

Environmental Science Atmospheres

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Md. K. A. Howlader, S. U. Zaman, Md. A. Hossen, Md. A. Kiber and A. Salam, *Environ. Sci.: Atmos.*, 2026, DOI: 10.1039/D6EA00036C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Environmental Significance Statement

Traffic intersections in rapidly growing megacities represent critical pollution hotspots with intense human exposure. This study quantifies seasonal variations of particulate matter ($PM_{1.0}$, $PM_{2.5}$, PM_{10}) and gaseous pollutants (NO_2 , CO_2) at major intersections in Dhaka, Bangladesh. Winter concentrations of PM were several times higher than pre-monsoon levels, and health risk assessment revealed hazard quotient (HQ) values >1 for $PM_{2.5}$, PM_{10} , and NO_2 , particularly for infants. Strong correlations between pollutants and vehicle density highlight traffic emissions as a dominant source. These findings emphasize the urgent need for targeted traffic management and emission-control strategies to reduce human exposure at urban intersection hotspots.



View Article Online
DOI: 10.1039/D6FA00036C

Traffic-Induced Air Pollution at Traffic Intersections in Dhaka: Seasonal Patterns and Associated Health Implications

Md. Khorshed Alam Howlader^a, Shahid Uz Zaman^{b*}, Md. Al-Amin Hossen^a, Md. Adnan Kiber^c, Abdus Salam^a

^aDepartment of Chemistry, Faculty of Science, University of Dhaka, Dhaka-1000, Bangladesh

^bDepartment of Chemistry, Faculty of Science, Bangladesh University of Engineering and Technology, Dhaka-1000, Bangladesh

^cDepartment of Electrical and Electronics Engineering, Faculty of Engineering and Technology, University of Dhaka, Dhaka-1000, Bangladesh

*Corresponding author's email: suzaman@chem.buet.ac.bd; shahiduzzamanadil@gmail.com



Abstract

Air pollution, poses a serious environmental and health challenges in Bangladesh. It is necessary to understand the concentration and seasonal variation of particulate matter (PM) and gaseous pollutants at traffic intersections to assess their impact on public health. This study was conducted to measure the levels of PM (PM_{1.0}, PM_{2.5} and PM₁₀) and trace gases (CO₂, NO₂) at five traffic intersections in Dhaka, Bangladesh, with the aim of evaluating seasonal variations and associated health risks. Sampling was conducted during the winter (December 2023-January 2024) and pre-monsoon (April-May 2024). Results revealed higher PM concentrations during winter: PM_{1.0}, PM_{2.5} and PM₁₀ averaged 108.3 ± 14.0 , 336.1 ± 68.3 and 449.2 ± 98.7 $\mu\text{g}/\text{m}^3$ respectively, compared to lower values in the pre-monsoon (41.7 ± 8.6 , 93.1 ± 20.1 and 151.9 ± 33.3 $\mu\text{g}/\text{m}^3$). NO₂ concentrations were higher during winter (0.13 ± 0.01 ppm) with oscillating diurnal variation and declined in pre-monsoon (0.10 ± 0.02 ppm), which exhibits a clear rising trend. CO₂ levels remained steady around 790 ± 30 ppm throughout both seasons. Health risk assessment showed Hazard Quotient (HQ) values above 1 for NO₂ (between 1.06 to 1.58), PM_{2.5} (between 0.89 to 4.78) and PM₁₀ (between 0.82 to 5.31) posing severe risks to infants. The Hazard Ratio (HR) for CO₂ ranged between 0.7 and 0.85, indicating no direct health effects. This study emphasizes the immediate need for specific mitigation strategies at traffic intersections to protect public health.

Keywords: Seasonal variations, Particulate matter, Gaseous pollutants, Traffic emission, Hazard Quotient, Hazard Ratio



1. Introduction

Air pollution is now regarded as a major global threat due to its negative effects on the environment, the economy and human health¹⁻⁴ and is also responsible worldwide for 9 to 12 million deaths annually^{5,6}. It is commonly acknowledged that transportation is one of the major and growing global sources of air pollution^{2,7-9} and traffic intersections are the centers of air pollution and create complicated traffic activities involving numerous pedestrians¹⁰⁻¹³. The emission of traffic-related air pollutants at traffic intersections can be two to three times greater than the ambient background resulting from vehicles continuous acceleration, slowdown and idling^{14,15}. The emissions of PM and gases (NO_x, SO_x, CO₂, VOCs etc.) from engine exhaust, as well as secondary air pollutants, are some of the ways that road traffic contributes to air pollution^{14,16,17}. Non-exhaust emissions include resuspended dust and wear on the tires, brakes and road surface^{18,19}.

Low-Middle Income Countries (LMICs) like Bangladesh are seeing a sharp rise in air pollution from brick kilns, mining, automobiles, industry and agriculture²⁰⁻²⁵. Air pollutants, including PM and gaseous pollutants have accumulated as a result of these activities²⁶. Dhaka is the commercial and industrial hub of Bangladesh with a population of more than 21 million and is expanding quickly despite all the issues that come with being a megacity^{16,27-29}. Among the many issues brought on by the city's explosive growth is traffic congestion, which has grown to be a significant burden for commuters and locals alike. The rapid growth in motorized vehicles has resulted from Dhaka's fast urbanization and population development, greatly increasing urban air pollution^{25,26,30}. The country's inadequate transportation system contributes to the high levels of PM pollution on the majority of its roadways, which has a



detrimental health effect on the city people^{31–33}. Recent studies have demonstrated increased illness and mortality among drivers, passengers and those residing close to major roadways as a result of traffic congestion^{16,34}. Respiratory ailments are more common in Dhaka's urban dwellers due to traffic-related air pollution and hospitalization for respiratory disorders is closely linked to air pollutants exposure^{21,35,36}. The high level of air pollution has particularly impacted children and the elderly³⁷. Road users experience high to extreme levels of stress due to traffic congestion^{38,39}.

Monitoring air quality at traffic intersections is essential to determine its impact on human health and how seasonal variations influence daily activity, yet empirical data remains limited. Mullick⁴⁰ along with Hashem⁴¹ has studied a cross-sectional survey regarding the knowledge, practices and respiratory health effects of air pollution among the traffic police exposed to vehicular emissions in Dhaka City, Bangladesh and Rahman⁴² have studied the self-reported health effects of air pollution on urban residents in Dhaka City, Bangladesh. Hoque⁴³ analyzed the spatiotemporal distribution and seasonal variations of air pollutants in fifteen different locations of Dhaka city, Bangladesh; no study has yet directly focused on air pollution at traffic intersections. Furthermore, the correlation between the air pollutants and vehicles at the traffic intersections also needs to be monitored to assess the contribution of air pollution due to traffic.

This study showed the temporal and seasonal variations of air quality at traffic intersections with their health effects. The relationship between average traffic number and air pollutants, which directly affect all types of intersection users, as well as a deeper understanding of the origin of PM, CO₂ and NO₂ is also studied. The study's findings focus insight into current implementation and policy gaps and offer suggestions for possible ways to lower emissions



from pollution sources generated by humans.

View Article Online
DOI: 10.1039/D6EA00036C

2. Methodology

2.1. Meteorology of Bangladesh

Bangladesh's weather pattern is characterized by its tropical monsoon, which is marked by high temperatures, humidity and clear seasonal fluctuations^{32,44}. From a climatic perspective, there are four distinct seasons: (1) the hot summer Pre-monsoon season, which lasts from March to May; (2) the dry Winter season, which lasts from December to February; (3) the Wet monsoon season, which lasts from June to September and (4) the Post monsoon fall season, which lasts from October to November^{27,31,44}. The usual pre-monsoon temperature ranges from 27 to 29°C, whereas the average winter temperature ranges from 18 to 21°C^{45,46}. The relative humidity ranges from 65 to 79% during pre-monsoon and has a mean of 74% during winter⁴⁷.

2.2. Sampling Sites

Five traffic intersections (Figure 1) in Bangladesh's capital, Dhaka (latitude 23°76' N, longitude 90°38' E, 8 m above sea level)⁴⁸, were sampled to guarantee representativeness and diversity in traffic conditions. Mirpur-10 (23°48'24.9"N 90°22'06.6"E) crossroads, an important intersection in Mirpur, often used by both public and private transportation. It is notoriously congested during rush hour because of its advantageous location that links several roads. Traffic intersection Mirpur-1 (23°46'54.4"N 90°21'06.4"E), an entrance to large urban districts, this crossroads is another important one in Mirpur, with high traffic density, especially from local transportation services and business vehicles. Bijoy Sarani (23°45'52.1"N 90°23'19.4"E) crossroads, ideally situated close to important business and administrative areas experience a



mixture of vehicles including motorcyclists, buses and cars that make up the intense commuting traffic. Gulistan ($23^{\circ}43'40.2''\text{N } 90^{\circ}24'37.8''\text{E}$) crossroads, a busy and central Dhaka traffic center, characterized by a high volume of traffic including private vehicles, rickshaws and buses. The intersection of Banani ($23^{\circ}47'49.1''\text{N } 90^{\circ}24'06.0''\text{E}$) situated in an affluent residential and business district, has heavy traffic because it's close to schools, shopping malls and business offices.




Fig. 1 Five sampling locations (Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani) chosen for particulate and gaseous pollutants sampling in Dhaka city, Bangladesh.

2.3. Sampling Procedure

The study employed a comprehensive temporal sampling strategy to monitor air quality parameters across multiple sites during two distinct seasonal periods: the winter season, which was from December 2023 to January 2024 and the pre-monsoon period from April 2024 to



May 2024. The sampling protocol for PM, NO₂ and CO₂ was systematically designed to capture diurnal variations by conducting measurements at three specific time intervals throughout the day: morning time (9:00 AM to 10:00 AM), afternoon time (1:00 PM to 2:00 PM) and evening (5:00 PM to 6:00 PM)^{43,49,50}. The number of vehicles was counted manually throughout the sample time in the traffic intersections. To ensure temporal consistency between traffic volume and pollutant concentration data, trained field personnel recorded all vehicles passing while also sampling air pollutants. To generate a representative estimate of traffic intensity, vehicle counts were averaged during the whole sample period at each site. This approach was used to establish a correlation between traffic volume and pollutant concentrations (PM, CO₂ and NO₂), as vehicles are the primary source of these pollutants in the research area.

Sampling of PM, NO₂ and CO₂ was done for three successive days in each location. The study spanned a total of 30 sampling days with 15 days in each season. Gaseous pollutants NO₂ and CO₂ were monitored by the AEROQUAL 500 SERIES (Aeroqual Ltd., Auckland, New Zealand) successively in one hour and with interval of 15 minutes. PM concentrations were recorded by AEROCET 531S (Met One Instrument, Washington, USA) in every 3 minutes, which allowed to achieve high temporal resolution of particle size distribution^{31,37,43,49,51,52}.

2.4. Health Risk Assessment due to PM and NO₂ Exposure

NO₂ and PM (PM_{2.5} and PM₁₀) are categorized as a Group 1 carcinogen to humans as per the IARC^{53,54}. NO₂ is classified as a component of air pollution and is associated with combustion processes, such as from power plants or vehicles. There is sufficient evidence of the human carcinogenic risk of exposure to NO₂ demonstrated by this categorization^{54,55}. Only the non-



carcinogenic risk was identified for PM and NO₂ using the USEPA-recommended technique (USEPA, 2009). The lifetime average daily dosage (LADD) is determined using the following Equation ^{37,56,57}:

$$\text{LADD} = \frac{CA \times IR \times ED \times EF}{BW \times AT} \dots\dots\dots (1)$$

Where, CA = Contaminant concentration (µg/m³), IR = Inhalation rate (m³/h), ED = Exposure duration (years), EF = Exposure frequency (days/year), BW = Body weight (kg) and AT = Average time for a lifetime (days) (Table 1).


The HQ is determined by the ratio of LADD to REL ⁵⁸:

$$\text{HQ} = \frac{\text{LADD}}{\text{REL}} \dots\dots\dots (2)$$

Where, REL = Reference Exposure Level. For PM_{2.5} and PM₁₀ the values for REL are 50 and 75 µg/m³ respectively ^{21,59,60} and according to USEPA (2008), REL value of NO₂ is 100 µg/m³. REL is the threshold at which exposed groups may experience negative health impacts in comparison to unexposed groups ⁶¹.

Table 1 Health risk assessment data for different age ^{21,58,60}

| Factors | Infant (0-1 year) | Children (6-12 year) | Adult (21-70 year) |
|-------------------------------------|-------------------|----------------------|--------------------|
| Inhalation rate (m ³ /h) | 6.8 | 13.8 | 16 |
| Body weight (kg) | 11.3 | 45.3 | 80 |
| Exposure frequency (days/year) | 350 | 350 | 350 |



| Exposure duration (years) | 1 | 12 | 30 |
|------------------------------------|-----|------|-------|
| Average time for a lifetime (days) | 365 | 4380 | 10950 |

View Article Online
DOI: 10.1039/D6EA00036C

2.5. Health Risk Assessment of CO₂

Global average atmospheric CO₂ concentration has been increasing and it has crossed 420 ppm now (USEPA). Although CO₂ is a normal constituent of the earth's atmosphere, high concentration can have several impacts on human health and thus should be evaluated. Using Formula (3), the Hazard Ratio (HR) was computed by dividing the average CO₂ concentration by the reference concentration ⁶²:

$$HR = \frac{C}{R_f C} \dots\dots\dots (3)$$

Where:

C = Average observed CO₂ concentration

R_fC = Reference concentration (1000 ppm)

When the HR is less than 1, non-carcinogenic health consequences are not taken into account.

On the other hand, if HR is more than 1, the exposure may be associated with non-cancer risks.

2.6. Quality Control and Quality Assurance

AEROQUAL 500 SERIES (Aeroqual Ltd., Auckland, New Zealand) implements a nondispersive infrared (NDIR) approach with a precision of $\leq \pm 10 \text{ ppm} + 5\%$ for the measurement of CO₂ concentration. NO₂ concentrations are detected with a precision of $\leq \pm 0.02 \text{ ppm} + 10\%$ using gas-sensitive electrochemical (GSE) sensors. PM concentrations were recorded by AEROCET 531S (Met One Instrument, Washington, USA) in μg/m³ with a ±10%



accuracy^{31,37,43,49,51,52}. Additionally, a standard filter-based device termed SIBATA (Model-090860-504, Saitama, Japan) was used to compare the devices³¹ and the results were in good agreement within 10%. The AEROCET 531S works on the principle of scattered light, which is studied as aerosols pass through a laser beam. Volume and distribution of the scattered light are therefore followed in order to understand size and concentration of particles within the sample. This makes it possible to monitor aerosol particles in real-time while the size range is from the nanometer scale to the micrometer scale⁶³.

2.7. Statistical Analysis

The correlation coefficient (R^2), Pearson's correlation and Analysis of Variance (ANOVA) test were determined by Origin 2025 and Microsoft Excel respectively to identify the pollutants primary sources, correlation among the pollutants and evaluate the significance of contributing variables.

2.7.1 Principal Component Analysis (PCA)

PCA is a multivariate statistical technique that simplifies the complexity of large datasets by representing most information with a smaller set of variables^{64,65}. This method is based on the hypothesis that limited numbers of the primary components, obtained as linear combinations of the original variables, can adequately capture the fundamental patterns of the data⁶⁶. The primary components are uncorrelated, therefore lowering redundancy in the obtained information⁶⁷. In PCA, the following dimensionless standard equation is established based on the chemical dataset in order to normalize the variables:

$$Z_{ij} = \frac{c_{ij} - \bar{c}_j}{\sigma_j} \dots \dots \dots (4)$$



In this analysis, i denotes the individual samples, ranging from 1 to n ; while j indicates the components under consideration, varying from 1 to m . (C_{ij}) is the presentation of component j 's concentration in sample i . The arithmetic mean concentration of component j (C_j) and its standard deviation (σ_j) are used to standardize the data.

3. Results and Discussion

3.1. Temporal and seasonal distribution of PM in Dhaka city

The average levels of PM in Mirpur-10, Gulistan, Bijoy Sarani, Mirpur-1 and Banani across three size categories: $PM_{1.0}$, $PM_{2.5}$ and PM_{10} were shown in Figure 2. PM concentrations during winter: $PM_{1.0}$, $PM_{2.5}$ and PM_{10} averaged 108.3 ± 14.0 , 336.1 ± 68.3 and $449.2 \pm 98.7 \mu\text{g}/\text{m}^3$, respectively, compared to lower values in the pre-monsoon season, roughly about one-third to half of winter levels (41.7 ± 8.6 , 93.1 ± 20.1 and $151.9 \pm 33.3 \mu\text{g}/\text{m}^3$), indicating improved air quality than winter season. Traffic and construction influenced areas Mirpur-1 and Banani showed higher PM levels than relatively less impacted traffic intersections. PM_{10} was significantly influenced by vehicles, road dust and construction activities. The intermediate source of $PM_{2.5}$, which was mostly produced by combustion processes in factories and automobiles, had a value of between 270 to 300 $\mu\text{g}/\text{m}^3$ during winter and 70 to 110 $\mu\text{g}/\text{m}^3$ during pre-monsoon across five traffic intersections. Observations in 2017 showed that $PM_{2.5}$ was 29.6 $\mu\text{g}/\text{m}^3$ and PM_{10} was 56.6 $\mu\text{g}/\text{m}^3$ ²⁹. Between 2016 and 2022, $PM_{2.5}$ values ranged from 4.94 to 351.2 $\mu\text{g}/\text{m}^3$, with an observed range of 346.3 $\mu\text{g}/\text{m}^3$ ⁶⁸ and the winter months (December-February) had significantly higher $PM_{2.5}$ concentrations, which exceeded 250 $\mu\text{g}/\text{m}^3$ ^{69,70}. Throughout the dry season, the average mass concentration of $PM_{2.5}$ and PM_{10} were



27 ± 8 and 109±38 µg/m³ respectively at an industrial site in Nigeria⁷¹. The lowest concentrations were seen in PM_{1.0}, the minimal particle size, which was between 90 to 120 µg/m³ during winter and 30 to 50 µg/m³ during pre-monsoon season. PM_{1.0} might penetrate deeper lung tissues and blood arteries, making it extremely harmful to human health^{72,73}. During winter, PM concentrations peaked in the morning at all five traffic intersections due to rush-hour vehicular emissions and a shallow planetary boundary layer. A distinct diurnal trend with high PM levels in the morning hours has been observed for Dhaka, principally impacted by traffic emissions and time-of-day climatic influences^{74,75}. During the afternoon, concentrations gradually decreased as solar heating extended the boundary layer, increasing vertical mixing and atmospheric dilution^{76,77}. Pre-monsoon PM concentrations were significantly lower, with a weaker diurnal structure due to stronger winds, higher temperatures that promote air mixing and increased wet deposition. Strong winds during the pre-monsoon disperse PM concentrations and meteorological conditions explain up to 76% of daily PM variability⁷⁸.







Fig. 2 Seasonal distribution of PM concentration during winter and pre-monsoon seasons at Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani in Dhaka city, Bangladesh.

3.2. Temporal and seasonal distribution of NO_2 in Dhaka city

NO_2 levels were consistently higher during the winter than pre-monsoon at all locations (Figure 3). Depending on the region under study, average winter NO_2 concentrations ranged from 0.12 to 0.16 ppm. Concentrations dropped to 0.08 to 0.12 ppm during the pre-monsoon. The winter season is characterized by low ultra-violet radiation intensity, high relative humidity and a



significant decrease in the height of the atmospheric boundary layer. As a result, the concentration of NO_2 released by vehicles is concentrated in a shallow surface layer, increasing the concentration at ground level. Gulistan and Bijoy-Sarani had the greatest NO_2 concentrations in winter and pre-monsoon seasons respectively, possibly as a result of industrialization and vehicle density. The comparatively lower results from two stations, Banani and Mirpur-10, suggested a smaller contribution from local emissions. Excessive coal burning in the brickfields close to Dhaka city during the winter may be linked to high NO_2 concentrations in the atmosphere⁷⁹. A similar pattern of NO_2 concentrations was observed in Dhaka city in 2003 to 2019^{22,29,80}, suggesting a consistent trend over the decade. The average NO_2 concentration was 5.8 ppm, with a range of 5.6 to 6.1 ppm during December 2021 and January 2022 in Dhaka⁴³. Similar trends were noted in India and China as well, suggesting a more widespread regional uniformity in the levels of NO_2 pollution in metropolitan areas^{81,82}. These results also imply vehicle emissions continue to be the primary causes of NO_2 concentrations in densely populated urban areas^{81–85}. Low temperatures and slow airflow brought on by cold-season weather patterns might be the origin of this seasonal trend, which confines pollutants to the atmosphere. In the pre-monsoon, high spatial and temporal variability showed, whereas it is considerably larger around midday and afternoon during the winter, showing higher variability. Later in the day, converging concentrations are visible at all five traffic intersections during winter season. NO_2 concentrations remain consistently higher and more fluctuating during the winter, whereas pre-monsoon seasons exhibit a clear rising trend, indicating stronger diurnal variation across all traffic intersections. NO_2 concentrations increase with progression of the day, indicating vehicular emissions as a significant



source^{43,79}. Greater air mixing and increased solar energy radiation, which cause pollutant fluctuation, are the causes of the pre-monsoon's more significant diurnal variation. Pollutants remain trapped close to the surface during the winter months due to temperature inversions and slow winds, which keep concentrations mostly constant throughout the day^{77,86,87}.




Fig. 3 Seasonal distribution of NO_2 concentration during winter and pre-monsoon at Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani in Dhaka city, Bangladesh.

3.3. Temporal and seasonal distribution of CO_2 in Dhaka city

CO_2 levels (Figure 4) are generally greater in winter than in the pre-monsoon season. A similar seasonal variation in CO_2 concentration was also observed in Ahmedabad, India, during the period from 2014 to 2015, although the overall levels were lower. Specifically, concentration averaged 424.8 ± 17 ppm in winter and 398.8 ± 2.8 ppm in the pre-monsoon season⁸⁸. CO_2 concentrations varied from 537 to 568 ppm (average: 554 ppm) during December 2021 and January 2022 in Dhaka⁴³. Nearly identical levels of CO_2 concentration were suggested by each of the five traffic intersections. All of the location's CO_2 levels stayed between 700 and 900 ppm, indicating that CO_2 levels were often high in metropolitan areas. Concentrations typically peak in the early morning during both seasons, fall during the middle of the day and then



slightly increase again in the evening which also evident in China and Korea^{89–91}. Heavy traffic emissions during morning rush hour and an insignificant boundary layer that restricts the dispersion of pollutants near the surface are the main causes of the elevated CO₂ concentrations seen in the early morning. Similar diurnal pattern was seen during the pre-monsoon season, however there was less variability and a lower peak concentration than during the winter. Bijoy Sarani typically has the lowest CO₂ levels, whereas Mirpur-10 and Banani have the highest amounts as a result of increased traffic or local emissions. These patterns show how urban activity and weather patterns affect the distribution of CO₂ throughout the day. It was evident that diverse sources of emissions were responsible for the equal amounts of CO₂ concentrations.




Fig. 4 Seasonal distribution of CO₂ concentration during pre-monsoon and winter at Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani in Dhaka city, Bangladesh.

3.4. Correlation among the pollutants

Pearson's correlation (r) and ANOVA test were carried out to analyze the relationship among the various pollutants (CO₂, NO₂, PM_{1.0}, PM_{2.5} and PM₁₀). The Correlation Coefficient (R^2) was determined to relate to the air pollutants with the average vehicle number at different traffic intersections in Dhaka City. Pearson's correlation analysis results are presented in Table S2.



The correlation analysis showed positive correlations among NO_2 , CO_2 , $\text{PM}_{1.0}$, $\text{PM}_{2.5}$ and PM_{10} concentrations. The Pearson correlation coefficients of pollutants during winter were 0.96 (NO_2 and CO_2), 0.53 ($\text{PM}_{1.0}$ and $\text{PM}_{2.5}$) and 0.78 ($\text{PM}_{2.5}$ and PM_{10}) and during pre-monsoon were 0.58 ($\text{PM}_{1.0}$ and $\text{PM}_{2.5}$), 0.95 ($\text{PM}_{2.5}$ and PM_{10}) and 0.64 (CO_2 and PM_{10}) ($p < 0.05$). Positive correlations indicate a strong linear correlation among the pollutants. According to the results of the ANOVA test, the connection between PM and gaseous pollutants is statistically significant ($p < 0.05$) (Table S2). Strongly positive correlation also observed in Dhaka⁴³, Chattogram⁹², Bangladesh and Southern China and California^{79,93}. In contrast, during the COVID-19 lockdown, a weakly positive correlation was observed between the pollutants in Wuhan, China⁹⁴. Vehicular emissions are a primary source of NO_2 and CO_2 both of which contribute to the formation of PM^{83} . The present study analysis showed a statistically significant ($p < 0.001$) moderately positive correlation ($R^2 = 0.40$) between $\text{PM}_{1.0}$ concentration and the average vehicle number during the pre-monsoon season. This correlation indicates fossil fuel combustion from vehicles is a dominant contributor to $\text{PM}_{1.0}$ pollution in traffic intersections. $\text{PM}_{2.5}$ and PM_{10} show strong positive correlations, while positive associations were also found among NO_2 , CO_2 , $\text{PM}_{1.0}$, $\text{PM}_{2.5}$ and PM_{10} ^{95–97}. These findings suggested pollutants were emitted from the similar sources, primarily linked to vehicular activity, emphasizing the need for stringent vehicular emission controls and enhanced traffic management to alleviate air pollution in traffic intersections.

3.4.1 Principal Component Analysis (PCA)

Principal Component Analysis (PCA) was employed to identify the major sources contributing to particulate matter and gaseous pollutants at the traffic intersections during winter and the



pre-monsoon season in Dhaka City, Bangladesh. The analysis employed a Varimax rotation to improve the interpretability of factor loadings by maximizing coefficient variance. This allows for a clearer identification of groups influencing particulate matter and gaseous pollutants.




Fig. 5 The PCA biplot highlighting the relationships between particulate matter and gaseous pollutants during winter and pre-monsoon seasons at Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani in Dhaka city, Bangladesh.

During the winter season, the first two principal components (PC1 and PC2) exhibited eigenvalues of 2.61 and 2.26 respectively, accounting for 97.38% of the total variance, showing that they sufficiently reflect the dataset (Table S4). PC1 accounted for 52.27% of the total variance and exhibited a significant correlation with particulate matter fractions, specifically $PM_{1.0}$ (0.583), $PM_{2.5}$ (0.562) and PM_{10} (0.540) (Table S5). The substantial positive loadings of these variables indicate that PC1 is a particulate matter-dominated source, most likely due to resuspended road dust, construction activities and traffic-related emissions⁹⁸. PC2 indicated a very distinct loading pattern, with strong positive loadings for CO_2 (0.629) and NO_2 (0.625)



(Table S5). This bipolar arrangement is consistent with a gaseous combustion emission factor, which is mostly caused by industrial burning, vehicle exhaust and energy generation activities that release CO₂ and NO₂ at the same time. The first two principal components (PC1 and PC2) exhibited eigenvalues of 3.193 and 0.865 respectively, cumulatively explaining 81.16% during pre-monsoon season (Table S6). PC1 is a general air pollution factor that reflects the overall intensity of anthropogenic emission activity, such as fuel combustion and recorded high and approximately uniform positive loadings across all five pollutants: NO₂ (0.356), CO₂ (0.356), PM₁₀ (0.485), PM_{2.5} (0.521) and PM_{1.0} (0.490) (Table S7). PC2 showed a unique loading pattern, with a substantial positive loading for NO₂ (0.784), indicating NO₂ from industrial and vehicle combustion. PC3 was dominated by an unusually high loading for CO₂ (0.908). This component captures industrial emissions that are uncorrelated with the major emission complex. These results are consistent with prior PCA based source apportionment studies conducted in urban environments, which consistently identify different gaseous and particle emission sources as the main causes of air pollution variance^{43,99–101}.

3.5. Health Risk Assessment

Air pollution had a major chronic impact on all age groups, with infants and children suffering the most^{102–106}. HQ was used to evaluate the non-cancer hazards related to air pollution^{107,108}. While there was no risk of adverse health consequences (HQ < 1) for the pollutants in the case of adults, the HQ values posed a considerable health threat (HQ > 1) for infants and children. Low breathing heights with other air pollution sources may be the primary cause of this problem in youngsters, who are especially vulnerable to ground-level pollution¹⁰⁹. Infants are exposed to concentrations of fine particles that are up to 44% greater than those of adults,



according to research done at a school by Sharma and Kumar ¹¹⁰.

3.5.1 Health Risk Assessment of PM

At every site, the HQ values of PM_{2.5} and PM₁₀ (Figure 6(a) and 6(b)) for every age group were greater throughout the winter than they were during the pre-monsoon season. This could happen as a result of wintertime weather conditions prevent PM from reducing ^{111,112}. Although HQ levels continue to decline during the pre-monsoon season, newborns were particularly at risk and in certain clusters, the HQ stays over the allowed limits. Because young populations were more susceptible to particulate matter exposure due to the physiological immaturity of their respiratory tracts and higher inhalation rates of particulate matter per unit body weight, newborns had much higher HQ values than children and adults ^{113–115}. Mirpur-1 and Banani showed the highest HQ values for all age groups, with infants exceeding a hazard quotient of 4. According to Bijoy Sarani's appendix, it had lower HQ values than any other place for both adults and children, indicating less exposure to PM or cleaner air. Similar to the winter, the HQ values in Banani and Mirpur-1 were higher during pre-monsoon. In locations like Bijoy Sarani and Banani, pre-monsoon PM levels remained hazardous despite being lower, particularly for young children. PM_{2.5} can enter the lungs¹¹⁶, whereas PM_{1.0} can penetrate blood vessels through alveoli, capturing the entire body^{117,118}. In addition, PM_{1.0} have a greater surface to mass ratio and can contain a high concentration of potentially toxic anthropogenic components¹¹⁹. Short-term exposure to PM_{1.0} has been regarded as the cause of death due to stroke, while long-term inhalation can raise the risk of neurological diseases^{120–122}. It also noted that babies who grow up in places with PM concentrations higher than the upper limit were more likely to experience harmful health consequences, including respiratory ailments and heart problems ^{123–125}. The



population most vulnerable to environmental pollution, particularly children because of their developing organs and immune systems, was at risk for serious health impacts from PM₁₂₆

128




Fig. 6(a) HQ variation of PM_{2.5} during winter and pre-monsoon seasons for different age at Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani in Dhaka city, Bangladesh.

PM causes numerous adverse health consequences, including a higher incidence of myocardial infarction, diabetes, respiratory diseases, cardiovascular and cerebrovascular diseases and lung cancer^{5,123,129,130}. In addition to negative health effects, including bronchial asthma, pneumonia and pulmonary disease^{131–134}.




Fig. 6(b) HQ variation of PM_{10} during winter and pre-monsoon for different age at Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani in Dhaka city, Bangladesh.

3.5.2 Health Risk Assessment of NO_2

The overall HQ values (Figure 7) for all age groups and all locations were greater throughout the winter than they were during pre-monsoon. This might be due to wintertime weather conditions, which limit the concentration of pollutants to areas near the surface (e.g., low mixing heights)^{135,136}. Infants exhibited the greatest HQ values, followed by children and adults. Infants were exposed to more air pollution than adults, making them far more vulnerable due to their developing organ systems. Mirpur-10, Gulistan and Bijoy Sarani have higher HQ levels than the Banani and Mirpur-1 areas. These discrepancies might have resulted from variations in the industrial activity and vehicle emissions that caused urban traffic congestion in various regions. Wintertime HQ values for infants at Mirpur-10, Gulistan and Bijoy Sarani were seen to increase to levels near or over the crucial level of 1. Therefore, NO_2 exposure was found to provide a non-carcinogenic health risk for newborns exposed to several places during the winter, as indicated by HQ values > 1 (USEPA). Airway inflammation and obstruction of airflow are caused by the NO_2 exposure^{137–141}. Annually ambient NO_2 is responsible for 3.52 million new cases of asthma in children and adolescents¹⁴².






Fig. 7 HQ variation of NO_2 during winter and pre-monsoon for different age at Mirpur-10, Mirpur-1, Bijoy Sarani, Gulistan and Banani in Dhaka city, Bangladesh.

3.5.3 Health Risk assessment of CO_2

The HR of CO_2 for the pre-monsoon and winter seasons for five traffic intersections was shown in Table 2. The HR values range from 0.7 to 0.85. With minor seasonal fluctuations noted at various places, the HR values were generally steady across both seasons. As CO_2 concentrations raised, the mean raw scores for seven of the nine decision-making performance scales consistently fell^{143,144} and cognitive impacts¹⁴⁵ were observed in comparison to 600 to 1000 ppm ($0.6 < \text{HR} < 1.0$).

Table 2 Hazard Ratio (HR) data for Health Risk Assessment of CO_2

| Sampling Sites | Hazard Ratio (HR) | |
|----------------|-------------------|--------|
| | Pre-monsoon | Winter |
| Mirpur-10 | 0.7705 | 0.8069 |
| Gulistan | 0.7781 | 0.8086 |
| Mirpur-1 | 0.7856 | 0.7694 |



| | | 0.7928 |
|--------------|--------|--------|
| Banani | 0.8074 | 0.7928 |
| Bijoy Sarani | 0.7666 | 0.7832 |

View Article Online

DOI: 10.1039/D6EA00036C

The short sampling duration, fewer traffic intersections and non-simultaneous pollution measurements are the limitations of this study. Future research should include simultaneous measurements, more traffic intersections and prolonged sampling period throughout several seasons (winter, pre-monsoon, monsoon and post-monsoon) to better capture the spatial and temporal variability in traffic-related air pollution.

4. Conclusion

This study investigated the seasonal and temporal variations of particulate matter (PM_{1.0}, PM_{2.5}, PM₁₀) and gaseous pollutants (NO₂, CO₂) at five major traffic intersections in Dhaka during winter (December 2023-January 2024) and pre-monsoon (April-May 2024). Results showed substantially higher pollutant levels in winter (PM_{1.0} = 108.3 ± 14.0 µg/m³, PM_{2.5} = 336.1 ± 68.3 µg/m³ and PM₁₀ = 449.2 ± 98.7 µg/m³, NO₂ = 0.13 ± 0.01 ppm) with Gulistan, Mirpur-10, Banani and Mirpur-1 emerging as hotspots for different pollutants. Health risk assessment revealed HQ values greater than one for NO₂, PM_{2.5} and PM₁₀, indicating serious risks, particularly for infants, while CO₂ levels (790 ± 30 ppm) mainly highlighted climatic rather than direct health impacts. The Pearson correlation analyses demonstrated strong associations among pollutants and a moderate link between PM_{1.0} and vehicle numbers (p < 0.001, R² = 0.40) during the pre-monsoon season, underscoring the central role of traffic emissions. The PCA analysis indicated that traffic emissions are one of the major sources of PM, NO₂ and CO₂. The findings of this study indicate the urgent need for sustained assessment and effective



intervention strategies to mitigate traffic-related air pollution in Dhaka. Future research should focus on expanding spatial and temporal coverage, including more intersections and continuous year-round monitoring, to capture long-term trends and the influence of meteorology.

View Article Online
DOI: 10.1039/D6EA00036C

Acknowledgement

The authors express their gratitude to the Department of Chemistry, University of Dhaka.

Funding

This research has received no external funding.

Author's contribution

Md. Khorshed Alam Howlader: Conceptualization, Sampling, Methodology, Data analysis, Writing original draft, Review & editing

Shahid Uz Zaman: Review & editing, Supervision

Md. Al-Amin Hossen: Review & editing

Md. Adnan Kiber: Review & editing

Abdus Salam: Conceptualization, Review & editing, Supervision

Conflict of interest statement


The authors declared that they had no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

5. References

1. Feng T, Sun Y, Shi Y, Ma J, Feng C, Chen Z. Air pollution control policies and impacts: A review. *Renewable and Sustainable Energy Reviews*. 2024;191.
2. Faisal A Al, Rahman MM, Haque S. Retrieving spatial variation of aerosol level over urban mixed land surfaces using Landsat imageries: Degree of air pollution in Dhaka Metropolitan Area. *Physics and Chemistry of the Earth* 2022;126.




3. Arfan Ali M, Bilal M, Wang Y, et al. Spatiotemporal changes in aerosols over Bangladesh using 18 years of MODIS and Reanalysis data. 2022;
4. Qiu W, Zhou Y, He H, et al. Short-term effects of air pollution on liver function among urban adults in China. *Atmos Environ* 2021;245.
5. Henning RJ. Particulate Matter Air Pollution is a Significant Risk Factor for Cardiovascular Disease. *Curr. Probl. Cardiol.* 2024;49(1).
6. Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 2015;525(7569):367–371.
7. Islam FAS. International Journal of Applied and Natural Sciences A Comprehensive Analysis of Air Pollution in Dhaka City, Bangladesh, and the Application of Artificial Intelligence and Machine Learning for Enhanced Management and Forecasting. 2025; Available from: <https://creativecommons.org/licenses/by/4.0/>
8. Patton AP, Boogaard H, Vienneau D, et al. Assessment of long-term exposure to traffic-related air pollution: An exposure framework. *J Expo Sci Environ Epidemiol* 2025;35(3):493–501.
9. Ilenič A, Pranjić AM, Zupančič N, Milačić R, Ščančar J. Fine particulate matter (PM_{2.5}) exposure assessment among active daily commuters to induce behaviour change to reduce air pollution. *Science of the Total Environment* 2024;912.
10. Zheng T, Xiang S, Zhang S, Wu Y. Variability of traffic-related air pollutants at two- and four-phase intersections. *Atmos Pollut Res* 2023;14(12).
11. Collado JT, Abalos JG, los Reyes I de, et al. Spatiotemporal Assessment of PM_{2.5} Exposure of a High-risk Occupational Group in a Southeast Asian Megacity. *Aerosol Air Qual Res* 2023;23(1).
12. Xiang S, Zhang S, Yu YT, et al. Evaluating Ultrafine Particles and PM_{2.5} in Microenvironments with Health Perspectives: Variability in Concentrations and



Pollutant Interrelationships. *Aerosol Air Qual Res* 2023;23(9).

View Article Online
DOI: 10.1039/D6EA00036C

13. Alobaidi MK, Badri RM, Salman MM. Evaluating the Negative Impact of Traffic Congestion on Air Pollution at Signalized Intersection. In: *IOP Conference Series: Materials Science and Engineering*. Institute of Physics Publishing, 2020;
14. Shen Y, Zhang Q, Wang D, et al. Evaluation of a cost-effective roadside sensor platform for identifying high emitters. *Science of the Total Environment* 2022;816.
15. Jia C, Li W, Wu T, He M. Road traffic and air pollution: Evidence from a nationwide traffic control during coronavirus disease 2019 outbreak. *Science of the Total Environment* 2021;781.
16. Islam ARMT, Awadh M Al, Mallick J, et al. Estimating ground-level PM_{2.5} using subset regression model and machine learning algorithms in Asian megacity, Dhaka, Bangladesh. *Air Qual Atmos Health* 2023;16(6):1117–1139.
17. Jin MY, Gallagher J, Li XB, Lu KF, Peng ZR, He H Di. Characterizing the distribution pattern of traffic-related air pollutants in near-road neighborhoods. *Environ Monit Assess* 2024;196(8).
18. Harrison RM, Vu T Van, Jafar H, Shi Z. More mileage in reducing urban air pollution from road traffic. *Environ Int* 2021;149.
19. Wei P, Brimblecombe P, Yang F, et al. Determination of local traffic emission and non-local background source contribution to on-road air pollution using fixed-route mobile air sensor network. *Environmental Pollution* 2021;290.
20. Zaman SU, Riad Sarkar Pavel M, Rani RI, et al. Aerosol climatology characterization over Bangladesh using ground-based and remotely sensed satellite measurements. *Elementa* 2022;10(1).
21. Salam A, Jeba F, Riad M, Pavel S, Zaman SU. Health Risk Assessment of High-Level Particulate Matter Exposure in Different Environments in Mega City Dhaka, Bangladesh [Homepage on the Internet]. 2021; Available from: www.baasbd.org



22. Pavel MRS, Zaman SU, Jeba F, Islam MS, Salam A. Long-Term (2003–2019) Air Quality, Climate Variables, and Human Health Consequences in Dhaka, Bangladesh. *Frontiers in Sustainable Cities* 2021;3.
23. Islam MM, Sharmin M, Ahmed F. Predicting air quality of Dhaka and Sylhet divisions in Bangladesh: a time series modeling approach. *Air Qual Atmos Health* 2020;13(5):607–615.
24. Rana MM, Sulaiman N, Sivertsen B, Khan MF, Nasreen S. Trends in atmospheric particulate matter in Dhaka, Bangladesh, and the vicinity. *Environmental Science and Pollution Research* 2016;23(17):17393–17403.
25. Begum BA, Hopke PK, Markwitz A. Air pollution by fine particulate matter in Bangladesh. *Atmos Pollut Res* 2013;4(1):75–86.
26. Safiur Rahman M, Khan MDH, Jolly YN, Kabir J, Akter S, Salam A. Assessing risk to human health for heavy metal contamination through street dust in the Southeast Asian Megacity: Dhaka, Bangladesh. *Science of the Total Environment* 2019;660:1610–1622.
27. Salam A, Mamoon H AI, Ullah MB, Ullah SM. Measurement of the atmospheric aerosol particle size distribution in a highly polluted mega-city in Southeast Asia (Dhaka-Bangladesh). *Atmos Environ* 2012;59:338–343.
28. Hossain I, Rahman S, Sattar S, Haque M. Environmental Overview of Air Quality Index (AQI) in Bangladesh: Characteristics and Challenges in Present Era [Homepage on the Internet]. 2021; Available from: <https://www.ijresm.com>
29. Mukta TA, Hoque MMM, Sarker ME, Hossain MN, Biswas GK. Seasonal variations of gaseous air pollutants (SO₂, NO₂, O₃, CO) and particulates (PM_{2.5}, PM₁₀) in Gazipur: An industrial city in Bangladesh. *Advances in Environmental Technology* 2020;6(4):195–209.
30. Salam A, Bauer H, Kassin K, Ullah SM, Puxbaum H. Aerosol chemical characteristics of a mega-city in Southeast Asia (Dhaka-Bangladesh). *Atmos Environ* 2003;37(18):2517–2528.



31. Uz Zaman S, Yesmin M, Riad Sarkar Pavel M, Jeba F, Salam A. Indoor air quality indicators and toxicity potential at the hospitals' environment in Dhaka, Bangladesh. Available from: <https://doi.org/10.1007/s11356-021-13162-8>
32. Salam A, Ullah MB, Islam MS, Salam MA, Ullah SM. Carbonaceous species in total suspended particulate matters at different urban and suburban locations in the Greater Dhaka region, Bangladesh. *Air Qual Atmos Health* 2013;6(1):239–245.
33. Begum BA, Biswas SK, Hopke PK. Key issues in controlling air pollutants in Dhaka, Bangladesh. In: *Atmospheric Environment*. Elsevier Ltd, 2011; p. 7705–7713.
34. Zheng T, Xiang S, Zhang S, Wu Y. Variability of traffic-related air pollutants at two- and four-phase intersections. *Atmos Pollut Res* 2023;14(12).
35. Cohen AJ, Brauer M, Burnett R, et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *The Lancet* 2017;389(10082):1907–1918.
36. Rahman M, Rashid F, Kumar D, Habib MA, Ullah A. Dataset of air pollutants (PM_{2.5}, PM₁₀, CO) concentrations in the export processing area of Dhaka, Bangladesh. *Data Brief* 2024;55.
37. Roy S, Zaman SU, Joy KS, Jeba F, Kumar P, Salam A. Impact of fine particulate matter and toxic gases on the health of school children in Dhaka, Bangladesh. *Environ Res Commun* 2023;5(2).
38. Fattah MA, Morshed SR, Kafy A Al. Insights into the socio-economic impacts of traffic congestion in the port and industrial areas of Chittagong city, Bangladesh. *Transportation Engineering* 2022;9.
39. Rizwan SA, Nongkynrih B, Gupta SK. Air pollution in Delhi: Its Magnitude and Effects on Health. *Indian Journal of Community Medicine* 2013;38(1):4–8.
40. Mullick DAR. Traffic Air Pollution and Respiratory Health: A Cross-Sectional Study among Traffic Police in Dhaka City (Bangladesh). *Journal of Medical Science And*

View Article Online
DOI: 10.1039/C5EA00036C



- clinical Research [homepage on the Internet] 2021;09(05). Available from: <http://jmscr.igmpublication.org/v9-i5/17%20jmscr.pdf>
41. Hashem RB, Siddique AB, Rasel SM, Hossain MS. Assessment of knowledge, attitudes, and practices regarding air pollution among traffic polices in Dhaka city, Bangladesh: a cross-sectional study. *BMC Public Health* 2024;24(1).
 42. Rahman MM, Hasanuzzaman ABM, Chisty MA, Alam E, Islam MK, Islam ARMT. Perceived-air pollution and self-reported health status: a study on air pollution-prone urban area of Bangladesh. *Front Public Health* 2025;13.
 43. Hoque MMdM, Mizan MdMR, Hossain MdN, et al. Spatial distribution and diurnal variation of particulate matters and gaseous air pollutants of Dhaka city, Bangladesh. *Discover Atmosphere* 2025;3(1).
 44. Salam A, Hossain T, Siddique MNA, Shafiqul Alam AM. Characteristics of atmospheric trace gases, particulate matter, and heavy metal pollution in Dhaka, Bangladesh. *Air Qual Atmos Health* 2008;1(2):101–109.
 45. Mainuddin M, Peña-Arancibia JL, Karim F, Hasan MM, Mojid MA, Kirby JM. Long-term spatio-temporal variability and trends in rainfall and temperature extremes and their potential risk to rice production in Bangladesh. *PLOS Climate* 2022;1(3).
 46. Sakib SN, Islam ARMT, Azad MAK, et al. Seasonality of meteorological factors influencing the COVID-19 era in coastal and inland regions of bangladesh. *Geocarto Int* 2023;38(1).
 47. Mortuza MR, Selmi S, Khudri MM, Ankur AK, Rahman MM. Evaluation of temporal and spatial trends in relative humidity and dew point temperature in Bangladesh. *Arabian Journal of Geosciences* 2014;7(12):5037–5050.
 48. Fattah MA, Morshed SR, Kafy A Al, Rahaman ZA, Rahman MT. Wavelet coherence analysis of PM_{2.5} variability in response to meteorological changes in South Asian cities. *Atmos Pollut Res* 2023;14(5).




49. Bani Khalifi N, Platymesi K, Vlachos S, Bartzanas T, Despoina Avgoustaki D. Quantifying the Scaling Effects of Urban Green Infrastructure on Air Quality and Greenhouse Gas Dynamics: Insights from a Multi-Site Evaluation in Athens, Greece. 2025; Available from: <https://doi.org/10.3390/su>
50. Blaszczyk-Boxe C, Karle NN, Ideki O, et al. Temporal variations in NO_x, O₃, and CO₂ and meteorological influences in New York and New Jersey urban atmospheres. *Discover Environment* 2025;3(1).
51. Lin C, Gillespie J, Schuder MD, Duberstein W, Beverland IJ, Heal MR. Evaluation and calibration of Aeroqual series 500 portable gas sensors for accurate measurement of ambient ozone and nitrogen dioxide. *Atmos Environ* 2015;100:111–116.
52. Delgado-Saborit JM. Use of real-time sensors to characterise human exposures to combustion related pollutants. *Journal of Environmental Monitoring* 2012;14(7):1824–1837.
53. Loomis D, Huang W, Chen G. The International Agency for Research on Cancer (IARC) evaluation of the carcinogenicity of outdoor air pollution: Focus on China. *Chin. J. Cancer*. 2014;33(4):189–196.
54. Yu G, Cui Y, Kang R, Wu J, Ge W, Han J. Air Pollution and the Risk of Liver Cancer Incidence and Mortality: A Systematic Review and Meta-Analysis. *Liver Int* 2025;45(11):e70409.
55. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 2014;
56. Riad M, Pavel S, Tarannum M, Zaman SU, Jeba F, Salam A. Health Risk Assessment of High-Level Particulate Matter Exposure in Different Environments in Mega City Dhaka, Bangladesh [Homepage on the Internet]. Available from: www.baasbd.org
57. Yunesian M, Rostami R, Zarei A, Fazlzadeh M, Janjani H. Exposure to high levels of PM_{2.5} and PM₁₀ in the metropolis of Tehran and the associated health risks during 2016–2017. *Microchemical Journal* 2019;150.

View Article Online
DOI: 10.1039/D5EA00036C



58. Kim H, Kang K, Kim T. Measurement of particulate matter (PM_{2.5}) and health risk assessment of cooking-generated particles in the kitchen and living rooms of apartment houses. *Sustainability (Switzerland)* 2018;10(3).
59. Othman M, Latif MT, Yee CZ, et al. PM_{2.5} and ozone in office environments and their potential impact on human health. *Ecotoxicol Environ Saf* 2020;194.
60. Matooane M, Diab R. Health risk assessment for sulfur dioxide pollution in South Durban, South Africa. *Arch Environ Health* 2003;58(12):763–770.
61. Mayowa Morakinyo O, Stephen Adebawale A, Ingrid Mokgobu M, Stanley Mukhola M. Health risk of inhalation exposure to sub-10 μm particulate matter and gaseous pollutants in an urban-industrial area in South Africa: an ecological study. *BMJ Open* [homepage on the Internet] 2017;7:13941. Available from: <http://dx.doi.org/10.1136/bmjopen-2016-013941>
62. Kumar P, Hama S, Abbass RA, et al. CO₂ exposure, ventilation, thermal comfort and health risks in low-income home kitchens of twelve global cities. *Journal of Building Engineering* 2022;61.
63. Akther T, Ahmed M, Shohel M, Ferdousi FK, Salam A. Particulate matters and gaseous pollutants in indoor environment and Association of ultra-fine particulate matters (PM₁) with lung function. *Environmental Science and Pollution Research* 2019;26(6):5475–5484.
64. Liu Q, Yang P, Hu Z, Shu Q, Chen Y. Identification of the sources and influencing factors of the spatial variation of heavy metals in surface sediments along the northern Jiangsu coast. *Ecol Indic* 2022;137.
65. Aziz S, Zaman SU, Roy S, et al. Comprehensive analysis of heavy metals in indoor PM_{2.5}: Source identification and health risk assessment in Dhaka, Bangladesh. *Atmos Environ X* 2025;27.
66. Aidoo EN, Appiah SK, Awashie GE, Boateng A, Darko G. Geographically weighted principal component analysis for characterising the spatial heterogeneity and




- connectivity of soil heavy metals in Kumasi, Ghana. *Heliyon* 2021;7(9). DOI: 10.1039/D6EA00036C
67. Feng W, Zhang Y, Huang L, et al. Source apportionment of environmentally persistent free radicals (EPFRs) and heavy metals in size fractions of urban arterial road dust. *Process Safety and Environmental Protection* 2022;157:352–361.
68. Hasan K, Rahman M, Akhter M, Mohinuzzaman M, Kayes I, Rahman S. A new dynamic approach using data-driven and machine learning models for forecasting particulate matter in Dhaka megacity. *Environmental Pollution and Management* 2024;1:235–247.
69. Billah SMM, Haq HM, Rahman A, Biswas H, Islam MI. Meteorological drivers of fine particulate matter variability in Dhaka City with a seasonal analysis of temperature, humidity, rainfall, and wind speed. *Discover Environment* 2026;4(1).
70. Zaman SU, Roy S, Foy B de, et al. Comparison of PM2.5 trends and source factors in urban and rural locations in Bangladesh. *Atmos Pollut Res* 2026;17(2).
71. Kolawole TO, Fomba KW, Ezeh GC, et al. Chemical composition, sources, and health risks assessment of PM 10 and PM 2.5 -bound metals at an industrial site in Nigeria . *Environmental Science: Atmospheres* 2026;
72. Mariga ST. Indoor Particulate Matter and Gaseous Phase Particles' Dynamics of Fuels-Burning 2 Emissions, in Slum Dwellings, Kampala [Homepage on the Internet]. Available from: <https://ssrn.com/abstract=5276169>
73. Chen T, Chen F, Wang K, et al. Acute respiratory response to individual particle exposure (PM1.0, PM2.5 and PM10) in the elderly with and without chronic respiratory diseases. *Environmental Pollution* 2021;271.
74. Rahman M, Meng L. Examining the Spatial and Temporal Variation of PM2.5 and Its Linkage with Meteorological Conditions in Dhaka, Bangladesh. *Atmosphere (Basel)* 2024;15(12).
75. Zaman SU, Budhavant K, Salam A. The influence of meteorological factors on wintertime black carbon and PM2.5 pollution in Dhaka, Bangladesh. *International*

View Article Online

DOI: 10.1039/D6EA00036C



- Journal of Environmental Science and Technology 2025;22(9):8225–8234.
76. Miao Y, Li J, Miao S, et al. Interaction Between Planetary Boundary Layer and PM_{2.5} Pollution in Megacities in China: a Review. *Curr. Pollut. Rep.* 2019;5(4):261–271.
 77. Foy B de, Edwards R, Joy KS, Zaman SU, Salam A, Schauer JJ. Interpretable machine learning tools to analyze PM_{2.5} sensor network data so as to quantify local source impacts and long-range transport. *Atmos Res* 2024;311.
 78. Islam N, Toha TR, Islam MM, Ahmed T. The association between particulate matter concentration and meteorological parameters in Dhaka, Bangladesh. *Meteorology and Atmospheric Physics* 2022;134(4).
 79. Hu M, Wang Y, Wang S, Jiao M, Huang G, Xia B. Spatial-temporal heterogeneity of air pollution and its relationship with meteorological factors in the Pearl River Delta, China. *Atmos Environ* 2021;254.
 80. Meteorological Influences on Seasonal Variations of Air Pollutants (SO₂, NO₂, O₃, CO, PM_{2.5}, and PM₁₀) in the Dhaka Megacity. *American Journal of Pure and Applied Biosciences* 2020;15–23.
 81. Xiao K, Wang Y, Wu G, Fu B, Zhu Y. Spatiotemporal characteristics of air pollutants (PM₁₀, PM_{2.5}, SO₂, NO₂, O₃, and CO) in the Inland Basin City of Chengdu, Southwest China. *Atmosphere (Basel)* 2018;9(2).
 82. Gaur A, Tripathi SN, Kanawade VP, Tare V, Shukla SP. Four-year measurements of trace gases (SO₂, NO_x, CO, and O₃) at an urban location, Kanpur, in Northern India. *J Atmos Chem* 2014;71(4):283–301.
 83. Zuraski K, Harkins C, Peischl J, et al. On-Road Measurements of Nitrogen Oxides, CO, CO₂, and VOC Emissions in Two Southwestern U.S. Cities. *ACS ES&T Air* 2025;2(4):589–598.
 84. Munir S, Mayfield M, Coca D. Understanding spatial variability of no₂ in urban areas using spatial modelling and data fusion approaches. *Atmosphere (Basel)* 2021;12(2):1–



- 20.
85. Voiculescu M, Constantin DE, Condurache-Bota S, Călmuc V, Roșu A, Bălănică CMD. Role of meteorological parameters in the diurnal and seasonal variation of no₂ in a Romanian urban environment. *Int J Environ Res Public Health* 2020;17(17):1–15.
86. Kendrick CM, Koonce P, George LA. Diurnal and seasonal variations of NO, NO₂ and PM_{2.5} mass as a function of traffic volumes alongside an urban arterial. *Atmos Environ* 2015;122:133–141.
87. Pancholi P, Kumar A, Bikundia DS, Chourasiya S. An observation of seasonal and diurnal behavior of O₃–NO_x relationships and local/regional oxidant (OX = O₃ + NO₂) levels at a semi-arid urban site of western India. *Sustainable Environment Research* 2018;28(2):79–89.
88. Chandra N, Lal S, Venkataramani S, Patra PK, Sheel V. Temporal variations of atmospheric CO₂ and CO at Ahmedabad in western India. *Atmos Chem Phys* 2016;16(10):6153–6173.
89. Huang X, Wang T, Talbot R, et al. Temporal characteristics of atmospheric CO₂ in urban Nanjing, China. *Atmos Res* 2015;153:437–450.
90. Hong JW, Lee SD, Lee K, Hong J. Seasonal variations in the surface energy and CO₂ flux over a high-rise, high-population, residential urban area in the East Asian monsoon region. *International Journal of Climatology* 2020;40(10):4384–4407.
91. Wei C, Wang M, Fu Q, Dai C, Huang R, Bao Q. Temporal characteristics of greenhouse gases (CO₂ and CH₄) in the megacity Shanghai, China: Association with air pollutants and meteorological conditions. *Atmos Res* 2020;235.
92. Masum MMH, Islam MB, Hossen MA, Nath H, Hoque A. Navigating pollution: The impact of transportation modes on air and noise quality in Chattogram City, Bangladesh. *J Transp Health* 2025;44.
93. Sun Y, Li X, Benmarhnia T, et al. Exposure to air pollutant mixture and gestational



- diabetes mellitus in Southern California: Results from electronic health record data of a large pregnancy cohort. *Environ Int* 2022;158.
94. Sulaymon ID, Zhang Y, Hopke PK, Zhang Y, Hua J, Mei X. COVID-19 pandemic in Wuhan: Ambient air quality and the relationships between criteria air pollutants and meteorological variables before, during, and after lockdown. *Atmos Res* 2021;250.
 95. Saha M, Kafy A Al, Bakshi A, et al. The urban air quality nexus: Assessing the interplay of land cover change and air pollution in emerging South Asian cities. *Environmental Pollution* 2024;361.
 96. Suthar G, Singh S, Kaul N, Khandelwal S. Diurnal variation of air pollutants and their relationship with land surface temperature in Bengaluru and Hyderabad cities of India. *Remote Sens Appl* 2024;35.
 97. Calama-González CM, Redondas D, Sabariego-Moreno K, Barbero-Barrera M del M. Statistical correlation analysis on indoor air high-priority pollutants in Spanish public primary schools. *Journal of Building Engineering* 2025;101.
 98. Putaud JP, Dingenen R Van, Alastuey A, et al. A European aerosol phenomenology - 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe. *Atmos Environ* 2010;44(10):1308–1320.
 99. Yoo HJ, Kim J, Yi SM, Zoh KD. Analysis of black carbon, particulate matter, and gaseous pollutants in an industrial area in Korea. In: *Atmospheric Environment*. Elsevier Ltd, 2011; p. 7698–7704.
 100. Shihab AS. Identification of Air Pollution Sources and Temporal Assessment of Air Quality at a Sector in Mosul City Using Principal Component Analysis. *Pol J Environ Stud* 2022;31(3):2223–2235.
 101. Ielpo P, Paolillo V, Gennaro G de, Dambruoso PR. PM10 and gaseous pollutants trends from air quality monitoring networks in Bari province: Principal component analysis and absolute principal component scores on a two years and half data set. *Chem Cent J* 2014;8(1).



102. Lina Thabethe ND, Engelbrecht JC, Wright CY, Oosthuizen MA. Human health risks posed by exposure to PM10 for four life stages in a low socio-economic community in South Africa. *Pan African Medical Journal* 2014;18.
103. Fu J, Lin Q, Ai B, et al. Associations between maternal exposure to air pollution during pregnancy and trajectories of infant growth: A birth cohort study. *Ecotoxicol Environ Saf* 2024;269.
104. Comotti A, Alberti I, Spolidoro GCI, et al. Air pollution and hospitalization risk in infants with bronchiolitis: A systematic review and meta-analysis. *Pediatric Allergy and Immunology*. 2025;36(5).
105. Berberian AG, Perera F, Arunachalam S, et al. Children's health impacts from a proposed decarbonization policy in the transportation sector in the Eastern United States. *Environmental Research Letters* 2024;19(4).
106. Cheng B, Ma Y, Qin P, et al. Characterization of air pollution and associated health risks in Gansu Province, China from 2015 to 2022. *Sci Rep* 2024;14(1).
107. Ahmadian F, Rajabi S, Maleky S, Baghapour MA. Spatiotemporal analysis of airborne pollutants and health risks in Mashhad metropolis: enhanced insights through sensitivity analysis and machine learning. *Environ Geochem Health* 2025;47(2).
108. Mondal A, Mondal S, Ghosh P, Das P. Analyzing the interconnected dynamics of domestic biofuel burning in India: unravelling VOC emissions, surface-ozone formation, diagnostic ratios, and source identification. *RSC Sustainability*. 2024;2(8):2150–2168.
109. Meo SA, Salih MA, Alkhalifah JM, Alsomali AH, Almushawah AA. Environmental pollutants particulate matter (PM2.5, PM10), Carbon Monoxide (CO), Nitrogen dioxide (NO2), Sulfur dioxide (SO2), and Ozone (O3) impact on lung functions. *J King Saud Univ Sci* 2024;36(7).
110. Sharma A, Kumar P. Quantification of air pollution exposure to in-pram babies and mitigation strategies. *Environ Int* 2020;139.

View Article Online

DOI: 10.1039/C4EA00036C



111. Liu T, Zhu Q, Wei J, et al. The Interactive and Joint Associations of Ambient PM_{2.5} and Temperature on the Onset of Acute Coronary Syndrome: Findings from The Chinese Cardiovascular Association (CCA) Database-Chest Pain Center Registry. *Environ Sci Technol* 2024;58(50):21978–21988.
112. Ye C, Tan Z, Li X, Ma X, Lu K, Zhang Y. Persistent Nitrogen Oxide Reductions Unmask Worsening Wintertime Photochemical Pollution in Urban Environments. *Environ Sci Technol* 2025;59(33):17383–17386.
113. Carabin A, Delpla I, Rodriguez M, Guilherme S, Dorea C. Subchronic Exposure to Regulated Trihalomethanes and Haloacetic Acids: Do We Need to Refine Our Exposure Assessment? A Full-Scale Case Study in a Canadian Municipality. *Environ Sci Technol* 2025;59(2):1100–1111.
114. Wang R, Lu H, Kang N, et al. Estimating under-five mortality attributable to fine particulate matter brought by wind-blown dust in 100 low- and middle-income countries in 2000–2017. *Natl Sci Rev* 2025;12(10).
115. Lu ZH, Liu C, Chen YJ, et al. Gestational Exposure to PM_{2.5} and Specific Constituents, Meconium Metabolites, and Neonatal Neurobehavioral Development: A Cohort Study. *Environ Sci Technol* 2024;58(23):9980–9990.
116. Ciarelli G, Colette A, Schucht S, et al. Long-term health impact assessment of total PM_{2.5} in Europe during the 1990–2015 period. *Atmos Environ X* 2019;3.
117. Akther T, Ahmed M, Shohel M, Ferdousi FK, Salam A. Particulate matters and gaseous pollutants in indoor environment and Association of ultra-fine particulate matters (PM₁) with lung function. *Environmental Science and Pollution Research* 2019;26(6):5475–5484.
118. Alam ZB, Mohiuddin KABM. Micro-characterization of Dust and Materials of Dust Origin at a Cement Industry Located in Bangladesh. *Aerosol Air Qual Res* 2023;23(1).
119. Trippetta S, Sabia S, Caggiano R. Fine aerosol particles (PM₁): natural and anthropogenic contributions and health risk assessment. *Air Qual Atmos Health*



2016;9(6):621–629.

View Article Online
DOI: 10.1039/D6EA00036C

120. Mainka A. Children health risk assessment of metals in total suspended particulate matter (Tsp) and pm1 in kindergartens during winter and spring seasons. *Atmosphere (Basel)* 2021;12(9).
121. Saju JA, Bari QH, Mohiuddin KABM, Strezov V. Measurement of ambient particulate matter (PM1.0, PM2.5 and PM10) in Khulna City of Bangladesh and their implications for human health. *Environmental Systems Research* 2023;12(1).
122. Liu T, Jiang Y, Hu J, et al. Association of ambient PM1 with hospital admission and recurrence of stroke in China. *Science of the Total Environment* 2022;828.
123. Lin F, Li G, Wang Y, et al. Impacts of air pollutions on cardiovascular and cerebrovascular diseases through inflammation: a comprehensive analysis of one million Chinese and half million UK individuals. *J Transl Med* 2025;23(1).
124. Miller MR. Air pollution and myocardial infarction in Poland. *The Lancet Regional Health - Europe*. 2024;41.
125. Ji JS, Dominici F, Gouveia N, Kelly FJ, Neira M. Air pollution interventions for health. *Nat. Med.* 2025;31(9):2888–2900.
126. Amnuaylojaroen T, Parasin N. Pathogenesis of PM2.5-Related Disorders in Different Age Groups: Children, Adults, and the Elderly. *Epigenomes*. 2024;8(2).
127. Chong-Neto HJ, Filho NAR. How does air quality affect the health of children and adolescents? *J. Pediatr. (Rio. J)*. 2025;101:S77–S83.
128. Ma J, Chiu YF, Kao CC, et al. Fine particulate matter manipulates immune response to exacerbate microbial pathogenesis in the respiratory tract. *European Respiratory Review*. 2024;33(173).
129. Hasnain MG, Garcia-Esperon C, Tomari YK, et al. Effect of short-term exposure to air pollution on daily cardio- and cerebrovascular hospitalisations in areas with a low level of air pollution. *Environmental Science and Pollution Research* 2023;30(46):102438–



- 102445.
130. Lepeule J, Laden F, Dockery D, Schwartz J. Chronic exposure to fine particles and mortality: An extended follow-up of the Harvard six cities study from 1974 to 2009. *Environ Health Perspect* 2012;120(7):965–970.
131. Ścibor M, Malinowska-Cieślik M. The association of exposure to PM10 with the quality of life in adult asthma patients. *Int J Occup Med Environ Health* 2020;33(3):311–324.
132. Young MT, Sandler DP, DeRoo LA, Vedal S, Kaufman JD, London SJ. Ambient air pollution exposure and incident adult asthma in a nationwide cohort of U.S. women. *Am J Respir Crit Care Med* 2014;190(8):914–921.
133. Xing YF, Xu YH, Shi MH, Lian YX. The impact of PM2.5 on the human respiratory system. *J. Thorac. Dis.* 2016;8(1):E69–E74.
134. Daellenbach KR, Uzu G, Jiang J, et al. Sources of particulate-matter air pollution and its oxidative potential in Europe. *Nature* 2020;587(7834):414–419.
135. Xin J, Chen H, Sun B, Ji Y, Yan L, Feng G. Response of air quality changes and spatio-temporal distribution characteristics on the energy development changes in China. *Environ Dev Sustain* 2025;
136. Guan C, Liu M, Shi J, Li Y. Temporal and spatial heterogeneity of tropospheric O3 and NO2 and health impact analysis in Shaanxi, Gansu, and Ningxia regions of China. *Environ Monit Assess* 2025;197(4).
137. Fukuto JM, Carrington SJ, Tantillo DJ, et al. Small molecule signaling agents: The integrated chemistry and biochemistry of nitrogen oxides, oxides of carbon, dioxygen, hydrogen sulfide, and their derived species. *Chem. Res. Toxicol.* 2012;25(4):769–793.
138. Kim YM, Kim YS, Jeon SG, Kim YK. Immunopathogenesis of allergic asthma: More than the Th2 hypothesis. *Allergy Asthma Immunol Res* 2013;5(4):189–196.
139. Stevens EL, Rosser F, Forno E, Peden D, Celedón JC. Can the effects of outdoor air pollution on asthma be mitigated? *Journal of Allergy and Clinical Immunology*.



2019;143(6):2016-2018.e1.

View Article Online
DOI: 10.1039/D6EA00036C

140. Yang S, Li M, Guo C, et al. Associations of long-term exposure to nitrogen oxides with all-cause and cause-specific mortality. *Nature Communications* 2025;16(1).
141. Zhou K, Tang L, Zhu C, et al. Recent Advances in Structure Design and Application of Metal Halide Perovskite-Based Gas Sensor. *ACS Sens.* 2024;9(9):4425–4449.
142. Chowdhury S, Haines A, Klingmüller K, et al. Global and national assessment of the incidence of asthma in children and adolescents from major sources of ambient NO₂. *Environmental Research Letters* 2021;16(3).
143. Dritsaki M, Dritsaki C. The Relationship Between Health Expenditure, CO₂ Emissions, and Economic Growth in G7: Evidence from Heterogeneous Panel Data. *Journal of the Knowledge Economy* 2024;15(1):4886–4911.
144. Satish U, Mendell MJ, Shekhar K, et al. Is CO₂ an indoor pollutant? direct effects of low-to-moderate CO₂ concentrations on human decision-making performance. *Environ Health Perspect* 2012;120(12):1671–1677.
145. Jacobson TA, Kler JS, Hernke MT, Braun RK, Meyer KC, Funk WE. Direct human health risks of increased atmospheric carbon dioxide. *Nat. Sustain.* 2019;2(8):691–701.



- 1 The data supporting this article have been included as part of the Supplementary Information

