

Environmental Science Atmospheres

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



	<p>Physics, Physic Gordon, Hamish; Carnegie Mellon University, Chemical Engineering; Carnegie Mellon University, Center for Atmospheric Particulate Studies Granzin, Manuel; Goethe University Frankfurt, Institute for Atmospheric and Environmental Sciences Große Schute, Lena; Goethe University Frankfurt Institute for Atmospheric and Environmental Sciences Hartl, Herbert; University of Helsinki Institute for Atmospheric and Earth System Research; Airmodus Oy Hirvensalo, Ella; University of Helsinki Institute for Atmospheric and Earth System Research Jacobshagen, Aenne; University of Vienna, Faculty of Physics Judmaier, Bernhard; University of Innsbruck, Institute for Applied and Ion Physics Kaniyodical Sebastian, Milin; Karlsruhe Institute of Technology, IMKAAF Klebach, Hannah; Goethe-Universität Frankfurt am Main, Institute for Atmospheric and Environmental Sciences Koemets, Paap; University of Tartu, Institute of Physics, Faculty of Science and Technology; Airel Ltd Kohl, Matthias ; Max-Planck-Institut für Chemie, Atmospheric Chemistry Department Konrat, Ruth; University of Vienna, Aerosol Physics and Environmental Physics Krüger, Timm; Goethe-Universität Frankfurt am Main, Institut für Atmosphäre und Umwelt Kunkler, Felix; Max Planck Institute for Chemistry, Atmospheric Chemistry Kuerten, A.; Goethe-Universität Frankfurt am Main, Institute for Atmospheric and Environmental Sciences Leiminger, Markus; Ionicon Analytik GesmbH Lietzke, Clara; University of Colorado Boulder, Chemistry; University of Colorado Boulder, Cooperative Institute for Research in Environmental Sciences Liu, Lu; Paul Scherrer Institut PSI, PSI Center for Energy and Environmental Sciences Mauldin, Roy; Carnegie Mellon University, Department of Chemistry; University of Colorado Boulder, Department of Atmospheric and Oceanic Sciences Mentler, Bernhard; University of Innsbruck, Institute for Ion Physics and Applied Physics Morawiec, Aleksandra; University of Vienna, Faculty of Physics Müller, Markus; Ionicon Analytik GesmbH Petäjä, Tuukka; University of Helsinki, Institute for Atmospheric and Earth System Research (INAR) Pignatelli, Alessia; Paul Scherrer Institut PSI, Center for Energy and Environmental Sciences Rato, Pedro; CERN; Goethe-Universität Frankfurt am Main, Institute for Atmospheric and Environmental Sciences Reinecke, Tobias; Ionicon Analytik GesmbH Richter, Sarah; Goethe University Frankfurt, Institute for Atmospheric and Environmental Sciences Rörup, Birte; University of Helsinki, Institute for Atmospheric and Earth System Research/Physics Ruhl, Samuel; Max Planck Institute for Chemistry, Atmospheric chemistry department Russell, Douglas; Goethe-Universität Frankfurt am Main, Institute for Atmospheric and Environmental Sciences Scholz, Wiebke; University of Innsbruck, Institute of Ion Physics and Applied Physics Shen, Jiali; University of Helsinki, Institute for Atmospheric and Earth System Research/Physics, Faculty of Science</p>
--	--

Stinchfield, Alexandria; Carnegie Mellon University, Department of Chemical Engineering; Carnegie Mellon University, Center for Atmospheric Particle Studies

Thakur, Roseline; University of Helsinki, Institute for Atmospheric and Earth System Research/Physics

Tong, Yandong; University of Colorado Boulder, Department of Chemistry; University of Colorado Boulder, Cooperative Institute for Research in Environmental Sciences

Top, Jens; Paul Scherrer Institut PSI, Center for Energy and Environmental Sciences;; University of Helsinki, Faculty of Science, Institute for Atmospheric and Earth System Research/Physics

Umo, Nsikanabasi Silas; University of North Carolina Wilmington, Department of Chemistry and Biochemistry

Weissbacher, Jakob; University of Innsbruck, Institute for ion and applied physics

Yang, Boxing; Paul Scherrer Institut PSI, Laboratory of Atmospheric Chemistry

Zauner-Wieczorek, Marcel; Goethe University Frankfurt, Institute for Atmospheric and Environmental Sciences

Zhang, Jiangyi; University of Helsinki, Institute for Atmospheric and Earth System Research/Physics

Amorim, Antonio; LIP; Universidade de Lisboa, Faculdade de Ciencias

Baltensperger, Urs; Paul Scherrer Institut PSI, Laboratory of Atmospheric Chemistry

Christoudias, Theodoros; The Cyprus Institute, Climate and Atmosphere Research Centre

Curtius, Joachim; Goethe-Universitat Frankfurt am Main, Institute for Atmospheric and Environmental Sciences

Donahue, Neil; Carnegie Mellon University, Department of Chemical Engineering; Carnegie Mellon University, Department of Chemistry; Carnegie Mellon University, Center for Atmospheric Particulate Studies; Carnegie Mellon University, Department of Engineering & Public Policy

Flagan, Richard; California Institute of Technology Division of Chemistry and Chemical Engineering,

El Haddad, Imad; Paul Scherrer Institut PSI, Center for Energy and Environmental Sciences; PKU, College of Environmental Sciences and Engineering

Hansel, Armin; University of Innsbruck, Institute for ion and applied physics

Harder, Hartwig; Max Planck Institute for Chemistry, Atmospheric Chemistry

He, Xucheng; University of Helsinki, Institute for Atmospheric and Earth System Research/ Physics, Faculty of Science

Junninen, Heikki; University of Tartu, Laboratory of Atmospheric and Environmental Sciences

Kulmala, Markku; University of Helsinki, Faculty of Science, Institute for Atmospheric and Earth System Research/Physics

Lehtipalo, Katrianne; University of Helsinki, Faculty of Science, Institute for Atmospheric and Earth System Research/Physics

Lelieveld, Jos; Max Planck Institute for Chemistry, Atmospheric Chemistry

Moehler, Ottmar; Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research

Schobesberger, Siegfried ; University of Eastern Finland, Department of Technical Physics

Tomé, António; Universidade da Beira Interior, IDL

Volkamer, Rainer; University of Colorado Boulder, Department of Chemistry; University of Colorado Boulder, Cooperative Institute for Research in Environmental Sciences

Worsnop, Doug; University of Helsinki, Faculty of Science, Institute for Atmospheric and Earth System Research/Physics; Aerodyne Research Inc

Onnela, Antti; CERN



	Winkler, Paul; University of Vienna, Faculty of Physics Kirkby, Jasper; CERN, EP Department; Goethe University Frankfurt, Institute for Atmospheric and Environmental Sciences



Open Access Article. Published on 09 June 2026. Downloaded on 6/13/2026 4:16:05 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Accurate representation of new particle formation from sulfuric acid and from oxidized organics is crucial for capturing aerosol effects in global climate models. Existing parameterizations are based on limited experimental conditions and do not accurately capture combined acid/organic nucleation effects or their temperature dependence. Here, we report chamber measurements of nucleation rates from α -pinene oxidation products and sulfuric acid in the near-absence of base vapours such as ammonia, under conditions representative of the cool boundary layer. We show that their synergistic interaction enhances particle formation beyond either component alone. Based on these results, we develop a parameterization accounting for vapour concentrations, ionisation and temperature. This framework improves the representation of a major aerosol formation source in global models.



Cite this: DOI: 00.0000/xxxxxxxxxx

Nucleation of α -pinene oxidation products with sulfuric acid

Eva Sommer^{*ab}, João Almeida^{ac}, Wenjuan Yu^d, Mario Simon^e, Felix Möller^f, Christos Xenofontos^g, Andrea Pozzer^{gh}, Zhensen Zheng^{ij}, Nirvan Bhattacharyya^e, Dina Alfaouri^d, Samira Atabakhsh^k, Rima Baalbaki^g, Hannah Beckmann^f, Moritz Berntheusel^e, Pia Bogert^l, Mattia Busato^a, Manjula Canagaratna^m, Luía Caudillo-Plath^e, Anouck Chassaing^{no}, Romulo Cruz-Simbron^{pq}, Lubna Dada^r, Jenna DeVivost, Jonathan Duplissy^{du}, Hamish Gordon^{vt}, Manuel Granzin^e, Lena Große Schute^e, Herbert G. Hartl^{dw}, Ella Hirvensalo^d, Aenne Jacobshagen^b, Bernhard Judmaierⁱ, Milin Kaniyodical Sebastian^l, Hannah Klebach^e, Paap Koemets^{fx}, Matthias Kohl^h, Ruth Konrat^b, Timm Krüger^e, Felix Kunkler^h, Andreas Kürten^e, Markus Leiminger^j, Clara J. Lietzke^{pq}, Lu Liu^r, Roy Mauldin^{sy}, Bernhard Mentlerⁱ, Aleksandra Morawiec^b, Markus Müller^j, Tuukka Petäjä^d, Alessia Pignatelli^r, Pedro Rato^{ae}, Tobias Reinecke^j, Sarah Richter^e, Birte Rörup^d, Samuel Ruhl^h, Douglas M. Russell^e, Wiebke Scholzⁱ, Jiali Shen^{du}, Alexandria Stinchfield^{vt}, Roseline C. Thakur^d, Yandong Tong^{pq}, Jens Toprd, Nsikanabasi Silas Umo^z, Jakob Weissbacherⁱ, Boxing Yang^r, Marcel Zauner-Wieczorek^e, Jiangu Zhang^d, Antonio Amorim^{cα}, Urs Baltensberger^r, Theodoros Christoudias^g, Joachim Curtius^e, Neil M. Donahue^{vstβ}, Imad El Haddad^{rδ}, Richard C. Flagan^γ, Armin Hanselⁱ, Hartwig Harder^h, Xu-Cheng He^d, Heikki Junninen^ε, Markku Kulmala^d, Katrianne Lehtipalo^{dζ}, Jos Lelieveld^{hg}, Ottmar Möhler^l, Siegfried Schobesbergerⁿ, António Tomé^θ, Rainer Volkamer^{pq}, Douglas R. Worsnop^{dm}, Antti Onnela^a, Paul M. Winkler^b and Jasper Kirkby^{*ae}

Received Date
Accepted Date

DOI: 00.0000/xxxxxxxxxx

Particle nucleation from trace atmospheric vapours is important for climate since it gives rise to more than half of global cloud condensation nuclei. Sulfuric acid (H_2SO_4) has long been recognised to drive particle nucleation in the atmosphere and, more recently, highly oxygenated products of biogenic vapours—in particular monoterpenes such as α -pinene ($\text{C}_{10}\text{H}_{16}$)—have also been shown to nucleate under atmospheric conditions, without requiring additional vapours. This raises the question of whether a nucleation synergy exists between α -pinene oxygenated organic molecules (AP-OOM) and H_2SO_4 , as has been suggested by early studies. Here we report new particle formation from AP-OOM and H_2SO_4 in the absence of base vapours such as ammonia (NH_3), measured in experiments performed with the CERN CLOUD (Cosmics Leaving Outdoor Droplets) chamber at cool boundary layer temperatures of $-10\text{ }^\circ\text{C}$ and $+5\text{ }^\circ\text{C}$. We find that AP-OOM nucleation rates increase strongly when H_2SO_4 concentrations exceed around 10^6 cm^{-3} . The enhancement is synergistic and cannot be explained as a simple linear addition of independent chemical systems. Above this threshold, the nucleation rate depends approximately linearly on H_2SO_4 concentration, in contrast with the strong sensitivity to H_2SO_4 for H_2SO_4 - NH_3 nucleation. Nucleation rates are 10–100-fold higher in the presence of ions from galactic cosmic rays or from the CERN pion beam. Based on these measurements, we have parameterised a temperature-dependent H_2SO_4 -AP-OOM nucleation rate in the absence of base vapours and implemented it in the EMAC (ECHAM/MESSy Atmospheric Chemistry) Earth system model. In comparison with a parameterisation developed in an earlier study [Riccobono et al., *Science*, 2014, **344**, 717–721.], the new parameterisation indicates sharply reduced nucleation rates in the boundary layer over warm regions, and increased rates over northern boreal forests.



1 Introduction

Aerosol particles and aerosol-cloud interactions are the largest source of uncertainty in global climate projections¹. It is estimated that more than half of all cloud condensation nuclei (CCN) arise from new particle formation (NPF)². NPF involves the formation of molecular clusters from low-volatility vapours, a nucleation process. Once these clusters exceed a diameter of 1–2 nm

they become stable against evaporation and, if not lost to pre-existing aerosol particles or other surfaces, may grow above sizes of around 50 nm where they can act as CCN³.

Sulfuric acid (H₂SO₄) has long been recognised as a key driver of atmospheric new particle formation across a wide range of environments^{4–6}. Via autoxidation⁷, biogenic vapours such as α -pinene (C₁₀H₁₆) can also rapidly give rise to highly oxygenated compounds with extremely low volatility. These compounds can condense onto particles and drive their growth, forming secondary organic aerosol and CCN over extensive continental regions⁸. Highly oxygenated organic molecules from α -pinene (AP-OOM) have been shown to nucleate even in the absence of H₂SO₄⁹. Alpha-pinene is the most abundant monoterpene, and its low-volatility oxidation products play an important role in global CCN production, especially in the pristine pre-industrial atmosphere². While isoprene (C₅H₈) suppresses particle formation in the boundary layer¹⁰, a recent study at the CERN CLOUD (Cosmics Leaving Outdoor Droplets) chamber showed that isoprene oxygenated organic molecules (IP-OOM) can drive NPF at the cold temperatures of the upper-troposphere¹¹. The nucleation rates from IP-OOM were enhanced by about two orders of magnitude in the presence of trace amounts of H₂SO₄¹¹. This raises the question of whether a nucleation synergy also exists between AP-OOM and H₂SO₄ in the absence of base vapours, as suggested by previous experiments^{12,13,14}.

Global climate models require parameterisations of nucleation rates in order to evaluate the climate impact of NPF and CCN¹⁵. A temperature- and ionisation rate-dependent parameterisation of H₂SO₄(-NH₃)-H₂O nucleation was developed from experiments at the CERN CLOUD chamber by Dunne *et al.*⁶ and has been implemented in several global climate models. Kirkby *et al.*⁹ parameterised pure AP-OOM nucleation rates at +5 °C. The mixed system of H₂SO₄ with oxygenated organics and no added base vapours was investigated at CLOUD by Riccobono *et al.*¹², while Lehtipalo *et al.*¹⁴ investigated the system of H₂SO₄ and oxygenated organics with different levels of NO_x and NH₃ at +5 °C.

Riccobono *et al.*¹² used a quantity referred to as BioOxOrg as a proxy for α -pinene oxygenated organic molecules. The BioOxOrg was produced from pinanediol (PD; C₁₀H₁₈O₂), a first-generation oxidation product of α -pinene. BioOxOrg was defined as the total reaction products from PD oxidation by hydroxyl (OH) radicals, so it included compounds with a wide range of oxygen content. A subset of BioOxOrg comprised the same highly-oxygenated AP-OOM compounds measured in the present study that drive nucleation.

Riccobono *et al.*¹² made an important advance at the time by revealing rapid formation of highly-oxygenated α -pinene products with sufficiently low volatility that, together with H₂SO₄, could account for nucleation rates observed in the atmosphere. However, the experiments were limited in several respects: pure biogenic nucleation was unknown, and no experiments were performed with H₂SO₄ below 10⁶ cm⁻³; a pinanediol proxy was used instead of α -pinene so the BioOxOrg did not include the high yield of extremely low volatility OOM from ozone-initiated autoxidation of α -pinene; contaminant ammonia, which strongly affects nucleation rates from H₂SO₄, could not be excluded below

^a CERN, European Organization for Nuclear Research, 1211 Geneva, Switzerland

^b Faculty of Physics, University of Vienna, Strudlhofgasse 4, 1090 Vienna, Austria

^c LIP - Laboratory for Instrumentation and Experimental Particle Physics, Av. Prof. Gama Pinto, n.2, 1649-003 Lisboa, Portugal

^d Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, 00014 Helsinki, Finland

^e Goethe University Frankfurt, Institute for Atmospheric and Environmental Sciences, Altenhoferallee 1, 60438 Frankfurt am Main, Germany

^f Institute of Physics, Faculty of Science and Technology, University of Tartu, 50411, Tartu, Estonia

^g Climate and Atmosphere Research Center, The Cyprus Institute, 1645 Nicosia, Cyprus

^h Atmospheric Chemistry Department - Max Planck Institute for Chemistry, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

ⁱ Institute for ion and applied physics, University of Innsbruck, Technikerstraße 25, 6020 Innsbruck, Austria

^j IONICON Analytik GmbH, 6020 Innsbruck, Austria

^k Leibniz-Institute for Tropospheric Research, Permoser Straße 15, 04318 Leipzig, Germany

^l Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

^m Aerodyne Research, Billerica, MA 01821, USA

ⁿ Department of Environmental Science, Stockholm University, Stockholm, 10961 Sweden

^o Bolin Centre for Climate Research, Stockholm University, Stockholm, 10961 Sweden

^p Department of Chemistry, University of Colorado Boulder, Cristol Chemistry, 215 UCB, Boulder, CO, USA

^q CIRES, Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO 80309-0215, USA

^r PSI Center for Energy and Environmental Sciences, Paul Scherrer Institute, 5232 Villigen PSI, Switzerland

^s Department of Chemistry, Carnegie Mellon University, 5000 Forbes Ave Pittsburgh, PA 15213 USA

^t Center for Atmospheric Particulate Studies, Carnegie Mellon University, 5000 Forbes Ave Pittsburgh, PA 15213 USA

^u Helsinki Institute of Physics, University of Helsinki, Helsinki, Finland

^v Department of Chemical Engineering and Center for Atmospheric Particle Studies, Carnegie Mellon University, Pittsburgh, PA 15213, USA

^w Airmodus Oy., Erik Palménin aukio 1, 00560 Helsinki, Finland

^x Airel Ltd, Observatooriumi 5, 61602 Tõravere, Estonia

^y Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, Boulder, CO 80309, USA

^z Department of Chemistry and Biochemistry, University of North Carolina Wilmington, NC 28403, USA

^{aa} Faculdade de Ciências da Universidade de Lisboa, Edifício C8, Campo Grande, 1749-016 Lisboa, Portugal

^{ab} Department of Engineering and Public Policy, Carnegie Mellon University, 5000 Forbes Ave, Pittsburgh, PA 15213 USA

^{ac} California Institute of Technology, Pasadena, CA 91125, USA

^{ad} College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

^{ae} Laboratory of Atmospheric and Environmental Sciences, University of Tartu, W. Ostwaldi tn 1, 50411, Tartu, Estonia

^{af} Finnish Meteorological Institute, Helsinki, Finland

^{ag} Department of Technical Physics, University of Eastern Finland, PO Box 1627, 70211 Kuopio, Finland

^{ah} IDL-Universidade da Beira Interior, Rua Marquês D'Ávila e Bolama 6201-001 Covilhã, Portugal



around 30 pptv; and the experiments were carried out at a single temperature (5 °C). The resulting parameterisation of the nucleation rate is a simple power law, $J = k [\text{H}_2\text{SO}_4]^p [\text{BioOxOrg}]^q$, with no dependence on ionisation rates or temperature¹².

Here we address these limitations in experiments performed at the CERN CLOUD chamber⁴ to investigate nucleation of AP-OOM with trace concentrations of H_2SO_4 and in the absence of base vapours (< 2 pptv NH_3 contaminant level). We report nucleation rates from AP-OOM measured under cool boundary layer conditions between -10 °C and +5 °C, both in the presence and absence of H_2SO_4 . We also quantify the contribution of ion-induced nucleation. The measurements allow us to characterise the synergy between AP-OOM and H_2SO_4 and to parameterise the nucleation rate as a function of vapour concentrations, ionisation rate and temperature.

2 Methods

2.1 The CERN CLOUD experiment

We conducted all measurements at the CERN CLOUD experiment⁴. The CLOUD chamber is a 26 m³ continuously-stirred stainless-steel reactor. The chamber has ultra-low contaminants, so controlled experiments can be performed with trace gases at atmospheric concentrations. New particle formation experiments are performed under steady-state conditions, where the loss rates of gases and particles balance their injection or production rates. Trace gas concentrations, particle size distributions, and chamber conditions are monitored using a suite of state-of-the-art instruments that continuously sample the contents of the chamber. A continual injection of around 380 L min⁻¹ humidified ultrapure air plus precursor vapours compensates for the sampling losses.

2.2 Chamber conditions

We conducted the experiments during the CLOUD15, CLOUD16, and CLOUD17 campaigns in autumn 2022, 2023, and 2024, respectively. We investigated new particle formation from oxygenated organic molecules derived from α -pinene and H_2SO_4 under conditions representative of the cool boundary layer. Experiments were performed at temperatures of -10 °C and +5 °C.

We injected α -pinene into the chamber from an evaporator and oxidised it via ozonolysis to form AP-OOM. H_2SO_4 was produced *in situ* by oxidation of SO_2 with OH radicals generated from photolysis of O_3 with ultraviolet radiation below 320 nm¹⁶.

We measured nucleation rates for AP-OOM concentrations ranging from < 10⁵ cm⁻³ (instrument detection limit) to 10⁸ cm⁻³, and for H_2SO_4 concentrations between < 10⁴ cm⁻³ (instrument detection limit) and approximately 10⁷ cm⁻³.

At selected vapour concentrations, we measured nucleation rates under three ionisation conditions: a) *neutral* conditions, where all ions were removed from the chamber using an electric clearing field, b) *GCR* conditions, where the chamber was exposed to ground level ionisation from galactic cosmic rays, and c) *beam* conditions, where elevated ionisation rates were generated using a π^+ beam from the CERN Proton Synchrotron, simulating ionisation at higher altitudes. These three ionisation conditions are referred to as *neutral*, *GCR*, and *beam*, respectively.

We maintained the relative humidity at ~ 60 % at both temperatures. Ozone (O_3) concentrations were ~ 80 ppbv during the -10 °C experiments and ~ 40 ppbv during the +5 °C experiments. All experiments in CLOUD16 and CLOUD17, except for one reference run per campaign, were conducted in the presence of UV illumination¹⁶. No NH_3 was added to the chamber during or immediately before the experiments. The chamber background NH_3 concentration is estimated to be below 2 pptv, based on API-TOF¹⁷ measurements showing a very low abundance of NH_3 -containing H_2SO_4 clusters under comparable conditions¹⁸. The CERN CLOUD chamber is operated as a continuously stirred steady-state reactor with a constant flow. The total air flow in our experiments was 370-380 L min⁻¹, corresponding to a trace-gas residence time of ~ 70 min. Two mixing fans mounted at the top and bottom of the chamber ensured uniform trace gas distributions. All nucleation and growth rates reported here were obtained under steady-state conditions. We studied the mixed H_2SO_4 -AP-OOM system during the CLOUD16 and CLOUD17 campaigns and the pure biogenic system during CLOUD15. For the development of the H_2SO_4 -AP-OOM parameterisation, we additionally include nucleation rates for the H_2SO_4 -AP-OOM system reported previously by Kirkby *et al.*⁹.

2.3 Precursor gas and condensable vapour measurement

We measured α -pinene concentrations using proton-transfer-reaction mass spectrometry (PTR-MS) during all campaigns. During CLOUD15 and CLOUD17, α -pinene was measured with a H_3O^+ -PTR-MS (STOF)^{19,20}. The lower limit of detection of the STOF instrument for α -pinene is ~ 150 pptv. During CLOUD16, a FUSION PTR-MS²¹, with a sensitivity for α -pinene down to <1 pptv, was used as the primary α -pinene measurement, while the STOF was measuring in parallel to allow an instrument comparison. Both instruments were regularly calibrated using gas standards, and instrumental backgrounds were subtracted. We used the measured α -pinene concentrations from both instruments to develop an instrument- and campaign-independent relationship allowing us to calculate α -pinene concentrations in the chamber directly from the injection rate (i.e. mass flow controller, MFC, settings). A more detailed description of this approach is provided in the supplementary information.

H_2SO_4 and AP-OOM were measured using a nitrate chemical ionisation atmospheric pressure interface time-of-flight mass spectrometer (NO_3^- -CI-API-TOF, hereafter referred to as nitrate-CIMS)^{22,23}. H_2SO_4 concentrations were calibrated in each campaign using a well-established procedure^{24,25}, yielding absolute concentrations. We assume the same detection efficiency for AP-OOM as for H_2SO_4 . This is appropriate for the more highly oxygenated molecules (HOM)²⁶ with $\text{O} \geq 6$, which are the subset of AP-OOM that drive NPF. The overall uncertainty of nitrate-CIMS measurements is a factor of two. A more detailed instrument description is provided in the supplementary information.

2.4 Definition of AP-OOM

Previous CLOUD studies have shown that AP-OOM nucleation and early growth is driven by exceedingly low volatility com-



pounds^{27–29}, which are known as extremely-low-volatility- and ultra-low-volatility organic compounds (ELVOC and ULVOC, respectively) in the framework of the Volatility Basis Set, VBS^{30,31}. For this work, AP-OOM are defined as the entire subset of oxygenated compounds from α -pinene that are measured by a nitrate-CIMS, as previously used by Wang *et al.*³²; no additional requirements (number of carbon atoms, O:C ratio or mass range) are applied.

Nitrate chemical ionisation is sensitive to AP-OOM with more than around 6 oxygen atoms³³. Since all our experiments are without NO_x , the detected OOM consist exclusively of $\text{C}_x\text{H}_y\text{O}_z$ compounds. In the temperature range of our study, nitrate chemical ionisation therefore efficiently detects ELVOC and ULVOC, while discriminating against less-volatile compounds³⁴. We measure a mean AP-OOM yield per ozonolysis reaction (including secondary OH reactions) of 6.0 %. Within experimental differences, this is consistent with the ELVOC yields for α -pinene oxidation (3.4 % from O_3 and 0.44 % from OH) reported by Jokinen *et al.*³⁵ using a nitrate-CIMS, which are used in the EMAC model simulations reported here.

To ensure consistent AP-OOM concentrations across campaigns, we compared the measured nitrate-CIMS signal to the α -pinene ozonolysis rate, using the measured O_3 concentration and the α -pinene concentration derived from MFC settings, after calibration with mass spectrometers. We established separate temperature-dependent relationships for -10°C and $+5^\circ\text{C}$ and applied them to the datasets from CLOUD15–17. For earlier data from Kirkby *et al.*⁹, direct AP-OOM measurements were not available. We therefore estimate AP-OOM from the nitrate-CIMS C_{10} organic signal, which correlates strongly with AP-OOM in CLOUD15–17 ($R^2 = 0.99$). Further details are provided in the supplementary information and Fig. S1.

2.5 Aerosol particle and charged cluster measurement

We measured naturally charged clusters with an Aerodyne Research Inc. APi-TOF operated in negative-ion mode¹⁷. Particles up to ~ 3 nm were measured with an Airmodus Ltd. nano-condensation nucleus counter (nCNC) consisting of a particle size magnifier (PSM)³⁶ coupled to a butanol condensation particle counter (CPC). We monitored total particle number concentrations above 2.5 nm with a butanol CPC (TSI 3776). We measured particle number size distributions from 2–66 nm and 18–533 nm with two Scanning Mobility Particle Sizer (SMPS) systems³⁷: a nano-SMPS (TSI 3938) coupled to a butanol CPC (TSI 3776) and a long-SMPS (TSI 3082) coupled to a butanol CPC (TSI 3775), respectively. We used an Airel Ltd. Neutral cluster and Air Ion Spectrometer (NAIS)³⁸, to measure the naturally charged particle number size distribution from ~ 1 to 41 nm, as well as the total particle number size distribution from 2 to 42 nm in both positive and negative polarity.

We determined particle growth rates (GRs) from 3.2 to 8 nm using the appearance-time method described in Dada *et al.*³⁹. In CLOUD15, we obtained the input size distributions for the growth-rate calculation with a DMA-train configured as described by Stolzenburg *et al.*⁴⁰. In CLOUD16 and CLOUD17, we obtained

the size distributions for the growth rate calculation from NAIS measurements.

We calculated the nucleation rates of particles at 1.7 nm diameter threshold ($J_{1.7}$) following the method described in Dada *et al.*³⁹. Ideally, when measuring and parameterising nucleation rates, one would choose the size threshold to correspond to the critical size, i.e. the size at which the growth rate of a particle begins to exceed its evaporation rate. However, the critical size depends on the chemical system and temperature. CLOUD has therefore chosen a practical solution to quote particle nucleation rates at 1.7 nm mobility diameter, which is safely above the critical size for all chemical systems that we have studied so far. For example, in the case of sulfuric acid, it corresponds around 9 H_2SO_4 molecules, whereas the critical cluster contains 2–4 H_2SO_4 molecules equivalent to 1.1–1.3 nm mobility diameter. However, with this choice, under slow GR conditions the condensation sink (CS) can reduce the survival probability of the particle in growing from the critical size to 1.7 nm. At CLOUD the CS under standard conditions is $2.2 \cdot 10^{-3} \text{ s}^{-1}$, equal to the wall loss rate. Our studies have shown that, for a GR equivalent to above around $3 \cdot 10^6 \text{ cm}^{-3} \text{ H}_2\text{SO}_4$, i.e. 0.6 nm/h, in the CLOUD chamber, the difference between $J_{1.7}$ and the nucleation rate at the critical size is relatively small⁴¹.

2.6 EMAC global model implementation

Based on our measurements, as well as the inorganic H_2SO_4 - NH_3 parameterisation from Dunne *et al.*⁶, we derived a new parameterisation for the mixed H_2SO_4 -AP-OOM nucleation mechanism and implemented it in a global model. We performed all simulations with the ECHAM/MESy Atmospheric Chemistry (EMAC) model, which integrates the ECHAM5 general circulation model⁴² with the MESy2 framework⁴³ to represent meteorology, trace gases, and aerosol processes. Further information on the model configuration, as well as a full description of the H_2SO_4 -AP-OOM parameterisation developed in this study, is provided in the supplementary information.

3 Experimental results

3.1 H_2SO_4 -AP-OOM nucleation

Fig. 1 illustrates a representative experiment. We first establish pure biogenic nucleation from AP-OOM and subsequently introduce SO_2 to produce H_2SO_4 and quantify its effect on the nucleation rate. The experiment shown in Fig. 1 was conducted at -10°C during the CLOUD16 campaign under galactic cosmic ray (GCR) ionisation with ionisation rate $Q \approx 2 \text{ cm}^{-3} \text{ s}^{-1}$, at approximately 60 % relative humidity (RH), ~ 80 ppbv O_3 , and continuous UV illumination¹⁶. Prior to each experiment, we cleaned the chamber by operating the mixing fans at maximum rotation speed to increase the loss rates of aerosol particles and condensable vapours. During these cleaning periods, we establish stable conditions of temperature, humidity, O_3 , and α -pinene concentrations. For all experiments reported here, no NH_3 was added to the chamber during or prior to the measurements. The low abundance of NH_3 -containing H_2SO_4 clusters observed in the APi-TOF is consistent with contaminant NH_3 concentrations be-



low 2 pptv¹⁸.

The first vertical dashed line in Fig. 1 indicates the start of the experiment, where we reduced the mixing fan speed from 100 to 12 %. This reduces the loss rate of condensable vapours and increases their lifetime in the chamber. AP-OOM reach a new equilibrium concentration of around $3 \cdot 10^7 \text{ cm}^{-3}$ (Fig. 1 panel a). The new steady-state AP-OOM concentration initiated new particle formation with particles emerging directly from the ion band (Fig. 1c), which is consistent with the strong ion-induced contribution previously observed for pure biogenic nucleation⁹.




Fig. 1 Evolution of an example nucleation experiment at $-10 \text{ }^\circ\text{C}$ under GCR conditions. a) AP-OOM and H_2SO_4 concentrations (left axis) and the corresponding nucleation rate at 1.7 nm , $J_{1.7}$ (right axis). b) Total particle number size distribution between 2 and 40 nm , measured by the NAIS. The dashed black curves show particle growth from ~ 3 to 8 nm , together with their fitted growth rates. c) Naturally-charged size distribution of negative ions and charged particles between 1 and 40 nm , measured by the NAIS. Vertical dashed lines indicate when steady-state gas concentrations were changed: first an increase of AP-OOM and then, in two steps, an increase of SO_2 to produce H_2SO_4 . The experimental conditions are 80 ppbv O_3 , $<2 \text{ pptv NH}_3$, $120 \text{ pptv } \alpha\text{-pinene}$, $0\text{--}100 \text{ pptv SO}_2$, 60 \% RH and $-10 \text{ }^\circ\text{C}$.

After the AP-OOM-driven nucleation rate reached a steady state, we introduced SO_2 , which is rapidly oxidised by OH to form H_2SO_4 . We increased the H_2SO_4 concentration in two steps (Fig. 1a), reaching approximately $2 \cdot 10^6 \text{ cm}^{-3}$. At this relatively low H_2SO_4 concentration, the nucleation rate doubled from ~ 1.2 to $2.5 \text{ cm}^{-3} \text{ s}^{-1}$. Taking only the $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation mechanism from Dunne *et al.*⁶ into account, we would expect $J_{1.7} \sim 10^{-5} \text{ cm}^{-3} \text{ s}^{-1}$ at 2 pptv contaminant NH_3 . Even if we assume an unrealistically high NH_3 concentration of 100 pptv , the $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation rate predicted by Dunne *et al.*⁶ would remain below $4 \cdot 10^{-4} \text{ cm}^{-3} \text{ s}^{-1}$. We therefore attribute the observed increase in nucleation rate in Fig. 1 to a synergistic $\text{H}_2\text{SO}_4\text{-AP-OOM}$ nucleation pathway rather than to an independent contribution from $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation. Within experimental errors, the particle growth rate remained unchanged following the ad-

dition of $2 \cdot 10^6 \text{ cm}^{-3} \text{ H}_2\text{SO}_4$ (Fig. 1b), consistent with the small expected increase ($\sim 0.2 \text{ nm h}^{-1}$)⁴⁴.

3.2 Charged cluster composition

Fig. 2 shows APi-TOF mass-defect plots of negatively charged clusters for four steady-state nucleation events at $-10 \text{ }^\circ\text{C}$, spanning a range of H_2SO_4 and AP-OOM relative concentrations. Panels a)-d) are ordered by increasing $\text{H}_2\text{SO}_4\text{:AP-OOM}$ ratio. In each panel, the mass defect (Th) is plotted against the nominal mass-to-charge ratio (Th), such that clusters with similar elemental composition form characteristic bands^{17,27,45}. The symbol colour indicates the number of sulfur atoms in the cluster, and the symbol radius is proportional to the logarithm of the signal intensity.

Fig. 2a) represents pure biogenic AP-OOM nucleation and exhibits the characteristic C_{10} monomer (Th $\approx 240\text{--}420$) and C_{20} dimer (Th $\approx 400\text{--}620$) bands of AP-OOM²⁷. No sulfur-containing clusters are detected; instead, under these conditions the AP-OOM species are clustered with nitrate anions (NO_3^-) and AP-OOM⁻ ions. As the $\text{H}_2\text{SO}_4\text{:AP-OOM}$ ratio increases (panels b) to d), sulfur-containing clusters become increasingly abundant. In panels b) and c), the contribution of HSO_4^- anions appears and grows. Moreover, the organic bands include a substantial fraction containing two sulfur atoms, indicating that neutral H_2SO_4 is also present in the mixed clusters, and demonstrating synergistic nucleation of $\text{H}_2\text{SO}_4\text{-AP-OOM}$.

Clusters with negative mass defect correspond to H_2SO_4 and $\text{H}_2\text{SO}_4\text{-NH}_3$ clusters. NH_3 -containing clusters are not observed until the $\text{H}_2\text{SO}_4\text{:AP-OOM}$ ratio exceeds ~ 0.1 (Fig. 2c). The average number of NH_3 molecules per cluster remains well below two, even in the most sulfur-rich cases. This holds true even for the largest clusters containing up to seven H_2SO_4 molecules. This is consistent with chamber NH_3 contamination below $\sim 2 \text{ pptv}$ ¹⁸. In the H_2SO_4 -dominated experiment (panel d), some organic contaminants are visible near the nitrate-CIMS detection limit for AP-OOM ($\sim 1 \cdot 10^5 \text{ cm}^{-3}$). These background organics account for the high nucleation rates observed under the lowest AP-OOM conditions shown in Fig. 3.

3.3 Nucleation rates versus H_2SO_4 and AP-OOM

Fig. 3 shows nucleation rates at 1.7 nm , $J_{1.7}$, measured during CLOUD15–17 as well as those reported in Kirkby *et al.*⁹. Panels a) and b) show $J_{1.7}$ as a function of H_2SO_4 coloured by AP-OOM concentrations at $-10 \text{ }^\circ\text{C}$ and $+5 \text{ }^\circ\text{C}$, respectively. Panels c) and d) show the same data, but with AP-OOM on the horizontal axis, coloured by H_2SO_4 concentration. For these AP-OOM concentrations, the particle growth rates of all experiments exceed around 2 nm h^{-1} (see Fig. S2 in the supplementary information) so the $J_{1.7}$ measurements closely represent the nucleation rate at the critical size. The solid lines indicate $J_{1.7}$ predicted by the $\text{H}_2\text{SO}_4\text{-AP-OOM}$ parameterisation developed in this study. All nucleation rates shown in Fig. 3 were obtained under GCR conditions with ionisation rates of $Q \approx 2 \text{ cm}^{-3} \text{ s}^{-1}$ (additional data under neutral or beam conditions are not shown for clarity). In panels a) and b), data points without added H_2SO_4 are plotted at 10^4 cm^{-3} , corresponding to the instrumental lower limit of detection for H_2SO_4 .






Fig. 2 Molecular composition of negatively charged clusters during H_2SO_4 -AP-OOM nucleation events at -10°C . Mass defect (difference from integer mass) versus m/z during H_2SO_4 -AP-OOM nucleation events measured with an APi-TOF in negative mode. Panels are ordered by increasing ratio of H_2SO_4 to AP-OOM: a) < 0.0007 , b) 0.02 , c) 0.1 , and d) > 2 . All experiments were performed with comparable levels of contaminant vapours. The symbol radius is proportional to the logarithm of the signal intensity, and the colour indicates the number of sulfur atoms in the cluster. Clusters with no sulfur atoms generally contained nitrate ions (NO_3^-). Square outlines indicate clusters containing one NH_3 , and triangle outlines denote clusters containing two NH_3 molecules; no clusters were measured with more than two NH_3 molecules. The experimental conditions are 80 ppbv O_3 , $< 2\text{ pptv NH}_3$, $< 10\text{--}120\text{ pptv } \alpha\text{-pinene}$, $60\% \text{ RH}$ and -10°C .

The actual H_2SO_4 concentrations were likely to be substantially lower. The dashed lines in panels a) and b) indicate the nucleation rates expected for H_2SO_4 - NH_3 , calculated according to Dunne *et al.*⁶

Panels a) and b) show, at both temperatures, that nucleation rates at constant AP-OOM are insensitive to increasing H_2SO_4 below a certain H_2SO_4 concentration threshold, spanning the region where nucleation is purely from AP-OOM. This region extends up to H_2SO_4 concentrations of approximately 10^5 cm^{-3} at -10°C and 10^6 cm^{-3} at $+5^\circ\text{C}$. Below these thresholds, H_2SO_4 has no measurable effect on $J_{1.7}$, but once these thresholds are exceeded, the addition of H_2SO_4 leads to a pronounced increase in nucleation rates. For example, at -10°C , increasing H_2SO_4 to $\sim 3 \cdot 10^6\text{ cm}^{-3}$ in the presence of $\sim 1.5 \cdot 10^7\text{ cm}^{-3}$ AP-OOM enhances $J_{1.7}$ by a factor of approximately seven, while the same increase at $+5^\circ\text{C}$ raises $J_{1.7}$ by a factor of about five. These changes of H_2SO_4 would result in nucleation rates around 5 orders-of-magnitude lower for H_2SO_4 - NH_3 ⁶. We therefore attribute the observed enhancement to synergistic H_2SO_4 -AP-OOM nucleation. The near-

linear dependency of $J_{1.7}$ on acid concentration above the threshold is similar to the findings of Shen & Russell *et al.*¹¹ for H_2SO_4 -IP-OOM, but the acid enhancement is stronger for IP-OOM. A similar near-linear dependency of $J_{1.7}$ on AP-OOM is seen above the threshold for H_2SO_4 -AP-OOM nucleation (Figs. 3c and d).

We also observe that extremely low concentrations of AP-OOM have a strong effect on nucleation rates from the inorganic system. The dark blue triangles in Figs. 3a and b show measurements of nucleation rates with only contaminant levels of AP-OOM in the chamber, close to the detection limit of the nitrate-CIMS. Under these conditions, organic vapours alone do not produce measurable nucleation rates, yet the observed rates substantially exceed those predicted for purely inorganic H_2SO_4 - NH_3 nucleation, indicated by the dashed lines.⁶ This demonstrates that even very low concentrations of AP-OOM efficiently activate a mixed H_2SO_4 -AP-OOM nucleation pathway, leading to a large increase in $J_{1.7}$, despite organic vapours being insufficient to drive nucleation on their own.






Fig. 3 Measured and modelled nucleation rates, $J_{1,7}$. The nucleation rates are shown as a function of a) and b) H₂SO₄ and c) and d) AP-OOM concentrations, and at -10 °C (panels a and c) and +5 °C (panels b and d). The symbols denote measured nucleation rates and the curves show those predicted by the H₂SO₄-AP-OOM parameterisation developed in this study. The colours indicate a) and b) AP-OOM, and c) and d) H₂SO₄ concentrations. In a) and b), AP-OOM concentrations below 10⁶ cm⁻³ are represented with a single colour due to experimental uncertainties. The inverted triangles indicate measurements from Kirkby *et al.*⁹, while upright triangles represent measurements from the CLOUD15–17 campaigns. The black dashed lines in panels a) and b) indicate nucleation rates expected from H₂SO₄-NH₃⁶ assuming 2 pptv contaminant NH₃. The dark blue triangles in a) and b) correspond to nucleation rates measured in the presence of contaminant AP-OOM near the detection limit. All data shown here were obtained under GCR ionisation conditions with ionisation rates $Q \approx 2$ cm⁻³s⁻¹. The experimental conditions are a) and c) -10 °C, 70–90 ppbv O₃, b) and d) +5 °C, 40–50 ppbv O₃, and a–d) <2 pptv NH₃, 0–4000 pptv α -pinene and 60 % RH.

3.4 Nucleation rates versus ion concentrations

Fig. 4 shows nucleation rates $J_{1,7}$ measured at +5 °C with AP-OOM concentrations of 1.0–1.3 · 10⁷ cm⁻³ over a range of H₂SO₄ concentrations and ionisation conditions, together with the corresponding rates predicted by the H₂SO₄-AP-OOM parameterisation (see section 3.5), represented by the coloured bands. Neutral conditions ($Q = 0$ cm⁻³s⁻¹) are shown as circles, galactic cosmic ray (GCR) background ionisation as triangles, and pion-beam conditions as stars. Experiments without added H₂SO₄ are set to 10⁴ cm⁻³, corresponding to the instrumental lower limit of detection for H₂SO₄.

Fig. 4 shows a strong ion-induced enhancement of $J_{1,7}$ for both AP-OOM and H₂SO₄-AP-OOM and for all ionisation rates from 2 to 100 ion pairs cm⁻³s⁻¹. Moreover, for all ion conditions, the

addition of H₂SO₄ above $\sim 10^6$ cm⁻³ leads to a substantial increase in nucleation rates relative to pure AP-OOM nucleation. This demonstrates that the H₂SO₄-AP-OOM synergy is present for both charged (ion-induced) and neutral molecular clusters. Under neutral conditions, the addition of $\sim 5 \cdot 10^7$ cm⁻³ H₂SO₄ increases $J_{1,7}$ by more than a factor of 20, which is comparable to the enhancement of GCR compared with neutral conditions.

In Fig. 4 the neutral nucleation rates measured during CLOUD15–17 are systematically lower than those reported by Kirkby *et al.*⁹. The neutral channel is especially sensitive to small changes in background impurities in the chamber, which can cause systematic discrepancies between campaigns. Our parameterisation includes all data in the fitted values and, despite these systematic uncertainties, reproduces the measured nucle-






Fig. 4 Nucleation rates under different ionisation conditions at +5 °C. The symbols show measured nucleation rates versus SA at fixed AP-OOM of $(1.0\text{--}1.3) \cdot 10^7 \text{ cm}^{-3}$. The coloured bands indicate the modelled nucleation rates based on the parameterisation developed in this study, and their width reflects the range of AP-OOM and ion concentrations in the experimental data. The circles and blue shading correspond to neutral conditions (ion pair production rate, $Q = 0 \text{ cm}^{-3}\text{s}^{-1}$), the triangles and orange shading correspond to GCR conditions ($Q = 1.8\text{--}12 \text{ ion pairs cm}^{-3}\text{s}^{-1}$, depending on the presence of background muons from the CERN Proton Synchrotron), and the stars and red shading correspond to pion beam conditions ($50\text{--}100 \text{ ion pairs cm}^{-3}\text{s}^{-1}$). The hollow symbols denote measurements from Kirkby *et al.*⁹, and coloured symbols are measurements from the CLOUD15–17 campaigns. The experimental conditions are 40–50 ppbv O_3 , <2 pptv contaminant NH_3 , 60 % RH and +5 °C.

ation rates well across the full range of conditions (see section 4).

4 Parameterisation of nucleation rates

We have developed a temperature-dependent parameterisation of H_2SO_4 –AP-OOM nucleation which follows a similar approach as that for H_2SO_4 – NH_3 nucleation⁶, in which neutral and ion-induced nucleation are represented separately. The purely inorganic H_2SO_4 – NH_3 pathways retain the same functional form as Dunne *et al.*⁶ whereas the AP-OOM and new H_2SO_4 –AP-OOM pathways are refitted in the new parameterisation.

4.1 Treatment of oxidised organic vapours

Our experimental data involve oxidised organic molecules originating exclusively from α -pinene, which are therefore referred to as AP-OOM. For the parameterisation and its evaluation in the EMAC (ECHAM/MESSy Atmospheric Chemistry) Earth system model, we use the more general variable OOM to represent oxidised organic molecules derived from *all* monoterpenes. Our measured yield of AP-OOM from α -pinene ozonolysis is 6.0 % at

+5 °C and 5.7 % at -10 °C, while OOM in the EMAC model are derived from monoterpenes with a mean yield of ~ 5 %.

The good correspondence between the AP-OOM yields measured in CLOUD and the monoterpene-derived oxidised organic yields used in EMAC supports our simple definition of AP-OOM for the present study (see sections 2.3 and 2.4, and also the Supplementary Information). Future parameterisations of nucleation rates involving OOM will need further development to account for the influence of other organic precursors, such as isoprene or sesquiterpenes ($\text{C}_{15}\text{H}_{24}$), as well as vapours such as NO_x and HO_2 , which shift the final products to higher volatility, and reduce the nucleation rates^{7,46,47}.

4.2 Parameterisation

The new parameterisation developed in this study replaces the Dunne *et al.*⁶ H_2SO_4 – NH_3 parameterisation, the Kirkby *et al.*⁹ pure biogenic parameterisation and the Riccobono *et al.*¹² H_2SO_4 –BioOxOrg parameterisation. The new parameterisation calculates the total nucleation rate, J_{total} ($\text{cm}^{-3}\text{s}^{-1}$), as the sum of rates from five distinct channels, representing additive, independent contributions to the total nucleation rate:

- 1) neutral nucleation of H_2SO_4 ($J_{b,n}$)⁶;
- 2) neutral nucleation of H_2SO_4 – NH_3 ($J_{t,n}$)⁶;
- 3) neutral nucleation of pure OOM (J_0);
- 4) positive ion-induced nucleation of pure OOM plus pure H_2SO_4 (J_+); and
- 5) negative ion-induced nucleation of H_2SO_4 – NH_3 ⁶ plus H_2SO_4 –OOM (J_-):

$$J_{total} = J_{b,n} + J_{t,n} + J_0 + J_+ + J_- \quad (1)$$

(Water is implicit in all these chemical systems.)

The parameters were fitted to 93 nucleation rates measured during the CLOUD15–17 campaigns together with 54 nucleation rates measured by Kirkby *et al.*⁹. The dataset contains measurements at temperatures of -10 °C and +5 °C, ~ 60 % RH, H_2SO_4 concentrations between 10^4 and $4 \cdot 10^7 \text{ cm}^{-3}$, AP-OOM concentrations between 10^4 and $6 \cdot 10^8 \text{ cm}^{-3}$, NH_3 levels below 2 pptv, and three ionisation conditions (neutral, GCR, and pion beam). Temperature-dependent terms in the mixed H_2SO_4 –OOM pathways are therefore constrained by measurements at only two temperatures, and their application outside this range involves large uncertainties.

Because NH_3 contaminants were very low (<2 pptv), the parameterisation does not include an explicit NH_3 dependence for the H_2SO_4 –OOM pathway, although a strong NH_3 influence is expected from earlier studies¹⁴. For the present study, the influence of NH_3 remains confined to the H_2SO_4 – NH_3 channel inherited from Dunne *et al.*⁶. Future laboratory studies spanning a wider temperature range and including higher NH_3 concentrations will be required to account for H_2SO_4 – NH_3 –OOM nucleation over the full range of atmospheric conditions.

Fig. 5 shows the modelled nucleation rates from the H_2SO_4 –AP-OOM parameterisation versus all 147 measured nucleation rates used in the fit. The parameterisation explains more than 90 % of the variance in $\log_{10}(J_{1.7})$. It reproduces over 76 % of the mea-






Fig. 5 Modelled versus measured nucleation rates at $-10\text{ }^{\circ}\text{C}$ and $+5\text{ }^{\circ}\text{C}$. The circles correspond to neutral conditions, the triangles to GCR conditions, and the stars to pion beam conditions. The colours correspond to different datasets, as indicated in the legend. The solid line shows a 1:1 relationship, and the dotted lines indicate deviations by a factor 2.

sured nucleation rates within a factor of two and more than 93 % within a factor of three. A full description of the parameterisation equations, fitted coefficients, and model implementation details is provided in the supplementary information.

5 Impact on global nucleation rates

In Fig. 6 we show a comparison of our nucleation rate parameterisations with observations in the boundary layer. The parameterisations with organic vapours are shown as coloured bands, and assume $-10\text{ }^{\circ}\text{C}$, and $1\text{--}2 \cdot 10^7\text{ cm}^{-3}$ OOM/HOM/BioOxOrg as representative of the cool boundary layer over boreal forests^{48,49}. The parameterisation for $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation⁶ is also shown (solid black line), assuming 100 pptv NH_3 and $-10\text{ }^{\circ}\text{C}$. The atmospheric observations^{50–52} are indicated by small coloured circles and correspond to diverse experimental and environmental conditions. In particular, the Tecamac (Mexico City) measurements⁵⁰ of very rapid nucleation are likely to involve H_2SO_4 -amine nucleation⁵³. Variable condensation sinks and concentrations of base vapours and NO_x can account for the large scatter of the nucleation rates measured in less polluted environments¹⁴.

Fig. 6 shows that the earlier parameterisation of H_2SO_4 -BioOxOrg¹² (dark blue band) underestimates nucleation at low H_2SO_4 since it accounts neither for pure biogenic nucleation⁹ (light blue band with no dependency on H_2SO_4) nor the rise in H_2SO_4 -AP-OOM synergy at low H_2SO_4 . On the other, hand, the H_2SO_4 -BioOxOrg parameterisation overestimates nucleation at high H_2SO_4 compared with the present study. The addition of NH_3 to the H_2SO_4 -AP-OOM system will tend to shift the threshold for the H_2SO_4 -AP-OOM(-NH_3) synergy to lower H_2SO_4 con-

centrations, and raise the nucleation rate above that threshold. The importance of OOM in accounting for nucleation rates in the boundary layer can be readily seen by a comparison with the rates expected for $\text{H}_2\text{SO}_4\text{-NH}_3$ alone (black line).

To estimate its impact on global nucleation rates, we have embedded the new parameterisation in the EMAC (ECHAM/MESSy Atmospheric Chemistry) Earth system model. Fig. 7 compares global nucleation rates ($J_{1.7}$) predicted by the new parameterisation with those obtained using the previously implemented scheme combining the H_2SO_4 -BioOxOrg parameterisation of Riccobono *et al.*¹², the pure biogenic Kirkby *et al.*⁹ and the inorganic $\text{H}_2\text{SO}_4\text{-NH}_3$ parameterisation of Dunne *et al.*⁶. The analysis is restricted to the Northern Hemisphere at ground level, where monoterpene emissions are highest. In the model representation, OOM are produced only from oxidation of monoterpenes; any influence from other biogenic precursors such as isoprene is not included. Fig. 7a) shows ground-level nucleation rates calculated using the new parameterisation with H_2SO_4 -OOM, which is built on the Dunne *et al.*⁶ framework and reverts to this inorganic scheme under purely inorganic conditions. Panel b) shows nucleation rates obtained using the previously implemented H_2SO_4 -BioOxOrg scheme¹² in combination with the pure biogenic parameterisation of Kirkby *et al.*⁹, and panel c) displays the ratio of the nucleation rates (new parameterisation over previous).

The new parameterisation introduces several key improvements compared with the previous parameterisations. First, consistent with the treatment of organic vapours in the EMAC model, it represents oxidised organic vapours based on monoterpene oxidation, rather than relying on proxy species like BioOxOrg¹².

Second, it includes a temperature dependence of the mixed organic-inorganic nucleation pathway, which was not available previously. Nevertheless, we caution that the temperature dependence is constrained by measurements at only two temperatures ($-10\text{ }^{\circ}\text{C}$ and $+5\text{ }^{\circ}\text{C}$) so its application outside this range is uncertain.

The new parameterisation predicts higher nucleation rates over boreal forests, where monoterpene-derived OOM and H_2SO_4 coexist in the boundary layer under cool conditions (Fig. 7c). When the effects of NH_3 are also included, the nucleation rates are expected to be further enhanced¹⁴. In these boreal forest regions, the H_2SO_4 -BioOxOrg parameterisation¹² underestimated NPF rates. This can be seen in Fig. 6 at $1\text{--}5 \cdot 10^6\text{ cm}^{-3}$ H_2SO_4 and is even more pronounced at lower AP-OOM/BioOxOrg.

On the other hand, the new parameterisation predicts lower nucleation rates at lower latitudes of the Northern Hemisphere, where higher temperatures suppress nucleation despite the presence of precursor vapours. The earlier scheme, without any temperature dependence, strongly overestimates nucleation rates at lower latitudes. The strength of the decrease seen in Fig. 7c) varies regionally since it depends on not only the local temperature but also the H_2SO_4 , OOM and NH_3 concentrations. The largest reductions occur where OOM and H_2SO_4 coexist at appreciable concentrations under warm conditions, for which the previous temperature-independent H_2SO_4 -BioOxOrg parameterisation predicts rapid nucleation. Once more, this can be seen in Fig. 6 at H_2SO_4 concentrations above about 10^7 cm^{-3} . The dis-






Fig. 6 Comparison nucleation rate parameterisations with observations in the boundary layer. The nucleation rates, J ($\text{cm}^{-3}\text{s}^{-1}$), are shown versus H_2SO_4 concentration. The atmospheric observations^{50–52} are indicated by small coloured circles, and correspond to diverse experimental and environmental conditions. The coloured bands show the parameterisation of the present work (H_2SO_4 -AP-OOM; red band) together with those from earlier studies: Kirkby *et al.*, 2016⁹ (AP-HOM; light blue) and Riccobono *et al.*, 2014¹² (H_2SO_4 -BioOxOrg; dark blue), assuming $1\text{--}2 \cdot 10^7 \text{ cm}^{-3}$ OOM/HOM/BioOxOrg and $-10 \text{ }^\circ\text{C}$. The black line shows the parameterisation of Dunne *et al.*, 2016⁶ for H_2SO_4 - NH_3 at 100 pptv NH_3 and $-10 \text{ }^\circ\text{C}$.




Fig. 7 Annually-averaged nucleation rates, $J_{1,7}$, at the surface, simulated with the EMAC model. The nucleation rates are calculated using a) the new parameterisation with H_2SO_4 -OOM developed in the present study, and b) the H_2SO_4 -BioOxOrg parameterisation of Riccobono *et al.*¹² in combination with the pure biogenic parameterisation of Kirkby *et al.*⁹. Both simulations include H_2SO_4 - NH_3 nucleation calculated according to the parameterisation of Dunne *et al.*⁶. Panel c) shows the ratio of the nucleation rates in panel a) over those in panel b).



crepancies are smaller where OOM or H_2SO_4 are limiting; for these conditions the nucleation rate largely depends on the pure inorganic or pure biogenic systems, respectively, which are reasonably well parameterised.

Recent aircraft observations have shown that biogenic monoterpenes can persist at mixing ratios of a few tens of pptv in the upper troposphere, for example over the Amazon basin⁵⁴. In that study, new particle formation was associated with highly oxidised organic products from isoprene. These observations indicate that biogenic organic precursors and their oxidation products may be present well above the surface and can contribute to particle formation at high altitudes. In regions where monoterpene emissions dominate, such as boreal forests, oxidation products from monoterpenes may therefore also contribute to new particle formation through mixed organic- H_2SO_4 pathways, especially at cold, high altitudes, even when the concentrations of the individual vapours alone would be insufficient to produce substantial nucleation rates.

6 Conclusions

We have investigated new particle formation from α -pinene oxidation products (AP-OOM) and H_2SO_4 in the CERN CLOUD chamber. Through controlled experiments at atmospheric AP-OOM and H_2SO_4 concentrations, we have quantified the synergistic interaction between these two vapours in a near NH_3 -free environment (NH_3 contamination below 2 pptv). We conducted measurements at -10°C and $+5^\circ\text{C}$ under variable ionisation conditions representative of the boundary layer and the free troposphere.

Our results demonstrate that small additions of H_2SO_4 strongly enhance AP-OOM-driven nucleation and, similarly, that small AP-OOM concentrations strongly enhance H_2SO_4 -driven nucleation, where “small” implies levels that would produce negligible nucleation alone. This behaviour cannot be explained by inorganic H_2SO_4 - NH_3 nucleation, indicating a synergistic interaction between AP-OOM and H_2SO_4 rather than simply the addition of two independent nucleation mechanisms. The enhancement is more pronounced at lower temperatures, reflecting increased cluster stability under colder conditions. Across the full tropospheric ionisation conditions, nucleation rates are higher with increasing ionisation rates, indicating the importance of ion-induced nucleation for H_2SO_4 -AP-OOM, as well as for pure AP-OOM and H_2SO_4 (- NH_3).

Based on our measurements, we have developed a new parameterisation describing neutral and ion-induced nucleation in the H_2SO_4 -AP-OOM system, built on the H_2SO_4 (- NH_3) parameterisation of Dunne *et al.*⁶ at low AP-OOM. The parameterisation reproduces the measured nucleation rates within a factor of around two and captures their dependence on temperature, organic vapour concentrations, and ionisation.

When implemented in the EMAC global model, the new parameterisation predicts enhanced nucleation rates over high-monoterpene regions of the Northern Hemisphere compared with the earlier H_2SO_4 -BioOxOrg parameterisation of Riccobono *et al.*¹². Recent observations indicating the presence of monoterpenes at high altitudes suggest that the mixed H_2SO_4 -AP-OOM

pathway identified here may also be relevant at higher altitudes.

The present experiments involve H_2SO_4 -AP-OOM nucleation under ultra-low NH_3 conditions and at -10°C and $+5^\circ\text{C}$. Further experiments spanning a wider temperature range, relative humidities and atmospheric NH_3 concentrations will be required to extend this work to encompass the four-component H_2SO_4 -AP-OOM- NH_3 - H_2O system. Moreover, future parameterisations of nucleation rates involving OOM will need to account for more complex chemical environments that affect the volatility spectrum, combining monoterpenes with isoprene, sesquiterpenes, NO_x and HO_2 .

Author contributions

E.S., J.A., M.S., L.D., T.P., D.M.R., U.B., J.C., N.M.D., I.E.H., H.J., M.Ku., K.L., S.S., D.R.W. and J.K. planned the experiments; E.S., J.A., M.S., R.B., J.Du., P.K., A.Pi., P.R., S.Ru., A.A., H.J. and A.T. developed software; C.X., A.Po., M.Ko. and T.C. conducted the model implementation; E.S., J.A., W.Y., M.S., F.M., C.X., Z.Z., L.C.-P., M.L., L.L., M.M., T.R., D.M.R., Y.T., T.C. and J.K. analysed the data; E.S., J.A., W.Y., M.S., F.M., C.X., Z.Z., N.B., D.A., S.A., R.B., H.B., M.Be., P.B., M.Bu., M.C., L.C.-P., A.C., R.C.-S., L.D., J.De., J.Du., M.G., L.G.S., H.G.H., E.H., A.J., B.J., M.K.S., H.K., P.K., R.K., T.K., F.K., M.L., C.J.L., L.L., R.M., B.M., A.M., M.M., A.Pi., P.R., T.R., S.R., B.R., S.Ru., D.M.R., W.S., J.S., A.S., R.C.T., Y.T., J.T., N.S.U., J.W., B.Y., M.Z.-W., J.Z., T.C., H.H., X.-C.H., H.J., R.V., P.M.W. and J.K. conducted the experiments; E.S. and J.K. wrote the manuscript; E.S., J.A., W.Y., M.S., C.X., A.P., Z.Z., N.B., J.De., H.G., H.G.H., H.K., M.K., F.K., W.S., U.B., J.C., N.M.D., I.E.H., R.C.F., X.-C.H., K.L., J.L., S.S., P.M.W. and J.K. commented on and edited the manuscript; E.S., J.A., F.M., X.C. and T.C. visualised the data; H.G., A.K., R.M., T.C., J.C., A.H., H.J., M.Ku., K.L., J.L., S.S., R.V., P.M.W. and J.K. provided supervision; E.S., T.P., J.C., N.M.D., I.E.H., X.-C.H., H.J., M.Ku., R.V., A.O. and J.K. managed the project; J.Du., H.G., T.P., U.B., T.C., J.C., N.M.D., I.E.H., R.C.F., A.H., H.J., M.Ku., K.L., J.L., O.M., S.S., R.V. A.O. and J.K. acquired funding.

Conflicts of interest

There are no conflicts to declare.

Data availability

Data for all figures in the main text and supplementary materials are available at the Zenodo repository (DOI:10.5281/zenodo.19350026).

Acknowledgements

We would like to thank CERN for supporting CLOUD with important technical and financial resources, and for providing a particle beam from the CERN Proton Synchrotron. This work has been supported by European Union's MSCA Doctoral Network CLOUD-DOC 101073026, European Union's Horizon 2020 research and innovation programme under grant agreement No. 856612, the Research Council of Finland (grant no. 1325656, 302958, 311932, 316114, 325647, 325681, 334792, 337549, 337550, 337552, 346371, 347782, 349659, 352431, 357905, 359331, 364229, 371185), the Jane and Aatos Erkkö Founda-



tion ("Quantifying carbon sink, CarbonSink+ and their interaction with air quality", grant no. 220043), the Wihuri foundation ("Gigacity"), the European Research Council (ERC) (742206), the European Union via Non-CO2 Forcers and their Climate, Weather, Air Quality and Health Impacts (FOCI), German Federal Ministry of Research, Innovation and Space through the project CLOUD-22 (grant nos. 01LK2201A, 01LK2201B, and 01LK2201C), the German BMFTR SPP 1294 HALO project "CAFE-Pacific" 502266535, the US National Science Foundation (216181, AGS-2215522, AGS-2431817, AGS-2528506, AGS2431817, AGS2442132, ATM-2528507, ATM-2215527), the Austrian National Bank donation 2023-5223, the Estonian Research Council (projects PRG2738, RVTT3) and the Swiss National Science Foundation (grant no. 200021, 213071, 216181).

7 Environmental Significance Statement

Accurate representation of new particle formation from sulfuric acid and from oxidized organics is crucial for capturing aerosol effects in global climate models. Existing parameterisations are based on limited experimental conditions and do not accurately capture combined acid/organic nucleation effects or their temperature dependence. Here, we report chamber measurements of nucleation rates from α -pinene oxidation products and sulfuric acid in the near-absence of base vapours such as ammonia, under conditions representative of the cool boundary layer. We show that their synergistic interaction enhances particle formation beyond either component alone. Based on these results, we develop a parameterisation accounting for vapour concentrations, ionisation and temperature. This framework improves the representation of a major aerosol formation source in global models.

References

- 1 K. Armour, P. Forster, T. Storelvmo, W. Collins, J.-L. Dufresne, D. Frame, D. Lunt, T. Mauritsen, M. Palmer, M. Watanabe, M. Wild, H. Zhang, K. Alterskjaer and C. Smith, AGU Fall Meeting Abstracts, 2021, pp. U13B-07.
- 2 H. Gordon, J. Kirkby, U. Baltensperger, F. Bianchi, M. Breitenlechner, J. Curtius, A. Dias, J. Dommen, N. M. Donahue, E. M. Dunne, J. Duplissy, S. Ehrhart, R. C. Flagan, C. Frege, C. Fuchs, A. Hansel, C. R. Hoyle, M. Kulmala, A. Kürten, K. Lehtipalo, V. Makhmutov, U. Molteni, M. P. Rissanen, Y. Stozhkov, J. Tröstl, G. Tsagkogeorgas, R. Wagner, C. Williamson, D. Wimmer, P. M. Winkler, C. Yan and K. S. Carslaw, *Journal of Geophysical Research: Atmospheres*, 2017, **122**, 8739–8760.
- 3 M. Kulmala, T. Petäjä, M. Ehn, J. Thornton, M. Sipilä, D. Worsnop and V.-M. Kerminen, *Annual review of physical chemistry*, 2014, **65**, 21–37.
- 4 J. Kirkby, J. Curtius, J. Almeida, E. Dunne, J. Duplissy, S. Ehrhart, A. Franchin, S. Gagné, L. Ickes, A. Kürten, A. Kupc, A. Metzger, F. Riccobono, L. Rondo, S. Schobesberger, G. Tsagkogeorgas, D. Wimmer, A. Amorim, F. Bianchi, M. Breitenlechner, A. David, J. Dommen, A. Downard, M. Ehn, R. C. Flagan, S. Haider, A. Hansel, D. Hauser, W. Jud, H. Junninen, F. Kreissl, A. Kvashin, A. Laaksonen, K. Lehtipalo, J. Lima, E. R. Lovejoy, V. Makhmutov, S. Mathot, J. Mikkilä, P. Minginette, S. Mogo, T. Nieminen, A. Onnela, P. Pereira, T. Petäjä, R. Schnitzhofer, J. H. Seinfeld, M. Sipilä, Y. Stozhkov, F. Stratmann, A. Tomé, J. Vanhanen, Y. Viisanen, A. Vrtala, P. E. Wagner, H. Walther, E. Weingartner, H. Wex, P. M. Winkler, K. S. Carslaw, D. R. Worsnop, U. Baltensperger and M. Kulmala, *Nature*, 2011, **476**, 429–433.
- 5 M. Kulmala, J. Kontkanen, H. Junninen, K. Lehtipalo, H. E. Manninen, T. Nieminen, T. Petäjä, M. Sipilä, S. Schobesberger, P. Rantala, A. Franchin, T. Jokinen, E. Järvinen, M. äijälä, J. Kangasluoma, J. Hakala, P. P. Aalto, P. Paasonen, J. Mikkilä, J. Vanhanen, J. Aalto, H. Hakola, U. Makkonen, T. Ruuskanen, R. L. Mauldin, J. Duplissy, H. Vehkamäki, J. Bäck, A. Kortelainen, I. Riipinen, T. Kurtén, M. V. Johnston, J. N. Smith, M. Ehn, T. F. Mentel, K. E. J. Lehtinen, A. Laaksonen, V.-M. Kerminen and D. R. Worsnop, *Science*, 2013, **339**, 943–946.
- 6 E. M. Dunne, H. Gordon, A. Kürten, J. Almeida, J. Duplissy, C. Williamson, I. K. Ortega, K. J. Pringle, A. Adamov, U. Baltensperger, P. Barmet, F. Benduhn, F. Bianchi, M. Breitenlechner, A. Clarke, J. Curtius, J. Dommen, N. M. Donahue, S. Ehrhart, R. C. Flagan, A. Franchin, R. Guida, J. Hakala, A. Hansel, M. Heinritzi, T. Jokinen, J. Kangasluoma, J. Kirkby, M. Kulmala, A. Kupc, M. J. Lawler, K. Lehtipalo, V. Makhmutov, G. Mann, S. Mathot, J. Merikanto, P. Miettinen, A. Nenes, A. Onnela, A. Rap, C. L. S. Reddington, F. Riccobono, N. A. D. Richards, M. P. Rissanen, L. Rondo, N. Sarnela, S. Schobesberger, K. Sengupta, M. Simon, M. Sipilä, J. N. Smith, Y. Stozhkov, A. Tomé, J. Tröstl, P. E. Wagner, D. Wimmer, P. M. Winkler, D. R. Worsnop and K. S. Carslaw, *Science*, 2016, **354**, 1119–1124.
- 7 F. Bianchi, T. Kürten, M. Riva, C. Mohr, M. P. Rissanen, P. Roldin, T. Berndt, J. D. Crouse, P. O. Wennberg, T. F. Mentel, J. Wildt, H. Junninen, T. Jokinen, M. Kulmala, D. R. Worsnop, J. A. Thornton, N. Donahue, H. G. Kjaergaard and M. Ehn, *Chemical reviews*, 2019, **119**, 3472–3509.
- 8 M. Ehn, J. A. Thornton, E. Kleist, M. Sipilä, H. Junninen, I. Pullinen, M. Springer, F. Rubach, R. Tillmann, B. Lee, F. Lopez-Hilfiker, S. Andres, I.-H. Acir, M. Rissanen, T. Jokinen, S. Schobesberger, J. Kangasluoma, J. Kontkanen, T. Nieminen, T. Kurtén, L. B. Nielsen, S. Jørgensen, H. G. Kjaergaard, M. Canagaratna, M. D. Maso, T. Berndt, T. Petäjä, A. Wahner, V.-M. Kerminen, M. Kulmala, D. R. Worsnop, J. Wildt and T. F. Mentel, *Nature*, 2014, **506**, 476–479.
- 9 J. Kirkby, J. Duplissy, K. Sengupta, C. Frege, H. Gordon, C. Williamson, M. Heinritzi, M. Simon, C. Yan, J. Almeida, J. Tröstl, T. Nieminen, I. K. Ortega, R. Wagner, A. Adamov, A. Amorim, A.-K. Bernhammer, F. Bianchi, M. Breitenlechner, S. Brilke, X. Chen, J. Craven, A. Dias, S. Ehrhart, R. C. Flagan, A. Franchin, C. Fuchs, R. Guida, J. Hakala, C. R. Hoyle, T. Jokinen, H. Junninen, J. Kangasluoma, J. Kim, M. Krapf, A. Kürten, A. Laaksonen, K. Lehtipalo, V. Makhmutov, S. Mathot, U. Molteni, A. Onnela, O. Peräkylä, F. Piel, T. Petäjä, A. P. Praplan, K. Pringle, A. Rap, N. A. D. Richards, I. Riipinen, M. P. Rissanen, L. Rondo, N. Sarnela, S. Schobes-



- berger, C. E. Scott, J. H. Seinfeld, M. Sipilä, G. Steiner, Y. Stozhkov, F. Stratmann, A. Tomé, A. Virtanen, A. L. Vogel, A. C. Wagner, P. E. Wagner, E. Weingartner, D. Wimmer, P. M. Winkler, P. Ye, X. Zhang, A. Hansel, J. Dommen, N. M. Donahue, D. R. Worsnop, U. Baltensperger, M. Kulmala, K. S. Carslaw and J. Curtius, *Nature*, 2016, **533**, 521–526.
- 10 M. Heinritzi, L. Dada, M. Simon, D. Stolzenburg, A. C. Wagner, L. Fischer, L. R. Ahonen, S. Amanatidis, R. Baalbaki, A. Baccarini, P. S. Bauer, B. Baumgartner, F. Bianchi, S. Brilke, D. Chen, R. Chiu, A. Dias, J. Dommen, J. Duplissy, H. Finkenzeller, C. Frege, C. Fuchs, O. Garmash, H. Gordon, M. Granzin, I. El Haddad, X. He, J. Helm, V. Hofbauer, C. R. Hoyle, J. Kangasluoma, T. Keber, C. Kim, A. Kürten, H. Lamkaddam, T. M. Laurila, J. Lampilahti, C. P. Lee, K. Lehtipalo, M. Leiminger, H. Mai, V. Makhmutov, H. E. Manninen, R. Marten, S. Mathot, R. L. Mauldin, B. Mentler, U. Molteni, T. Müller, W. Nie, T. Nieminen, A. Onnela, E. Partoll, M. Passananti, T. Petäjä, J. Pfeifer, V. Pospisilova, L. L. J. Quéléver, M. P. Rissanen, C. Rose, S. Schobesberger, W. Scholz, K. Scholze, M. Sipilä, G. Steiner, Y. Stozhkov, C. Tauber, Y. J. Tham, M. Vazquez-Pufleau, A. Virtanen, A. L. Vogel, R. Volkamer, R. Wagner, M. Wang, L. Weitz, D. Wimmer, M. Xiao, C. Yan, P. Ye, Q. Zha, X. Zhou, A. Amorim, U. Baltensperger, A. Hansel, M. Kulmala, A. Tomé, P. M. Winkler, D. R. Worsnop, N. M. Donahue, J. Kirkby and J. Curtius, *Atmospheric Chemistry and Physics*, 2020, **20**, 11809–11821.
- 11 J. Shen, D. M. Russell, J. DeVivo, F. Kunkler, R. Baalbaki, B. Mentler, W. Scholz, W. Yu, L. Caudillo-Plath, E. Sommer, E. Ahongshangbam, D. Alfaouri, J. Almeida, A. Amorim, L. J. Beck, H. Beckmann, M. Berntheusel, N. Bhattacharyya, M. R. Canagaratna, A. Chassaing, R. Cruz-Simbron, L. Dada, J. Duplissy, H. Gordon, M. Granzin, L. Große Schute, M. Heinritzi, S. Iyer, H. Klebach, T. Krüger, A. Kürten, M. Lampimäki, L. Liu, B. Lopez, M. Martinez, A. Morawiec, A. Onnela, M. Peltola, P. Rato, M. Reza, S. Richter, B. Rörup, M. K. Sebastian, M. Simon, M. Surdu, K. Tamme, R. C. Thakur, A. Tomé, Y. Tong, J. Top, N. S. Umo, G. Unfer, L. Vettikkat, J. Weissbacher, C. Xenofontos, B. Yang, M. Zauner-Wieczorek, J. Zhang, Z. Zheng, U. Baltensperger, T. Christoudias, R. C. Flagan, I. El Haddad, H. Junninen, O. Möhler, I. Riipinen, U. Rohner, S. Schobesberger, R. Volkamer, P. M. Winkler, A. Hansel, K. Lehtipalo, N. M. Donahue, J. Lelieveld, H. Harder, M. Kulmala, D. R. Worsnop, J. Kirkby, J. Curtius and X.-C. He, *Nature*, 2024, **636**, 115–123.
- 12 F. Riccobono, S. Schobesberger, C. E. Scott, J. Dommen, I. K. Ortega, L. Rondo, J. Almeida, A. Amorim, F. Bianchi, M. Breitenlechner, A. David, A. Downard, E. M. Dunne, J. Duplissy, S. Ehrhart, R. C. Flagan, A. Franchin, A. Hansel, H. Junninen, M. Kajos, H. Keskinen, A. Kupc, A. Kürten, A. N. Kvashin, A. Laaksonen, K. Lehtipalo, V. Makhmutov, S. Mathot, T. Nieminen, A. Onnela, T. Petäjä, A. P. Praplan, F. D. Santos, S. Schallhart, J. H. Seinfeld, M. Sipilä, D. V. Spracklen, Y. Stozhkov, F. Stratmann, A. Tomé, G. Tsagkogeorgas, P. Vaattovaara, Y. Viisanen, A. Vrtala, P. E. Wagner, E. Weingartner, H. Wex, D. Wimmer, K. S. Carslaw, J. Curtius, N. M. Donahue, J. Kirkby, M. Kulmala, D. R. Worsnop and U. Baltensperger, *Science*, 2014, **344**, 717–721.
- 13 A. Metzger, B. Verheggen, J. Dommen, J. Duplissy, A. S. H. Prevot, E. Weingartner, I. Riipinen, M. Kulmala, D. V. Spracklen, K. S. Carslaw and U. Baltensperger, *Proceedings of the National Academy of Sciences*, 2010, **107**, 6646–6651.
- 14 K. Lehtipalo, C. Yan, L. Dada, F. Bianchi, M. Xiao, R. Wagner, D. Stolzenburg, L. R. Ahonen, A. Amorim, A. Baccarini, P. S. Bauer, B. Baumgartner, A. Bergen, A.-K. Bernhammer, M. Breitenlechner, S. Brilke, A. Buchholz, S. B. Mazon, D. Chen, X. Chen, A. Dias, J. Dommen, D. C. Draper, J. Duplissy, M. Ehn, H. Finkenzeller, L. Fischer, C. Frege, C. Fuchs, O. Garmash, H. Gordon, J. Hakala, X. He, L. Heikkinen, M. Heinritzi, J. C. Helm, V. Hofbauer, C. R. Hoyle, T. Jokinen, J. Kangasluoma, V.-M. Kerminen, C. Kim, J. Kirkby, J. Kontkanen, A. Kürten, M. J. Lawler, H. Mai, S. Mathot, R. L. Mauldin, U. Molteni, L. Nichman, W. Nie, T. Nieminen, A. Ojdanic, A. Onnela, M. Passananti, T. Petäjä, F. Piel, V. Pospisilova, L. L. J. Quéléver, M. P. Rissanen, C. Rose, N. Sarnela, S. Schallhart, S. Schuchmann, K. Sengupta, M. Simon, M. Sipilä, C. Tauber, A. Tomé, J. Tröstl, O. Väisänen, A. L. Vogel, R. Volkamer, A. C. Wagner, M. Wang, L. Weitz, D. Wimmer, P. Ye, A. Ylisirniö, Q. Zha, K. S. Carslaw, J. Curtius, N. M. Donahue, R. C. Flagan, A. Hansel, I. Riipinen, A. Virtanen, P. M. Winkler, U. Baltensperger, M. Kulmala and D. R. Worsnop, *Science advances*, 2018, **4**, eaau5363.
- 15 J. Merikanto, D. Spracklen, G. Mann, S. Pickering and K. Carslaw, *Atmospheric Chemistry and Physics*, 2009, **9**, 8601–8616.
- 16 A. Kupc, A. Amorim, J. Curtius, A. Danielczok, J. Duplissy, S. Ehrhart, H. Walther, L. Ickes, J. Kirkby, A. Kürten, J. Lima, S. Mathot, P. Minginette, A. Onnela, L. Rondo and P. Wagner, *Journal of Aerosol Science*, 2011, **42**, 532–543.
- 17 H. Junninen, M. Ehn, T. Petäjä, L. Luosujärvi, T. Kotiaho, R. Kostianen, U. Rohner, M. Gonin, K. Fuhrer, M. Kulmala and D. R. Worsnop, *Atmospheric Measurement Techniques*, 2010, **3**, 1039–1053.
- 18 S. Schobesberger, A. Franchin, F. Bianchi, L. Rondo, J. Duplissy, A. Kürten, I. K. Ortega, A. Metzger, R. Schnitzhofer, J. Almeida, A. Amorim, J. Dommen, E. M. Dunne, M. Ehn, S. Gagné, L. Ickes, H. Junninen, A. Hansel, V.-M. Kerminen, J. Kirkby, A. Kupc, A. Laaksonen, K. Lehtipalo, S. Mathot, A. Onnela, T. Petäjä, F. Riccobono, F. D. Santos, M. Sipilä, A. Tomé, G. Tsagkogeorgas, Y. Viisanen, P. E. Wagner, D. Wimmer, J. Curtius, N. M. Donahue, U. Baltensperger, M. Kulmala and D. R. Worsnop, *Atmospheric Chemistry and Physics*, 2015, **15**, 55–78.
- 19 M. Graus, M. Müller and A. Hansel, *Journal of the American Society for Mass Spectrometry*, 2011, **21**, 1037–1044.
- 20 E. Canaval, N. Hyttinen, B. Schmidbauer, L. Fischer and A. Hansel, *Frontiers in Chemistry*, 2019, **7**, 191.
- 21 T. Reinecke, M. Leiminger, A. Jordan, A. Wisthaler and M. Müller, *Analytical Chemistry*, 2023, **95**, 11879–11884.
- 22 T. Jokinen, M. Sipilä, H. Junninen, M. Ehn, G. Lönn, J. Hakala, T. Petäjä, R. Mauldin Iii, M. Kulmala and



- D. Worsnop, *Atmospheric Chemistry and Physics*, 2012, **12**, 4117–4125.
- 23 N. Hyttinen, O. Kupiainen-Maatta, M. P. Rissanen, M. Muuronen, M. Ehn and T. Kurtén, *The Journal of Physical Chemistry A*, 2015, **119**, 6339–6345.
- 24 A. Kürten, L. Rondo, S. Ehrhart and J. Curtius, *Atmospheric measurement techniques*, 2011, **4**, 437–443.
- 25 A. Kürten, L. Rondo, S. Ehrhart and J. Curtius, *The Journal of Physical Chemistry A*, 2012, **116**, 6375–6386.
- 26 D. Li, D. Wang, L. Caudillo, W. Scholz, M. Wang, S. Tomaz, G. Marie, M. Surdu, E. Eccli, X. Gong, L. Gonzalez-Carracedo, M. Granzin, J. Pfeifer, B. Rörup, B. Schulze, P. Rantala, S. Perrier, A. Hansel, J. Curtius, J. Kirkby, N. M. Donahue, C. George, I. El-Haddad and M. Riva, *Atmospheric Measurement Techniques*, 2024, **17**, 5413–5428.
- 27 M. Simon, L. Dada, M. Heinritzi, W. Scholz, D. Stolzenburg, L. Fischer, A. C. Wagner, A. Kürten, B. Rörup, X.-C. He, J. Almeida, R. Baalbaki, A. Baccharini, P. S. Bauer, L. Beck, A. Bergen, F. Bianchi, S. Bräkling, S. Brilke, L. Caudillo, D. Chen, B. Chu, A. Dias, D. C. Draper, J. Duplissy, I. El-Haddad, H. Finkenzeller, C. Frege, L. Gonzalez-Carracedo, H. Gordon, M. Granzin, J. Hakala, V. Hofbauer, C. R. Hoyle, C. Kim, W. Kong, H. Lamkaddam, C. P. Lee, K. Lehtipalo, M. Leiminger, H. Mai, H. E. Manninen, G. Marie, R. Marten, B. Mentler, U. Molteni, L. Nichman, W. Nie, A. Ojdanic, A. Onnela, E. Partoll, T. Petäjä, J. Pfeifer, M. Philippov, L. L. J. Quéléver, A. Ranjithkumar, M. P. Rissanen, S. Schallhart, S. Schobesberger, S. Schuchmann, J. Shen, M. Sipilä, G. Steiner, Y. Stozhkov, C. Tauber, Y. J. Tham, A. R. Tomé, M. Vazquez-Pufleau, A. L. Vogel, R. Wagner, M. Wang, D. S. Wang, Y. Wang, S. K. Weber, Y. Wu, M. Xiao, C. Yan, P. Ye, Q. Ye, M. Zauner-Wieczorek, X. Zhou, U. Baltensperger, J. Dommen, R. C. Flagan, A. Hansel, M. Kulmala, R. Volkamer, P. M. Winkler, D. R. Worsnop, N. M. Donahue, J. Kirkby and J. Curtius, *Atmospheric Chemistry and Physics*, 2020, **20**, 9183–9207.
- 28 L. Dada, D. Stolzenburg, M. Simon, L. Fischer, M. Heinritzi, M. Wang, M. Xiao, A. L. Vogel, L. Ahonen, A. Amorim, R. Baalbaki, A. Baccharini, U. Baltensperger, F. Bianchi, K. R. Daellenbach, J. DeVivo, A. Dias, J. Dommen, J. Duplissy, H. Finkenzeller, A. Hansel, X.-C. He, V. Hofbauer, C. R. Hoyle, J. Kangasluoma, C. Kim, A. Kürten, A. Kvashnin, R. Mauldin, V. Makhmutov, R. Marten, B. Mentler, W. Nie, T. Petäjä, L. L. J. Quéléver, H. Saathoff, C. Tauber, A. Tome, U. Molteni, R. Volkamer, R. Wagner, A. C. Wagner, D. Wimmer, P. M. Winkler, C. Yan, Q. Zha, M. Rissanen, H. Gordon, J. Curtius, D. R. Worsnop, K. Lehtipalo, N. M. Donahue, J. Kirkby, I. El Haddad and M. Kulmala, *Science advances*, 2023, **9**, eadi5297.
- 29 J. Kirkby, A. Amorim, U. Baltensperger, K. S. Carslaw, T. Christoudias, J. Curtius, N. M. Donahue, I. E. Haddad, R. C. Flagan, H. Gordon, A. Hansel, H. Harder, H. Junninen, M. Kulmala, A. Kürten, A. Laaksonen, K. Lehtipalo, J. Lelieveld, O. Möhler, I. Riipinen, F. Stratmann, A. Tomé, A. Virtanen, R. Volkamer, P. M. Winkler and D. R. Worsnop, *Nature Geoscience*, 2023, **16**, 948–957.
- 30 N. M. Donahue, S. A. Epstein, S. N. Pandis and A. L. Robinson, *Atmospheric Chemistry and Physics*, 2011, **11(7)**, 3303–3318.
- 31 N. M. Donahue, J. H. Kroll, S. N. Pandis and A. L. Robinson, *Atmospheric Chemistry and Physics*, 2012, **12(2)**, 615–634.
- 32 Y. Wang, P. Clusius, C. Yan, K. Dällenbach, R. Yin, M. Wang, X.-C. He, B. Chu, Y. Lu, L. Dada, J. Kangasluoma, P. Rantala, C. Deng, Z. Lin, W. Wang, L. Yao, X. Fan, W. Du, J. Cai, L. Heikkinen, Y. J. Tham, Q. Zha, Z. Ling, H. Junninen, T. Petäjä, M. Ge, Y. Wang, H. He, D. R. Worsnop, V.-M. Kerminen, F. Bianchi, L. Wang, J. Jiang, Y. Liu, M. Boy, M. Ehn, N. M. Donahue and M. Kulmala, *Environmental Science & Technology*, 2021, **56**, 770–778.
- 33 M. Riva, P. Rantala, J. E. Krechmer, O. Peräkylä, Y. Zhang, L. Heikkinen, O. Garmash, C. Yan, M. Kulmala, D. Worsnop and M. Ehn, *Atmospheric Measurement Techniques*, 2019, **12**, 2403–2421.
- 34 D. Stolzenburg, L. Fischer, A. L. Vogel, M. Heinritzi, M. Schervish, M. Simon, A. C. Wagner, L. Dada, L. R. Ahonen, A. Amorim, A. Baccharini, P. S. Bauer, B. Baumgartner, A. Bergen, F. Bianchi, M. Breitenlechner, S. Brilke, S. Buenrostro Mazon, D. Chen, A. Dias, D. C. Draper, J. Duplissy, I. El Haddad, H. Finkenzeller, C. Frege, C. Fuchs, O. Garmash, H. Gordon, X. He, J. Helm, V. Hofbauer, C. R. Hoyle, C. Kim, J. Kirkby, J. Kontkanen, A. Kürten, J. Lampilahti, M. Lawler, K. Lehtipalo, M. Leiminger, H. Mai, S. Mathot, B. Mentler, U. Molteni, W. Nie, T. Nieminen, J. B. Nowak, A. Ojdanic, A. Onnela, M. Passananti, T. Petäjä, L. L. J. Quéléver, M. P. Rissanen, N. Sarnela, S. Schallhart, C. Tauber, A. Tomé, R. Wagner, M. Wang, L. Weitz, D. Wimmer, M. Xiao, C. Yan, P. Ye, Q. Zha, U. Baltensperger, J. Curtius, J. Dommen, R. C. Flagan, M. Kulmala, J. N. Smith, D. R. Worsnop, A. Hansel, N. M. Donahue and P. M. Winkler, *Proceedings of the National Academy of Sciences*, 2018, **115**, 9122–9127.
- 35 T. Jokinen, T. Berndt, R. Makkonen, V.-M. Kerminen, H. Junninen, P. Paasonen, F. Stratmann, H. Herrmann, A. B. Guenther, D. R. Worsnop, M. Kulmala, M. Ehn and M. Sipilä, *Proceedings of the National Academy of Sciences*, 2015, **112**, 7123–7128.
- 36 J. Vanhanen, J. Mikkilä, K. Lehtipalo, M. Sipilä, H. Manninen, E. Siivola, T. Petäjä and M. Kulmala, *Aerosol Science and Technology*, 2011, **45**, 533–542.
- 37 S. C. Wang and R. C. Flagan, *Aerosol Science and Technology*, 1990, **13**, 230–240.
- 38 S. Mirme and A. Mirme, *Atmospheric Measurement Techniques*, 2013, **6**, 1061–1071.
- 39 L. Dada, K. Lehtipalo, J. Kontkanen, T. Nieminen, R. Baalbaki, L. Ahonen, J. Duplissy, C. Yan, B. Chu, T. Petäjä, K. Lehtinen, V.-M. Kerminen, M. Kulmala and J. Kangasluoma, *Nature protocols*, 2020, **15**, 1013–1040.
- 40 D. Stolzenburg, G. Steiner and P. M. Winkler, *Atmospheric Measurement Techniques*, 2017, **10**, 1639–1651.
- 41 A. Kürten, C. Williamson, J. Almeida, J. Kirkby and J. Curtius, *Atmospheric Chemistry and Physics*, 2015, **15(8)**, 4063–4075.



- 42 E. Roeckner, R. Brokopf, M. Esch, M. Giorgetta, S. Hagemann, L. Kornbluh, E. Manzini, U. Schlese and U. Schulzweida, *Journal of Climate*, 2006, **19**, 3771–3791.
- 43 P. Jöckel, A. Kerckweg, A. Pozzer, R. Sander, H. Tost, H. Riede, A. Baumgaertner, S. Gromov and B. Kern, *Geoscientific Model Development*, 2010, **3**, 717–752.
- 44 D. Stolzenburg, M. Simon, A. Ranjithkumar, A. Kürten, K. Lehtipalo, H. Gordon, S. Ehrhart, H. Finkenzeller, L. Pichelstorfer, T. Nieminen, X.-C. He, S. Brilke, M. Xiao, A. Amorim, R. Baalbaki, A. Baccarini, L. Beck, S. Bräkling, L. Caudillo Murillo, D. Chen, B. Chu, L. Dada, A. Dias, J. Dommen, J. Duplissy, I. El Haddad, L. Fischer, L. Gonzalez Carracedo, M. Heinritzi, C. Kim, T. K. Koenig, W. Kong, H. Lamkaddam, C. P. Lee, M. Leiminger, Z. Li, V. Makhmutov, H. E. Manninen, G. Marie, R. Marten, T. Müller, W. Nie, E. Partoll, T. Petäjä, J. Pfeifer, M. Philippov, M. P. Rissanen, B. Rörup, S. Schobesberger, S. Schuchmann, J. Shen, M. Sipilä, G. Steiner, Y. Stozhkov, C. Tauber, Y. J. Tham, A. Tomé, M. Vazquez-Pufleau, A. C. Wagner, M. Wang, Y. Wang, S. K. Weber, D. Wimmer, P. J. Wlasits, Y. Wu, Q. Ye, M. Zauner-Wieczorek, U. Baltensperger, K. S. Carslaw, J. Curtius, N. M. Donahue, R. C. Flagan, A. Hansel, M. Kulmala, J. Lelieveld, R. Volkamer, J. Kirkby and P. M. Winkler, *Atmospheric Chemistry and Physics*, 2020, **20**, 7359–7372.
- 45 S. Schobesberger, H. Junninen, F. Bianchi, G. Lönn, M. Ehn, K. Lehtipalo, J. Dommen, S. Ehrhart, I. K. Ortega, A. Franchin, T. Nieminen, F. Riccobono, M. Hutterli, J. Duplissy, J. Almeida, A. Amorim, M. Breitenlechner, A. J. Downard, E. M. Dunne, R. C. Flagan, M. Kajos, H. Keskinen, J. Kirkby, A. Kupc, A. Kürten, T. Kurtén, A. Laaksonen, S. Mathot, A. Onnela, A. P. Praplan, L. Rondo, F. D. Santos, S. Schallhart, R. Schnitzhofer, M. Sipilä, A. Tomé, G. Tsagkogeorgas, H. Vehkamäki, D. Wimmer, U. Baltensperger, K. S. Carslaw, J. Curtius, A. Hansel, T. Petäjä, M. Kulmala, N. M. Donahue and D. R. Worsnop, *Proceedings of the National Academy of Sciences*, 2013, **110**, 17223–17228.
- 46 C. Yan, W. Nie, A. L. Vogel, L. Dada, K. Lehtipalo, D. Stolzenburg, R. Wagner, M. P. Rissanen, M. Xiao, L. Ahonen, L. Fischer, C. Rose, F. Bianchi, H. Gordon, M. Simon, M. Heinritzi, O. Garmash, P. Roldin, A. Dias, P. Ye, V. Hofbauer, A. Amorim, P. S. Bauer, A. Bergen, A.-K. Bernhammer, M. Breitenlechner, S. Brilke, A. Buchholz, S. B. Mazon, M. R. Canagaratna, X. Chen, A. Ding, J. Dommen, D. C. Draper, J. Duplissy, C. Frege, C. Heyn, R. Guida, J. Hakala, L. Heikkinen, C. R. Hoyle, T. Jokinen, J. Kangasluoma, J. Kirkby, J. Kontkanen, A. Kürten, M. J. Lawler, H. Mai, S. Mathot, R. L. Mauldin, U. Molteni, L. Nichman, T. Nieminen, J. Nowak, A. Ojdanic, A. Onnela, A. Pajunoja, T. Petäjä, F. Piel, L. L. J. Quéléver, N. Sarnela, S. Schallhart, K. Sengupta, M. Sipilä, A. Tomé, J. Tröstl, O. Väisänen, A. C. Wagner, A. Ylisirniö, Q. Zha, U. Baltensperger, K. S. Carslaw, J. Curtius, R. C. Flagan, A. Hansel, I. Riipinen, J. N. Smith, A. Virtanen, P. M. Winkler, N. M. Donahue, V.-M. Kerminen, M. Kulmala, M. Ehn and D. R. Worsnop, *Science advances*, 2020, **6**, eaay4945.
- 47 L. Yang, W. Nie, C. Yan, M. Ehn, Y. Liu, P. Roldin, X. Qi, V.-M. Kerminen, N. M. Donahue, D. Worsnop, M. Kulmala and A. Ding, *Nature Communications*, 2025, **17**, 302.
- 48 T. Jokinen, K. Lehtipalo, R. C. Thakur, I. Ylivinkka, K. Neitola, N. Sarnela, T. Laitinen, M. Kulmala, T. Petäjä and M. Sipilä, *Atmospheric Chemistry and Physics*, 2022, **22**, 2237–2254.
- 49 J. Sulo, N. Sarnela, J. Kontkanen, L. Ahonen, P. Paasonen, T. Laurila, T. Jokinen, J. Kangasluoma, H. Junninen, M. Sipilä, T. Petäjä, M. Kulmala and K. Lehtipalo, *Atmospheric Chemistry and Physics*, 2021, **21**, 695–715.
- 50 C. Kuang, P. H. McMurry, A. V. McCormick and F. Eisele, *Journal of Geophysical Research: Atmospheres*, 2008, **113**, 209.
- 51 S.-L. Sihto, M. Kulmala, V.-M. Kerminen, M. Dal Maso, T. Petäjä, I. Riipinen, H. Korhonen, F. Arnold, R. Janson, M. Boy, A. Laaksonen and K. E. J. Lehtinen, *Atmospheric Chemistry and Physics*, 2006, **6**, 4079–4091.
- 52 P. Paasonen, T. Nieminen, E. Asmi, H. E. Manninen, T. Petäjä, C. Plass-Dülmer, H. Flentje, W. Birmili, A. Wiedensohler, U. Hörrak, A. Metzger, A. Hamed, A. Laaksonen, M. C. Facchini, V.-M. Kerminen and M. Kulmala, *Atmospheric Chemistry and Physics*, 2010, **10**, 11223–11242.
- 53 J. Almeida, S. Schobesberger, A. Kürten, I. K. Ortega, O. Kupiainen-Määttä, A. P. Praplan, A. Adamov, A. Amorim, F. Bianchi, M. Breitenlechner, A. David, J. Dommen, N. M. Donahue, A. Downard, E. Dunne, J. Duplissy, S. Ehrhart, R. C. Flagan, A. Franchin, R. Guida, J. Hakala, A. Hansel, M. Heinritzi, H. Henschel, T. Jokinen, H. Junninen, M. Kajos, J. Kangasluoma, H. Keskinen, A. Kupc, T. Kurtén, A. N. Kvashin, A. Laaksonen, K. Lehtipalo, M. Leiminger, J. Leppä, V. Loukonen, V. Makhmutov, S. Mathot, M. J. McGrath, T. Nieminen, T. Olenius, A. Onnela, T. Petäjä, F. Riccobono, I. Riipinen, M. Rissanen, L. Rondo, T. Ruuskanen, F. D. Santos, N. Sarnela, S. Schallhart, R. Schnitzhofer, J. H. Seinfeld, M. Simon, M. Sipilä, Y. Stozhkov, F. Stratmann, A. Tomé, J. Tröstl, G. Tsagkogeorgas, P. Vaattovaara, Y. Viisanen, A. Virtanen, A. Vrtala, P. E. Wagner, E. Weingartner, H. Wex, C. Williamson, D. Wimmer, P. Ye, T. Yli-Juuti, K. S. Carslaw, M. Kulmala, J. Curtius, U. Baltensperger, D. R. Worsnop, H. Vehkamäki and J. Kirkby, *Nature*, 2013, **502**, 359–363.
- 54 J. Curtius, M. Heinritzi, L. J. Beck, M. L. Pöhlker, N. Tripathi, B. E. Krumm, P. Holzbeck, C. M. Nussbaumer, L. Hernández Pardo, T. Klimach, K. Barmounis, S. T. Andersen, R. Bardakov, B. Bohn, M. A. Cecchini, J.-P. Chaboureaud, T. Dauhut, D. Dienhart, R. Dörich, A. Edtbauer, A. Giez, A. Hartmann, B. A. Holanda, P. Joppe, K. Kaiser, T. Keber, H. Klebach, O. O. Krüger, A. Kürten, C. Mallaun, D. Marno, M. Martinez, C. Monteiro, C. Nelson, L. Ort, S. S. Raj, S. Richter, A. Ringsdorf, F. Rocha, M. Simon, S. Sree Kumar, A. Tsokankunku, G. R. Unfer, I. D. Valenti, N. Wang, A. Zahn, M. Zauner-Wieczorek, R. I. Albrecht, M. O. Andreae, P. Artaxo, J. N. Crowley, H. Fischer, H. Harder, D. L. Herdies, L. A. T. Machado, C. Pöhlker, U. Pöschl, A. Possner, A. Pozzer, J. Schneider, J. Williams and J. Lelieveld, *Nature*, 2024, **636**, 124–130.



Data availability statement for “Nucleation of α -pinene oxidation products with sulfuric acid”

[View Article Online](#)
DOI: 10.1039/D6EA00046K

Data for all figures in the main text and supplementary materials are available at the Zenodo repository (DOI:10.5281/zenodo.19350026).

