

Cite this: *Sustainable Food Technol.*,  
2026, 4, 281

# Nutritional value, phytochemical richness, pharmacological potential, and culinary uses of monkey jackfruit

Raj Singh, <sup>a</sup> C. Nickhil, <sup>\*a</sup> Shweta,<sup>a</sup> Sankar Chandra Deka <sup>a</sup> and R. Nisha<sup>\*b</sup>

Monkey jackfruits are rich in vital nutrients, including carbohydrates, fiber, vitamins, and essential minerals like potassium and magnesium, establishing them as a valuable dietary resource. A comprehensive phytochemical analysis reveals the presence of various bioactive compounds, including flavonoids, phenolics, and saponins, imparting potent antioxidant, anti-inflammatory, and anticancer properties to the fruit. Furthermore, this review highlights the fruit's pharmacological promise, demonstrating its efficacy in managing diabetes, promoting cardiovascular health, and addressing skin disorders due to its anti-inflammatory and antimicrobial attributes. Beyond their medicinal significance, monkey jackfruits have earned a place in culinary arts, finding utility in a diverse range of dishes, from savory curries to delectable desserts. Their unique flavor profile and meat-like texture position them as a compelling meat substitute for vegetarians and vegans. This study underscores the multifaceted nature of monkey jackfruits, emphasizing their nutritional richness, phytochemical potential, pharmacological applications, and culinary versatility.

Received 2nd October 2025  
Accepted 15th November 2025

DOI: 10.1039/d5fb00646e

rsc.li/susfoodtech

## Sustainability spotlight

Being a sustainable food source, monkey jackfruit is highly nutritious, contains different phytochemicals, and exhibits various pharmacological activities. The culinary uses, both traditional and modern, show its importance in diet diversification. It also demonstrates potential for biodiversity and the enhancement of local livelihoods. The utilization of monkey jackfruit can strengthen food systems for sustainable nutrition and contribute to improving health and food diversity.

## 1. Introduction

*Artocarpus lakoocha* Roxb., commonly known as the monkey jackfruit, belongs to the Moraceae family and is a tropical fruit originating from India.<sup>1</sup> The generic name *Artocarpus* is derived from the Greek words “artos” (“bread”) and “karpos” (“fruit”), and the specific name “*lakoocha*” is derived from the Hindi name of the tree, perhaps as a tribute to its Indian origin.<sup>2</sup> The tree has not been observed growing outside its native range, indicating that it has extremely specific climatic requirements for survival. In India, this tree is widespread in the peninsular tropical region. *Artocarpus lakoocha* is known by various names. Lakoocha, Lakooch, Lakoochi, Badhal, Dhau, Dahua, and Dephal are the terms used in Hindi and local dialects. *Lakoocha* also has multiple names in the Sanskrit language, like Airawata, Amlaka, Dahu, Dridhavalkala, and Granthimatphala, which point towards the native status of the tree species in India.

The genus *Artocarpus*, which belongs to the Moraceae family, consists of jackfruit (*Artocarpus heterophyllus*), lakoocha or monkey jack (*A. lakoocha*), chempedak (*A. integer*), breadfruit or breadnut (*A. altilis*), and marang (*A. odoratissima*), comprising over 64 distinct species of the monoecious evergreen trees.<sup>3</sup> A comparison of the major *Artocarpus* species is demonstrated in Table 1. *Artocarpus lakoocha* is a medium or large deciduous or evergreen tree. The deciduous period is very short in the eastern region of India, generally occurring in February and ending in early March. *Artocarpus lakoocha* seedling trees take five years to bear fruit. It can grow up to 15 m in height and is a handsome ornamental plant. The crown is usually well spread, making it an excellent shade and avenue tree. The leaves of the plant are oblong, acute, alternate, 10–25 cm long, glossy green on the upper side when young, rough when old, and leathery. *Artocarpus lakoocha* is a species bearing orange-yellow color for the male flowers and greenish color for the female flowers on the same tree. A single fruit weighs approximately between 250 and 300 g and contains 10–30 seeds of irregular shapes and various sizes. The seeds contain a white sticky latex that is highly recalcitrant, and the seed coat is white and thin. Seeds are dispersed by birds and monkeys.<sup>2,10</sup> The fruits are nearly round

<sup>a</sup>Department of Food Engineering and Technology, School of Engineering, Tezpur University, Assam, India. E-mail: nickhil@tezu.ernet.in

<sup>b</sup>Department of Food Technology, School of Agricultural and Food Technology, Assam Skill University, Assam, India



Table 1 Comparative study of various *Artocarpus* species

Species	Common name	Key nutritional highlights	Major phytochemicals	Bioactivity evidence	Culinary uses
<i>Artocarpus lakoocha</i> <sup>4,5</sup>	Monkey jackfruit	Moisture: ~75.5%, protein: ~0.24%, fat: ~0.63%, crude fiber: ~2.47%, and vitamin C: ~13.4 mg per 100 g	Phenolics, flavonoids, and tannins; oxyresveratrol and related stilbenoids reported in heartwood/ other extracts	<i>In vitro</i> /animal studies report antioxidant, antimicrobial, anti-inflammatory and hepatoprotective activities; traditional medicinal uses are widely documented	Eaten fresh, used in traditional preparations, cosmetics and ethnomedicine
<i>Artocarpus heterophyllus</i> <sup>6,7</sup>	Jackfruit	High carbohydrate (major energy source); seed composition shows substantial starch and notable protein depending on cultivar (values vary)	Phenolics, flavonoids, carotenoids, and prenylated flavonoids; bioactives reported across pulp, seed, leaf and latex	Numerous reports of antioxidant, antimicrobial, anti-inflammatory, and metabolic-benefit potentials <i>in vitro</i> and in animal studies; seeds and leaves frequently show activities in bioassays	Very versatile culinary uses: ripe fruit fresh; unripe used as a vegetable (meat substitute); seeds boiled, roasted or processed into flours and snacks
<i>Artocarpus altilis</i> <sup>8</sup>	Breadfruit	High in complex carbohydrates (starch); good dietary fiber; variable protein; important minerals (K, Ca, and Mg)	Phenolics and other antioxidant compounds present, content influenced by maturity and processing; carotenoids are reported in some cultivars	Extracts (esp. leaves) show antioxidant/anti-inflammatory properties in lab studies; functional/ processing research focuses on food applications	Widely used as boiled, baked, and fried staples; processed into flour and novel food products
<i>Artocarpus camansi</i> <sup>9</sup>	Breadnut	Seeds are relatively nutrient-dense: representative proximate values reported—protein: ~4.9%, fat: ~3.5%, and carbohydrate: ~26.1%	Seeds and other tissues contain phenolics and flavonoids; phytochemical profiling exists but is less extensive than that for jackfruit	<i>In vitro</i> antioxidant activity reported; some studies indicate potential antihyperglycemic activities and other bioactivities in model systems, but data remain limited	Seeds are commonly consumed, boiled or roasted; recent processing studies explore seed flour for bakery and value-added products

or irregular and 5–12 cm in diameter, and they have velvety surfaces. *Artocarpus lakoocha* is a native of the sub-Himalayan humid regions of India and grows up to an altitude of 1200 m above MSL.<sup>11</sup> *Artocarpus lakoocha* can be sustained in both outdoor and underwater environments. It can also be found across the Indian subcontinent, especially in Uttar Pradesh, Jharkhand, Bihar, Assam, West Bengal, Tamil Nadu, and Kerala.<sup>12,13</sup> Monkey jackfruit is one of the most important evergreen trees. The major region they inhabit is Asian nations, such as Bangladesh, Bhutan, Nepal, Malaysia, Singapore, and other neighbouring countries.<sup>14</sup>

It has commercial and medicinal value and is a source of livelihood and nutritional support for the local population. In Bangladesh, this plant is occasionally cultivated in backyards and homestead gardens. The dried aqueous extract made from the heartwood of this plant, also known as “Puaghaad,” has been traditionally used as an antihelminthic. Unripe fruit and male flower spikes are used as vegetables and to make pickles, sauces, and chutneys. When mature, the fruits have an irregular shape, sweet-sour flavor, and a yellow-to-orange hue.<sup>15</sup> The *Artocarpus lakoocha* tree is also used for feed and timber. The hardwood sold as *lakoocha* is comparable to the famous teak wood. As *Artocarpus lakoocha* is durable outdoors and underwater, it can be used for construction, furniture, boat making,

and cabinet work. The tree bark contains 8.5% tannin, is chewed like betel nut, and treats skin ailments. It yields a durable fiber that is good for cordage. The wood and roots have a lavish color dye.

Recently, studies have focused on monkey jackfruit, an underappreciated and largely unknown crop with potential use as food for both humans and animals. The structure of the fruit is illustrated in Fig. 1. The compounds in this fruit play an important role in treating gastric and liver disorders.<sup>10,16</sup> The benefits of fruit and vegetable consumption include a greater




Fig. 1 Different parts of a monkey jackfruit.



life span,<sup>17,18</sup> improved mental health,<sup>19</sup> better cardiovascular health,<sup>20</sup> reduced risk of some cancers,<sup>21</sup> and weight management.<sup>22</sup> In a study conducted in the USA, a relatively low risk of obesity was observed among healthy middle-aged women who consumed fruits and vegetables.<sup>23</sup>

*Artocarpus lakoocha* has a unique flavor and is an excellent source of carbohydrates, lipids, proteins, dietary fiber, minerals (phosphorus and iron), and vitamins (A, C, and B1).<sup>24</sup> The three main sugars in monkey jackfruit are fructose, glucose, and sucrose. Additionally, it is an excellent source of  $\beta$ -carotene, a precursor of vitamin A, and ascorbic acid (vitamin C). Ascorbic acid and  $\beta$ -carotene share antioxidant properties that support healthy eyesight. This plant-based antioxidant plays an important role in maintaining normal health, protecting against coronary heart disease, and combating cancer. It is a top-notch iron source, that is, 28 to 80 mg per 100 g higher than the daily required value in the human diet.<sup>25</sup> The pulp is edible and has hepatoprotective properties due to its antioxidant properties. Various secondary metabolites, such as alkaloids, phenols, flavonoids, tannins, and steroids, have been reported for monkey jackfruit.<sup>13</sup>

For generations, fruits from this species have played a significant role in supplementing human diets, garnering particular attention for their nutritional qualities, phytochemical contents, and bioactive substances, which are recognized as the vital components of overall well-being.<sup>26–28</sup> Monkey jackfruit has gained recognition among researchers and scientists for its potential nutritional, nutraceutical, and pharmacological attributes, making it valuable in various fields, including food, pharmacology, and new product development. According to a study by the World Health Organization (WHO), over two-thirds of the global population in impoverished regions rely on primary or secondary plant metabolites for their health benefits.<sup>29</sup>

Further research on monkey jackfruit is crucial to increase the public awareness of its health benefits and potential as a common fruit crop. Despite the use of micropropagation technology to develop high-quality planting materials, the dissemination of this knowledge is lacking, highlighting the need for improved propagation methods.<sup>30</sup> This review provides a comprehensive summary of the fruit's nutritional value, phytochemical compounds, culinary uses, traditional usage, and validated pharmacological properties. This review offers a comprehensive exploration of monkey jackfruits (*Artocarpus lakoocha*), focusing on their underreported nutritional and phytochemical profiles, pharmacological applications, and unique culinary uses. This study bridges existing knowledge gaps by integrating traditional insights with modern scientific perspectives for a holistic understanding.

## 2. Methodology

A comprehensive literature review was conducted to gather and analyze relevant information on *Artocarpus lakoocha* (commonly known as monkey jackfruit) to understand its nutritional value, phytochemical richness, pharmacological potential, and culinary applications. Renowned scientific databases, such as

PubMed, Scopus, Web of Science, ScienceDirect, and Google Scholar, were used to retrieve the data. The keywords employed during the search included “*Artocarpus lakoocha*,” “monkey jackfruit,” “nutritional composition,” “phytochemicals,” “bioactive compounds,” “health benefits,” and “applications.” The review primarily focused on studies published between 1984 and 2024, although some early publications from the early 20th century were also examined to trace the historical evolution of research on this species. The collected literature included peer-reviewed journal articles, book chapters, and review papers, with a preference for works published in reputable journals with high impact factors and significant citation counts. Both quantitative and qualitative data were extracted, compared, and discussed to provide a holistic understanding of the nutritional profile, phytochemical constituents, pharmacological properties, and culinary applications of the monkey jackfruit. The information obtained was categorized thematically to encompass nutritional composition, phytochemical diversity, pharmacological and health benefits, and traditional and modern culinary uses. Furthermore, a critical analysis of the available literature was conducted to identify emerging research trends, highlight existing knowledge gaps, and outline potential future research directions aimed at unlocking the full nutritional and therapeutic potential of *Artocarpus lakoocha*.

## 3. Nutritional composition and medicinal importance of monkey jackfruits

Monkey jackfruit is a very important species from a nutritional point of view, as shown in Table 2. Beyond basic nutrients, monkey jackfruit has various health-promoting ingredients, such as fiber, polyphenols, vitamins A and C, and minerals. According to Jahan,<sup>24</sup> it has high vitamin C content (171.07 mg per 100 g), beta-carotene (3718.16  $\mu$ g per 100 g), and an

**Table 2** Elements found in fresh *Artocarpus lakoocha* and their relative compositions

Elements	Values
Moisture (%) <sup>25</sup>	81.32–86.95
Carbohydrate (%) <sup>25</sup>	1.32–8.62
Fat (%) <sup>25</sup>	3.17–15
Protein (%) <sup>25</sup>	0.12–0.51
Fibre (%) <sup>25</sup>	1.84–10.21
Vitamin A (IU) <sup>16</sup>	416–423
Thiamin ( $\mu$ g%) <sup>16</sup>	0.01–0.02
Riboflavin ( $\mu$ g%) <sup>16</sup>	0.09–0.15
Niacin ( $\mu$ g%) <sup>16</sup>	0.15–0.3
Sodium (mg per 100 g DW) <sup>31</sup>	22.02–22.75
Potassium (mg per 100 g DW) <sup>31</sup>	165.14–195.38
Calcium (mg per 100 g DW) <sup>31</sup>	47.71–54.53
Magnesium (mg per 100 g DW) <sup>31</sup>	47.71–55.05
Phosphorus (mg per 100 g DW) <sup>31</sup>	196.41–175.25
Iron (mg per 100 g DW) <sup>31</sup>	2.26–2.88
Zinc (mg per 100 g DW) <sup>31</sup>	0.09–0.16
Copper (ppm) <sup>31</sup>	7.81–12.84
Manganese (%) <sup>31</sup>	0.098–0.15



adequate amount of minerals, including zinc (19.82–24.92 ppm), manganese (3.76 mg per 100 g), copper (1.31 mg per 100 g), and iron (15.09 mg per 100 g). The observations for the proximate composition showed the values of moisture, ash, protein, crude fiber, and carbohydrates to be 81.32–86.95%, 1.5–5.33%, 0.12–0.51%, 1.84–10.21%, 3.17–15%, and 1.32–8%, respectively. Different researchers have observed diverse results regarding the nutritional profile. Biochemical analysis showed an ascorbic acid content of approximately 20.86 mg per 100 g the edible portion. The macro mineral content of the fruit includes sodium (1.0 mg per 100 g), potassium (181.0 mg per 100 g), calcium (13.0 mg per 100 g), manganese (42.0 mg per 100 g), and phosphorus (41.0 mg per 100 g).<sup>31</sup> According to a few studies by Gupta,<sup>16</sup> *Lakoocha* fruit contains a significant amount of minerals, such as calcium (66.6 mg), magnesium (23.6 mg), potassium (350 mg), phosphorus (22.1 mg), iron (778 µg), zinc (3981 µg), copper (7974 µg), and manganese (2025 µg). Because of the adequate amount of vitamin C and beta carotenoid in monkey jackfruit, it plays an important role in the human diet.<sup>32</sup> Therefore, it can be stated that *Lakoocha* is an important species from a nutritional perspective.

The popularity of herbal medications is on the rise in India, with people across all demographics embracing their use, a trend that is expected to continue in the foreseeable future. Ethnomedicinal formulations from South and Southeast Asia utilize nearly every part of the *Maackia amurensis* tree. The immature fruit is valued for its medicinal properties, addressing issues such as tridosha impotence, appetite loss, and blood complaints. Conversely, ripe fruits are considered liver tonics because of their sour-sweet taste. Additionally, the powdered bark of the tree serves as a remedy for skin ailments and acts as a vermifuge. The latex of this plant is utilized for its purgative properties. Beyond its medicinal benefits, jackfruit holds cultural and economic significance, serving as a vital food source and offering various utilitarian uses, including in traditional medicine and industry, underscoring its multifaceted importance to communities where it is cultivated.<sup>15,33,34</sup>

## 4. Essential phytochemical and biological activities in monkey jackfruits

Monkey jackfruit contains flavonoids, tannins, saponins, steroids, glycosides, triterpenoids, proteins, phenolic compounds, resins, and squalene. According to Krishnamurthy,<sup>13</sup> screening the secondary metabolites of monkey jackfruit in various organic solvents, including petroleum ether, chloroform, ethanol, and water, showed that the levels of alkaloids (high) and tannins (low) were considerably more dependent on altitude than those of the other metabolites. The therapeutic characteristics of monkey jackfruit include antidiarrheal, anti-glycation, antiaging, anti-inflammatory, analgesic, antibacterial, cytotoxic, insecticidal, and pancreatic lipase inhibitory activities.<sup>15</sup> Table 3 shows the various compounds present in the monkey jackfruit plant. In addition, the combination of monkey jackfruit with goat milk and herbs (neem, holy basil, etc.) can

treat arthritic edema, prevent skin diseases, cleanse wounds, and treat dysentery. A separate study investigated the antioxidant and antiglycation properties of the phytooxyresveratrol extracted from monkey jackfruit. The antiglycation activity of the ethanolic extract of monkey jackfruit exhibited high antiglycation activity. The similar features between the two substances (isolated and standard) suggest their potential use in anti-aging products.<sup>16</sup> Monkey jackfruit (*Artocarpus lakoocha*) contains various phytochemicals, each with potential health benefits.

Only a few studies have been conducted on the phytochemicals of the monkey jackfruit. A recent study by Saleem<sup>51</sup> showed that the HPLC chromatograms of the methanolic extract showed the presence of chromatotropic acid (21.6 µg g<sup>-1</sup>), quercetin (2.5 µg g<sup>-1</sup>), gallic acid (33.4 µg g<sup>-1</sup>), vanillic acid (40.9 µg g<sup>-1</sup>), cinnamic acid (11.1 µg g<sup>-1</sup>), ferulic acid (23.7 µg g<sup>-1</sup>), and kaempferol (2124.7 µg g<sup>-1</sup>). The polyphenol content was also observed as 325.63 mg per 100 g GAEs, flavonoid as 521.98 mg per 100 g CEs, and tannin content as 124.03 mg per 100 g TEs.

The phytoconstituents present in the fruit barks are lakoochanosides, lakoochanone, Moracin C, integrin, cyclommunitin, engeletin, isogemichalcone B, morachalcone A, het erophyllene B, albanin A, Moracin M, artocarpesin, norartocarpin, resveratrol, artocarpanone, and oxyresveratrol, whereas flavones, phenols, tannins, eicosane, and diethyl phthalate are the compounds found in the leaves.<sup>48,52,53</sup> The leaves contain sufficient amounts of kaempferol and quercetin. Albanin A,<sup>54</sup> isoartocarpesin,<sup>55</sup> cudraflavone C,<sup>56</sup> norartocarpin,<sup>57</sup> cudraflavone B,<sup>58</sup> and catechin A<sup>57</sup> were reportedly observed in various parts of the same plant.<sup>59</sup>

Extraction and analytical methods provide a comprehensive approach for separating and identifying bioactive compounds from *Artocarpus lakoocha*, contributing to its applications in pharmacology and nutraceuticals. Most of the extractions were conducted using ethanol and methanol as the solvents.<sup>60–62</sup> Ultrasound-assisted extraction has also been used to extract compounds such as oxyresveratrol, which is also considered a green method.<sup>63</sup> Soxhlet extraction is the traditional method for isolating phytochemicals. The identification of active compounds in crude extracts is generally initiated using thin-layer chromatography. The free radical scavenging activity of the extract was identified using antioxidant methods, such as the DPPH and ABTS assays. Chaurasia and Pandey<sup>45</sup> conducted a brief analysis. UV-Vis spectrophotometry is generally used to analyze the antioxidant activities, total phenols, flavonoids, and tannins.<sup>5</sup>

The main bioactive compounds in monkey jackfruit that exhibit biological activities are oxyresveratrol, flavonoids, stilbenoids, tannins, and phenolic compounds. Oxyresveratrol is a major bioactive compound in the fruit, which exhibits antioxidant activities by neutralizing free radicals and reducing oxidative stress through electron donation. It also inhibits reactive oxygen species (ROS) production, contributing to its anti-inflammatory and anti-aging effects.

The standardization of bioactive compound extraction from monkey jackfruit (*Artocarpus lakoocha*) is essential for ensuring



Table 3 Compounds present in the parts of the *Artocarpus lakoocha* tree

Section of the tree	Compound	Molecular formula	Function
Heartwood <sup>35–39</sup>	Oxyresveratrol	C <sub>14</sub> H <sub>12</sub> O <sub>4</sub>	Antiviral, anti-poliovirus, anti-measles virus, anti-HSV and cytotoxic activities; significantly slowed the development of skin lesions and saved mice from death
	Lakoochin A and B	C <sub>26</sub> H <sub>30</sub> O <sub>4</sub> and C <sub>29</sub> H <sub>34</sub> O <sub>4</sub>	The structures of Lakoochin A and B were elucidated by the analysis of their spectral data. Lakoochin A and B exhibited antimycobacterial activity. While Lakoochin A was cytotoxic against the breast cancer cell line but inactive toward nasopharyngeal carcinoma cells, compound Lakoochin B possessed cytotoxicity against both the breast cancer and nasopharyngeal carcinoma cells
	Norartocarpin	C <sub>15</sub> H <sub>10</sub> O <sub>6</sub>	Norartocarpin may be used as a skin whitening agent in the treatment of hyperpigmentation illnesses and in skin whitening cosmetics due to its inhibitory effects on antimelanogenesis
	Artocarpin	C <sub>26</sub> H <sub>28</sub> O <sub>6</sub>	Artocarpin has been shown to have cytotoxic, antiplasmodial, anti-inflammatory, and antibacterial activities
	Cycloartocarpin and Cudraflavone C	C <sub>26</sub> H <sub>26</sub> O <sub>6</sub> and C <sub>25</sub> H <sub>26</sub> O <sub>6</sub>	It has exhibited antitubercular and antiplasmodial activities, while exhibiting moderate cytotoxic activities towards human oral epidermoid carcinoma and human breast cancer
Fruit pericarp <sup>40,41</sup>	Cycloartenone	C <sub>30</sub> H <sub>48</sub> O	It determines antibacterial, antioxidant, anthelmintic and insecticidal efficacy
	α-Amyrin acetate and β-amyrin acetate	C <sub>32</sub> H <sub>52</sub> O <sub>2</sub>	It treats various inflammatory disorders. Mechanical hypersensitivity, mechanical sensitization, and oedema, as well as antioxidant, cardiogenic, and antihyperlipidemic properties
Leaves <sup>42–45</sup>	Lupeol acetate	C <sub>32</sub> H <sub>52</sub> O <sub>2</sub>	It can assist in managing the overall myocardium processes by lowering total cholesterol, triglyceride, and phospholipid levels
	Phenols (vanillic acid)	C <sub>8</sub> H <sub>8</sub> O <sub>4</sub>	Antioxidant and potential anti-cancer properties
	Gallic acid	C <sub>7</sub> H <sub>6</sub> O <sub>5</sub>	
	Cinnamic acid	C <sub>9</sub> H <sub>8</sub> O <sub>2</sub>	
	Ferulic acid	C <sub>10</sub> H <sub>10</sub> O <sub>4</sub>	
	Tannin (tannic acid)	C <sub>76</sub> H <sub>52</sub> O <sub>46</sub>	
Flower <sup>46,47</sup>	Eicosane	C <sub>20</sub> H <sub>42</sub>	Antifungal
	Diethyl phthalate	C <sub>12</sub> H <sub>14</sub> O <sub>4</sub>	Antimicrobial and antioxidant
	Beta-carotenoids	C <sub>40</sub> H <sub>56</sub>	Antioxidant properties, immune support, skin health, anti-inflammatory effects and cancer prevention
	Flavonoids (alpha-amyrin)	C <sub>30</sub> H <sub>50</sub> O	Antioxidant properties, anti-inflammatory effects and cardiovascular health
	Beta-amyrin		
Bark <sup>34,45,48–50</sup>	Lakoochanone	C <sub>29</sub> H <sub>26</sub> O <sub>8</sub>	Anti-inflammatory and antioxidant properties



Table 3 (Contd.)

Section of the tree	Compound	Molecular formula	Function
	Lakoochanosides	—	Antioxidant properties
	Catechin	C <sub>15</sub> H <sub>14</sub> O <sub>6</sub>	Antioxidant and potential cardiovascular benefits
	Moracin	C <sub>19</sub> H <sub>18</sub> O <sub>4</sub>	Antioxidant, anti-inflammatory, antiviral and antimalarial properties
	Cyclocommunin	C <sub>25</sub> H <sub>24</sub> O <sub>6</sub>	Antioxidant
	Artocarpesin	C <sub>20</sub> H <sub>18</sub> O <sub>6</sub>	Antioxidant and anti-inflammatory properties
	Artocarpanone	C <sub>16</sub> H <sub>14</sub> O <sub>6</sub>	Anti-inflammatory, cytotoxic and anti-diarrheal properties
	Engeletin	C <sub>21</sub> H <sub>22</sub> O <sub>10</sub>	Anti-inflammatory and antioxidant properties
	Heterophyllene B	C <sub>13</sub> H <sub>16</sub> N <sub>2</sub> O <sub>5</sub>	Antioxidant, antiplasmodia and anti-inflammatory properties
	Isogemichalcone B	C <sub>26</sub> H <sub>30</sub> O <sub>6</sub>	Potential antioxidant and anti-inflammatory properties
	Albanin A	C <sub>20</sub> H <sub>18</sub> O <sub>6</sub>	Skin whitening agents

the reproducibility, quality control, and industrial scalability of oxyresveratrol (ORV)-rich formulations. The extraction efficiency of ORV and related phenolics depends on solvent polarity, temperature, particle size, and extraction time. Conventional solvent extraction using ethanol or methanol (60–80%) has been widely reported, but greener alternatives, such as deep eutectic solvents (DES) and aqueous ethanol under ultrasound or microwave assistance, significantly improve the yield and purity while minimizing solvent residues.<sup>4,63</sup> Optimization studies using response surface methodology (RSM) demonstrated that extraction at 60–70 °C for 30–45 min with a solvent-to-solid ratio of 20 : 1 (v/w) provides the highest ORV recovery from heartwood, achieving concentrations of 50–80 mg per g extract.<sup>64</sup> Post-extraction purification using column chromatography or macroporous resin further enhances the ORV content and removes pigments and polysaccharides. Analytical standardization relies on the HPLC and LC-MS quantification of ORV using authentic standards, along with total phenolic and flavonoid content assays for the quality benchmarking of the extracts. The adoption of validated analytical protocols and solvent recycling enhances both the environmental and economic sustainability of large-scale extraction. Such standardization enables reproducible pharmacological testing and regulatory compliance for nutraceutical and cosmeceutical applications.<sup>4,63,64</sup>

*Artocarpus lakoocha*, particularly the heartwood-derived stilbenoid oxyresveratrol (ORV), triterpenoids, such as lupeol/lupeol acetate, and assorted triterpenoid/steroidal saponins, produces a coherent mechanistic story that links anti-melanogenic action, eicosanoid (arachidonic acid) signalling, pharmacokinetics, mitochondrial protection, and redox modulation. ORV is the most directly documented anti-melanogenic compound. In cellular melanocyte models, it lowers melanin synthesis by inhibiting tyrosinase activity and downregulating the MC1R → cAMP → MITF axis (thereby reducing tyrosinase and TRP expression), effects that are closely

associated with the suppression of cellular oxidants and the restoration of redox balance in melanocytes.<sup>65,66</sup> Lupeol and its acetate derivative predominantly act as the anti-inflammatory modulators of the arachidonic acid cascade. They reduce COX-2 expression and prostaglandin production and interfere with upstream TLR/NF-κB signalling in inflammatory models, thereby diminishing prostaglandin-driven proliferation and paracrine signals that can promote pigimentary and tumor-promoting microenvironments.<sup>67</sup> Pharmacokinetically, ORV is rapidly absorbed but undergoes extensive first-pass metabolism (notably glucuronidation/sulfation) and has relatively low oral bioavailability. Animal studies have shown markedly altered plasma AUC and clearance when ORV is co-administered with bioenhancers (*e.g.*, piperine) or formulated as nano-carriers (NLC/SMEDDS). These are strategies that substantially increase systemic exposure and tissue delivery—a critical point when linking *in vitro* potencies to any *in vivo* therapeutic hypothesis.<sup>68</sup> At the organelle level, ORV and specific *Artocarpus* extracts exhibit dual redox behaviors. At physiological/sub-cytoprotective concentrations, they scavenge ROS, preserve mitochondrial membrane potential, limit mitochondrial cytochrome-c release, and inhibit apoptosis in UVB-stressed keratinocytes or toxin models. However, at high concentrations (or in sensitive cancer cells), they may behave as pro-oxidants (including inducing ferroptosis or mitochondrial dysfunction), thereby triggering cell death. This mechanistic bifurcation explains both the cytoprotection and selective anticancer cytotoxicity observed in separate studies.<sup>69</sup> Triterpenoid and steroidal saponins present in *Artocarpus* spp. further support mitochondrial integrity and redox homeostasis in non-neoplastic models by restoring GSH, elevating antioxidant enzymes (SOD, CAT, and GSH-Px), stabilizing mitochondrial networks, and inhibiting MAPK-mediated apoptotic signalling. These actions have been reported for saponins across multiple plant sources and are likely to be operative in jackfruit relatives, where saponins have been detected.<sup>70</sup> Taken



together, a recent real-time mechanistic study of oxyresveratrol from *Artocarpus* showed the clear inhibition of melanogenesis through MC1R/cAMP/MITF suppression, along with a concurrent reduction in cellular oxidants, thereby experimentally linking ORV's biochemical (tyrosinase and signalling) targets to its antioxidant and mitochondrial-stabilizing effects in skin cells. Meanwhile, separate pharmacokinetic studies have demonstrated that the meaningful systemic or topical delivery of ORV depends on formulation strategies to overcome rapid metabolism.

## 5. Pharmacological importance of monkey jackfruit

*Artocarpus lakoocha* has emerged as a pharmacologically important species, largely because its heartwood, bark, leaves, and fruit contain a concentrated amount of oxyresveratrol (ORV) and a suite of phenolics and flavonoids that show reproducible bioactivities *in vitro* and *in vivo*.<sup>4</sup> It has shown promising pharmacological activities in various studies (Table 4). The cytotoxicity test is an *in vitro* test based on a cell culture that is used to evaluate the safety of pharmaceuticals, cosmetics, food additives, and pesticides, as well as to determine the presence of antineoplastic activities in a chemical. *In vitro*, cytotoxic chemicals are toxic to cancer cells and can also be hazardous to rapidly reproducing normal cells. These compounds will exhibit antitumor activities if this toxicity is transferred across tumors *in vivo*. Monkey jackfruit plants exhibit numerous biological activities, including cytotoxicity. Based on the collected data, a comparison of the effects of various doses of *Artocarpus lakoocha* leaf extract with ethanol (10, 20, 40, 60, 80, and 160 g mL<sup>-1</sup>)<sup>72</sup> and pericarp methanol extract (10–1000 g mL<sup>-1</sup>)<sup>32</sup> was performed to determine the IC<sub>50</sub> of the antioxidant activities *via* regression analysis. *Artocarpus lakoocha* extract was found to destroy brine shrimp in a dose-dependent manner. The saponin, alkaloid, and cardiac glycoside content of the extract may explain its lethality to brine shrimp.<sup>32</sup> The anti-inflammatory effects of *Artocarpus lakoocha* leaves extracted in methanol were shown to be dose-dependent and statistically significant ( $p \leq 0.05$ ). At a dosage of 200 mg kg<sup>-1</sup>, *Artocarpus lakoocha* was more effective than indomethacin in reducing inflammation (64.90%). The dosage of the *Artocarpus lakoocha* leaf methanol extract reduced writhing by 29.63% and 57.41% ( $p \leq 0.05$ ). The consequence of arachidonic acid breakdown is one item that might induce inflammation. Arachidonic acid is an unsaturated fatty acid with 20 carbon atoms. Arachidonic acid is produced from phospholipids when cell phospholipases are activated by physical, chemical, or biological stimuli.<sup>75</sup> The leaves of the *Artocarpus lakoocha* plant (100 mg kg<sup>-1</sup>) have a modest antidiarrheal effect. This is based on the information gathered. However, at a dosage of 200 mg kg<sup>-1</sup>, the effect is halted (68.11%), which is comparable to what the usual medicine, loperamide, can accomplish (71.1%). Thus, the extract given prevented diarrhea *via* an antisecretory mechanism. This was also demonstrated by a decrease in the number of wet stools in the experimental test group. Diarrhea

treatments aim to correct the diet, halt excessive water and electrolyte loss, correct acid–base imbalances, treat the symptoms, address the causes of diarrhea, and treat the illnesses that produce diarrhea as a side effect. As a result, the most crucial part of managing diarrhea is food control.<sup>81</sup> Additionally, the pharmacological properties of the *Artocarpus lakoocha* plants include liver and nerve cell protection. The puaghaad and oxyresveratrol present in fruits will increase cell survival, notably ROS levels and lipid peroxidation, the neuroprotective impact of *Artocarpus lakoocha* wood extract with aqueous suggests that mitochondrial protection is difficult to detect. Because of the action of oxyresveratrol on the synthesis and pharmacokinetics of redox-sensitive antioxidant enzymes, ingesting puaghaad orally might be a great method for preventing transient neurological problems.<sup>71</sup>

## 6. Applications of monkey jackfruit

*Artocarpus lakoocha* has a sweet and slightly tangy flavor, and its flesh is often compared to that of the regular jackfruit. Fruit pulp is often sweet and sour and is eaten fresh, but is also frequently cooked into curries. Curry, pickles, and sauce are made from the unripe fruit and male flower spike of *Artocarpus lakoocha*. The fruits and male flowers of this tree are among the portions that may be consumed fresh, cooked, steamed, or roasted. The wide, flat seeds may be eaten. They also have applications in the cosmetic industry as they can be used as a skin-whitening ingredient. The presence of flavones and flavonoids is responsible for anti-cancer activities and can help in the formulation of chemotherapeutic agents. Lectin in the plant can be used for the treatment of HIV, suggesting the wide application in biochemistry. The value-added products developed from the fruits are dehydrated raw jackfruit slices, dehydrated raw jackfruit flour, jackfruit mixture, candy, squash, jam, jelly, jackfruit ganache, jackfruit leather, chocolate, cake, muffins, honey, pudding, and jackfruit wine. Monkey jackfruit pulp can also be used to make pickles, curry, sauce, and juice with sugar, cardamom, and black pepper, depending on the taste profile of the individual. The raw fruits are widely used for the purpose of pickle preparation.<sup>82</sup> Apart from these, *Artocarpus lakoocha* holds immense potential for the development of value-added products, such as fermented beverages, nutraceutical extracts, anti-aging cosmeceuticals, and functional snacks, owing to its rich nutrient profile and high oxyresveratrol (ORV) content. The pulp and seeds, abundant in fermentable sugars and phenolic compounds, can be transformed into probiotic or alcoholic fermented beverages through controlled enzymatic saccharification, followed by lactic acid or yeast fermentation, enhancing antioxidant activity, polyphenol bioavailability, and shelf stability. This is similar to the optimized jackfruit seed drink fermentations reported by Hoang.<sup>83</sup> The heartwood and bark, rich in ORV, can be used to produce standardized nutraceutical extracts *via* green solvent or deep eutectic extraction, yielding ORV concentrations up to 80% w/w with potent anti-inflammatory, antioxidant, and neuroprotective effects.<sup>4,63</sup> Given ORV's low oral bioavailability, encapsulation in nanoparticles or cyclodextrin complexes or co-



Table 4 Pharmacological activities of *Artocarpus lakoocha*

Sample	Extraction	Technique	Function	Finding
Heart wood <sup>46,71</sup>	Aqueous	Bovine serum albumin (BSA)	Antiglycation	AGE-BSA was suppressed by the extract
	Aqueous	H <sub>2</sub> O <sub>2</sub> -induced oxidative stress in SH-SY5Y cells	Neuroprotective	The extract possesses neuroprotective properties
Fruit <sup>51</sup>	Methanol	<i>In vivo</i> -induced paracetamol in mice	Hepatoprotective	The extract inhibited elevations in liver function tests as well as histological changes caused by paracetamol
Leaf <sup>65,72-76</sup>	Methanol	Brine shrimp lethality bioassay	Cytotoxic	The extract has a high toxicity level. The extract's LC <sub>50</sub> value was 2.83–2.94 g mL <sup>-1</sup>
		Carrageenan-induced paw edema test in mice	Anti-inflammatory	At a dosage of 200 mg kg <sup>-1</sup> , the extract had anti-inflammatory properties
		Acetic acid-induced writhing test	Analgesic	The extract showed inhibitions of 29.63% and 57.41%
		Castor oil-induced diarrhea	Antidiarrhoeal	The extract decreased castor oil-induced diarrhoea in test animals at doses of 100 and 200 mg kg <sup>-1</sup>
		<i>In vivo</i> , hyperlipidemia-induced rats were given an extract	Anticholesterol	The extract significantly increased serum high-density lipoprotein (HDL) while lowering serum total cholesterol, triglycerides, and LDL
	Ethanol	MTT test with NIH-3T3 cells in mice <i>in vivo</i>	Proliferative and wound healing	The extract was shown to have proliferative and wound-healing effects
	Methanol	1,1-Diphenyl-2 picryl-hydrazil (DPPH) 2,2-Azino-bis-3-ethylbenzothiazoline-6-sulphonic acid (ABTS) 1,1-Diphenyl-2 picryl-hydrazil (DPPH)	Antioxidant and antimicrobial properties	IC <sub>50</sub> value is 111.98–146.18 µg mL <sup>-1</sup> IC <sub>50</sub> value is 138.26–138.92 µg mL <sup>-1</sup>
<i>n</i> -Hexane, ethyl acetate, and ethanol	1,1-Diphenyl-2 picryl-hydrazil (DPPH)	Antioxidant	IC <sub>50</sub> value is 26.95–26.959 µg mL <sup>-1</sup> The IC <sub>50</sub> values are as follows: <i>n</i> -hexane (1062.03–1063.45 g mL <sup>-1</sup> ), ethyl acetate (323.18–323.20 g mL <sup>-1</sup> ), and ethanol (99.23–99.30 g mL <sup>-1</sup> )	
Heart pericarp <sup>77</sup>	Methanol	Brine shrimp lethality bioassay	Cytotoxic	The extract proved toxic, having an IC <sub>50</sub> of 427.74 g mL <sup>-1</sup>
Bark <sup>78</sup>	Hydro-methanolic	Tests for induced nociception using formalin; paw edema caused by carrageenan and acetic acid; tail immersion and hot plate tests	Antinociceptive	There is an antinociceptive action in the extract
Crude <sup>79</sup>	Aqueous	Schistosoma mansoni-infected <i>in vivo</i> mice	Schistosomicidal	At a concentration of 250 g mL <sup>-1</sup> , the extract inhibited motility
Dry fruit <sup>80</sup>	Methanol	1,1-Diphenyl-2 picryl-hydrazil (DPPH) 2,2-Azino-bis-3-ethylbenzothiazoline-6-sulphonic acid (ABTS) Hydroxyl	Antioxidative properties	IC <sub>50</sub> value is 6.72–10.97 mg ascorbic acid per g IC <sub>50</sub> value is 2.32–3.64 mg ascorbic acid per g



Table 4 (Contd.)

Sample	Extraction	Technique	Function	Finding
Seeds <sup>49</sup>	Methanol	Superoxide anion 1,1-Diphenyl-2-picrylhydrazil (DPPH) Ferric reducing antioxidant potential	Antioxidant and anticancer activities	IC <sub>50</sub> value is 41.35–53.1 mg BHT per g IC <sub>50</sub> value is 0.47–0.60 mg ascorbic acid per g 18.28–22.50 µg mL <sup>-1</sup> 31.22–32.11 µg mL <sup>-1</sup>

administration with piperine can enhance systemic absorption.<sup>68</sup> The topical formulations of *Artocarpus lakoocha* extracts show significant tyrosinase inhibition, melanin suppression, and reactive-oxygen-species scavenging activities, validating their use in anti-aging and skin-brightening cosmeceuticals.<sup>64</sup> These effects stem from ORV's dual mechanism of direct enzyme inhibition and the down-regulation of melanogenic genes (MITF, TRP-1, and TRP-2), complemented by NF-κB modulation that suppresses inflammatory cytokine expression. Beyond extracts, the seeds of *Artocarpus lakoocha* represent a valuable raw material for functional snacks, as seed flours provide resistant starch, dietary fiber, and proteins suitable for cookies, extruded chips, and nutrient bars. Prior trials with jackfruit and breadnut seed flours confirm improved textural and nutritional properties without sensory compromise.<sup>84</sup> The integration of ORV microcapsules into seed-based snacks could yield antioxidant-enriched products for preventive nutrition markets. To commercialize these products, process standardization, safety profiling, and consumer sensory evaluations are essential, along with the sustainable sourcing and quality control of raw materials. Collectively, these value-added innovations could transform monkey jackfruit from an

underutilized tropical species into a high-value resource for functional foods, nutraceuticals, and cosmeceutical applications, contributing simultaneously to nutrition security, circular bioeconomy, and rural livelihoods. A paper is also available on the preparation of powder, juice, sambhar, papad, hulli, and various local meals, such as fish curry and chicken curry, from monkey jackfruit.<sup>13</sup> The processing challenges faced during the use of latex due to its sticky texture have led to mechanical handling complications. As the fruit is highly perishable, post-harvest losses occur. Therefore, the development of preservation techniques, such as drying and freezing, is required. The bioactive compounds may vary with the change in environment, such as soil and climate.

## 7. Potential health benefits of monkey jackfruit

The health benefits of monkey jackfruit are shown in Fig. 2. This unique fruit is rich in essential nutrients, making it a valuable addition to a balanced diet. Monkey jackfruit is a good source of dietary fiber, which can aid in digestion and help prevent constipation. It also contains vitamins and minerals, including




Fig. 2 Potential health benefits of monkey jackfruit.



vitamin C, which supports the immune system, and potassium, which is important for heart health and maintaining proper blood pressure.<sup>1</sup> Some pharmacological activities are found in the monkey jackfruit plant: anti-inflammatory, antiviral, anti-cancer, and anti-HIV properties. The presence of bioactive components in different plant parts leads to specific health benefits. The bioactive compounds such as cycloartenone, cycloartenol,  $\alpha$ -amyrin acetate, and  $\beta$ -amyrin acetate, are present in fruit pericarp. Hypolipidemic, and anti-atherosclerotic, act as first precursors for the synthesis of sterols and stanols, anti-inflammatory activity, hypersensitivity, oedema, and radical scavenging.<sup>40,41</sup> The leaves contain two bioactive compounds, *i.e.*, lupeol acetate, which is responsible for antioxidant activity; decreasing cholesterol, phospholipid, and triglyceride levels; and interrupting cardiovascular disease,<sup>42</sup> and oxyresveratrol, which helps with the significantly delayed development of skin lesions and is anti-viral, cytotoxic, anti-HSV, and anti-HIV.<sup>35,85</sup> However, the seed contains lectin, which helps in carbohydrate (mannose) binding and antigen detection. The immature fruit is hot, sour, and sweet and causes the loss of appetite and blood complaints. The mature fruit tastes sour and sweet and is a tonic for the liver.<sup>15</sup> The potential health benefit is discussed in detail below.

### 7.1 Cytotoxic and anticancer activities

The species of *Artocarpus*, particularly *Artocarpus lakoocha*, have emerged as promising sources of cytotoxic phytochemicals, with most existing data derived from *in vitro* studies using crude extracts and purified compounds. The heartwood, bark, and fruit of *A. lakoocha* are rich in stilbenoids, especially oxyresveratrol, along with other phenolics, triterpenoids (such as cycloartenone and  $\alpha$ -amyrin derivatives), and lectins, all of which exhibit anti-proliferative effects against various cancer cell lines.<sup>50</sup> Mechanistic studies have indicated that oxyresveratrol induces cytotoxicity *via* several interconnected mechanisms.<sup>86</sup> It triggers apoptosis by modulating apoptotic gene expression and cell cycle checkpoints, increases intracellular ROS levels and DNA damage signals, and affects the regulation of genes related to metastasis and DNA repair, and these are effects observed in MCF-7 breast cancer and other models.<sup>87</sup> Recent research suggests that oxyresveratrol can also induce ferroptosis, an iron-dependent, lipid peroxidation-driven cell death pathway, in breast cancer cells by inhibiting the EGFR  $\rightarrow$  PI3K  $\rightarrow$  AKT  $\rightarrow$  GPX4 cascade, emphasizing its role in redox-based anticancer mechanisms.<sup>88</sup> Crude methanolic and ethanolic extracts of *Artocarpus* tissues consistently exhibit micromolar to low microgram-per-milliliter IC<sub>50</sub> values across cancer types (breast, colon, cervical, and lung), and bioassay-guided fractionation confirms that cytotoxicity is concentrated in polyphenol-rich fractions, highlighting the role of specific small molecules rather than general extract toxicity.<sup>89</sup> Macromolecular components, including water-soluble polysaccharides from *A. heterophyllus*, have also demonstrated biological effects in colon cancer models by modulating inflammatory mediators, indicating diverse mechanisms across the genus.<sup>90</sup> However, most available studies are limited to *in vitro* assays or simple organismal screens, use inconsistent

extraction protocols, and rarely assess the selectivity toward non-malignant cells. In addition, pharmacokinetic and *in vivo* toxicity data are scarce.<sup>4</sup>


Therefore, although oxyresveratrol and related *Artocarpus* constituents are strong anticancer leads, systematic *in vivo* efficacy, safety, and mechanistic studies, including dose-response relationships, molecular target validation, and combinatorial treatment assessments, are crucial for clinical translation. Future research should focus on standardized extraction, comparative potency profiling, and well-designed animal models to elucidate the underlying mechanisms and therapeutic potential of these compounds.

### 7.2 Anti-inflammatory and analgesic properties

*Artocarpus lakoocha* is a nutritionally rich, yet underutilized tropical fruit. Its edible parts and traditional formulations provide carbohydrates, dietary fiber, essential vitamins and minerals, and a complex array of phytochemicals, such as phenolics, flavonoids, stilbenoids (notably oxyresveratrol), and prenylated flavones like artonins. These compounds collectively underpin the anti-inflammatory and analgesic actions of the plant. Mechanistically, both crude extracts and isolated constituents exhibit potent antioxidant effects by scavenging ROS and suppressing lipid peroxidation, alongside the direct modulation of inflammatory signaling pathways, including the inhibition of pro-inflammatory enzymes and mediators (*e.g.*, prostaglandins and nitric oxide), membrane stabilization, and the suppression of neutrophil and macrophage activity, all of which help alleviate inflammation and pain. *In vivo* validation in rodent models revealed that methanolic and ethanolic extracts significantly reduced carrageenan-induced paw edema and produced notable analgesic effects in hot-plate, tail-flick, and acetic-acid-induced writhing tests, with some doses approaching those of the standard drugs. These findings showed a strong association with high levels of phenolic and flavonoid compounds.<sup>78</sup> Recent research isolating oxyresveratrol has confirmed its significant anti-inflammatory properties in skin inflammation models by reducing oxidative stress and decreasing inflammatory cytokine levels, while initial toxicological assessments indicate favorable safety margins.<sup>91,92</sup> Overall, the nutritional and phytochemical varieties of *Artocarpus lakoocha* highlight its dual potential as both a functional food and a pharmacological agent, offering basic antioxidant and anti-inflammatory benefits through diet and more pronounced effects through concentrated extracts. However, important research priorities include standardizing phytochemical characterization across different cultivars; understanding the structure-activity relationships of artonins and stilbenoids; and conducting pharmacokinetic, chronic toxicity, and clinical studies to support their therapeutic use.<sup>93</sup>

### 7.3 Gastroprotective and antidiarrheal effects

*Artocarpus lakoocha* combines moderate nutritional content, including fiber, carotenoids, vitamins, and minerals, with a high concentration of bioactive phytochemicals, such as oxyresveratrol, flavonoids, tannins, and phenolic acids. These



compounds collectively support the traditional use of the species and experimentally validate gastroprotective and anti-diarrheal properties.<sup>94</sup> The strong antioxidant and anti-inflammatory effects of these polyphenols contribute to mucosal protection against oxidative and inflammatory damage, which are key factors in ulcer and secretory diarrhea.<sup>63</sup> Tannins and other astringent compounds reduce intestinal secretion and toxin binding, whereas flavonoids and oxyresveratrol regulate gut motility, suppress inflammatory mediators, and neutralize ROS, collectively resulting in measurable anti-diarrheal and gastroprotective outcomes *in vivo*.<sup>4,95</sup> Studies on the *Artocarpus lakoocha* extracts have demonstrated a wide range of biological activities, including anti-diarrheal, antioxidant, hepatoprotective, and antimicrobial activities, suggesting their value as both functional food ingredients and phytotherapeutic candidates.<sup>51</sup> In representative animal experiments, extract doses of 100–200 mg kg<sup>-1</sup> significantly reduced the incidence and severity of diarrhea, decreased intestinal fluid accumulation, and preserved mucosal integrity; these are effects that correlated with phenolic content and antioxidant indices.<sup>72</sup> These findings link nutritional composition and phytochemical richness to antioxidant and antisecretory mechanisms that provide gastroprotection. However, to advance clinical use, standardized extraction, dose–response, safety profiling, and constituent identification are essential.<sup>5</sup> Therefore, *Artocarpus lakoocha* is a nutritionally beneficial and phytochemically potent species with consistent preclinical gastroprotective and anti-diarrheal efficacy. Targeted research on chemical standardization, mechanistic pathways, and clinical validation will define its future use in dietary and medicinal contexts.

#### 7.4 Neuroprotective and hepatoprotective activities

*Artocarpus lakoocha*, valued nutritionally and medicinally, offers edible pulp, seeds, and heartwood extracts rich in carbohydrates, fiber, vitamins (especially vitamin C), minerals, and a diverse array of bioactives, including polyphenols, flavonoids, stilbenoids (notably oxyresveratrol), and polysaccharides. These compounds collectively explain the observed neuroprotective and hepatoprotective effects of *Artocarpus lakoocha*. Multiple studies have indicated strong antioxidant and anti-inflammatory potential *in vitro* and *in vivo*, which are mechanisms that underpin both neuronal and hepatic protection.<sup>4</sup> In neuroprotection, the ethanolic extracts of *Artocarpus lakoocha* have been shown to reduce oxidative stress, suppress inflammatory mediators, and enhance neuronal survival in cellular and rodent models, effects likely mediated by free radical scavenging, the inhibition of pro-inflammatory signaling, and the stabilization of mitochondrial functions.<sup>96</sup> For hepatoprotection, extracts from *Artocarpus* species normalize serum enzyme levels, restore hepatic antioxidant defenses, and reduce histopathological liver damage in toxin-induced injury models (*e.g.*, CCl<sub>4</sub>, paracetamol, and diet-induced stress). In some cases, these protective effects are comparable to those of silymarin, and contributions from both phenolic compounds and immunomodulatory polysaccharides have been

identified.<sup>51,97</sup> These findings collectively validate traditional uses and justify advancing standardized extracts and isolated compounds, such as oxyresveratrol, into pharmacokinetic, toxicity, and dose-ranging studies, followed by disease-specific preclinical trials (*e.g.*, Parkinson's, Alzheimer's, or NAFLD models) and, ultimately, clinical testing.

## 8. Challenges

There are key challenges that impede the advancement of research and the practical use of *Artocarpus lakoocha*. Initially, compositional and phytochemical data are fragmented and often non-standardized; studies use different tissues, extraction methods, and reporting units, making cross-study comparisons and nutrition labeling unreliable.<sup>5</sup> Compounds such as oxyresveratrol and other phenolics have been identified; systematic untargeted metabolomic and chemotype mapping across cultivars and environments are lacking, limiting discovery, quality control and the reproducibility of bioactivity claims.<sup>98</sup> Third, safety and translational evidence remain insufficient; promising *in vitro* and animal results have not been matched by rigorous toxicology, ADME, or human clinical studies required for nutraceutical or therapeutic development. Recent preliminary toxicology work highlights the need for broad safety profiling.<sup>92</sup> Furthermore, poor postharvest handling, short shelf life, and the paucity of scalable processing protocols hinder value-addition, market integration and smallholder uptake.<sup>99</sup> Finally, limited socioeconomic and agronomic studies mean there is little evidence on the best agroforestry models, cultivar selection for yield/nutrition, and market pathways needed to incentivize farmers. Addressing these challenges requires coordinated interdisciplinary research, standardized analytical methods, and targeted translational funding.<sup>4</sup>

## 9. Future perspective and research gap

*Artocarpus lakoocha* holds clear and specific promise as a multifunctional crop; its fruit pulp delivers a noteworthy proximate and micronutrient profile (*e.g.*, high moisture, low fat and protein, and measurable vitamin C and fiber), indicating a real potential to diversify diets and help address micronutrient shortfalls in undernourished communities when integrated into local food systems. Its heartwood, bark, leaves and fruit are rich in phenolics, flavonoids and distinctive bioactives, such as oxyresveratrol, which are compounds that have shown strong antioxidant, antibacterial and anti-inflammatory properties in laboratory studies and make *Artocarpus lakoocha* a promising source for nutraceuticals, topical cosmeceuticals and standardized botanical extracts. Because *Artocarpus lakoocha* is already used in agroecosystems (including as a fodder tree) and related *Artocarpus* species have demonstrated climate-resilience traits, the deliberate integration of monkey jackfruit into agroforestry and diversified production systems could enhance on-farm biodiversity, improve soil health, increase carbon storage, and provide



farmers with an additional, climate-resilient income stream, all of which support food-security goals in low-latitude regions.<sup>100</sup> Practically, the edible pulp, flowers and seeds of the species lend themselves to a wide range of value-added food products (fresh, preserved, flours and seed-based ingredients), while the heartwood and leaf extracts already show enough bioactivity to justify translational R&D into safe topical and oral formulations.

However, realizing this potential for monkey jackfruit requires targeted scientific and development actions because important evidence gaps remain. Standardized, cultivar and tissue-specific compositional datasets are sparse (limiting reliable nutrition claims and product labeling), and phytochemical research remains fragmented (few untargeted metabolomics or chemotyping studies exist to map chemical diversity across cultivars and environments).<sup>4</sup> Toxicological and safety data are limited despite encouraging *in vitro* and animal results; recent toxicology and *in vivo* assessments underscore the urgent need for systematic acute/chronic toxicology, ADME profiling, and dose-finding work before human applications proceed. Likewise, mechanistic pharmacology and well-designed human clinical studies are essentially absent, and applied research on postharvest handling, shelf-life extension, sensory optimization and low-cost value-addition is insufficient.<sup>99</sup> To unlock monkey jackfruit's role in nutrition, sustainability and rural livelihoods, an interdisciplinary program is needed that combines standardized compositional and metabolomic mapping across cultivars and tissues, rigorous toxicology and translational pharmacology, phased human trials for priority indications, applied postharvest and product-development studies tailored to low-resource contexts, and agroecological and socioeconomic research to design farmer-friendly agroforestry and value-chain models.<sup>92,100</sup>

## 10. Consumer studies and toxicity

Comprehensive research on the consumer acceptance, toxicity, and dosage control of *Artocarpus lakoocha* remains limited, yet emerging evidence provides valuable insights into its nutritional and pharmacological potential. Consumer acceptance studies focusing exclusively on *Artocarpus lakoocha* are scarce; however, regional evaluations from India, Thailand, and Bangladesh indicate encouraging results when the fruit pulp is incorporated into blended food products, such as jams, beverages, or fermented foods, improving sensory scores for color, flavor, and mouthfeel. These studies, though preliminary, highlight the potential of *Artocarpus lakoocha* to be developed into functional and value-added foods, provided that standardized sensory evaluation protocols and broad demographic testing are employed.<sup>101,102</sup> Comparable sensory work on related *Artocarpus* species (e.g., *A. heterophyllus* and *A. altilis*) demonstrates that texture, sweetness, and aroma strongly influence consumer preference, suggesting similar drivers for *Artocarpus lakoocha*-based products.

Toxicological studies on *Artocarpus lakoocha* extracts reveal relatively low acute toxicity but highlight the need for systematic dose-response and chronic toxicity evaluations. Recent

investigations on ethanolic and ethyl acetate extracts from *Artocarpus lakoocha* bark and leaves show no mortality or significant biochemical alterations in rodents at doses up to 2000 mg kg<sup>-1</sup>, suggesting a wide safety margin in acute exposure. However, subchronic and genotoxicity data remain scarce, and the pharmacokinetic behaviour of its major bioactive compound, oxyresveratrol, shows moderate oral absorption (~10–15%) and rapid systemic clearance, indicating a need for controlled dosing and formulation optimization to enhance bioavailability.<sup>103</sup> *In vitro* studies demonstrate that oxyresveratrol exerts strong antioxidant and anti-inflammatory activities but may exhibit pro-oxidant or genotoxic effects at high concentrations or under metal-rich conditions, necessitating careful safety evaluations before clinical applications.<sup>68,104</sup>

No standardized human dosage guidelines currently exist for *Artocarpus lakoocha* or its bioactive constituents. Translational studies should therefore determine the no-observed-adverse-effect levels (NOAELs) and lowest-observed-adverse-effect levels (LOAELs) in well-designed animal trials, followed by human pharmacokinetic and tolerability assessments. Establishing controlled dose ranges is vital for potential nutraceutical or therapeutic formulations to ensure safety and reproducibility. Furthermore, standardized extraction methods and compositional characterization are essential to maintain consistency between batches and across studies. Until robust clinical data become available, recommending safe intake levels remains premature.<sup>105</sup> Therefore, it is a potentially nutritious and bioactive food resource and is safe for human consumption.

## 11. Conclusion

Monkey jackfruits, with their impressive nutritional content and adaptability, have emerged as a crucial resource in addressing global challenges. Rich in essential vitamins, minerals, and fiber, they combat malnutrition and diet-related health issues while offering a sustainable solution for food security and climate resilience. As a meat substitute, they support the shift towards plant-based diets, and their historical use in traditional medicine hints at numerous health benefits, from antioxidant to potential cancer-fighting properties. This intersection of nutrition, sustainability, and health underscores their potential as innovative pharmaceuticals and nutraceuticals, requiring continued research, investment, and awareness to fully leverage their benefits in building a more resilient world.

## Declarations

This review paper, titled “Nutritional Value, Phytochemical Richness, Pharmacological Potential, and Culinary Uses of Monkey Jackfruit”, has been prepared solely by the author(s), and all contributors have been appropriately credited.

## Author contributions

Raj Singh: conceptualization, data curation and writing-review and editing; C. Nickhil: investigation, writing and editing;



Shweta: writing and editing; Sankar Chandra Deka: investigation and final checking of the manuscript; R. Nisha: investigation and final checking of the manuscript.

## Conflicts of interest

The authors declare that there is no conflict of interest.

## Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## References

- 1 M. Yadav, K. Srilekha and K. U. Maheswari, Potential health benefit of underutilized fruits: a review, *J. Pharmacogn. Phytochem.*, 2018, 7(5), 1417–1420.
- 2 C. Orwa, A. Mutua, R. Kindt, R. Jamnadass, S. Anthony and L. Arbutus unedo, *Agroforestry Database: A Tree Reference and Selection Guide*, Version 4, 2009.
- 3 R. A. Drew, The application of biotechnology to the conservation and improvement of tropical and subtropical fruit species, *Seed and Plant Genetic Resources Service*, Food and Agriculture Organization of the United Nations, 1997.
- 4 P. Sitorus, J. M. Keliat, V. Asfianti, M. Muhammad and D. Satria, A literature review of *Artocarpus lacucha* focusing on the phytochemical constituents and pharmacological properties of the plant, *Molecules*, 2022, 27(20), 6940.
- 5 E. Bhattacharya, R. Dutta, S. Chakraborty and S. M. Biswas, Phytochemical profiling of *Artocarpus lakoocha* leaf methanol extract and its antioxidant and antimicrobial activities, *Asian Pac. J. Trop. Biomed.*, 2019, 9(11), 484–492.
- 6 D. Pertiwi, R. Hartati, E. Julianti and I. Fidrianny, Antibacterial and antioxidant activities in various parts of *Artocarpus lacucha* Buch.-Ham ethanolic extract, *Biomed. Rep.*, 2024, 20(4), 66.
- 7 R. A. S. N. Ranasinghe, S. D. T. Maduwanthi and R. A. U. J. Marapana, Nutritional and health benefits of jackfruit (*Artocarpus heterophyllus*)—a review, *Int. J. Food Sci.*, 2019, 2019, 4327183.
- 8 K. A. Mehta, Y. C. R. Quek and C. J. Henry, Breadfruit (*Artocarpus altilis*): Processing, nutritional quality and food applications, *Front. Nutr.*, 2023, 10, 1156155.
- 9 R. O. Adeleke and O. A. Abiodun, Nutritional composition of breadnut seeds (*Artocarpus camansi*), *Afr. J. Agric. Res.*, 2010, 5(11), 1273–1276.
- 10 S. M. Islam, F. M. Hasan, M. Ali, M. Robbani and T. M. Hossain, Socioeconomic potential of Monkey Jack: an underutilized fruit in Bangladesh, *Int. J. Innovative Res.*, 2016, 1(1), 40–44.
- 11 D. H. Dwivedi, V. Mishra, N. Singh and S. K. Dwivedi, Genetic Variability Studies in Barhal (*Artocarpus lakoocha* Roxb.), in *Central Uttar Pradesh. In XXVIII International Horticultural Congress on Science and Horticulture for People (IHC2010): III International Symposium on 918*, 2010, vol. 22, pp. 419–425.
- 12 S. K. Bishnoi, J. B. Tomar, K. K. Saini and P. Kumar, *Artocarpus lakoocha*: a potential tree species for forest and agricultural diversification in Jharkhand, India, in *3rd National Symposium on Agricultural Production and Protection in Context of Climate Change*, 2012, pp. 3–5.
- 13 S. R. Krishnamurthy and P. Sarala, Phytochemical studies of *Artocarpus gomezianus* Wall. ex Trecul. var. *lakoocha* Roxb. fruits collected from various altitudes of Central Western Ghats, *Indian J. Nat. Prod. Resour.*, 2013, 4(4), 398–411.
- 14 M. F. Hossain, M. A. Islam, S. Akhtar and S. M. Numan, Nutritional value and medicinal uses of Monkey Jack fruit (*Artocarpus lakoocha*), *Int. Res. J. Biol. Sci.*, 2016, 5(1), 60–63.
- 15 P. Gautam and R. Patel, *Artocarpus lakoocha* Roxb.: An overview, *Eur. J. Complementary Altern. Med.*, 2014, 1(1), 10–14.
- 16 A. K. Gupta, U. Pathak, M. Medhi, A. Mastinu, M. S. Sikarwar and P. Mishra, Botanical, chemical and pharmacological properties of *Artocarpus lakoocha* (monkey fruit): a review, *Agric. Rev.*, 2020, 41(4), 305–316.
- 17 R. Singh, C. Nickhil, R. Nisha, K. Upendar, B. Jithender and S. C. Deka, A comprehensive review of advanced deep learning approaches for food freshness detection, *Food Eng. Rev.*, 2024, 1–34.
- 18 A. Bellavia, S. C. Larsson, M. Bottai, A. Wolk and N. Orsini, Fruit and vegetable consumption and all-cause mortality: a dose–response analysis, *Am. J. Clin. Nutr.*, 2013, 98(2), 454–459.
- 19 T. S. Conner, K. L. Brookie, A. C. Carr, L. A. Mainvil and M. C. Vissers, Effect of fruit and vegetable consumption on psychological well-being in young adults, *PLoS One*, 2017, 12(2), e0171206.
- 20 O. Oyeboode, V. Gordon-Dseagu, A. Walker and J. S. Mindell, Fruit and vegetable consumption and all-cause, cancer and CVD mortality: analysis of Health Survey for England data, *J. Epidemiol. Community Health*, 2014, 68(9), 856–862.
- 21 P. Boffetta, E. Couto, J. Wichmann, P. Ferrari, D. Trichopoulos, H. B. Bueno-de-Mesquita, *et al.*, Fruit and vegetable intake and cancer risk in EPIC, *J. Natl. Cancer Inst.*, 2010, 102(8), 529–537.
- 22 B. J. Rolls, J. A. Ello-Martin and B. C. Tohill, What can intervention studies tell us about the relationship between fruit and vegetable consumption and weight management?, *Nutr. Rev.*, 2004, 62(1), 1–17.
- 23 K. He, F. B. Hu, G. A. Colditz, J. E. Manson, W. C. Willett and S. Liu, Changes in intake of fruits and vegetables and risk of obesity and weight gain among middle-aged women, *Int. J. Obes.*, 2004, 28(12), 1569–1574.
- 24 S. Jahan, T. Gosh, M. Begum and B. K. Saha, Nutritional profile of some tropical fruits in Bangladesh, *Bangladesh J. Med. Sci.*, 2011, 10(2), 95–100.
- 25 P. Sarala and S. R. Krishnamurthy, Monkey jack: Underutilized edible medicinal plant, nutritional attributes and traditional foods of Western Ghats,



- Karnataka, India, *Indian Journal of Traditional Knowledge*, 2014, **13**(3), 508–518.
- 26 A. Kumar, M. Premoli, F. Aria, S. A. Bonini, G. Maccarinelli, A. Gianoncelli, *et al.*, Cannabimimetic plants: New cannabinoidergic modulators?, *Planta*, 2019, **249**, 1681–1694.
- 27 A. Mastinu, S. A. Bonini, W. Rungratanawanich, F. Aria, M. Marziano, G. Maccarinelli, *et al.*, Gamma-oryzanol prevents LPS-induced brain inflammation and cognitive impairment in mice, *Nutrients*, 2019, **11**(4), 728.
- 28 W. S. D. Yamika, N. Aini and B. Waluyo, *Physalis peruviana* L. growth, yield and phytochemical content: a review, *Agric. Rev.*, 2019, **40**(4), 324–328.
- 29 S. Khalid, N. Khalid, R. S. Khan, H. Ahmed and A. Ahmad, A review on chemistry and pharmacology of Ajwa date fruit and pit, *Trends Food Sci. Technol.*, 2017, **63**, 60–69.
- 30 R. Singh, R. Nisha, R. Naik, K. Upendar, C. Nickhil and S. C. Deka, Sensor fusion techniques in deep learning for multimodal fruit and vegetable quality assessment: a comprehensive review, *J. Food Meas. Charact.*, 2024, **18**(9), 8088–8109.
- 31 M. M. Hossain, M. A. Rahim and M. R. Haque, Biochemical properties of some important underutilized minor fruits, *J. Agric. Food Res.*, 2021, **5**, 100148.
- 32 T. R. Prashith Kekuda, H. L. Raghavendra, N. Mallikarjun, T. M. Venugopal and H. S. Anil Kumar, Elemental composition, anticariogenic, pancreatic lipase inhibitory and cytotoxic activity of *Artocarpus lakoocha* Roxb pericarp, *Int. J. Drug Dev. Res.*, 2012, **4**(1), 330–336.
- 33 C. Nickhil, Development of Mechanical Tool to Separate Bulbs from Whole Jackfruit, Master's thesis, University of Agricultural Sciences, Bengaluru, GKVK, 2015.
- 34 A. Hari, K. G. Revikumar and D. Divya, *Artocarpus*: A review of its phytochemistry and pharmacology, *Journal of Pharma Search*, 2014, **9**(1), 7–12.
- 35 T. Chuanasa, J. Phromjai, V. Lipipun, K. Likhitwitayawuid, M. Suzuki, P. Pramyothin, *et al.*, Anti-HSV-1 activity of oxresveratrol and therapeutic efficacy in mice, *Antiviral Res.*, 2008, **80**(1), 62–70.
- 36 A. Puntumchai, P. Kittakoop, S. Rajviroongit, S. Vimuttipong, K. Likhitwitayawuid and Y. Thebtaranonth, Lakoochins A and B: New antimycobacterial stilbene derivatives from *Artocarpus lakoocha*, *J. Nat. Prod.*, 2004, **67**(3), 485–486.
- 37 H. H. Ko, Y. T. Tsai, M. H. Yen, C. C. Lin, C. J. Liang, T. H. Yang, *et al.*, Norartocarpetin from *Artocarpus communis* acts as a melanogenesis inhibitor without cytotoxicity, *BMC Complementary Altern. Med.*, 2013, **13**, 348.
- 38 M. H. Tsai, J. F. Liu, Y. C. Chiang, S. C. S. Hu, L. F. Hsu, Y. C. Lin, *et al.*, Artocarpin, an isoprenyl flavonoid, induces p53-dependent or independent apoptosis via ROS-mediated MAPKs and Akt activation in non-small cell lung cancer cells, *Oncotarget*, 2017, **8**(17), 28342.
- 39 M. S. Sikarwar, B. J. Hui, K. Subramaniam, B. D. Valeisamy, L. K. Yean and K. Balaji, A review on *Artocarpus altilis* (Parkinson) Fosberg (breadfruit), *J. Appl. Pharm. Sci.*, 2014, **4**(8), 91–97.
- 40 M. S. Kumar, M. R. Kumar, A. C. Bharath, H. V. Kumar, T. P. Kekuda, K. C. Nandini, *et al.*, Screening of selected biological activities of *Artocarpus lakoocha* Roxb fruit pericarp, *J. Basic Clin. Pharm.*, 2010, **1**(4), 239.
- 41 K. Krishnan, L. E. Mathew, N. R. Vijayalakshmi and A. Helen, Anti-inflammatory potential of  $\beta$ -amyryn, a triterpenoid isolated from *Costus igneus*, *Inflammopharmacology*, 2014, **22**, 373–385.
- 42 N. Han and M. Bakovic, Biologically active triterpenoids and their cardioprotective and anti-inflammatory effects, *J. Bioanal. Biomed.*, 2015, **S12**, 005.
- 43 T. Z. Shoshe, *Quantification of total phenol, in vitro investigation of antioxidant activity, cytotoxic effect and phytochemical screening of the leaf extract of Artocarpus lacucha*, 2017.
- 44 M. T. Islam Shajib, M. E. M. Ferdous, A. Naz Shah, U. Habiba and S. N. Islam, Monkey Jack (*Artocarpus lakoocha*), in *Handbook of Phytonutrients in Indigenous Fruits and Vegetables*, 2022, pp. 311–322.
- 45 S. Chaurasia and A. Pandey, Phytochemistry and pharmacology of genus *Artocarpus*, *Russ. J. Bioorg. Chem.*, 2023, **49**(3), 481–514.
- 46 S. Pandey and A. Poonia, Monkey jackfruit (*Artocarpus lakoocha* Roxb.): The lesser-known fruit, *Indian Food Ind.*, 2021, **6**, 38–45.
- 47 M. Vanajakshi, J. H. Virupaksha and S. Maria, A review on pharmacological actions of *Artocarpus lacoocha* Roxb, *Res. J. Pharmacol. Pharmacodyn.*, 2016, **8**(4), 181–184.
- 48 S. Boonyaketguson, V. Rukachaisirikul, S. Phongpaichit and K. Trisuwan, Deoxybenzoin and flavan derivatives from twigs of *Artocarpus lakoocha*, *Phytochem. Lett.*, 2019, **31**, 96–100.
- 49 T. Ahamad, M. A. Khan, W. A. Ansari and M. F. Khan, Antioxidant and anticancer activities of selected Indian medicinal plants viz., *Artocarpus lakoocha*, *Kigelia pinnata*, and *Amaranthus viridis*, *Era's J. of Med. Res.*, 2020, **7**(1), 46–51.
- 50 S. Krupa, C. S. Karigar and K. S. Murthy, Pharmacology and phytochemistry of *Artocarpus* family: A review, *Indo Global J. Pharm. Sci.*, 2020, **10**(3), 48–55.
- 51 M. Saleem, A. Asif, M. F. Akhtar and A. Saleem, Hepatoprotective potential and chemical characterization of *Artocarpus lakoocha* fruit extract, *Bangladesh J. Pharmacol.*, 2018, **13**(1), 90–97.
- 52 P. Hankittichai, P. Buacheen, P. Pitchakarn, M. Na Takuathung, N. Wikan, D. R. Smith, *et al.*, *Artocarpus lakoocha* extract inhibits LPS-induced inflammatory response in RAW 264.7 macrophage cells, *Int. J. Mol. Sci.*, 2020, **21**(4), 1355.
- 53 A. K. Buddhisuharto, H. Pramastya, M. Insanu and I. Fidrianny, Updated review of phytochemicals and pharmacological activities of *Artocarpus* genus, *Biointerface Res. Appl. Chem.*, 2021, **11**(6), 14898–14905.
- 54 F. Ferrari, I. Messana and M. D. C. M. de Araujo, Structures of three new flavone derivatives from *Brosimopsis oblongifolia*, *Planta Med.*, 1989, **55**(1), 70–72.



- 55 Z. P. Zheng, K. W. Cheng, J. T. K. To, H. Li and M. Wang, Isolation of tyrosinase inhibitors from *Artocarpus heterophyllus* and use of its extract as antibrowning agent, *Mol. Nutr. Food Res.*, 2008, **52**(12), 1530–1538.
- 56 Y. Hano, Y. Yamagami, M. Kobayashi, R. Isohata and T. Nomura, Artonins E and F, two new prenylflavones from the bark of *Artocarpus communis* Forst, *Heterocycles*, 1990, **31**(5), 877–882.
- 57 E. T. Arung, K. Shimizu and R. Kondo, Inhibitory effect of artocarpanone from *Artocarpus heterophyllus* on melanin biosynthesis, *Biol. Pharm. Bull.*, 2006, **29**(9), 1966–1969.
- 58 T. Fujimoto, Y. Hano, T. Nomura and J. Uzawa, Components of root bark of *Cudrania tricuspidata*: two new isoprenylated flavones, *Planta Med.*, 1984, **50**(2), 161–163.
- 59 S. Maneechai, W. De-Eknamkul, K. Umehara, H. Noguchi and K. Likhitwitayawuid, Flavonoid and stilbenoid production in callus cultures of *Artocarpus lakoocha*, *Phytochemistry*, 2012, **81**, 42–49.
- 60 U. Neog, P. Dhar, T. Kumari, C. Nickhil, S. C. Deka and R. Pandiselvam, Optimization of microwave-assisted extraction of phytochemicals from norabogori fruit (*Prunus persica* L. Batsch) and application in fruit leather, *Biomass Convers. Biorefin.*, 2023, 1–15.
- 61 B. D. Maibam, S. Chakraborty, C. Nickhil and S. C. Deka, Effect of *Euryale ferox* seed shell extract on starch digestibility and predicted glycemic index of wheat bread, *Int. J. Biol. Macromol.*, 2023, **226**, 1066–1078.
- 62 B. Jithender, D. M. Vyas, C. Nickhil and P. J. Rathod, Development of juice extraction process for pomegranate (*Punica granatum*): evaluation and physicochemical aspects, *Indian J. Agric. Sci.*, 2021, **91**(6), 936–938.
- 63 K. Saesue, P. Thanomrak, W. Prompan, W. Punan, N. Khorana, W. Juprasert, *et al.*, Development of a ready-to-use oxyresveratrol-enriched extract from *Artocarpus lakoocha* Roxb. using greener and deep eutectic solvents for whitening applications, *Cosmetics*, 2024, **11**(2), 58.
- 64 K. Likhitwitayawuid, Oxyresveratrol: Sources, production, biological activities, pharmacokinetics and delivery systems, *Molecules*, 2021, **26**(14), 4212.
- 65 J. Zhang, S. Liu, W. Guo, Y. Huang and N. Li, Oxyresveratrol suppressed melanogenesis, dendrite formation and melanosome transport in melanocytes via regulation of the MC1R/cAMP/MITF pathway, *Sci. Rep.*, 2025, **15**(1), 20400.
- 66 T. Rodboon, P. Puchadapirom, S. Okada and P. Suwannalert, Oxyresveratrol from *Artocarpus lakoocha* Roxb. inhibits melanogenesis in B16 melanoma cells by regulating cellular oxidants, *Walailak J. Sci. Technol.*, 2016, **13**(4), 261–270.
- 67 D. L. Lucetti, E. C. Lucetti, M. A. M. Bandeira, H. N. Veras, A. H. Silva, L. K. A. Leal, *et al.*, Anti-inflammatory effects and mechanism of action of lupeol acetate isolated from *Himatanthus drasticus*, *J. Inflammation*, 2010, **7**(1), 60.
- 68 D. Junsang, T. Anukunwithaya, P. Songvut, B. Sritularak, K. Likhitwitayawuid and P. Khemawoot, Comparative pharmacokinetics of oxyresveratrol alone and in combination with piperine as a bioenhancer in rats, *BMC Complementary Altern. Med.*, 2019, **19**(1), 235.
- 69 K. Malaniyom, P. Ratanachamnong, P. Namchaiw, U. Namdaung, S. Suksamrarn and Y. Jaisin, Suppression of inflammatory response by oxyresveratrol from *Artocarpus lakoocha* root bark against UV-B-induced keratinocytes, *Heliyon*, 2024, **10**(20), e38962.
- 70 Y. Cui, F. Li, X. Zhu, J. Xu, A. Muhammad, Y. Chen, *et al.*, Alfalfa saponins inhibit oxidative stress-induced apoptosis via MAPK pathway, *Redox Rep.*, 2022, **27**(1), 1–8.
- 71 W. O. M. Hasriadi, P. Lapphanichayakool and N. Limpeanchob, Neuroprotective effect of *Artocarpus lakoocha* extract and oxyresveratrol against hydrogen peroxide-induced toxicity in SH-SY5Y cells, *Int. J. Pharm. Sci.*, 2017, **9**, 229–233.
- 72 M. L. Nesa, S. Munira, A. S. Bristy, M. M. Islam and H. Chayan, Cytotoxic, anti-inflammatory, analgesic, CNS depressant and anti-diarrhoeal activities of methanolic extract of *Artocarpus lakoocha* leaves, *World. J. Pharm. Sci.*, 2015, 167–174.
- 73 S. Seiki and W. H. Frishman, Pharmacologic inhibition of squalene synthase and other downstream enzymes of the cholesterol synthesis pathway: a new therapeutic approach to treatment of hypercholesterolemia, *Cardiol. Rev.*, 2009, **17**(2), 70–76.
- 74 N. Nazliniwaty, O. A. Hanafiah, D. Pertiwi, M. Muhammad and D. Satria, Ethanolic extract of *Artocarpus lacucha* and *Anredera cordifolia* leaves improves wound healing in NIH-3T3 cells, *Open Access Maced. J. Med. Sci.*, 2022, **10**(A), 807–811.
- 75 M. G. Uddin, M. M. Islam, A. Ahmed, S. A. Siddiqui and A. Debnath, Evaluation of anthelmintic, antioxidant and anti-inflammatory potential of methanolic extract of *Artocarpus lacucha* leaves, *Discovery Phytomed.*, 2020, **7**(1), 27–32.
- 76 P. A. Z. Hasibuan, Antioxidant activity of n-hexane, ethyl acetate, and ethanol extracts from *Artocarpus lacucha* leaves using DPPH method, *Indones. J. Pharm. Clin. Res.*, 2018, **1**(2), 41–47.
- 77 H. Raghavendra, N. Mallikarjun, T. Venugopal and H. S. Anil Kumar, Elemental composition, anticariogenic, pancreatic lipase inhibitory and cytotoxic activity of *Artocarpus lakoocha* Roxb pericarp, *Int. J. Drug Dev. Res.*, 2012, **4**(1), 330–336.
- 78 S. Islam, M. S. Shajib, R. B. Rashid, M. F. Khan, M. A. Al-Mansur, B. K. Datta, *et al.*, Antinociceptive activities of *Artocarpus lacucha* and its isolated phenolic compound catechin in mice, *BMC Complementary Altern. Med.*, 2019, **19**, 1–13.
- 79 N. Preyavichyapugdee, M. Sangfuang, S. Chaiyapum, S. Sriburin, Y. Pootaeng-on, P. Chusongsang, *et al.*, Schistosomicidal activity of crude extract of *Artocarpus lakoocha*, *Southeast Asian J. Trop. Med. Public Health*, 2016, **47**, 1–15.
- 80 S. Tarbiat, In vitro antioxidant, anti-inflammatory and lipid-lowering activities of *Artocarpus lakoocha* fruit extract



- and its implication on treatment of dyslipidemia, *Int. J. Pharm. Sci. Rev. Res.*, 2018, **52**(2), 75–82.
- 81 W. Iskandar, O. W. K. Handayani and W. H. Cahyati, Analysis of family income factors on diarrhea incidence through behavior in Tapalang, *Public Health Perspect. J.*, 2019, **4**(3), 206–213.
- 82 P. E. Thomas and B. Dharmapalan, Value added products from jackfruit (*Artocarpus heterophyllus*) fruit, *Acta Sci. Nutr. Health*, 2020, **4**(2), 105–110.
- 83 B. Q. Hoang, H. T. Nguyen and D. N. T. Duong, Development of lactic acid fermentation of jackfruit (*Artocarpus heterophyllus*) seed drink and its physicochemical and sensory properties, *J. Food Sci. Technol.*, 2024, **61**(6), 1180–1187.
- 84 S. Mohammed, P. K. Dubey, A. A. Mishra and S. Rahman, Valorisation of jackfruit seed flour in extrusion and bakery products: A review, *Food Sci. Biotechnol.*, 2024, **33**(14), 3167–3180.
- 85 S. Maneechai, K. Likhitwitayawuid, B. Sritularak, C. Palanuvej, N. Ruangrunsi and P. Sirisa-Ard, Quantitative analysis of oxyresveratrol in *Artocarpus lakoocha* and 'Puag-Haad', *Med. Princ. Pract.*, 2009, **18**(3), 223–227.
- 86 M. T. Islam, S. K. Barman, M. Saha, M. S. Islam, and M. I. Sahid, *Phytochemical Screening and Evaluation of Antioxidant Potential of Ethanolic Extract of Artocarpus lacucha Leaves In-vitro*, 2024, pp. 156–170.
- 87 S. Radapong, K. Chan, S. D. Sarker and K. J. Ritchie, Oxyresveratrol modulates genes associated with apoptosis, cell cycle control and DNA repair in MCF-7 cells, *Front. Pharmacol.*, 2021, **12**, 694562.
- 88 L. Xiang, Q. Li, Z. Guan, G. Wang, X. Yu, X. Zhang, *et al.*, Oxyresveratrol as a novel ferroptosis inducer exhibits anticancer activity against breast cancer via the EGFR/PI3K/AKT/GPX4 signalling axis, *Front. Pharmacol.*, 2025, **15**, 1527286.
- 89 M. J. Haruni, M. S. Parvin, S. Munira and M. E. Islam, Phytochemical profile and evaluation of antioxidant, anticancer, and antimicrobial potential of *Artocarpus lacucha* bark extracts: in vitro and in silico studies, *J. Food Biochem.*, 2025, **2025**(1), 8016598.
- 90 A. Wiater, R. Paduch, S. Trojnar, A. Choma, M. Pleszczyńska, P. Adamczyk, *et al.*, Effect of water-soluble polysaccharide from jackfruit (*Artocarpus heterophyllus* Lam.) on human colon carcinoma cells cultured in vitro, *Plants*, 2020, **9**(1), 103.
- 91 N. Wongkattiya, P. Sanguansermisri, I. Fraser, A. Shuayprom, S. Pongmuangmul, R. Koranee, *et al.*, Oxyresveratrol derived from *Artocarpus lakoocha* extract represents a potential natural agent for treating acne and skin inflammation, *J. Appl. Pharm. Sci.*, 2025, **15**(11), 50–57.
- 92 T. Banerjee, A. Sarkar, A. De, S. Sar, A. Mandal, P. Panda, *et al.*, Toxicological evaluation of *Artocarpus lacucha* ethyl acetate extract: in vitro and in vivo assessment, *J. Ethnopharmacol.*, 2025, **347**, 119804.
- 93 A. T. Shafaq, H. M. Ishaq and M. Shahzad, Pharmacological effects of *Artocarpus lakoocha* methanol extract on inhibition of squalene synthase and other downstream enzymes of the cholesterol synthesis pathway, *Pharm. Biol.*, 2022, **60**(1), 840–845.
- 94 S. Singhatong, D. Leelarungrayub and C. Chaiyasut, Antioxidant and toxicity activities of *Artocarpus lakoocha* Roxb. heartwood extract, *J. Med. Plants Res.*, 2010, **4**(10), 947–953.
- 95 L. Xu, C. Liu, W. Xiang, H. Chen, X. Qin and X. Huang, Advances in the study of oxyresveratrol, *Int. J. Pharmacol.*, 2014, **10**(1), 44–54.
- 96 P. Jantaharn, T. Sornda, N. Limpeanchob, T. Pitaksuteepong, W. Tojaroen, K. Chootip, *et al.*, Neuroprotective activity of *Artocarpus lacucha* extract in SH-SY5Y cells and lifespan extension in *C. elegans*, *Ind. Crops Prod.*, 2025, **237**, 122203.
- 97 S. Zeng, Y. Chen, C. Wei, L. Tan, C. Li, Y. Zhang, *et al.*, Protective effects of polysaccharide from *Artocarpus heterophyllus* Lam. (jackfruit) pulp on non-alcoholic fatty liver disease in high-fat diet rats via PPAR and AMPK signalling pathways, *J. Funct. Foods*, 2022, **95**, 105195.
- 98 D. K. Joung, S. H. Mun, S. H. Choi, O. H. Kang, S. B. Kim, Y. S. Lee, *et al.*, Antibacterial activity of oxyresveratrol against methicillin-resistant *Staphylococcus aureus* and its mechanism, *Exp. Ther. Med.*, 2016, **12**(3), 1579–1584.
- 99 M. N. Hossain, M. S. Hasan, M. I. Keya, M. G. Rabby, R. Tarannum, R. Parvin, *et al.*, Effects of *Artocarpus lacucha* fruit pulp extracts on physicochemical characteristics and microbial stability of raw chicken meatballs during refrigeration, *Appl. Food Res.*, 2025, 101253.
- 100 H. J. Borah, R. Singhal and S. Hazarika, *Artocarpus lakoocha* Roxb.: An untapped bioresource of resveratrol from NE India and its antioxidant activity, *Ind. Crops Prod.*, 2017, **95**, 75–82.
- 101 S. K. Bishnoi, R. Shinde and P. K. Sarkar, *Monkey Jack (Artocarpus Lakoocha Roxb.): Hope for Sustaining Livelihood*, 2017.
- 102 R. Verma, R. Sharma and S. Bhamra, Standardization, preparation and evaluation of dheu (*Artocarpus lakoocha*)-based toffee, *Himachal J. Agric. Res.*, 2021, 228–234.
- 103 D. Satria, P. Sitorus, A. Dalimunthe, S. B. Waruwu and V. Asfianti, Oral acute toxicity study of ethanol extract of Mobe leaves (*Artocarpus lacucha* Buch.-Ham) in Wistar rats, *Pharmacia*, 2024, **71**, 1–8.
- 104 S. Radapong, S. D. Sarker and K. J. Ritchie, Oxyresveratrol possesses DNA-damaging activity, *Molecules*, 2020, **25**(11), 2577.
- 105 N. Alam, H. Najnin, M. Islam, S. Shakya, I. M. Khan and R. Zaidi, Biochemical and histopathological analysis after sub-chronic administration of oxyresveratrol in Wistar rats, *Drug Chem. Toxicol.*, 2023, **46**(1), 166–175.

