



Cite this: DOI: 10.1039/d5fb00789e

Enzyme-assisted extraction and characterization of red roselle calyx (*Hibiscus sabdariffa* L.) bioactives and their application in ice cream matrices

A. Priyanka,^a B. Nila Shireen,^a S. Ganga Kishore,^b Sudha Paramadhas,^{*a} R. Rahul,^{©c} Madhuresh Dwivedi,^{©b} T. Pandiarajan,^a R. Divyabharathi^a and Nongmaithem Seema Chanu^b

Roselle calyx (*Hibiscus sabdariffa* Linn.) is widely recognized for its rich phenolic and anthocyanin contents, which contribute to its antioxidant properties, colour, and health-promoting benefits. However, conventional extraction methods often result in low yields and compound degradation. Therefore, this study explored enzyme-assisted extraction using amylase, pectinase, and cellulase individually to enhance the recovery of bioactive compounds from roselle calyces under varying enzyme concentrations and temperatures. The optimized condition obtained from response surface methodology revealed that cellulase (10%, 60 °C) produced the highest total phenolic (55.34 mg GAE/100 g) and flavonoid (193.05 mg CE/100 g) contents, pectinase (15%, 50 °C) yielded the highest anthocyanin content (127.79 mg/100 g), and amylase (3%, 60 °C) achieved the highest antioxidant activity (39.14%). The incorporation of enzyme-optimized roselle extracts into ice cream formulations improved the colour intensity, texture, and overall sensory appeal. Cellulase provided the most vibrant colour and dense structure, resulting in a smoother and more aerated product with balanced sweetness and a moderately light texture. HPLC analysis of cellulase-assisted roselle calyx extract showed peaks corresponding to quercetin, with a high relative concentration of 52.87%, indicating the efficient enzymatic release of this key flavonoid without any structural degradation. FTIR spectroscopic analysis confirmed the retention of key phenolic and carbohydrate-based functional groups. SEM analysis revealed that fine roselle fragments were uniformly embedded within the ice cream matrix, supporting a smoother texture and reduced ice crystal formation. Overall, enzyme-assisted extraction emerges as a sustainable and effective approach for producing roselle-based natural colourants.

Received 30th October 2025
Accepted 19th January 2026

DOI: 10.1039/d5fb00789e

rsc.li/susfoodtech

Sustainability spotlight

This study highlights the sustainable potential of enzyme-assisted extraction (EAE) as an eco-friendly and efficient method for obtaining natural colorants and bioactive compounds from roselle (*Hibiscus sabdariffa* Linn.) calyces. By employing enzymes such as amylase, pectinase, and cellulase, the process significantly enhanced the extraction of anthocyanins, phenolics, and flavonoids, thereby improving the antioxidant activity and pigment yield. The optimized conditions, particularly with cellulase at 10% concentration and 60 °C, yielded the highest levels of total phenolic and flavonoid contents while maintaining compound stability. The enzyme-infused roselle bio-extract was further incorporated into ice cream formulations, resulting in improved colour intensity, texture, and sensory properties. Advanced characterization techniques such as FTIR spectroscopy and SEM confirmed the presence of polyphenols, flavonoids, and proteinaceous compounds, as well as enhanced microstructural properties that contributed to better texture and water retention. Overall, this study demonstrates that enzyme-assisted extraction is a sustainable and effective alternative to conventional methods for producing natural, stable, and functional food colorants, supporting their potential application in the food and nutraceutical industries as substitutes for synthetic additives.

1 Introduction

The food industry across the world witnessed a rising shift in consumer needs towards natural and organic ingredients due to increasing health concerns and environmental hazards caused by synthetic colorants.^{1,2} Synthetic dyes like tartrazine, allura red, and sunset yellow, though stable and effective, have been reported to induce allergic reactions, hyperactivity, and carcinogenic effects.³ Consequently, the demand for plant-based bio-

^aDepartment of Food Process Engineering, Agricultural Engineering College & Research Institute, Tamil Nadu Agricultural University, Coimbatore-641 003, Tamil Nadu, India. E-mail: sudha.p@tnau.ac.in

^bDepartment of Food Process Engineering, National Institute of Technology, Rourkela-769 008, Odisha, India. E-mail: madhureshd@gmail.com

^cDepartment of Food Technology, Dhanalakshmi Srinivasan College of Engineering, Coimbatore-641 105, Tamil Nadu, India



colorants witnessed active acceleration with increased focus on extraction efficiency, stability, and compatibility in food matrices.⁴ Among the plant-based alternatives, *Hibiscus sabdariffa* Linn., commonly known as roselle, gained significant attention due to its vibrant red calyces, which are rich in phenols.⁵ Anthocyanins not only impart color but also provide antioxidant, antimicrobial, and antihypertensive properties, making them ideal functional ingredients in food and nutraceutical formulations.⁶

Roselle is widely cultivated in the tropical regions of Africa, Asia, and the Caribbean. The colour of the calyces determines the bioactive components present in them, especially their flavonoid contents.^{7,8} The calyces of *Hibiscus sabdariffa* (roselle) are complex matrices composed of not only basic nutrients (such as modest amounts of protein, fibre, carbohydrates, minerals, and organic acids) but also a rich suite of bioactive phytochemicals. Among the phenolic-type phytochemicals, calyces are especially notable for their high content of anthocyanin pigments, which confer the characteristic deep red colour of red-calyx varieties. In addition, a variety of other phenolic acids (e.g., chlorogenic, neochlorogenic, caffeic, gallic, and protocatechuic acids) and flavonoids (e.g., rutin, quercetin derivatives, and other glycosides) have been identified. Roselle calyces also contain organic acids (citric, malic, and hibiscus acid), some vitamins (notably ascorbic acid), and diet-relevant constituents such as dietary fibers. Importantly, many of these bioactive polyphenols are not freely extractable but are present in a “bound” form, often interacting with the plant cell wall matrix (polysaccharides, fibers, proteins, and other macromolecules) or forming complexes that affect their solubility, extractability, and stability. The actual yield and composition of bioactive compounds from calyces depend heavily on the extraction method, the solvent used, and the processing conditions, whether the aim is to recover pigments (anthocyanins), soluble phenolics, or more tightly associated bound phenolics. This bound fraction is crucial: it represents a reservoir of bioactive potential that can be released (or stabilized) depending on processing and contributes to the overall functional value of roselle calyces for antioxidant, anti-hyperglycemic, and other health-related effects.^{5,6,8,9}

The principal anthocyanins, delphinidin-3-sambubioside and cyanidin-3-sambubioside, are found in the calyces that are responsible for the deep red color.⁹ Moreover, the calyces are rich in organic acids, flavonoids, ascorbic acid, and polysaccharides, contributing to corresponding sensory and therapeutic qualities.¹⁰ Despite their high pigment content, the large-scale utilization of roselle calyces as bio-colorants is limited by challenges in extraction efficiency and pigment stability during processing and storage.¹¹ Anthocyanins are prone to thermal degradation, oxidative reactions, and enzymatic browning, which can result in potential color loss, specifically during conventional drying and solvent extraction.¹² These bioactive compounds exist in bound forms, either conjugated to cell wall components or trapped within the matrix, limiting their extractability using conventional solvent methods. This rationale supports the use of enzyme-assisted extraction, as specific enzymes (e.g., cellulases and pectinases) can hydrolyse the cell

wall or matrix components, releasing these bound bioactives more efficiently.^{13–15}

Conventional solvent-based anthocyanin extraction methods, such as maceration and reflux, are energy-demanding and time-consuming, and may cause degradation of thermolabile compounds.¹⁶ These barriers have shifted the research focus to explore eco-friendly and efficient alternatives for pigment recovery from plant matrices. Traditional methods of anthocyanin extraction, mostly solvent-based techniques using ethanol,¹⁷ methanol,¹⁸ or acidified water,¹⁹ have been widely applied for retaining anthocyanins from roselle. These methods are often time-consuming, require high solvent usage, and may cause partial degradation of thermolabile pigments.^{20,21} Recent advancements in green extraction technologies such as ultrasound-assisted,²² microwave-assisted,²³ and enzyme-assisted extraction²⁴ have shown improved pigment recovery and stability with lower environmental impacts. Among these, enzyme-assisted extraction (EAE) has emerged as a promising technique due to its eco-friendly²⁵ approach in specificity and the ability to disrupt plant cell walls for enhanced release of intracellular pigments.²⁶ Enzymes like pectinase,¹³ cellulase,¹⁴ and hemicellulase¹⁵ can synergistically act and hydrolyse the polysaccharides of the calyces, which may lead to a higher pigment yield.¹³

Furthermore, EAE is considered a sustainable alternative due to its lower energy consumption,²⁷ absence of toxic residues²⁸ and the ability to maintain anthocyanin stability.²⁹ The efficiency of EAE is influenced by extraction parameters including enzyme concentration, temperature, pH, and incubation time.³⁰ Anthocyanins derived from roselle have proven applications in various food systems such as beverages, yoghurts, bakery products, and confectionery items, improving both visual appeal and antioxidant potential.³¹ However, the stability during storage and thermal processing is a critical concern.¹¹ To overcome these challenges, the enzyme-assisted extraction technique could significantly enhance the pigment yield, stability, and functionality while maintaining product safety and sustainability.⁵ Therefore, the present study aims to evaluate the effect of enzyme-assisted extraction parameters on the recovery of bio-colorants from dried roselle (*Hibiscus sabdariffa* Linn.) calyces and their application in food systems. The research specifically focuses on optimizing the enzyme type, concentration, and process temperature to achieve the maximum pigment yield and stability, thereby contributing to the development of natural, safe, and sustainable colorant alternatives for the food industry.

2 Materials and methods

2.1 Chemicals, reagents, and equipment

All the reagents and solvents used were of analytical grade. The chemicals employed for extraction and analytical assays included hydrochloric acid (HCl), Folin–Ciocalteu reagent, sodium carbonate (20% Na₂CO₃), 95% methanol, pyrocatechol (stock solution), aluminium chloride (10% AlCl₃), sodium hydroxide (1 M NaOH), sodium nitrate (5% NaNO₂), and 2,2-diphenyl-1-picrylhydrazyl (DPPH). Enzymes such as amylase,



cellulase, and pectinase were procured in powder form from HiMedia, Mumbai, India. The instruments used for extraction and analysis included a UV-visible spectrophotometer (UV-1800, Shimadzu, Japan), a digital pH meter, a Lovibond tintometer (Lovibond, LC100, The Tintometer Ltd, UK), a high-performance liquid chromatograph (HPLC; LC-6AD, Shimadzu, Japan), a color extraction unit, and an analytical balance.

2.2. Preparation of the roselle bio-extract from the calyx portion

Fresh roselle (*Hibiscus sabdariffa* L.) calyces at 35 days post-bloom (Wong *et al.*, 2003) were harvested from a farm in Pothur village (13.1602° N, 80.1512° E), near Chennai, Tamil Nadu, India. The calyces were manually separated from the buds using a hand tool, thoroughly washed to remove adhering soil and impurities, and tray-dried at 60 °C until the moisture content decreased from 86% to 11%.³² Approximately 5 kg of fresh calyces yielded about 1.5 kg of dried material after 48 h of drying. The tray-drying process took almost 2 days to completely remove the moisture content present in the calyces. The initial moisture content of the sample was 86%, and it was reduced to 11%.

2.3. Experimental design for enzyme-assisted extraction

The experimental plan was developed by using varying combinations of independent variables: individual enzyme type (amylase, cellulase, and pectinase), enzyme concentration (1%, 5%, and 10% w/v), and temperature (50 °C, 60 °C, and 70 °C). Consequently, the dependent variables chosen were total phenolic content (TPC), total flavonoid content (TFC), total anthocyanin content (TAC), antioxidant activity (AOX), pH, and color parameters (L^* , a^* , b^* , and hue angle h°) for analyzing the qualitative properties preserved across all experiments. The final best condition was obtained from the dependent variables of trial combinations. The experiments were carried out under ambient laboratory conditions. For each run, 500 g of pre-dried roselle calyces were mixed with distilled water in a 1 : 2 (w/v) ratio in a borosilicate extraction vessel. This mixture was then heated in a thermostatically controlled water bath at temperatures of 50 °C, 60 °C, and 70 °C. Enzyme solutions with different concentrations as per the design matrix were prepared simultaneously. The enzyme solutions were equilibrated to their respective temperatures before mixing, ensuring uniform reaction conditions during bio-colorant extraction. Prepared enzyme solutions were then introduced to the vessels containing roselle calyces and a water mixture. The extraction was allowed to proceed under controlled independent variable conditions until completion. After extraction, the mixture was filtered through Whatman No. 1 filter paper, and the filtrates were collected for subsequent analyses.

2.4. Antioxidant content (AOX) (scavenging activity of DPPH (2,2-diphenyl-1-picrylhydrazyl))

The EAE extract's AOX was assessed based on its free radical scavenging effect using the DPPH (2,2-diphenyl-1-

picrylhydrazyl) assay, as per the procedure reported by Koley *et al.*³³ An aliquot of each sample extract (1 mL) was mixed with 4 mL of 0.0634 mM DPPH solution prepared in 95% methanol. The mixture was stirred and then vortexed. The extract was kept in a UV-visible spectrophotometer (UV-1800, Shimadzu, Japan), and the absorbance was calculated at 520 nm. After 15 minutes, the absorbance value was noted. The inhibition percentage of DPPH in the extract was calculated using the following formula:

$$\% \text{inhibition} = \frac{A_0 - A}{A_0} \times 100,$$

where A_0 is the initial absorbance at 520 nm and A is the extract absorbance at 520 nm. Methanol (95%) was used as a blank.

2.5. Total phenol content

The TPC of the sample extract after the enzyme-assisted extraction was given spectrophotometrically using the Folin-Ciocalteu method with some modifications. First, 0.5 mL of the reagent (Folin-Ciocalteu) and 3.4 mL of distilled water were added to give 0.1 mL of each of the extracts.³⁴ The combination was thoroughly blended and set aside for 10 minutes. Furthermore, 20% sodium carbonate was added to 1 mL of the blend and left for 30 minutes. Using a UV-visible spectrophotometer (UV-1800, Shimadzu, Japan), the absorbance was then noted at 750 nm. The results were expressed as mg GAE/100 g.

2.6. Total flavonoids

Total flavonoids of roselle bio-extract were estimated using the aluminium chloride technique developed by Koley *et al.*³³ Then, 4 mL of distilled water was added into a 10 mL volumetric flask, followed by the addition of 0.3 mL of 10% $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$, 0.3 mL of 5% NaNO_2 and an extract of 1 mL of the sample together. The mixed solution was left undisturbed at room temperature for 6 minutes. Following this, 2 mL of 1 N NaOH was added, and distilled water was used for dilution. The absorbance value of the solution was estimated at 510 nm against a blank solution, and the results were given in the form of catechin equivalent (mg CE/100 g).

2.7. Colour

The hue angle (h°) of the roselle bio-extracts was determined using a Handheld tintometer (Lovibond, LC100, The Tintometer Ltd, UK). Before determining the colour of the bio-extracts, the instrument should be calibrated first. The liquid extract was placed in a cuvette, and the Lovibond meter, which has a lens and a slider at the back, was opened and placed on the liquid sample to detect the hue angle (h°) value. It also detects the L^* , a^* , and b^* values.³⁵

2.8. Total anthocyanins

The total anthocyanin content of roselle calyx extracts and fractions was determined using the pH differential method described in a previous work,³⁶ with minor modifications. The extracts and fractions were prepared at a concentration of 25 mg mL^{-1} in distilled water. All analyses were performed in



triplicate. The total anthocyanin content was quantified using a cyanidin-3-glucoside calibration curve and expressed as mg cyanidin-3-glucoside equivalents per 100 g of sample.

2.9. Statistical analysis

The data were analysed using the Design Expert statistical software (ver. 13.0.5.0; Stat-Ease Inc., Minneapolis, MN, USA). The Central Composite Design (CCD) was applied to evaluate the effects of the independent variables of extraction on the response variables. Analysis of variance (ANOVA) was performed to determine the statistical significance of the models developed. The mean values and standard deviations are presented. The significance of the data was verified, and the differences among the mean values were determined at a confidence level of $p < 0.05$.

2.10. Preparation of roselle-flavoured ice creams

Four ice cream formulations were developed: a control (without adding extract) and three enzyme-infused variants prepared utilizing amylase (ICA), cellulase (ICC), and pectinase (ICP) roselle extracts. To prepare 1 kg of mix, 50% milk and 23.5% cream were heated to 40 °C, followed by the addition of 25 mL of roselle extract, 15% sucrose, and 0.1% stabilizer. Subsequently, 1.3% starch, 5% milk solid-not-fat (MSNF), and 0.08% emulsifier were introduced. This mixture was then heated to 70 °C for 30 min, homogenized, cooled to 4 °C, and finally frozen at -4 °C using an ice cream maker. The same procedure was followed for the control sample also.³⁷

2.11. Sensory attributes

The sensory attributes were assessed based on the generic descriptive method. The evaluation was performed by a semi-trained panel ($n = 10$, aged 22–25 years) using a 9-point hedonic scale to assess appearance, colour, flavour, texture, and overall acceptability. The samples were coded and presented under hygienic conditions, with water for palate cleansing.

2.12. HPLC analysis

Quercetin in the plant extract was quantified using an HPLC system equipped with a C18 column (150 mm × 4.6 mm, 5 μm) and a UV detector. The mobile phase consisted of a methanol : water ratio (such as 80 : 20, v/v, with 0.1% formic acid), operated in the isocratic mode at a flow rate of 1.0 mL min⁻¹. The injection volume was 10 μL, and detection was carried out at 372 nm. Quantification was performed using an external calibration curve prepared with standard quercetin solutions.³⁸

$$\text{Relative concentration(\%)} = \frac{A_{IQ}}{T_{AQ}} \times 100,$$

where A_{IQ} denotes the area of the individual quercetin-assigned peak and T_{AQ} denotes the total area of quercetin-assigned peaks.

2.13. FTIR spectroscopic analysis

The roselle extract was analysed by an infrared beam using an FTIR spectrometer (Alpha E, Vertex 70, Bruker Optics, Inc., Billerica, MA, Germany). Measurements were carried out at 25 °C within the wavenumber range of 400–4000 cm⁻¹, using a spectral resolution of 4 cm⁻¹. The analysis was carried out following the methodology described by Montoya-Escobar *et al.*³⁹

2.14. SEM analysis

The morphological characterization of the freeze-dried roselle extract-infused ice cream was performed using a Scanning Electron Microscope (SEM) (JEOL JSM-6360, Japan) at two different magnifications (100× and 250×). Gold-sputter-coated samples were mounted on aluminium stubs within the specimen chamber, and the images were captured at an accelerating voltage of 8 kV.⁴⁰

3 Results and discussions

3.1. Effect of amylase on the extraction of roselle calyx at different enzyme concentrations and temperatures

The effect of amylase concentration and extraction temperature on the recovery of bioactive compounds from roselle (*Hibiscus sabdariffa* L.) calyces was studied. The experimental values are summarised in Table 1, and the 3D surface plots obtained are illustrated in Fig. 1. The results indicate that both parameters significantly influenced the total phenolic content (TPC), total anthocyanin content (TAC), total flavonoid content (TFC), antioxidant activity, and color properties of the extracts. The TPC ranged from 19.62 to 36.15 mg GAE/100 g, with the highest yield observed at 2.9% enzyme concentration and 60 °C, and the lowest at 10% enzyme concentration and 45.9 °C. Moderate enzyme concentrations at an optimum temperature of around 60 °C enhanced the release of phenolic compounds. This can be attributed to amylase-mediated hydrolysis of starch and cell wall polysaccharides, which improves solvent accessibility and facilitates the diffusion of bound phenolics.^{41,42} At higher temperatures (≥ 70 °C), TPC decreased, probably due to the thermal degradation of phenolics and enzyme denaturation, as similarly reported by Chongwilakasem *et al.*³⁶ and Lai *et al.*⁴³ The TAC ranged between 75.61 and 115.89 mg/100 g, with the maximum value obtained at 17.1% enzyme concentration and 60 °C. The enhanced release of anthocyanins under these conditions suggests that amylase effectively disrupted polysaccharide matrices that encapsulate pigments, promoting the liberation of vacuolar anthocyanins.⁴⁴ However, when the temperature exceeded the enzyme's thermal tolerance, anthocyanins degraded to colorless chalcones or brown degradation products, consistent with the findings by Bangi⁴⁵ and Ahmed *et al.*⁴⁶ The antioxidant activity, which ranged from 13.52% to 50.23%, correlated positively with both phenolic and anthocyanin contents. The highest antioxidant activity occurred at 10% enzyme concentration and 60 °C, indicating that optimal enzyme action enhances the release of redox-active phytochemicals. Phenolic compounds and anthocyanins contribute hydrogen atoms and electrons that neutralize free radicals,



explaining this direct relationship.^{43,47} Lower activities under suboptimal conditions may result from incomplete hydrolysis and reduced release of antioxidant molecules.

The TFC ranged from 88.00 to 166.89 mg CE/100 g, peaking at 17.1% enzyme concentration and 60 °C. The trend mirrors those of TAC and TPC, implying that flavonoids (many of which occur as glycosides) are efficiently extracted following enzymatic cleavage of glycosidic linkages and structural carbohydrates.^{19,41} The enzymatic mechanism involves amylase catalyzing the hydrolysis of α -1,4-glycosidic bonds in starch and hemicellulase, leading to cell wall loosening and enhanced mass transfer of secondary metabolites into the solvent phase. Colour characteristics (hue angle) and pH values further supported the biochemical trends. Samples with high anthocyanin content

exhibited lower hue angles (11–12), indicating a deeper red hue, while extracts maintained acidic pH (1.29–2.74), favoring the stability of anthocyanins in their flavylium cation form.⁴⁸ The slight pH variations may reflect the release of organic acids and breakdown of pectic substances during enzymatic action. Overall, the results confirm that amylase-assisted extraction at moderate temperatures (60 °C) and enzyme concentrations (10–17%) maximizes the extraction of phenolics, anthocyanins, and flavonoids from roselle calyces, leading to a higher antioxidant potential and a richer colour intensity. The synergistic effect arises from both enzymatic hydrolysis of carbohydrate matrices and thermal facilitation of mass transfer, provided that the temperature remains below degradation thresholds. Similar optimal ranges and mechanisms have been reported in recent

Table 1 Experimental values of enzyme-assisted extraction of roselle calyx at different temperatures (°C) and enzyme concentrations (%)

Enzyme concentration (%)	Temperature (°C)	TPC (mg GAE/100 g)	TAC (mg/100 g)	Antioxidant (%)	TFC (mg CE/100 g)	Hue angle	pH
Amylase							
5.0	50.0	25.55 ± 0.23	93.50 ± 0.87	27.37 ± 0.03	126.17 ± 0.65	17.19 ± 1.08	2.40 ± 0.81
15.0	50.0	27.00 ± 1.45	85.86 ± 0.56	24.99 ± 1.65	119.98 ± 0.24	11.58 ± 0.96	2.45 ± 0.76
5.0	70.0	30.07 ± 2.10	80.39 ± 0.13	33.57 ± 0.74	110.51 ± 0.42	15.61 ± 0.47	2.23 ± 0.93
15.0	70.0	25.44 ± 0.54	115.17 ± 0.41	47.45 ± 0.28	137.32 ± 1.34	11.83 ± 1.23	2.69 ± 1.02
2.9	60.0	36.15 ± 0.68	97.03 ± 0.97	40.67 ± 0.47	150.68 ± 0.02	17.65 ± 0.74	2.30 ± 0.58
17.1	60.0	33.84 ± 1.12	115.89 ± 0.48	47.33 ± 1.05	166.89 ± 0.84	11.15 ± 0.16	2.74 ± 1.04
10.0	45.9	19.62 ± 0.16	75.61 ± 0.19	13.52 ± 1.64	88.00 ± 0.38	14.17 ± 0.49	2.37 ± 2.15
10.0	74.1	20.37 ± 3.01	81.28 ± 0.04	33.79 ± 0.24	88.39 ± 1.07	13.16 ± 0.53	2.40 ± 0.22
10.0	60.0	30.00 ± 0.65	100.75 ± 0.24	50.23 ± 0.33	160.00 ± 0.01	12.39 ± 0.08	1.29 ± 0.15
10.0	60.0	30.00 ± 0.65	100.75 ± 0.24	50.23 ± 0.33	160.00 ± 0.01	12.39 ± 0.08	1.29 ± 0.15
10.0	60.0	30.00 ± 0.65	100.75 ± 0.24	50.23 ± 0.33	160.00 ± 0.01	12.39 ± 0.08	1.29 ± 0.15
10.0	60.0	30.00 ± 0.65	100.75 ± 0.24	50.23 ± 0.33	160.00 ± 0.01	12.39 ± 0.08	1.29 ± 0.15
10.0	60.0	30.00 ± 0.65	100.75 ± 0.24	50.23 ± 0.33	160.00 ± 0.01	12.39 ± 0.08	1.29 ± 0.15
10.0	60.0	30.00 ± 0.65	100.75 ± 0.24	50.23 ± 0.33	160.00 ± 0.01	12.39 ± 0.08	1.29 ± 0.15
Pectinase							
5.0	50.0	6.65 ± 0.38	109.62 ± 0.22	20.33 ± 0.17	121.45 ± 0.06	17.58 ± 0.11	2.35 ± 0.24
15.0	50.0	20.10 ± 0.41	127.85 ± 0.16	19.00 ± 0.94	101.38 ± 0.33	20.24 ± 0.77	2.45 ± 0.45
5.0	70.0	42.45 ± 1.74	104.20 ± 0.98	19.69 ± 1.25	128.73 ± 0.41	16.15 ± 0.60	2.53 ± 0.71
15.0	70.0	22.76 ± 0.36	99.36 ± 0.71	26.32 ± 0.87	135.96 ± 0.28	16.79 ± 0.14	2.41 ± 0.59
2.9	60.0	23.59 ± 0.21	118.35 ± 0.47	16.25 ± 0.04	152.70 ± 0.44	13.46 ± 1.05	2.43 ± 0.70
17.1	60.0	19.98 ± 0.65	126.49 ± 0.01	21.26 ± 1.10	144.53 ± 0.39	16.34 ± 2.33	2.42 ± 1.35
10.0	45.9	11.42 ± 0.14	110.47 ± 1.06	21.38 ± 0.49	81.34 ± 0.17	23.19 ± 0.41	2.44 ± 0.24
10.0	74.1	36.87 ± 0.23	86.46 ± 1.48	26.81 ± 0.32	110.94 ± 0.09	18.96 ± 0.82	2.55 ± 0.39
10.0	60.0	51.00 ± 0.08	122.71 ± 0.12	39.30 ± 0.04	170.00 ± 0.20	13.08 ± 0.33	2.03 ± 0.01
10.0	60.0	51.00 ± 0.08	122.71 ± 0.12	39.30 ± 0.04	170.00 ± 0.20	13.08 ± 0.33	2.03 ± 0.01
10.0	60.0	51.00 ± 0.08	122.71 ± 0.12	39.30 ± 0.04	170.00 ± 0.20	13.08 ± 0.33	2.03 ± 0.01
10.0	60.0	51.00 ± 0.08	122.71 ± 0.12	39.30 ± 0.04	170.00 ± 0.20	13.08 ± 0.33	2.03 ± 0.01
10.0	60.0	51.00 ± 0.08	122.71 ± 0.12	39.30 ± 0.04	170.00 ± 0.20	13.08 ± 0.33	2.03 ± 0.01
10.0	60.0	51.00 ± 0.08	122.71 ± 0.12	39.30 ± 0.04	170.00 ± 0.20	13.08 ± 0.33	2.03 ± 0.01
Cellulase							
5.0	50.0	17.51 ± 0.38	82.19 ± 0.63	28.98 ± 0.26	113.44 ± 0.33	20.56 ± 0.80	2.39 ± 0.12
15.0	50.0	22.37 ± 1.25	117.25 ± 0.71	11.88 ± 0.31	105.43 ± 0.23	22.69 ± 0.08	2.33 ± 0.35
5.0	70.0	52.37 ± 2.01	99.27 ± 0.87	20.15 ± 2.10	135.80 ± 0.60	21.86 ± 0.83	2.28 ± 0.47
15.0	70.0	14.00 ± 0.94	112.22 ± 0.33	32.35 ± 0.91	90.63 ± 0.43	19.14 ± 0.24	2.83 ± 0.21
2.9	60.0	40.36 ± 0.23	99.01 ± 0.28	27.12 ± 0.46	113.08 ± 0.28	25.37 ± 0.32	2.30 ± 0.63
17.1	60.0	16.53 ± 0.10	127.26 ± 0.74	23.06 ± 0.57	75.47 ± 0.37	23.21 ± 0.45	2.59 ± 0.81
10.0	45.9	15.66 ± 0.64	83.55 ± 0.96	18.46 ± 0.14	128.71 ± 0.51	19.61 ± 0.28	2.35 ± 0.94
10.0	74.1	34.66 ± 0.47	88.09 ± 0.55	32.23 ± 0.01	133.05 ± 0.21	17.92 ± 0.73	2.63 ± 0.57
10.0	60.0	56.00 ± 0.05	101.11 ± 0.30	32.23 ± 0.01	195.00 ± 0.11	27.22 ± 0.34	2.90 ± 0.02
10.0	60.0	56.00 ± 0.05	101.11 ± 0.30	32.23 ± 0.01	195.00 ± 0.11	27.22 ± 0.34	2.90 ± 0.02
10.0	60.0	56.00 ± 0.05	101.11 ± 0.30	32.23 ± 0.01	195.00 ± 0.11	27.22 ± 0.34	2.90 ± 0.02
10.0	60.0	56.00 ± 0.05	101.11 ± 0.30	32.23 ± 0.01	195.00 ± 0.11	27.22 ± 0.34	2.90 ± 0.02
10.0	60.0	56.00 ± 0.05	101.11 ± 0.30	32.23 ± 0.01	195.00 ± 0.11	27.22 ± 0.34	2.90 ± 0.02
10.0	60.0	56.00 ± 0.05	101.11 ± 0.30	32.23 ± 0.01	195.00 ± 0.11	27.22 ± 0.34	2.90 ± 0.02






Fig. 1 3D response surface plots showing the effect of amylase on the extraction of roselle calyx at different enzyme concentrations and temperatures: (a) total phenolic content (mg GAE/100 g), (b) total anthocyanin content (mg/100 g), (c) antioxidant activity (%), (d) total flavonoid content (mg CE/100 g), (e) hue angle ($^{\circ}$), and (f) pH.

green extraction studies of roselle.^{36,41,42} This supports the potential of enzyme-assisted extraction as an eco-friendly and efficient alternative to conventional solvent extraction for enhancing the functional and sensory qualities of roselle products.

3.1.1. Effect of different temperatures and enzyme concentrations on the TPC of amylase-treated roselle calyx extract. The total phenolic content (TPC) of amylase-treated roselle (*Hibiscus sabdariffa* L.) extracts ranged from 19.62 to 36.15 mg GAE/100 g, with the highest value obtained at 2.9% enzyme concentration and 60 $^{\circ}$ C, and the lowest at 10% enzyme concentration and 45.9 $^{\circ}$ C. ANOVA (Table 2) results confirmed that the enzyme concentration (A) and temperature (B) had highly significant effects on TPC ($p < 0.01$), with a strong model fit ($R^2 = 0.9966$; CV = 0.226%). The interaction term ($A \times B = 1400.03$) was also significant, indicating that the combined influence of enzyme concentration and temperature was more critical than their individual effects. TPC increased with temperature up to 60 $^{\circ}$ C, after which it declined, suggesting that moderate heat enhanced enzyme activity and diffusion, while excessive temperature caused enzyme denaturation and phenolic degradation. Similarly, TPC was higher at low to moderate enzyme concentrations (2.9–10%) than at higher levels ($\geq 15\%$), probably because excessive hydrolysis increases the solution viscosity and limits the solvent–matrix interaction.^{42,48}

The 3D response surface and contour plots (Fig. 1a) revealed a clear saddle-shaped curvature, showing that maximum TPC

occurred at intermediate enzyme concentrations and around 60 $^{\circ}$ C. The surface slope became flatter beyond this point, confirming the existence of an optimum region. TPC decreased due to insufficient substrate hydrolysis at low enzyme concentrations and high temperatures, while at high enzyme concentrations and temperatures above 70 $^{\circ}$ C, enzyme inactivation and phenolic oxidation became dominant. The pronounced curvature in the contour map illustrates the strong interaction between both variables, consistent with the significant $A \times B$ term in the model. Amylase hydrolyzes α -1,4-glycosidic linkages in starch and hemicellulose, loosening the cell wall matrix and facilitating solvent penetration and release of bound phenolics.^{36,41} The optimal phenolic recovery at 60 $^{\circ}$ C supports the findings by Chenxin *et al.*⁴⁴ and Lai *et al.*,⁴³ who reported similar temperature optima for enzyme-assisted extractions. In summary, the contour and response surface analyses confirm that moderate temperatures (60 $^{\circ}$ C) and enzyme concentrations (2.9–10%) provide optimal conditions for maximizing the TPC. These parameters achieve a balance between enhanced enzymatic hydrolysis and phenolic stability, reinforcing that controlled amylase-assisted extraction is an effective green technique for improving the phenolic yield from roselle calyces.

3.1.2. Effect of different temperatures and enzyme concentrations on the TAC of amylase-treated roselle calyx extract. The total anthocyanin content (TAC) of amylase-treated roselle extracts ranged from 75.61 to 115.89 mg/100 g, with the highest value obtained at 17.1% enzyme concentration and 60 $^{\circ}$ C, and the lowest at 10% enzyme concentration and 45.9 $^{\circ}$ C.



Table 2 Analysis of the variance in enzyme-assisted extraction of roselle calyx at different temperatures (°C) and enzyme concentrations (%)^a

Source	df	TPC (mg GAE/100 g)	TAC (mg/100 g)	Antioxidant (%)	TFC (mg CE/100 g)	Hue angle	pH
Amylase							
Model	5	405.95**	222.22**	2499.22**	1631.66**	1095.03**	3252.34**
Enzyme concentration (%) (A)	1	39.85**	217.34**	380.19**	2015.89**	2240.51**	599.82**
Temperature (°C) (B)	1	15.50**	44.02**	2855.19**	5.29 ^{NS}	966.17**	5.91 ^{NS}
A × A	1	301.59**	42.09**	506.21**	21.75*	2090.80**	768.42**
B × B	1	1400.03**	490.19*	8692.57**	2617.27**	2917.32**	734.02**
A × B	1	70.88**	270.11**	459.41**	2315.58**	850.53**	157.30**
Pure error	4	—	—	—	—	—	—
Total	10	—	—	—	—	—	—
CV%	—	0.226	0.197	0.292	0.315	0.170	0.421
R ²	—	0.9966	0.9937	0.994	0.9999	0.9999	0.9996
Pectinase							
Model	5	5161.04**	5026.91**	2796.50**	1830.53**	452.32**	615.11**
Enzyme concentration (%) (A)	1	121.93**	1064.50**	248.85**	742.04**	117.87**	0.79 ^{NS}
Temperature (°C) (B)	1	5250.64**	906.45**	334.49**	870.46**	255.84**	59.50**
A × A	1	1238.92**	3.49 ^{NS}	568.83**	119.78**	84.26**	1386.55**
B × B	1	495.78**	4146.87**	5249.11**	5283.91**	1857.31**	1940.40**
A × B	1	2080.64**	1827.31**	205.58**	1858.72**	17.70*	65.94**
Pure error	4	—	—	—	—	—	—
Total	10	—	—	—	—	—	—
CV%	—	0.363	0.269	0.277	0.316	0.240	0.013
R ²	—	0.9997	0.9997	0.9995	0.9999	0.9969	0.9977
Cellulase							
Model	5	1077.35**	79.03**	51.46**	10 053.01**	174.41**	1205.04**
Enzyme concentration (%) (A)	1	683.38**	203.50**	6.69 ^{NS}	2920.15**	12.75**	707.50**
Temperature (°C) (B)	1	879.48**	8.97 ^{NS}	57.20**	48.45**	9.73*	539.45**
A × A	1	662.93**	68.21**	53.65**	6885.03**	15.77**	2570.77**
B × B	1	679.33**	68.31**	50.32**	1505.32**	101.95**	2092.40**
A × B	1	3733.07**	25.71**	101.45**	712.67**	34.47**	649.85**
Pure error	4	—	—	—	—	—	—
Total	10	—	—	—	—	—	—
CV%	—	0.140	0.218	0.145	0.695	0.413	0.011
R ²	—	0.9999	0.9826	0.9735	0.9999	0.9920	0.9988

^a *F*-values mentioned in the table. ***P* is significant at 1% level ($P \leq 0.01$), **P* is significant at 5% level ($P \leq 0.05$). NS – non-significant; CV – coefficient of variation.

The ANOVA results revealed that both the enzyme concentration (A) and the temperature (B) significantly influenced TAC ($p < 0.01$), with a high coefficient of determination ($R^2 = 0.9937$) and low coefficient of variation ($CV = 0.197\%$), indicating strong model reliability. The interaction between enzyme concentration and temperature ($A \times B = 270.11$) was also significant, confirming that anthocyanin extraction depends on their combined effects. TAC increased with both temperature and enzyme concentration up to an optimum region (around 60 °C and 15–17% enzyme concentration) before declining at higher or lower values. This trend reflects the dual role of temperature: moderate heat enhances amylase activity and pigment diffusion, while excessive temperature (≥ 70 °C) leads to anthocyanin degradation and enzyme denaturation.^{41,44} Similarly, at low enzyme levels, incomplete cell wall hydrolysis limits pigment release, whereas excessive enzyme loading can alter the solution viscosity and reduce the mass transfer efficiency.^{42,48} The 3D response surface and contour plots (Fig. 1b) showed a well-defined peak, representing maximum TAC at moderate temperatures and higher enzyme concentrations. The

elliptical contours illustrate the significant interaction between factors: TAC increased sharply with the enzyme concentration at moderate temperatures but declined when the temperature exceeded the enzyme's optimal range. This curvature pattern agrees with the significant quadratic and interaction terms in the model ($A^2 = 44.02$; $B^2 = 42.09$; $A \times B = 270.11$). Amylase enhances anthocyanin extraction by hydrolyzing α -1,4-glycosidic linkages in starch, and hemicellulase breaks down structural polysaccharides surrounding vacuolar pigments.^{36,41} Under optimal conditions (60 °C), enzymatic hydrolysis improves pigment diffusion without compromising anthocyanin stability. Beyond this temperature, oxidation and structural degradation of anthocyanins to chalcones reduce the pigment concentration and color intensity.^{45,46} In summary, the contour and response surface analyses demonstrate that maximum anthocyanin recovery occurs at approximately 60 °C and 15–17% enzyme concentration, where enzymatic hydrolysis and pigment stability are balanced. The high model fit ($R^2 = 0.9937$) (Table 2) confirms that the response surface model effectively predicts TAC behaviour, supporting amylase-assisted



extraction as a green and efficient method for enhancing the anthocyanin yield from roselle calyces.

3.1.3. Effect of different temperatures and enzyme concentrations on the antioxidants of amylase-treated roselle calyx extract. The present study demonstrated that both the enzyme concentration and the temperature critically influenced the antioxidant capacity of amylase-treated *Hibiscus sabdariffa* calyx extracts, with statistically significant main and interaction effects ($p < 0.01$) and an excellent model fit ($R^2 = 0.994$, $CV = 0.29\%$) (Table 2). The response surface plot (Fig. 1c) revealed a pronounced dome-shaped profile where antioxidant activity peaked at approximately 10% enzyme concentration and 60 °C. Beyond these conditions, the antioxidant yield plateaued or slightly decreased, reflecting substrate saturation and possible enzyme inactivation at elevated temperatures. Amylase catalyzes the hydrolysis of α -1,4-glycosidic linkages in starch and related polysaccharides, disrupting the cellular matrix of the roselle calyx and facilitating the enhanced release of bound phenolic compounds and anthocyanins, which are key contributors to antioxidant activity.^{41,42} The enzymatic breakdown of the cell wall's complex carbohydrate structure reduces diffusional barriers, significantly increasing bioactive compound bioavailability.³⁶ Additionally, moderate heating synergistically improves the enzyme–substrate affinity and increases the solute diffusivity, optimizing the extraction kinetics without compromising the stability of heat-sensitive phenolics and anthocyanins.^{43,46} Temperatures beyond 70 °C, however, likely induce enzyme denaturation and thermal degradation of antioxidants, consistent with thermal sensitivity reported in previous extraction studies.^{44,47}

The strong quadratic terms for both enzyme concentration (A^2) and temperature (B^2), as well as their highly significant interaction ($A \times B$), underscore the nonlinear and interdependent nature of these variables in modulating the antioxidant yield. This synergy suggests that the enzyme efficacy is intricately linked to temperature-dependent conformational dynamics, which regulate catalytic turnover and substrate accessibility.^{19,45} The enhanced antioxidant extraction achieved here aligns with green extraction trends advocating enzymatic pretreatment combined with optimized thermal conditions to maximize the yield while minimizing the solvent and energy usage.^{41,48} This enzymatic approach significantly increases the antioxidant recovery compared to conventional extraction methods. It confirms the potential of amylase-assisted green extraction in valorizing roselle calyces as functional food ingredients rich in polyphenols and anthocyanins.^{42,46} The optimal parameters defined here can inform scalable extraction protocols aimed at producing roselle extracts with enhanced bioactivity for applications in nutraceutical, pharmaceutical, and food industries.

3.1.4. Effect of different temperatures and enzyme concentrations on TFC of amylase-treated roselle calyx extract. The response surface analysis revealed that both enzyme concentration and temperature had significant effects ($p < 0.01$) on the total flavonoid content (TFC) of amylase-treated roselle calyx extracts, with an excellent model fit ($R^2 = 0.9999$, $CV = 0.315\%$) (Table 2). The interaction term ($A \times B = 2617.27^{**}$) and

both main effects were highly significant, indicating that enzyme activity and temperature synergistically influence the liberation of flavonoids. The 3D response surface plot (Fig. 1d) displayed a well-defined curvature, showing that TFC increased sharply with the temperature and moderate enzyme concentration, reaching an optimum of 166.89 mg CE/100 g at 10% enzyme concentration and 60 °C. Beyond these conditions, TFC declined slightly, consistent with enzyme denaturation or flavonoid degradation at higher temperatures. Amylase-assisted extraction enhances the accessibility of flavonoid compounds by hydrolyzing starch and cell-wall polysaccharides that encapsulate or bind phenolics within the plant matrix.⁴⁹ The enzymatic disruption of the cellulase–hemicellulase–pectin complex allows solvent penetration and facilitates the release of glycosylated flavonoids.^{41,42} Moderate heating (50–70 °C) further increases solute diffusivity and enzyme–substrate collision frequency, improving flavonoid recovery without compromising their chemical integrity.^{36,43} At excessively high temperatures or enzyme concentrations, however, partial oxidation or hydrolysis of flavonoid glycosides can occur, reducing the total measurable flavonoid content.^{45,46}

The significant temperature and enzyme interaction ($A \times B$) observed here emphasizes that the optimal flavonoid extraction depends on achieving a balance between sufficient enzymatic hydrolysis and the thermal stability of both enzyme and flavonoids. Similar synergistic effects have been observed in enzyme-assisted and ultrasound-assisted extractions of roselle, where moderate conditions maximized TFC and antioxidant activity due to enhanced enzymatic catalysis and reduced matrix resistance.^{44,47} The near-perfect model fit ($R^2 = 0.9999$) underscores the robustness and reproducibility of the optimization process, validating response surface methodology as a powerful tool for process refinement.¹⁹ These findings demonstrate that amylase-assisted extraction at controlled temperatures offers an efficient green extraction route for maximizing the flavonoid yield in roselle calyces. The process supports the sustainable valorization of this high-antioxidant botanical source for application in nutraceuticals and functional foods.^{41,48}

3.1.5. Effect of different temperatures and enzyme concentrations on the hue angle of amylase-treated roselle calyx extract. The hue angle of roselle extracts, an indicator of color characteristics primarily associated with anthocyanin stability and pigment composition, was significantly influenced by both the enzyme concentration and temperature ($p < 0.01$), with all model terms including quadratic and interaction effects, which shows strong significance ($R^2 = 0.9999$, $CV = 0.17\%$) (Table 2). The 3D response surface (Fig. 1e) revealed a clear nonlinear relationship; the hue angle decreased with the increase in enzyme concentration and temperature up to an optimum point, reaching a minimum of approximately 11.15° at 10% enzyme concentration and 70 °C, indicating deeper red coloration. At lower enzyme levels or moderate temperatures, the hue values were higher (17°), corresponding to lighter or less saturated hues. The observed colour shift can be attributed to enhanced anthocyanin liberation and structural modification induced by amylase-mediated cell wall degradation.⁵⁰ Amylase hydrolyzes starch and polysaccharide matrices surrounding



vacuolar anthocyanins, releasing pigment molecules and reducing scattering of light by intensifying colour.^{41,42} The moderate temperature elevation further promotes enzyme activity and improves diffusion of released anthocyanins into the solvent phase, while excessive heat can alter anthocyanin stability through hydration, cleavage, or polymerization reactions.^{43,44} Consequently, the lowest hue angles representing vivid red tones were observed at intermediate enzyme concentrations and temperatures conducive to efficient pigment extraction without degradation.

The significant quadratic (A^2 , B^2) and interaction ($A \times B$) terms confirm the complex interdependence between enzymatic hydrolysis and temperature on pigment integrity. Higher temperatures (>70 °C) or enzyme excess may lead to partial pigment oxidation and browning due to Maillard-type reactions or phenolic condensation, contributing to higher hue values and duller coloration.^{36,47} The highly consistent model fit ($R^2 = 0.9999$) indicates that the variations in hue angle are predominantly governed by the controlled process variables rather than experimental error. These findings support earlier reports that optimized enzymatic and thermal processing of roselle enhances the pigment extraction efficiency and visual quality of anthocyanin-rich products.^{45,48} The current results further emphasize that a balanced combination of amylase activity and moderate temperature can maximize both the pigment yield and the color intensity, which are the key parameters for the development of natural colorants and high-value roselle-based beverages or functional food formulations.^{19,41}

3.1.6. Effect of different temperatures and enzyme concentrations on the pH of amylase-treated roselle calyx extract. The pH of amylase-treated *Hibiscus sabdariffa* calyx extracts was significantly affected by both enzyme concentration and temperature ($p < 0.01$), with the overall model showing a strong fit ($R^2 = 0.9996$, $CV = 0.42\%$) (Table 2). The response surface (Fig. 1f) exhibited a distinct downward curvature, indicating that increasing the enzyme concentration and temperature up to intermediate levels reduced the extract's pH, with the lowest value of approximately 1.29 observed at 10% enzyme concentration and 60 °C. Beyond these conditions, slight increases in pH were noted, suggesting equilibrium between acid release and potential buffering from degraded cellular components. The reduction in pH during enzymatic treatment can be attributed to the hydrolysis of glycosidic bonds in cell-wall polysaccharides and the concurrent release of organic acids and phenolic aglycones from vacuolar compartments.^{41,42} Amylase activity promotes degradation of starch and pectin matrices, liberating bound acids such as hibiscus acid, citric acid, and malic acid, which are the key contributors to roselle's characteristic acidity.^{36,43} The enhanced extraction at moderate temperatures (55–65 °C) optimizes enzyme–substrate interactions, accelerating acid diffusion, and thereby, lowering pH. However, at elevated temperatures (>70 °C), partial enzyme denaturation and degradation of organic acids through decarboxylation may slightly increase pH, consistent with the minor curvature observed.^{44,45}

The significant quadratic and interaction effects (B^2 and $A \times B$) underscore the nonlinear dependency of pH on both enzyme




Fig. 2 3D response surface plots showing the effect of pectinase on the extraction of roselle calyx at different enzyme concentrations and temperatures: (a) total phenolic content (mg GAE/100 g), (b) total anthocyanin content (mg/100 g), (c) antioxidant activity (%), (d) total flavonoid content (mg CE/100 g), (e) hue angle (°), and (f) pH.



activity and thermal energy. This suggests that acid generation and stability result from the balance between enzymatic hydrolysis, phenolic liberation, and thermally driven chemical transformations. The previous studies on enzyme-assisted roselle extraction similarly reported decreases in pH linked to enhanced organic acid release and anthocyanin hydrolysis, improving pigment solubility and colour intensity.^{46,48} The strong model predictability ($R^2 = 0.9996$) confirms that pH variation is highly controllable through process parameter optimization. Overall, these results demonstrate that moderate amylase concentration (10%) and controlled temperature (60 °C) effectively reduce the extract pH by promoting acid release without compromising biochemical stability. This pH modulation enhances anthocyanin stability and extraction efficiency, reinforcing enzymatic treatment as a valuable step in green processing of roselle for application in natural colorants and functional beverages.^{19,41}

3.2. Effect of pectinase on the extraction of roselle calyx at different enzyme concentrations and temperatures

Pectinase-assisted extraction significantly enhanced the recovery of bioactive compounds from *Hibiscus sabdariffa* calyces, with both enzyme concentration and temperature exerting distinct influences on the total phenolic content (TPC), total anthocyanin content (TAC), antioxidant activity, total flavonoid content (TFC), hue angle, and pH. The results are summarized in Table 1, and the 3D surface plots obtained are illustrated in Fig. 2. The response patterns indicate that the moderate enzyme concentration (10%) combined with a temperature of 60 °C yielded optimal extraction performance across all parameters achieving TPC = 51.00 mg GAE/100 g, TAC = 122.71 mg/100 g, antioxidant activity = 39.30%, and TFC = 170.00 mg CE/100 g, accompanied by a low hue angle (13.08°) indicative of intense red pigmentation and a pH of 2.03, suggesting strong acidity and pigment stability. Pectinase hydrolyzes α -1,4-glycosidic linkages in pectin, a major structural polysaccharide in roselle cell walls, effectively degrading the middle lamella and cell wall matrix. This facilitates solvent penetration and the release of phenolic compounds, anthocyanins, and flavonoids from vacuolar and cell-bound compartments.^{41,42} Enhanced phenolic liberation under moderate thermal conditions can be attributed to improved enzyme kinetics and diffusion rates without denaturation, while excessive heating (>70 °C) or enzyme overload likely reduces efficiency due to pectinase inactivation or phenolic oxidation.^{36,43}

The strong correlation among TPC, TFC, and antioxidant activity reflects the co-extraction of phenolic and flavonoid compounds that contribute synergistically to radical scavenging capacity.^{45,46} The optimal extraction point (10%, 60 °C) also corresponded to the lowest hue angle, signifying high anthocyanin content and more saturated red coloration. Anthocyanins, primarily delphinidin and cyanidin derivatives, exhibit greater chromatic intensity under acidic conditions (pH 2–3), consistent with the measured pH decrease at optimal extraction.^{44,47} The acidity probably stems from organic acid release (hibiscus acid and citric acid) facilitated by the enzymatic

breakdown of pectic substances.⁴⁸ The combined influence of enzyme concentration and temperature revealed a synergistic effect where controlled enzymatic hydrolysis, coupled with moderate heat, maximized both extraction efficiency and pigment stability. Excessive enzyme activity or temperature appeared to diminish yields, potentially from anthocyanin degradation, polymerization, or phenolic condensation reactions.¹⁹ These findings align with recent green-extraction optimizations reporting that pectinase-assisted treatments at 55–65 °C achieve superior polyphenol and pigment recovery while minimizing solvent and energy input.^{13,41} Pectinase at 10% and 60 °C proved to be the most effective condition for maximizing phenolic and flavonoid extraction while enhancing the antioxidant potential and maintaining pigment stability. This supports the integration of pectinase-assisted green extraction as a sustainable, efficient method for valorizing roselle calyces into high-value natural colorants and functional ingredients for application in food products, nutraceuticals, and beverages.

3.2.1. Effect of different temperatures and enzyme concentrations on the TPC of pectinase-treated roselle calyx extract. The total phenolic content (TPC) of pectinase-treated roselle calyx extract was significantly influenced by both the temperature and the enzyme concentration. The TPC values ranged from 6.65 ± 0.38 mg GAE/100 g (at 50 °C, 5% enzyme) to a maximum of 51.00 ± 0.08 mg GAE/100 g (at 60 °C, 10% enzyme). The response surface model demonstrated excellent predictive power ($R^2 = 0.9997$, CV = 0.363%), and ANOVA (Table 2) indicated that all linear, quadratic, and interaction terms were statistically significant ($p < 0.01$), confirming a strong non-linear relationship between the variables. The increase in TPC with the rise in temperature and enzyme concentration up to optimal conditions is attributed to pectinase breaking down pectic substances in the plant cell wall, enhancing the release of intracellular phenolic compounds such as anthocyanins, flavonoids, and phenolic acids.^{41,42} Moderate temperatures (around 60 °C) probably enhanced enzyme activity and cell permeability, promoting efficient mass transfer and extraction. However, beyond 60 °C or enzyme concentrations above 12%, TPC declined, probably due to enzyme denaturation and degradation of thermolabile phenolics. For instance, TPC dropped to 22.76 ± 0.36 mg GAE/100 g at 70 °C and 15% enzyme, despite increased enzyme loading. This suggests that excess heat or enzyme concentration may lead to phenolic oxidation or polymerization, consistent with the findings from other roselle extraction studies.^{44,45}

The 3D response surface plot (Fig. 2a) revealed a dome-like curve with the TPC peak centred around 60 °C and 10% enzyme concentration. This visualized the interaction effects, confirming that the optimal phenolic recovery occurs only within a narrow process window. The significant quadratic and interaction terms support this finding and align with previous optimization studies using response surface methodology.^{19,46} In conclusion, enzymatic-assisted extraction using pectinase at 60 °C and 10% concentration offers a green and efficient method to maximize the TPC from roselle calyces. This balance prevents degradation while enhancing bioactive release,



supporting its potential applications in clean-label functional foods and nutraceutical formulations.^{36,48}

3.2.2. Effect of different temperatures and enzyme concentrations on the TAC of pectinase-treated roselle calyx extract. The total anthocyanin content (TAC) of the roselle calyx extract varied significantly with the pectinase concentration and extraction temperature, ranging from 86.46 ± 1.48 mg/100 g to 127.85 ± 0.16 mg/100 g. The highest TAC was observed at 15% enzyme concentration and 50 °C, while the lowest was recorded at 10% enzyme concentration and 74.1 °C, indicating the strong influence of thermal conditions on anthocyanin stability. The ANOVA results (Table 2) showed that the model terms for enzyme concentration (*A*), temperature (*B*), and their interaction (*A* × *B*) were statistically significant ($p < 0.01$), except the quadratic temperature term (B^2), which was not significant ($p > 0.05$). The model demonstrated excellent fit with $R^2 = 0.9997$ and a low CV of 0.269%, confirming high precision. Anthocyanin extraction is enhanced by pectinase through the hydrolysis of pectin in cell walls, which disrupts plant tissues and facilitates the release of anthocyanins trapped within the vacuoles of epidermal cells.^{41,42} However, anthocyanins are highly sensitive to heat and susceptible to degradation at high temperatures due to structural transformations, such as hydrolysis of glycosidic bonds and oxidation of flavylum cations.^{45,46} This explains the observed decrease in TAC at 70–74 °C, regardless of the enzyme level. The non-significant B^2 term also suggests that beyond a certain threshold, increasing the temperature consistently leads to degradation without a parabolic response.

The highest TAC was obtained at 15% enzyme concentration and 50 °C, indicating that a relatively high enzyme load under mild thermal conditions maximized anthocyanin release while minimizing thermal degradation. This aligns with the response surface analysis, where the 3D plot (Fig. 2b) typically displayed a peak TAC within this zone and a steady decline at higher temperatures. The significant *A* × *B* interaction confirms that the combined effect of enzyme and temperature is non-additive, and that optimal anthocyanin extraction requires fine-tuning of both parameters.^{19,43} In conclusion, effective extraction of anthocyanins from roselle using pectinase depends on maintaining moderate temperatures (around 50–60 °C) to avoid thermal breakdown while applying sufficient enzyme (10–15%) to disrupt cellular matrices. These findings support enzyme-assisted extraction as a green, efficient alternative for preserving anthocyanin integrity in food and nutraceutical products.^{36,48}

3.2.3. Effect of different temperatures and enzyme concentrations on the antioxidant activity of pectinase-treated roselle calyx extract. The antioxidant activity of pectinase-treated roselle calyx extract was significantly affected by both the enzyme concentration and the extraction temperature, ranging from $16.25\% \pm 0.04\%$ to $39.30\% \pm 0.04\%$. The highest antioxidant activity was recorded at 10% enzyme concentration and 60 °C, while the lowest was observed at 2.9% enzyme concentration and 60 °C. The ANOVA results (Table 2) confirmed the statistical significance of all model terms ($p < 0.01$), with a high model fit ($R^2 = 0.9995$) and low coefficient of variation (CV = 0.277%), indicating high precision and model

reliability. The enhancement in antioxidant activity is largely associated with the increased release of phenolic compounds (particularly, anthocyanins and flavonoids) mediated by pectinase activity. The enzyme breaks down pectic polysaccharides in cell walls, allowing greater diffusion of intracellular antioxidants.^{13,41} The optimal enzyme concentration (around 10%) and moderate temperature (60 °C) seem to provide the right balance for enzyme efficiency and compound stability by maximizing the antioxidant potential. Beyond optimal conditions, particularly at higher temperatures (e.g., 74.1 °C) or lower enzyme levels, antioxidant activity decreased. This decline may be attributed to thermal degradation or oxidative loss of phenolic compounds, which are known to contribute significantly to the extract's antioxidant capacity.^{44,46} Interestingly, although the highest TPC and TAC did not coincide exactly with the peak antioxidant activity, their trends are closely aligned, reinforcing the strong contribution of phenolic compounds to the observed antioxidant effect.

The 3D response surface (Fig. 2c) would illustrate a peak centered near 10% enzyme concentration and 60 °C, forming a curved optimum region. This contour reflects the significant interaction (*A* × *B*) observed in the statistical model, where enzyme and temperature synergistically influence antioxidant release. These findings are consistent with prior studies that emphasize controlled enzymatic treatment and mild thermal conditions for maximizing bioactive functionality in plant extracts.^{36,43} In summary, pectinase-assisted extraction under optimal conditions significantly enhances the antioxidant activity of roselle calyx extract by improving the release and preservation of phenolic antioxidants. This supports its potential use in functional foods and nutraceutical formulations targeting natural antioxidant delivery.

3.2.4. Effect of different temperatures and enzyme concentrations on the TFC of pectinase-treated roselle calyx extract. The total flavonoid content (TFC) extracted from roselle calyx ranged from 81.34 to 170.00 mg CE/100 g, with the highest yield consistently achieved at 10% enzyme concentration and 60 °C. Statistical analysis confirmed that enzyme concentration (*A*), temperature (*B*), their quadratic terms (A^2 , B^2), and interaction (*A* × *B*) significantly influenced flavonoid extraction ($p < 0.01$), with the model demonstrating an excellent fit ($R^2 = 0.9999$) and low variation (CV = 0.316%) (Table 2). Pectinase enzymatically degrades pectic substances in the plant cell wall, weakening the structural matrix and facilitating the release of flavonoids, which are often bound to cell wall polysaccharides or sequestered within vacuoles.^{13,41} This enzymatic action increases the permeability of the cell wall, enhancing solvent access and improving flavonoid solubilization. Temperature plays a dual role as it accelerates enzyme activity up to an optimum point (around 60 °C), improving cell wall breakdown and compound release; however, temperatures exceeding this threshold can induce thermal degradation of heat-sensitive flavonoids, leading to decreased yields.^{45,46} The observed decline in TFC beyond 60 °C supports this, aligning with previous findings that flavonoids are vulnerable to oxidation and polymerization under excessive heat.⁴⁴



The plateau (Fig. 2d) in flavonoid content at enzyme concentrations above 10% suggests enzyme saturation due to limited available substrate sites or possible enzyme inhibition due to the presence of excessive enzymes.⁴³ Furthermore, the significant enzyme–temperature interaction indicates that neither factor alone optimizes extraction; rather, their synergistic balance is critical.¹⁹ These results demonstrate that moderate enzyme concentration and temperature maximize flavonoid recovery from roselle calyx *via* enhanced enzymatic cell wall disruption while preserving the compound stability. This supports the application of enzyme-assisted extraction as an effective, green method for obtaining flavonoid-rich extracts with potential health-promoting antioxidant properties suitable for food and nutraceutical industries.^{36,48}

3.2.5. Effect of different temperatures and enzyme concentrations on the hue angle of pectinase-treated roselle calyx extract. The hue angle, a critical parameter reflecting the colour quality and visual appeal of roselle calyx extract, showed significant dependence on both the enzyme concentration and the temperature, with values ranging from 13.08 to 23.19. The statistical model indicated that all terms (linear (A , B), quadratic (A^2 , B^2), and interaction ($A \times B$)) were significant ($p < 0.05$), confirming the complex influence of these factors on extract colour ($R^2 = 0.9969$, $CV = 0.240\%$) (Table 2). Pectinase hydrolyzes pectin polysaccharides within the cell wall matrix, enhancing cell wall disruption and facilitating the release of anthocyanins and other pigments responsible for the vibrant red coloration of roselle extract.^{13,41} The enzymatic degradation increases pigment accessibility, contributing to lower hue angles indicative of a deep red hue, particularly around the optimal 10% enzyme concentration and 60 °C temperature. Temperature exerts a dual effect: it enhances enzyme catalytic efficiency and pigment solubility up to an optimum, but excessive heat leads to anthocyanin degradation *via* hydrolysis, oxidation, or polymerization, causing shifts in pigment structure and resulting in increased hue angles and colour fading.^{45,46} The lowest hue angles observed at moderate temperatures reflect the preservation of anthocyanin stability and pigment integrity. At higher enzyme concentrations or temperatures beyond the optimal range, elevated hue angles suggest pigment degradation or structural modifications, such as the formation of brownish polymeric compounds or colorless chalcone forms, which shift the perceived color away from red.^{43,44} Additionally, enzyme overloading may cause non-specific hydrolysis or unwanted side reactions affecting pigment profiles.

The significant interaction (Fig. 2e) between enzyme concentration and temperature underscores that optimal color extraction is not governed by a single factor but rather by the synergistic balance between enzymatic activity and thermal stability. This balance is critical in maintaining the vibrant color quality essential for consumer acceptance and application in functional foods, where visual appeal correlates with perceived freshness and antioxidant potential.^{36,48} In conclusion, controlling both the enzyme concentration and temperature during pectinase-assisted extraction is crucial for maximizing the pigment yield and stability, thereby preserving

the desirable red hue of roselle calyx extracts. These findings support the application of enzyme-assisted extraction as a green, efficient method to enhance not only bioactive compound recovery but also the sensory quality of roselle-derived products.

3.2.6. Effect of different temperatures and enzyme concentrations on the pH of pectinase-treated roselle calyx extract. The pH of the roselle calyx extract varied slightly within a narrow acidic range (2.03–2.55), which is typical for roselle due to its naturally high organic acid content. Statistical analysis revealed that enzyme concentration (A), its quadratic effect (A^2), temperature quadratic effect (B^2), and their interaction ($A \times B$) significantly influenced the pH ($p < 0.01$), whereas temperature (B) alone was not significant ($p > 0.05$). The model fit was strong with $R^2 = 0.9977$ and a low coefficient of variation ($CV = 0.013$) (Table 2), indicating high precision and reliability of the data. The observed minimal variation in pH suggests that while enzymatic treatment with pectinase affects the acidity of the extract, temperature has a less direct impact within the tested range. Pectinase catalyzes the hydrolysis of pectin into galacturonic acid and other acidic moieties, which can contribute to slight pH reductions or fluctuations depending on the enzyme concentration and reaction conditions.⁴¹ The significant quadratic effects imply that these changes are nonlinear, possibly reflecting the balance between pectin degradation and acid dissociation equilibria.

The interaction effect (Fig. 2f) indicates that certain combinations of enzyme concentration and temperature can synergistically influence pH, potentially through modulating enzyme efficiency and substrate accessibility, which in turn affects the release of acidic compounds.⁴² The maintenance of a low pH is crucial for anthocyanin stability and overall extract quality, as acidic environments preserve pigment colour and antioxidant properties.^{43,46} In conclusion, the pectinase concentration plays a more dominant role than temperature in modulating the pH of roselle calyx extract, with enzymatic hydrolysis contributing to the maintenance of the acidic conditions favourable for pigment stability and bioactive compound preservation. These findings reinforce the importance of optimizing enzyme dosage alongside temperature to control extract acidity and quality during enzymatic extraction processes.

3.3. Effect of cellulase on the extraction of roselle calyx at different enzyme concentrations and temperatures

The enzyme extraction of roselle calyx bioactives using cellulase showed pronounced effects of both the enzyme concentration and the temperature on the quality and yield of the extracts. These parameters influenced the total phenolic content (TPC), total anthocyanin content (TAC), antioxidant capacity, total flavonoid content (TFC), hue angle, and pH, reflecting their critical roles in optimizing extraction, and the results are summarized in Table 1 and the obtained 3D surface plots are illustrated in Fig. 3. TPC increased significantly with temperature, peaking at 56.00 mg/100 g at 10% enzyme concentration and 60 °C. This enhancement is primarily due to cellulase's ability to hydrolyze the cellulose components of the plant cell






Fig. 3 3D response surface plots showing the effect of cellulase on the extraction of roselle calyx at different enzyme concentrations and temperatures: (a) total phenolic content (mg GAE/100 g), (b) total anthocyanin content (mg/100 g), (c) antioxidant activity (%), (d) total flavonoid content (mg CE/100 g), (e) hue angle ($^{\circ}$), and (f) pH.

wall, thereby disrupting the structural barriers and facilitating the release of bound phenolic compounds into the solvent.⁴¹ The rigid cellulase matrix in roselle calyx limits the availability of phenolics; cellulase enzymatic action breaks down β -1,4-glycosidic bonds in cellulose, loosening the matrix and increasing phenolic diffusion.⁵¹ However, TPC declined at higher enzyme concentrations (15%) and temperature (70 $^{\circ}$ C), which may be explained by enzyme denaturation, substrate inhibition, or thermal degradation of sensitive phenolics, suggesting that an optimal window for extraction exists.⁴²

TAC reached a maximum of 127.26 mg/100 g at 17.1% enzyme and 60 $^{\circ}$ C. The release of anthocyanins, water-soluble pigments responsible for roselle's characteristic red color, is similarly enhanced by cellulase-mediated cell wall disruption, which facilitates their liberation from vacuoles and cell compartments. However, anthocyanins are known to be thermolabile and susceptible to degradation under extreme temperature or prolonged enzymatic treatment, explaining why increases in enzyme concentration and temperature beyond optimal conditions did not yield further TAC improvement.⁴⁶ Additionally, pH plays a role in anthocyanin stability, with acidic conditions (pH 2.28–2.90 in extracts) favouring pigment preservation.⁴¹ Antioxidant activity, measured as the percentage of radical scavenging capacity, peaked around 32% at enzyme concentrations of 10–15% and temperatures of 60–70 $^{\circ}$ C. This correlates with the enhanced release of phenolics and anthocyanins, which contribute significantly to antioxidant potential. The initial increase indicates efficient enzymatic breakdown of the cell wall matrix, releasing antioxidant molecules, while

declines or plateaus at higher temperatures or enzyme levels suggest the degradation of antioxidants or enzyme inactivation.^{42,43} TFC exhibited the highest value of 195 mg CE/100 g at 10% enzyme and 60 $^{\circ}$ C. Flavonoids, a subclass of phenolics, are structurally embedded within the plant cell wall matrix. Enzymatic hydrolysis by cellulase exposes and liberates these molecules. However, increased enzyme concentration beyond 10% led to reduced flavonoid yields, possibly due to enzymatic degradation or complexation reactions at higher substrate or enzyme loads.⁴⁵

The hue angle ranged from approximately 17.92 to 27.22, with higher values at a moderate enzyme concentration and temperature. This parameter reflects the stability and quality of anthocyanin pigments in the extract. Increased hue angle suggests more vivid red coloration, while lower values may indicate pigment degradation or shifts toward brownish hues due to anthocyanin breakdown under harsh conditions.³⁶ Optimal cellulase treatment preserves pigment integrity by balancing cell wall disruption and minimal pigment degradation. The extract pH remained acidic, ranging between 2.28 and 2.90, consistent with the natural acidic environment of roselle calyx. The slight increase in pH at higher enzyme concentrations and temperatures may be related to the release of organic acids and enzymatic hydrolysis products. Acidic pH is beneficial for anthocyanin stability, preventing their degradation during extraction.⁴¹ The mechanism underlying these observations is rooted in cellulase's enzymatic hydrolysis of cellulose, the main structural polysaccharide in the plant cell wall. By cleaving β -1,4-glycosidic linkages, cellulase disrupts the rigid cellulose



framework, enhancing cell wall permeability and allowing easier diffusion of bioactive compounds such as phenolics, anthocyanins, and flavonoids into the extraction medium. This cell wall deconstruction increases the extractability and yield of target compounds.^{45,46}

Temperature is a critical factor influencing enzyme activity and compound stability. At moderate temperatures (60 °C), cellulase exhibits optimal catalytic efficiency, maximizing cell wall breakdown without significant thermal denaturation or degradation of heat-sensitive compounds. However, temperatures above this threshold may denature the enzyme or degrade phenolic and pigment molecules, reducing the overall extract quality. Similarly, the enzyme concentration affects the availability of active sites for substrate hydrolysis. Low to moderate enzyme concentrations improve the extraction yield by increasing the hydrolysis rates, while excessive enzyme amounts can lead to substrate inhibition, enzyme aggregation, or nonspecific degradation of bioactives, resulting in diminished returns.⁴² In summary, cellulase-assisted extraction of roselle calyx bioactives is highly dependent on the fine-tuning of enzyme concentration and extraction temperature. Optimal conditions, particularly around 10% enzyme concentration and 60 °C, maximize the release and stability of phenolics, anthocyanins, flavonoids, and antioxidant capacity while preserving pigment color and pH. These findings highlight the importance of balancing enzymatic hydrolysis and thermal stability to enhance the functional quality of roselle extracts for application in food, nutraceutical, and pharmaceutical products.

3.3.1. Effect of different temperatures and enzyme concentrations on the TPC of cellulase-treated roselle calyx extract. The total phenolic content (TPC) of roselle calyx extract treated with cellulase ranged from 14.00 ± 0.94 to 56.00 ± 0.05 mg GAE/100 g, with significant dependence on both enzyme concentration and extraction temperature. The highest TPC was observed at 10% enzyme concentration and 60 °C, while the lowest occurred at 15% and 70 °C, suggesting that optimal conditions lie within moderate ranges of both variables. Cellulase facilitates TPC extraction by hydrolyzing β -1,4-glycosidic bonds in cellulose, thereby loosening the plant cell wall and enhancing the release of intracellular phenolic compounds.⁴¹ Moderate temperatures (55–65 °C) improve the enzymatic efficiency and solvent penetration, but excessive heat (>70 °C) can lead to enzyme denaturation or phenolic degradation, reducing yields.^{42,46}

The significant quadratic and interaction terms ($p < 0.01$) in the model suggest that the effects of enzyme concentration and temperature are non-linear and interdependent. For example, higher enzyme concentrations at elevated temperatures yielded lower TPC, probably due to substrate saturation and thermal instability of phenolics.^{43,45} These findings underscore the importance of balancing enzyme load and thermal conditions for efficient extraction. The response surface model showed excellent fit ($R^2 = 0.9999$; $CV = 0.140$) (Table 2), confirming the reliability of the results. The model further supports that cellulase-assisted extraction is highly tunable and efficient under optimized conditions. In conclusion, a moderate enzyme concentration (10%) combined with a controlled temperature

(60 °C) yields the highest TPC from roselle calyx. These conditions maximize phenolic release while minimizing degradation, offering a practical and sustainable approach for producing antioxidant-rich extracts for application in functional foods or nutraceuticals.

3.3.2. Effect of different temperatures and enzyme concentrations on the TAC of cellulase-treated roselle calyx extract. The total anthocyanin content (TAC) of roselle calyx extract treated with cellulase ranged from 82.19 ± 0.63 to 127.26 ± 0.74 mg/100 g, with a clear dependence on the enzyme concentration and extraction temperature. The highest TAC was achieved at 17.1% enzyme concentration and 60 °C, indicating that optimal cell wall degradation and anthocyanin release occurred under these conditions. The model showed a good fit ($R^2 = 0.9826$, $CV = 0.218$) (Table 2), with temperature (B), enzyme concentration (A), their interaction ($A \times B$), and the temperature quadratic term (B^2) all showing statistically significant effects ($p < 0.01$), while the enzyme quadratic term (A^2) was not significant. Cellulase promotes anthocyanin release by breaking down the cellulose structure of the plant cell walls, enabling the release of anthocyanins stored in vacuoles.⁴¹ The effectiveness of extraction is enhanced by moderate heating (around 60 °C), which improves cell wall permeability and enzyme activity, without triggering anthocyanin degradation. Anthocyanins are thermolabile; thus, higher temperatures (>70 °C) can lead to pigment decomposition, whereas lower temperatures may hinder sufficient enzyme performance.^{43,46}

A relatively high TAC (117.25 mg/100 g) was also recorded at 15% enzyme concentration and 50 °C, demonstrating that even at lower temperatures, a high enzyme loading can sufficiently degrade plant tissue to release anthocyanins while minimizing the thermal degradation. This suggests that optimal anthocyanin recovery is not solely temperature-dependent but requires a well-balanced enzyme-to-substrate interaction.³⁶ The significant $A \times B$ interaction reflects the synergistic role of enzyme concentration and temperature. For example, high enzyme concentrations are more effective at moderate temperatures but may not yield proportional improvements at elevated temperatures due to the possible anthocyanin breakdown or enzyme inactivation. This behaviour aligns with prior observations in enzyme-assisted extraction systems from other plant sources, where anthocyanin stability limited extraction efficiency at higher processing temperatures.^{44,45} In conclusion, cellulase-assisted extraction significantly improves anthocyanin yield from roselle calyces when applied under optimized conditions. A temperature of 60 °C and an enzyme concentration of 15–17.1% were most effective, achieving high pigment recovery while preserving anthocyanin stability. These findings are important for the development of clean-label natural colorants and antioxidant-rich functional ingredients derived from roselle.

3.3.3. Effect of different temperatures and enzyme concentrations on the antioxidant activity of cellulase-treated roselle calyx extract. The antioxidant activity of cellulase-treated roselle calyx extract varied from $11.88\% \pm 0.31\%$ to $32.35\% \pm 0.91\%$, indicating a substantial influence of enzyme concentration and extraction temperature on bioactive



compound release. The highest antioxidant activity was observed at 15% enzyme concentration and 70 °C, whereas the lowest was recorded at 15% and 50 °C. The regression model showed a strong fit ($R^2 = 0.9735$, $CV = 0.145$) (Table 2), with significant contributions from the linear term of enzyme concentration (A), the quadratic terms (A^2 and B^2), and their interaction ($A \times B$) ($p < 0.01$). However, the linear effect of temperature (B) alone was not statistically significant. The increase in antioxidant activity under optimal conditions can be attributed to the enhanced release of phenolic compounds and anthocyanins, both of which contribute significantly to radical scavenging capacity. Cellulase assists this release by disrupting the cellulase matrix in plant tissues, thereby freeing phenolic antioxidants that are otherwise bound within the cell walls.^{41,42} Interestingly, maximum antioxidant activity did not strictly align with the highest phenolic or anthocyanin contents, highlighting the complex interactions among multiple compounds contributing to antioxidant potential. For example, the antioxidant activity peaked at 15% and 70 °C, while the highest TPC was at 10% and 60 °C. This may be due to differences in the stability, synergistic effects, or bioactivity of individual phenolics and anthocyanins under different conditions.^{43,46}

The significant $A \times B$ interaction (Fig. 3c) indicates that the enzyme concentration modifies how temperature affects the antioxidant activity. For instance, higher enzyme levels may require elevated temperatures to achieve optimal antioxidant release, but excessive heat may also degrade some thermolabile compounds. The presence of a strong quadratic effect further suggests that both under- and over-processing can limit the antioxidant yield, reinforcing the need for a precisely controlled extraction environment.^{36,45} In conclusion, the antioxidant activity of roselle extract can be maximized using cellulase under optimized conditions, specifically at 15% enzyme concentration and 70 °C. While higher enzyme concentrations and temperatures generally enhance antioxidant recovery, a balanced interaction between these factors is critical to avoid compound degradation and ensure functional potency of the extract.

3.3.4. Effect of different temperatures and enzyme concentrations on the TFC of cellulase-treated roselle calyx extract. The total flavonoid content (TFC) of cellulase-treated roselle calyx extract showed significant variation, ranging from 75.47 ± 0.37 to 195.00 ± 0.11 mg CE/100 g, strongly influenced by both the temperature and the enzyme concentration. The optimal TFC was obtained at 10% enzyme concentration and 60 °C, while the lowest was recorded at 17.1% and 60 °C. The regression model demonstrated excellent fit and predictive power with $R^2 = 0.9999$ and $CV = 0.695$ (Table 2), and all terms in the model (A , B , A^2 , B^2 , and $A \times B$) were highly significant ($p < 0.01$). Cellulase enhances flavonoid extraction by hydrolyzing cellulose and weakening the plant cell wall, facilitating the release of vacuole-bound flavonoids.⁵² The increase in TFC at moderate enzyme concentrations (10%) and optimal temperatures (60 °C) suggest a balance between effective cell wall breakdown and compound stability. Higher enzyme concentrations, as observed at 17.1%, may lead to excessive breakdown or oxidation, resulting in lower TFC due to flavonoid degradation or structural alteration.^{41,42}

The strong quadratic effects (A^2 and B^2) indicate that both under- and over-processing (low or high enzyme levels and temperatures) can reduce the flavonoid yield. This aligns with previous reports where enzymatic overactivity or thermal stress degraded sensitive phenolic structures.^{43,44} Additionally, the interaction effect ($A \times B$) suggests that temperature modulates the enzyme's efficiency, where higher temperatures can boost enzyme activity to a point, but beyond optimal levels, the degradation of heat-sensitive flavonoids probably offsets the benefits. The plateau in TFC values at 195.00 mg CE/100 g across multiple replicates at 10% and 60 °C indicates a saturation point beyond which no additional release occurs, possibly due to extraction equilibrium.^{36,45} In conclusion, cellulase-assisted extraction significantly enhances flavonoid recovery from roselle calyces when the enzyme concentration and temperature are optimized. The best conditions observed were 10% cellulase at 60 °C, where the maximum TFC was consistently achieved. These findings are crucial for designing efficient, enzyme-based extraction protocols for natural antioxidants for application in functional foods and nutraceuticals.

3.3.5. Effect of different temperatures and enzyme concentrations on the hue angle of cellulase-treated roselle calyx extract. The hue angle, indicative of color perception and pigment stability in roselle extract, ranged from 17.92 ± 0.73 to 27.22 ± 0.34 across different temperatures and enzyme concentrations. The highest hue angle, reflecting a shift toward more orange tones, was recorded at 10% cellulase and 60 °C, while the lowest values were observed under higher enzyme concentrations and lower temperatures. The fitted quadratic model for hue angle demonstrated high predictive accuracy with $R^2 = 0.9920$ and $CV = 0.413\%$ (Table 2). Significant effects were observed from both linear and quadratic terms for the enzyme concentration and temperature ($p < 0.01$), as well as their interaction, highlighting a complex interplay between thermal and enzymatic factors in pigment stability and color expression. Higher temperatures generally promoted increased hue angle, possibly due to enhanced pigment solubilization and reduced co-pigmentation. However, excessive enzyme activity (particularly at >15% concentration) may have led to degradation or transformation of anthocyanins, contributing to less desirable colour changes. This is consistent with previous findings that indicate that the enzymatic breakdown of cell wall structures can release both stable and unstable anthocyanins, which may be further altered by thermal exposure.^{44,47}

The significant interaction between the temperature and the enzyme concentration ($A \times B$) suggests that the temperature amplifies the enzymatic impact on colour. At optimal moderate enzyme levels (10%) and elevated temperatures (60 °C), the hue angle peaked, potentially due to maximized anthocyanin release and stabilization. In contrast, either too low or too high enzyme levels, combined with suboptimal temperatures, resulted in hue reduction, likely linked to pigment oxidation or polymerization.³⁶ In conclusion, the hue angle of cellulase-treated roselle extract is significantly affected by extraction conditions. A moderate enzyme concentration (10%) combined with elevated temperatures (60 °C) preserves the most vivid coloration by



optimizing the anthocyanin extraction while minimizing degradation. These findings are crucial for industrial applications aiming to retain the visual appeal and functional colour properties of roselle-based products.

3.3.6. Effect of different temperatures and enzyme concentrations on the pH of cellulase-treated roselle calyx extract. The pH values of cellulase-treated roselle calyx extracts ranged from 2.28 ± 0.47 to 2.90 ± 0.02 , indicating the overall acidic nature of the extract, which is characteristic of roselle due to its organic acid content. The highest pH was observed at 10% cellulase and 60 °C, suggesting a potential buffering or reducing effect of enzymatic treatment on acidity under optimal thermal conditions. The fitted model showed high significance ($p < 0.01$) for both linear and quadratic terms of enzyme concentration and temperature, as well as for their interaction. The model's coefficient of determination ($R^2 = 0.9988$) and low coefficient of variation ($CV = 0.011\%$) (Table 2) confirm the reliability and accuracy of the data and model. Increased enzyme concentration initially led to a slight decrease in pH, possibly due to enhanced release of acidic compounds like organic acids or polyphenols during cell wall degradation. However, at an optimal combination of 10% enzyme concentration and 60 °C, the pH increased, probably because of the liberation of buffering components or due to dilution effects from higher extraction yields. This pattern aligns with previous findings showing that enzymatic treatment modifies the balance of acidic and non-acidic phytochemicals in plant extracts.^{41,43} Temperature played a synergistic role, where moderate to high heat levels likely enhanced the enzyme activity up to a threshold. Beyond that, enzyme denaturation or degradation of acidic compounds may have influenced pH stability. The significant interaction effect ($A \times B$) confirms that the combined influence of temperature and enzyme concentration critically dictates the final pH of the extract. In conclusion, both enzyme concentration and temperature significantly influence the pH of cellulase-treated roselle extract. Maintaining optimal conditions (10% enzyme, 60 °C) ensures balanced extraction while minimizing excessive acidity, which is beneficial for enhancing extract stability and improving sensory acceptance for application in food products and beverages.

3.4. Comparative analysis of amylase, cellulase, and pectinase under optimized conditions

The enzyme-assisted extraction of roselle calyx bioactives reveals that the nature of the enzyme and extraction conditions

critically influence the yield and profile of phenolic compounds, anthocyanins, antioxidant activity, flavonoids, color attributes, and pH. The comparative analysis results are listed (Table 3). Cellulase, applied at 10% concentration and 60 °C, yielded the highest total phenolic content (55.34 mg GAE/100 g) and total flavonoid content (193.05 mg CE/100 g). Cellulase hydrolyzes β -1,4-glycosidic bonds in cellulose, a major structural polysaccharide in plant cell walls. By breaking down cellulase microfibrils, cellulase disrupts the cell wall matrix by facilitating the release of intracellular and cell wall-bound phenolic and flavonoid compounds.^{41,42} This enzymatic degradation increases cell wall porosity and extractability, which is consistent with the enhanced phenolic recovery reported in green extraction methods.³⁶ The increase in flavonoids, a subclass of phenolics often bound to polysaccharides, further supports cellulase's role in liberating complexed bioactives. Pectinase at 15% and 50 °C was most effective in extracting anthocyanins (127.79 mg/100 g), pigments responsible for the characteristic red hue of roselle. Pectinase catalyzes the breakdown of pectin, a polysaccharide present in the middle lamella of plant cells, which acts as a cementing agent holding cells together.⁴⁴ The degradation of pectin loosens cell adhesion and enhances pigment release from vacuoles, which are the primary storage sites of anthocyanins. This mechanism explains the superior anthocyanin content and the lower hue angle observed with pectinase treatment, indicating a more vivid red coloration.^{45,48} Additionally, pectinase treatment can improve the juice yield and clarity, beneficial for industrial processing.⁴²

Amylase, used at 3% concentration and 60 °C, led to the highest antioxidant activity (39.14%). Amylase hydrolyzes starch molecules into simpler sugars such as maltose and glucose, which may contribute to increased antioxidant potential either by releasing antioxidant-bound polysaccharides or through the formation of Maillard reaction products during processing, known for antioxidant properties.^{43,46} Although amylase showed lower phenolic and flavonoid extraction than cellulase, its enhancement of antioxidant capacity suggests a complementary mechanism where starch breakdown supports the availability or stability of antioxidant compounds. The pH values observed in cellulase-treated extracts were higher (around 2.9), which could influence anthocyanin stability and color expression, as anthocyanins are pH-sensitive pigments.¹⁹ Pectinase-treated extracts exhibited lower pH values, potentially preserving anthocyanin stability and enhancing colour

Table 3 Comparative analysis of the effects of amylase, cellulase, and pectinase at optimized conditions in enhancing the enzyme-assisted extraction of roselle calyx^a

Enzymes	Enzyme concentration (%)	Temperature (°C)	TPC (mg GAE/100 g)	TAC (mg/100 g)	Antioxidant (%)	TFC (mg CE/100 g)	Hue angle	pH
Amylase	3	60	35.09 ± 1.20	99.65 ± 0.46	39.14 ± 0.11	151.88 ± 0.03	17.93 ± 1.16	2.37 ± 0.13
Pectinase	15	50	33.35 ± 0.34	127.79 ± 0.48	28.95 ± 0.41	133.07 ± 0.88	17.32 ± 1.02	2.24 ± 0.38
Cellulase	10	60	55.34 ± 0.17	102.36 ± 0.52	32.36 ± 0.07	193.05 ± 0.32	27.08 ± 0.53	2.91 ± 0.29

^a TPC – total phenolic content (mg GAE/100 g); TAC – total anthocyanin content (mg/100 g); TFC – total flavonoid content (mg CE/100 g); AOX – antioxidant activity (%); h° – hue angle (°); results are expressed as mean ± SD ($n = 3$).



intensity. In summary, in the differential enzymatic actions, cellulase breaks down cellulose, pectinase degrades pectin, and amylase hydrolyzes starch, and each facilitates the liberation of specific bioactive compounds from the roselle calyx matrix. This specificity underscores the importance of enzyme selection based on target compounds and desired extract qualities.⁴¹ Integrating these enzymes sequentially or simultaneously could synergistically improve the extraction efficiency, maximizing the yield and functional properties of Roselle extracts for applications in food, nutraceutical, and pharmaceutical products.

3.5. Comparative analysis of amylase, cellulase, and pectinase under optimized conditions infused in ice creams

The integration of amylase, cellulase, and pectinase enzymes into roselle-infused ice creams led to distinct differences in colour intensity, texture, structure, and air incorporation, which are visually represented (Table 4). These changes are attributed to the enzymatic breakdown of complex macromolecules (starch, cellulase, and pectin), which significantly influenced the physicochemical and sensory characteristics of the final product. The amylase-treated ice cream exhibited a light pink hue with a porous, aerated surface. Amylase acts on starches

present in dairy and plant-derived ingredients by hydrolyzing α -1,4-glycosidic bonds, converting polysaccharides into maltose and glucose. This reaction not only increases the sweetness of the base but also reduces the viscosity, leading to improved aeration and overrun during churning.⁵³ The visible fine air bubbles (Table 4) suggest enhanced emulsification due to reduced starch thickness, allowing better dispersion of fat and water phases. The degradation of starch reduces its water-holding capacity, enabling better freezing and a smoother melt-down profile. The increased sugar concentration due to amylolysis also acts as a cryoprotectant, influencing freezing point depression, which contributes to the softer texture observed.⁵⁴ However, excessive amylase activity may lead to overly thin consistency, which is why optimization is crucial.

The cellulase-infused sample showed a deep red, dense ice cream matrix with minimal air incorporation. This enzyme hydrolyzes β -1,4-glycosidic bonds in cellulase, producing glucose or cellobiose from plant-based fibers in the roselle calyces. The breakdown of cell wall components facilitates the release of intracellular pigments, including anthocyanins, which account for the pronounced red coloration.⁵⁵ This cellulolytic action improves colorant bioavailability and may enhance antioxidant retention. Additionally, the liberated soluble fibers increase water-binding and viscosity, contributing to a firmer, gel-like consistency with minimal overrun.³⁷ The limited presence of air bubbles suggests the reduced ability of the mixture to entrap air during freezing, probably due to increased viscosity, which restricts the formation of a lighter structure. The mechanism here reflects dual benefits: improved colour extraction from roselle tissues and structural enhancement due to fiber solubilization. However, excessive viscosity may lead to a heavy mouthfeel, which should be balanced for consumer acceptability.

The pectinase-treated ice creams exhibited a uniform pink tone and a visibly smoother surface, with evenly distributed micro-bubbles indicating excellent aeration and emulsification. Pectinase catalyzes the hydrolysis of α -1,4-glycosidic bonds in pectic substances, particularly polygalacturonic acid chains found in plant cell walls. This depolymerization facilitates the release of water-soluble anthocyanins and flavonoids, enhancing color uniformity.⁵⁶ The degradation of high-molecular-weight pectins lowers viscosity, improves flowability, and enhances the interaction between air, fat, and water during churning. The result is a creamier texture, better aeration, and a refined mouthfeel.⁵³ However, excessive pectin degradation can reduce the body and stability of the ice cream, though under this optimized condition, a balanced structure was maintained. Additionally, the enhanced diffusion of bioactive compounds improves the product's antioxidant profile and visual appeal, both essential for consumer preference and health-oriented formulations.⁵⁴

These findings are consistent with previous studies highlighting the benefits of enzyme-assisted extraction and modification in functional food systems. Gengatharan⁵⁶ and Bandara & Mahendran⁵⁵ showed that pectinase and cellulase, respectively, improve pigment extraction and product texture. Similarly, Homayouni Rad *et al.*⁵³ confirmed that enzyme-treated ice

Table 4 Ice cream samples containing different concentrations of roselle calyx extract




S. no.	Enzyme used	Flavoured ice cream images
1	Amylase-infused roselle bio-colourant	
2	Cellulase-infused roselle bio-colourant	
3	Pectinase-infused roselle bio-colourant	



Table 5 Colour values of the roselle-flavoured ice creams^a

Sample	L^*	a^*	b^*	WI	Hue (°)
Control	72.50 ^a ± 0.30	1.10 ^a ± 0.10	5.30 ^a ± 0.20	89.00 ^a ± 0.40	87.80 ^a ± 0.50
ICA	71.70 ^a ± 0.40	0.07 ^a ± 0.10	2.80 ^a ± 0.20	78.20 ^a ± 0.45	73.30 ^a ± 0.40
ICC	93.00 ^b ± 0.20	0.67 ^a ± 0.10	0.50 ^b ± 0.10	60.50 ^b ± 0.35	62.10 ^a ± 0.30
ICP	82.20 ^a ± 0.30	2.20 ^a ± 0.10	0.43 ^b ± 0.20	82.50 ^b ± 0.40	29.90 ^a ± 0.40

^a L^* – lightness; a^* – red-green value; b^* – yellow-blue value; WI – whiteness index; hue (°) – colour tone. Results are expressed as mean ± SD ($n = 3$). Different superscript letters within a column indicate significant differences ($p < 0.05$).

cream formulations using plant extracts yield better sensory and nutritional outcomes. The application of food-grade enzymes (amylase, cellulase, and pectinase) in roselle-colored ice creams not only improved the aesthetic quality (colour and uniformity), but also affected physical properties such as texture, viscosity, and aeration. The underlying biochemical mechanisms of each enzyme significantly influence the matrix behaviour, making them valuable tools in the formulation of clean-label, functional ice creams. Optimal enzyme use allows the tailoring of product attributes based on consumer preferences and nutritional goals. Overall, enzymatic infusion improved the colour extraction, texture, and visual appeal of roselle bio-colorant ice creams, with each enzyme offering distinct advantages. Cellulase provided the most vibrant colour and dense structure, with a smoother and more aerated product, which contributed to a balanced sweetness and moderately light texture.

3.5.1. Colour and sensory attributes of amylase, cellulase, and pectinase under optimized conditions infused in ice creams. The infusion of amylase (ICA), cellulase (ICC), and pectinase (ICP) into roselle-flavoured ice creams significantly altered both colour intensity and sensory perception compared to the untreated control (Tables 5 and 6). Instrumental colour analysis revealed distinct differences among treatments. The control sample showed moderate lightness ($L^* = 72.50$), mild redness ($a^* = 1.10$), and the highest whiteness index (WI = 89.00), indicating limited anthocyanin extraction. Amylase-treated ice creams (ICA) had slightly lower lightness ($L^* = 71.70$) and a very low a^* value (0.07), with a hue angle of 73.30°, indicating a pale pink tone. This is attributed to amylase's role

in hydrolyzing starch into maltose and glucose, which enhanced sweetness (score: 8.0) but had minimal effects on pigment release due to a lack of direct interaction with plant cell walls.⁵³ In contrast, cellulase-treated ice creams (ICC) exhibited the highest lightness ($L^* = 93.00$) but the lowest WI (60.50), suggesting a bright yet pigment-dense matrix. This is due to cellulase breaking down β -1,4 linkages in cellulose, facilitating anthocyanin release from roselle tissues, consistent with Bandara *et al.*⁵⁵ ICC also had improved redness ($a^* = 0.67$) and reduced b^* and hue angle values, contributing to a visually rich, deep red colour. These enhancements were reflected in sensory scores, with ICC receiving the highest ratings for appearance (8.0), texture (8.2), creaminess (8.0), and overall acceptability (8.0), indicating that enhanced colour release and fiber solubilization contributed to a smooth, dense mouthfeel.³⁷

Pectinase-treated ice creams (ICP) showed a strong red colour ($a^* = 2.20$) with the lowest hue angle (29.90°), indicating the optimal anthocyanin release. This aligns with the enzyme's ability to degrade pectic polysaccharides, loosening plant cell structures and enhancing pigment diffusion.⁵⁶ ICP also maintained high lightness ($L^* = 82.20$) and moderate WI (82.50), resulting in a uniformly coloured and creamy appearance. Sensory attributes such as texture (7.8), appearance (7.1), and sweetness (7.6) were favorable, though slightly lower creaminess (7.0) and milky perception (7.2) suggested that excessive pectin breakdown may have reduced stabilizing effects within the matrix. These findings demonstrate that cellulase had the most pronounced effect on overall quality, improving both pigment extraction and sensory texture. Amylase enhanced sweetness naturally but contributed least to colour development. Pectinase was most effective in enhancing red pigmentation, but slightly compromised the body. These enzyme-specific effects on starch, cellulase, and pectin structures directly influenced anthocyanin release, viscosity, texture, and flavour perception, supporting previous research on enzyme-assisted functional ice creams.^{37,53–56}

3.6. Spectroscopic and chromatographic analysis

3.6.1. HPLC analysis of cellulase-assisted roselle extract. The high-performance liquid chromatography (HPLC) analysis of the cellulase-assisted roselle (*Hibiscus sabdariffa*, L) extract revealed two prominent peaks with retention times of 4.001 and 4.019 minutes, respectively. Both peaks were identified as quercetin, confirmed by comparison with the quercetin standard, which exhibited a retention time of 4.019 minutes (Table

Table 6 Sensory attributes of the roselle-flavoured ice creams^a

Sample	Control	ICA	ICC	ICP
Appearance	6.9 ± 0.30	7.2 ± 0.42	8.0 ± 0.47	7.1 ± 0.30
Flavor	7.1 ± 0.30	7.0 ± 0.00	7.5 ± 0.00	7.0 ± 0.00
Texture	7.4 ± 0.52	8.1 ± 0.52	8.2 ± 0.42	7.8 ± 0.42
Creamy	7.1 ± 0.30	7.0 ± 0.00	8.0 ± 0.42	7.0 ± 0.00
Sweetness	8.0 ± 0.30	7.0 ± 0.00	7.0 ± 0.00	7.6 ± 0.50
Milky	8.4 ± 0.52	7.6 ± 0.50	7.4 ± 0.33	7.2 ± 0.47
Overall acceptability	7.8 ± 0.42	7.4 ± 0.50	8.0 ± 0.00	7.4 ± 0.00

^a ICA – amylase-infused ice cream; ICC – cellulase-infused ice cream; ICP – pectinase-infused ice cream. Results are expressed as mean ± SD ($n = 3$).



Table 7 HPLC analysis of the cellulase-assisted extraction of roselle calyx

Peak	Retention time (min)	Area	Height	Relative concentration (%)	Assignment
1	4.001	64 556 600	53 330	52.87	Quercetin
2	4.019	57 554 320	16 833	47.13	Quercetin standard

Table 8 FTIR peaks and corresponding functional groups in the roselle calyx extract

Wavenumber (cm ⁻¹)	Assigned functional group	Possible compound class
420.28	Aromatic skeletal vibration/ring deformation	Aromatic phenolic rings, anthocyanidin core structures, mineral-organic complexes
1633.00	C=O (amide I) stretching and C=C aromatic stretching	Protein amide groups, hydroxycinnamic acids, anthocyanins, conjugated phenolic structures
2163.55	Weak C≡N or C≡C-related overtone/composition band	Trace nitrogenous residues, minor plant secondary metabolites, or instrumental artifacts
3303.30	O-H and N-H stretching (broad, H-bonded)	Flavonoids, anthocyanins (glycosylated OH), polysaccharide hydroxyls, proteinaceous NH groups

7). The relative concentration of quercetin in the extract was approximately 52.87%, closely matching the 47.13% relative concentration of the quercetin standard peak. This high concentration highlights the efficiency of cellulase-assisted extraction in releasing quercetin from roselle calyces. The enzymatic treatment probably facilitated the breakdown of the plant cell wall matrix, thus enhancing the release and subsequent quantification of bioactive flavonoids like quercetin.^{41,42} The previous studies have demonstrated that enzyme-assisted extraction improves the yield and bioavailability of polyphenolic compounds in roselle by targeting polysaccharides and cellulase structures within the plant matrix.^{36,48} The retention times and peak characteristics observed are consistent with those reported for quercetin, a flavonoid known for its antioxidant, anti-inflammatory, and antibacterial properties.^{43,45} Furthermore, the similarity in peak areas and retention times between the sample and standard indicates minimal degradation or alteration of quercetin during the enzymatic extraction process. This aligns with the findings of Chenxin *et al.*,⁴⁴ who reported that optimized extraction conditions preserve anthocyanins and flavonoids in roselle extracts. The presence of high quercetin content corroborates the antioxidant potential reported in enzymatically extracted roselle calyx samples, contributing to the functional benefits attributed to roselle for application in food and pharmaceutical products.^{46,47} The cellulase-assisted extraction method thus represents a green and efficient technique to obtain high-quality bioactive compounds, supporting sustainability and potential commercial scalability.¹⁹ In summary, the HPLC analysis confirms that cellulase-assisted extraction effectively liberates quercetin from roselle calyces with high purity and concentration. This enzymatic method enhances the extraction yield of valuable flavonoids, supporting the development of functional foods and nutraceuticals based on roselle bioactives.

3.6.2. FTIR spectroscopic analysis of cellulase-assisted roselle extract. The Fourier transform infrared (FTIR) spectrum of the cellulase-assisted roselle extract revealed characteristic bands corresponding to several functional groups, indicative of the complex chemical composition of the extract (Table 8 and Fig. 4). The broad peak at 3303.30 cm⁻¹ corresponds to O-H and N-H stretching vibrations, typically attributed to alcohols, phenolic compounds, and amines. This broad band suggests the presence of hydroxyl groups, a hallmark of polyphenols and flavonoids in roselle extract.^{36,41} The absorption at 1633.00 cm⁻¹ can be assigned to C=O stretching (amide I) and C=C stretching in aromatic rings, which is characteristic of phenolic acids and residual proteinaceous components. This aligns with previous reports that enzymatic extraction can liberate phenolics and other associated bioactive compounds.^{43,44} The absorption features in the region of 1200–




Fig. 4 FTIR spectra of cellulase-assisted roselle extract infused ice cream, showing characteristic absorption bands.






Fig. 5 SEM images of the cellulase-assisted roselle extract-infused ice cream: (a) 100 \times magnification and (b) 250 \times magnification.

1000 cm^{-1} correspond to the C–O and C–O–C stretching vibrations, typical of carbohydrates, glycosides, and polysaccharide structures present in roselle. The weak band in the lower wavenumber region around 420.28 cm^{-1} represents skeletal vibrations of complex biomolecular matrices or mineral-associated signals commonly found in plant-based extracts.⁴⁵ Overall, the FTIR spectrum confirms that cellulase-assisted extraction of roselle calyces retains a diverse range of functional groups associated with phenolic, proteinaceous, and carbohydrate-based compounds. This molecular diversity supports the enhanced bioactivity of roselle extracts obtained through enzymatic treatment, as documented in related studies.^{19,47}

3.6.3. SEM analysis of cellulase-assisted roselle ice creams.

The scanning electron microscopy (SEM) analysis provides insights into the microstructure of ice creams infused with cellulase-extracted roselle extract. At 100 \times magnification (Fig. 5a), clusters of irregularly shaped plant-derived particles are visible within the ice cream matrix. These aggregates probably represent partially hydrolyzed roselle fibre fragments and phenolic-rich residues carried into the formulation through the extract. This may probably result from the increased soluble solids generated during cellulase-assisted hydrolysis.⁵⁷ Their distribution indicates the successful incorporation of the extract components into the frozen structure without visible aggregation. At 250 \times magnification (Fig. 5b), the particles appear more clearly defined, with smoother surfaces and spherical morphologies. The surrounding matrix appears continuous, suggesting that the roselle-derived micro-particles are well integrated into the ice cream structure. The smaller particle size produced through enzymatic pretreatment facilitates more uniform dispersion within the frozen matrix, preventing local clustering or structural discontinuities.⁵⁸ The presence of fine, evenly distributed fragments is consistent with cellulase-assisted extraction, which reduces plant materials into more dispersible structures before infusion.⁵⁹ Uniformly distributed fine fragments can also provide moderate local water mobility during freezing, thereby helping to reduce ice-crystal growth and attain a smoother texture in the final

product.⁶⁰ Although SEM does not provide chemical identification,⁶¹ the visible structures correspond to roselle-derived plant fragments embedded within the ice cream matrix, and their integration suggests that key bioactive components remain structurally retained during freezing.

4 Conclusion

This study demonstrated that enzyme-assisted extraction using amylase, pectinase, and cellulase significantly enhances the recovery of bioactive compounds from *Hibiscus sabdariffa* L. calyces compared to conventional techniques. Among the enzymes, cellulase produced the highest total phenolic and flavonoid yields, pectinase maximized anthocyanin recovery and colour intensity, while amylase enhanced antioxidant activity. Incorporating enzyme-treated roselle extracts into ice cream formulations improved the colour uniformity, texture, and sensory acceptability, demonstrating practical potential for developing functional and visually appealing food products. The spectroscopic and chromatographic analysis confirmed the preservation of key compounds such as quercetin, validating the efficiency of the optimized enzymatic process. Overall, the study highlights enzyme-assisted extraction as an eco-friendly, scalable, and energy-efficient approach for producing natural colorants and antioxidant-rich ingredients from roselle. The future research should focus on scaling up extraction for industrial applications, exploring synergistic enzyme combinations and assessing the stability and bioavailability of extracted compounds during product processing and storage. Additionally, integrating advanced green technologies such as ultrasound or microwave-assisted enzymatic extraction could further enhance the yield and sustainability, paving the way for broader use of roselle in functional foods, nutraceuticals, and natural product formulations.

Consent for publication

All authors agreed on the publication of this research work.



Author contributions

Priyanka A: conceptualization, methodology, writing – original draft. Nila Shireen B: validation, writing – original draft, formal analysis. Ganga Kishore: investigation, validation, writing – original draft, formal analysis. Sudha Paramadhas: conceptualization, investigation, supervision, writing – original draft, formal analysis. S. R. Rahul: writing – original draft, validation, formal analysis. Madhuresh Dwivedi: methodology, writing – original draft, review and editing, formal analysis. Pandiarajan T: supervision. Divyabharathi R: validation, investigation, formal analysis. Nongmaithem Seema Chanu: formal analysis.

Conflicts of interest

The authors declare no competing interest.

Abbreviations

<i>Hibiscus sabdariffa</i> L.	<i>Hibiscus sabdariffa</i> Linn.
HCl	Hydrochloric acid
HPLC	High-performance liquid chromatography
FTIR	Fourier transform infrared
SEM	Scanning electron microscope
EAE	Enzyme-assisted extraction
DPPH	2,2-Diphenyl-1-picrylhydrazyl
TPC	Total phenol content
TFC	Total flavonoid content
TAC	Total anthocyanin content
AOX	Antioxidant activity
CE	Catechin equivalent
MSNF	Milk solid-not-fat
ICA	Ice cream with amylase-treated extract
ICC	Ice cream with cellulase-treated extract
ICP	Ice cream with pectinase-treated extract
cyn-3-sam	Cyanidin-3-sambubioside
dp-3-glu	Delphinidin-3-glucoside
cyn-3-glu	Cyanidin-3-glucoside
CCD	Central composite design
ANOVA	Analysis of variance
FWHM	Full width at half maximum

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgements

Funded by the ICAR-AICRP-PHET scheme.

References

- 1 J. Golijan and B. Dimitrijević, *Acta Agric. Serb.*, 2018, **23**, 125–140.
- 2 L. T. Reynolds, *World Dev.*, 2004, **32**, 725–743.
- 3 R. Banc, L. Filip, A. Cozma-Petruț, D. Ciobârca and D. Miere, *Bull. Univ. Agric. Sci. Vet. Med. Cluj-Napoca. Food Sci. Technol.*, 2024, **81**, 1–17.
- 4 R. Bocker and E. K. Silva, *Food Chem.:X*, 2022, **15**, 100398.
- 5 N. Li, J. E. Simon and Q. Wu, *Food Chem.*, 2024, **461**, 140782.
- 6 H. A. Sindi, L. J. Marshall and M. R. A. Morgan, *Food Chem.*, 2014, **164**, 23–29.
- 7 T. Estiasih, J. E. Witoyo, G. K. M. Runtung, D. D. P. Moelyono, A. R. Murini, W. Setyaningsih and K. Ahmadi, *CyTA-J. Food*, 2024, **22**, 2356847.
- 8 A. A. Abou-Arab, F. M. Abu-Salem and E. A. Abou-Arab, *Am. J. Sci.*, 2011, **7**, 445–456.
- 9 P. J. Tsai, J. McIntosh, P. Pearce, B. Camden and B. R. Jordan, *Food Res. Int.*, 2002, **35**, 351–356.
- 10 N. Mahadevan, S. Shivali and P. K. Pradeep Kamboj, *Nat. Product. Radiance*, 2009, **8**, 77–83.
- 11 M. Cissé, P. Bohuon, F. Sambe, C. Kane, M. Sakho and M. Dornier, *J. Food Eng.*, 2012, **109**, 16–21.
- 12 M. Dyrby, N. Westergaard and H. Stapelfeldt, *Food Chem.*, 2001, **72**, 431–437.
- 13 M. Mardiah, N. Novidahlia, M. Khoirunnisa, H. Hanafi and A. Aminullah, *J. Teknol. Ind. Has. Pertan.*, 2021, **26**, 65–71.
- 14 N. Stanek-Wandzel, A. Krzyszowska, M. Zarębska, K. Gębura, T. Wasilewski, Z. Hordyjewicz-Baran and M. Tomaka, *Int. J. Mol. Sci.*, 2024, **25**, 13538.
- 15 M. Spagnuolo, C. Crecchio, M. D. R. Pizzigallo and P. Ruggiero, *Bioresour. Technol.*, 1997, **60**, 215–222.
- 16 F. Chemat, N. Rombaut, A. G. Sicaire, A. Meullemiestre, A. S. Fabiano-Tixier and M. Abert-Vian, *Ultrason. Sonochem.*, 2017, **34**, 540–560.
- 17 A. Escobar-Ortiz, E. Castaño-Tostado, N. E. Rocha-Guzmán, J. A. Gallegos-Infante and R. Reynoso-Camacho, *J. Sci. Food Agric.*, 2021, **101**, 110–119.
- 18 M. M. K. Chowdhury, S. Chowdhury, L. Bari and S. Yeasmin, *Dhaka Univ. J. Biol. Sci.*, 2024, **33**, 139–147.
- 19 K. N. Yulianda Setyawan and K. Kartini, *J. Appl. Biol. Biotechnol.*, 2023, **11**, 91–97.
- 20 A. A. Harahap, K. Letare S and N. Hendrianie, *AIP Conf. Proc.*, 2023, **2667**, 020003.
- 21 A. N. Nunes, A. Borges, A. A. Matias, M. R. Bronze and J. Oliveira, *Molecules*, 2022, **27**, 368.
- 22 G. Linares and M. L. Rojas, *Front. Nutr.*, 2022, **9**, 891462.
- 23 D. L. Nonglait and J. S. Gokhale, *Food Bioprocess Technol.*, 2024, **17**, 1681–1705.
- 24 I. Benucci, C. Lombardelli, C. Mazzocchi and M. Esti, *Compr. Rev. Food Sci. Food Saf.*, 2022, **21**, 2715–2737.
- 25 J. J. Liu, M. A. A. Gasmalla, P. Li and R. Yang, *Innovative Food Sci. Emerging Technol.*, 2016, **35**, 184–193.
- 26 K. Rafińska, O. Wrona, A. Krakowska-Sieprawska, J. Walczak-Skierska, A. Kielbasa, Z. Rafiński and B. Buszewski, *Ind. Crops Prod.*, 2022, **187**, 115424.
- 27 D. Alavarsa-Cascales, M. J. Aliaño-González, M. Palma, G. F. Barbero and C. Carrera, *Agronomy*, 2022, **12**, 2327.
- 28 M. José Aliaño González, C. Carrera, G. F. Barbero and M. Palma, *Food Chem.:X*, 2022, **13**, 100192.



- 29 H. T. Ngoc Nhon, N. T. Diem My, V. N. Tuong Vi, P. T. Kim Lien, N. T. Thao Minh, L. N. Doan Duy, L. T. Hong Anh, D. T. Anh Dao and J. Agric, *Food Res.*, 2022, **10**, 100408.
- 30 P. R. Amulya and R. ul Islam, *Food Chem.:X*, 2023, **18**, 100643.
- 31 K. Duangmal, B. Saicheua and S. Sueeprasan, *Proceedings of the AIC*, 2004, pp. 155–158.
- 32 M. V. de Paiva, M. Z. Sanches, L. G. Chies, A. L. Vieira, J. F. Nicoleti and L. S. Sakanaka, *Res. Soc. Dev.*, 2021, **10**, e12910212224.
- 33 T. K. Koley, C. Kaur, S. Nagal, S. Walia and S. Jaggi, *Arabian J. Chem.*, 2016, **9**, S1044–S1052.
- 34 L. Jakobek and P. Matić, *Croat. J. Food Sci. Technol.*, 2021, **13**, 176–183.
- 35 R. E. Wrolstad, R. W. Durst and J. Lee, *Trends Food Sci. Technol.*, 2005, **16**, 423–428.
- 36 N. Chongwilaikasem, P. Sithisarn, P. Rojsanga and P. Sithisarn, *J. Food Sci.*, 2024, **89**, 8819–8835.
- 37 T. M. Singo and D. Beswa, *Int. J. Food Prop.*, 2019, **22**, 42–53.
- 38 A. Al-Rifai, A. Aqel, A. Awaad and Z. A. AlOthman, *Commun. Soil Sci. Plant Anal.*, 2015, **46**, 1411–1418.
- 39 N. Montoya-Escobar, D. Ospina-Acero, J. A. Velásquez-Cock, C. Gómez-Hoyos, A. Serpa Guerra, P. F. Gañan Rojo, L. A. Vélez Acosta, C. Castro Herazo and R. Zuluaga Gallego, *Polymers*, 2022, **14**, 5199.
- 40 U. Pithanthanakul, S. Vatanyoopaisarn, B. Thumthanaruk, C. Puttanlek, D. Uttapap, B. Kietthanakorn and V. Rungsardthong, *Flavour Fragrance J.*, 2021, **36**, 365–373.
- 41 L. A. Cira-Chávez, L. E. Gassós-Ortega, J. Nuñez-Vega, G. A. Camelo-Méndez, R. Cañedo-Urias, A. K. Blanco-Rios and M. I. Estrada-Alvarado, *Appl. Food Res.*, 2025, **5**, 101149.
- 42 M. Mardiah, M. A. Fulazzaky, S. I. Rahmawati and M. Fulazzaky, *Next Res.*, 2025, **2**(3), 100536.
- 43 Y.-J. Lai, Y.-C. Chiang, Y.-S. Jhan, T.-Y. Song and M.-C. Cheng, *Foods*, 2024, **13**, 2172.
- 44 L. U. Chenxin, L. I. Yangxinhui, Y. Jinyi, G. Longyao and Z. Zhen, *Food Mach.*, 2025, **40**, 32–42.
- 45 U. K. M. Bangi, *Sains Malays.*, 2023, **52**, 3147–3162.
- 46 T. Ahmed, M. R. Rana, M. A. Hossain, S. Ullah and M. Suzauddula, *Biomass Convers. Biorefin.*, 2024, **14**, 28985–28999.
- 47 N. N. Aneke, W. I. Okonkwo, S. L. Ezeoha, G. I. Okafor and C. N. Anyanwu, *Res. Innov. Food Sci. Technol.*, 2023, **11**, 437–450.
- 48 F. D'Heureux-Calix and N. Badrie, *Food Serv. Technol.*, 2004, **4**, 141–148.
- 49 A. Rani, Y. Arfat, R. S. Aziz, L. Ali, H. Ahmed, S. Asim, M. Rashid and C. H. Hocart, *Environ. Technol. Innov.*, 2021, **23**, 101620.
- 50 F. Weber and L. R. Larsen, *Food Res. Int.*, 2017, **100**, 354–365.
- 51 H.-K. Tran, T.-T.-H. Nguyen, T. T. Huynh, N. M. T. Vo, H. O. Le, D.-H. Truong, H.-C. Nguyen and C. J. Barrow, *Int. J. Food Sci. Technol.*, 2025, **60**(1), DOI: [10.1093/ijfood/vvaf057](https://doi.org/10.1093/ijfood/vvaf057).
- 52 L. Wang, Y. Wu, Y. Liu and Z. Wu, *Molecules*, 2017, **22**, 1648.
- 53 A. Homayouni Rad, F. Karbalaee, M. A. Torbati, M. Moslemi, F. Shahraz, R. Babadi and M. Javadi, *J. Food Sci. Technol.*, 2022, **59**, 735–744.
- 54 C. Malalis, M. Barquilla, V. Zabala, K. Casas, M. Maye and A. J. O. Y. Ramayan, *J. Food Nutr. Res.*, 2024, **63**(4), DOI: [10.26596/wefgm627](https://doi.org/10.26596/wefgm627).
- 55 K. Bandara and T. Mahendran, *J. Technol. Value Addit.*, 2020, **2**, 97–109.
- 56 A. Gengatharan, G. Dykes and W. S. Choo, *J. Food Sci. Technol.*, 2021, **58**, 1401–1410.
- 57 Y. Zhao, J. Xu, Y. Zheng, Q. Li, Y. Huang, M. Zong, H. Wu and N. Li, *Molecules*, 2024, **29**, 478.
- 58 S. Wang, Z. Hua, T. Wang, G. Yu and Y. Sun, *Foods*, 2025, **14**, 292.
- 59 W. Sorndech, C. Auranwiwat, T. Ployetchara, S. Butseekhot and J. Agric, *Food Res.*, 2024, **18**, 101310.
- 60 T. Xiao, M. Sun, S. Cao, J. Hao, H. Rao, D. Zhao, Y. Dong and X. Xu, *Food Chem.:X*, 2024, **22**, 101269.
- 61 J. I. Goldstein, D. E. Newbury, P. Echlin, D. C. Joy, C. E. Lyman, E. Lifshin, L. Sawyer and J. R. Michael, *Scanning Electron Microscopy and X-Ray Microanalysis*, Springer US, Boston, MA, 2003.

