



Cite this: DOI: 10.1039/d5fb00935a

Optimization and greenness evaluation of extraction methods of bioactive compounds from cocoa bean shells (*Theobroma cacao* L.)

Erick Alvarez-Yanamango,^{id} *^{abc} Milagros Huaytalla-Ramirez,^{id} ^b
Nereyda Sarmiento-Perez,^{id} ^b Gerald Chumpitaz-Huanqui,^{id} ^{bc} Luis Napan-
Tacca,^{id} ^b Daniel Obregon,^{id} ^b and Alfredo Ibáñez,^{id} ^c

In this study, the efficiency of two green technologies, ultrasound-assisted extraction (CBS-UAE) and microwave-assisted extraction (CBS-MAE), was optimized and compared for the recovery of bioactive compounds from cocoa bean shells (CBSs). Response surface (I-optimal) experimental designs were applied, considering variables such as temperature, time, solid–liquid ratio, particle size, and power. Total phenol content (TPC) and antioxidant capacity (DPPH = 2,2-diphenyl-1-picrylhydrazyl, ABTS = 2,2'-azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid, and FRAP = ferric reducing antioxidant power) were analyzed. CBS-UAE showed higher antioxidant activity, as measured by the ABTS (265.8 ± 5.1 mg TE per g CBS extract b.s.) and FRAP (212.86 ± 4.95 mg TE per g CBS extract b.s.) assays, than CBS-MAE. Optimal conditions obtained for UAE were 65 °C, 45 min, a ratio of 70 mL g⁻¹, and a fine particle size (75–150 μm). While both methods yielded similar TPC and DPPH profiles, UAE achieved the highest antioxidant activity in the ABTS and FRAP assays, whereas MAE drastically reduced the extraction time. UHPLC–HRMS (Q-Exactive–MS/MS) analysis of the optimal CBS-UAE/CBS-MAE extracts mainly identified methylxanthines (theobromine and caffeine), amino acids (L-phenylalanine and arginine), and organic acids (mannitol). The environmental impact of the extraction technique was validated using AGREE metrics, confirming the ecological sustainability of MAE and UAE with AGREE scores of 0.62/1.00 and 0.55/1.00, respectively. These results validate the use of cocoa by-products *via* clean, sustainable technologies, thereby adding value to these residues.

Received 10th December 2025
Accepted 8th April 2026

DOI: 10.1039/d5fb00935a

rsc.li/susfoodtech

Sustainability spotlight

Cocoa bean shells (CBSs) are abundant agro-industrial residues whose valorization is essential to reduce waste and advance circular bioeconomy practices. This study optimizes two green extraction technologies—ultrasound (UAE) and microwave-assisted extraction (MAE)—to recover antioxidant compounds using energy-efficient, low-solvent processes. A response surface design enabled an efficient, robust evaluation of extraction variables, resulting in environmentally favorable conditions. Both methods showed positive AGREE scores (0.62 for MAE and 0.55 for UAE) and produced comparable extracts rich in methylxanthines, organic acids, and amino acids with potential nutraceutical applications. By enabling high-value recovery from cocoa waste and minimizing environmental impact, this work supports SDG 12 (Responsible Consumption and Production), SDG 9 (Industry, Innovation and Infrastructure), and SDG 3 (Good Health and Well-Being).

Introduction

In recent years, waste generated by the food industry has been valorized to develop circular-economy processes. In the chocolate industry, the most exploited fraction of cocoa is the cotyledon after the bean is subjected to fermentation, drying, roasting, and shelling, generating as waste the shell that

represents between 10 and 20% of the cocoa bean.^{1,2} According to the International Cocoa Organization (ICCO), in the period 2021/2022 cocoa bean production reached 4.8 million tons, and it is estimated that by the period 2022/2023 it will have reached 5 million tons of cocoa.³ Based on these data, the estimated global production of cocoa bean shells (CBSs) is between 480 000 and 960 000 tons per year (10–20%).

As shown in Fig. 1, CBSs are an important source of phytochemicals, including bioactive substances such as methylxanthines and polyphenolic compounds, as well as fiber, volatile cocoa compounds, proteins, minerals, and vitamins, which would allow it to be considered as an additive for innovative and functional foods.⁴ The polyphenols present in CBSs are notable

^aDepartamento de Ingeniería, Pontificia Universidad Católica Del Perú, San Miguel, Lima, Peru. E-mail: erick.alvarez@pucp.edu.pe

^bGrupo de Investigación en Tecnologías y Procesos Agroindustriales (ITEPA PUCP), Pontificia Universidad Católica Del Perú, San Miguel, Lima, Peru

^cInstituto de Ciencias Ómicas y Biotecnología Aplicada (ICOBA PUCP), Pontificia Universidad Católica Del Perú, San Miguel, Lima, Peru






Fig. 1 Whole cocoa bean shell treatment from grinding and sieving, green extraction, analysis, optimization, and AGREE validation.

for their high antioxidant activity, as they act as enzyme modulators and inhibitors. These compounds have demonstrated the ability to prevent the progression of various diseases, with potential antidiabetic, anti-inflammatory, and cardioprotective effects.^{5–7} In particular, methylxanthines are associated with improved cardiovascular function and a lower risk of neurodegenerative disorders, such as Alzheimer's and Parkinson's diseases, as well as metabolic conditions, including obesity and diabetes.^{8,9} Therefore, CBS extracts represent a promising source of bioactive compounds for the development of nutraceutical products, *i.e.*, concentrated forms of these phytochemicals intended to supplement the diet and provide health benefits beyond basic nutrition, particularly in the prevention and management of chronic diseases.^{4,10}

Traditional extraction techniques offer simplicity, widespread availability, and low cost; however, they are constrained by inefficient penetration of solvents into plant cell walls and the potential toxicity of the solvents used. Ultrasonic probe cavitation disrupts cell walls *via* microjets and shockwaves; however, scaling remains challenging because probes offer high intensity but limited volume and thermal control, whereas ultrasonic baths provide uniformity at lower intensity over large volumes. Meanwhile, microwave-assisted extraction (MAE) uses temperature-induced diffusion and localized heating to rupture cells. Both technologies thus represent effective, scalable

options for bioactive compound recovery, each with distinct advantages depending on the process requirements.¹¹

This study optimizes and compares two green extraction technologies—ultrasound-assisted extraction (UAE) and microwave-assisted extraction (MAE), for recovering bioactive compounds from cocoa bean shells (CBSs). Using response surface methodology, key process parameters were evaluated to maximize total phenolic content and antioxidant capacity. In addition to phytochemical profiling by UHPLC–HRMS, environmental sustainability was quantitatively assessed using AGREE metrics. The results enable selection of the most suitable technology based on the target outcome, maximizing antioxidant activity or minimizing processing time, supporting CBS valorization through clean, scalable processes.

Materials and methods

Chemical reagents

The chemicals used in the sample analysis were analytical grade chemicals, including Folin&Ciocalteu's phenol reagent, sodium carbonate ($\geq 99.5\%$), 2,2'-diphenyl-1-picryl hydrazyl (95%; DPPH), 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (97%; trolox), (+)-catechin hydrate ($>98\%$), methanol ($\geq 99.9\%$), hydrochloric acid (37%), aluminium chloride (99%), sodium nitrite ($\geq 99\%$), formic acid ($\geq 98\%$) and quercetin ($\geq 98.5\%$) acquired from Sigma-Aldrich (St. Louis, Missouri, USA);



isotopically labeled standards for LC-MS/MS (>99%), 3,7-dimethylxanthine-(dimethyl-D₆) and caffeine-(trimethyl-¹³C₃) were obtained from Merck (Sigma-Aldrich and Supelco, respectively). In addition, the chemicals used in the extraction process, including food-grade ethanol, were purchased from Colaromo SAC (Lima, Peru).

Raw material preparation

The CBS sample was obtained from fine-aroma cocoa of Amazonian origin (*Theobroma cacao* L.), the Criollo variety, provided by Kuyay Chocolates, located in the Jahuanga zone, Bagua Grande district, Utcubamba province, Amazonas region, Peru (5°46'19.6" S, 78°32'57.4" W, 605 m altitude). The cocoa bean shells were obtained after fermentation in handmade wooden crates, drying in the shade, and roasting with a heat ramp from 90 to 120 °C for 50 minutes, followed by mechanical shelling.

To obtain the study sample, the shells were ground to a powder using a knife mill (MKM6003, Bosch, Slovenia) for 30 s, and then passed through 60-, 80-, 100-, and 200-mesh sieves (ASTM E11, Retsch, Germany). The obtained fractions were used as samples with particle sizes of 75–150 μm and 180–250 μm. The samples were vacuum-packed in trilaminate bags and stored at –20 °C until further analysis (Fig. 1).

Green extraction process of CBS bioactives

In both cases, the UAE-and MAE-assisted extracts were separated only by centrifugation at 5000 rpm for 10 minutes. The obtained supernatant was made up to 40 mL and stored in amber vials at freezing temperatures. Before analysis, the supernatant was centrifuged again, and filtered through Whatman 41 paper (20–25 μm pore size), and then through a 0.22 μm syringe filter.

Ultrasonic-assisted extraction (UAE). All tests were performed at 40 kHz/70W (Branson, M1800H-E, USA) in an ethanol–water (50/50 v/v) solvent. The CBS samples were weighed according to the DOE, using 40 mL of solvent. The mixture was vortexed to homogenize, and then ultrasonic extraction was performed according to DOE operating parameters (Table 1). Vortex agitation was performed every 5 min.

Microwave-assisted extraction (MAE). The CBS sample was weighed according to the specified ratios in the DOE, using 40 mL ethanol–water (50/50 v/v) solvent volume based on preliminary tests. The mixture was vortexed to homogenize, and then microwave extraction was performed according to DOE operating parameters (Table 2).

Table 1 DOE parameters of the UAE response surface

Factor name	Type	Min	Max	Coded	
				Low	High
Temperature (A)	Numeric	60	80	–1	+1
Time (B)	Numeric	45	90	–1	+1
Ratio (C)	Numeric	60	80	–1	+1
Granulometry (D)	Categoric	A	B	–1	+1

Table 2 DOE parameters of the MAE response surface

Factor name	Type	Min	Max	Coded	
				Low	High
Ratio (E)	Numeric	60	80	–1	+1
Power (F)	Numeric	95	950	–1	+1
Time (G)	Numeric	2	4	–1	+1
Granulometry (H)	Categoric	A	B	–1	+1

Spectrophotometric assays

Absorbance was measured using a spectrophotometer, Thermo Scientific, Genesys 10S UV-VIS, USA, in all cases.

Total phenolic content. The total phenolic content (TPC) of the extracts was determined using the Folin–Ciocalteu reagent following the method proposed by Singleton & Rossi (1965).¹² In a test tube, 0.5 mL of CBS extract, 2.5 mL of Folin–Ciocalteu reagent (10%), and 2 mL of Na₂CO₃ (7.5%) were added. The mixture was then incubated in a water bath at 45 °C for 15 minutes. It was then cooled in the dark, and the absorbance was measured at 765 nm. TPC values were expressed as milligrams of gallic acid equivalents per gram of dry weight of CBS extract (mg GAE per g CBS d.w), based on a calibration curve plotted from 10 to 100 μg mL^{–1}.

Antioxidant *in vitro* assays. The antioxidant activity of the samples was evaluated using different antioxidant methodologies with minor modifications.

The DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging assay was performed as described by Brand William *et al.* (1995).¹³ The DPPH radical activity of the CBS extract was obtained after mixing 100 μL of CBS extract with 3.9 mL of 500 μM DPPH. The mixture was shaken and left to stand in the dark for 30 minutes, and then the absorbance was measured at 515 nm. The radical scavenging activity was calculated using eqn (1), where the control represents the absorbance of the DPPH reagent blank (without extract), and the sample is the absorbance of the DPPH solution in the presence of the CBS extract.

$$\text{Scavenging effect} = \left(\frac{\text{control} - \text{sample}}{\text{control}} \right) \times 100 \quad (1)$$

ABTS (2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid)) radical scavenging was carried out as proposed by Re *et al.* (1999).¹⁴ It consists of mixing 100 μL of CBS extract with 2 mL of ABTS solution (7 mM), incubating in the dark for 6 minutes, and measuring the absorbance at 734 nm. ABTS values were expressed as micrograms of trolox equivalents per gram of CBS extract on a dry-weight basis (mg TE per g CBS d.w), using a trolox standard curve at concentrations of 5 and 150 μg mL^{–1}.

The ferric reducing antioxidant power (FRAP) assay was performed following the methodology proposed by Benzie & Strain (1996).¹⁵ To 100 μL of CBS extract, 1.9 mL of FRAP solution is added, and the mixture is incubated in the dark for 30 minutes, with absorbance read at 593 nm. FRAP values were expressed in micrograms of trolox equivalent per gram of CBS extract on a dry weight basis (mg TE per g CBS d.w), using



a trolox standard curve at concentrations of 10 and 100 $\mu\text{g mL}^{-1}$, respectively.

DPPH, ABTS, and FRAP are widely used *in vitro* electron-transfer-based assays that evaluate chemical reducing power and radical-scavenging capacity rather than direct biological antioxidant activity.^{16,17} As these methods rely on synthetic radicals and nonphysiological conditions and do not account for bioavailability, their results should be considered preliminary.^{18,19} Therefore, further validation through cellular antioxidant assays (e.g., ORAC) or *in vivo* studies is required to confirm their physiological relevance.²⁰

Identification of metabolites by liquid chromatography-mass spectrometry (UHPLC-MS/MS)

The identification of phenolic compounds and methylxanthines in the MAE/UAE extract was carried out by UHPLC-Q-Orbitrap-MS/MS following the methodology adapted from a previous study.²¹ An ultra-high-performance liquid chromatograph (UHPLC Dionex UltiMate 3000, Thermo Fisher Scientific, Waltham, MA, USA) equipped with an Acclaim™ C18 reversed-phase HPLC column (5 μm , 4.6 \times 150 mm) was used, with a flow rate of 0.7 mL min^{-1} . The mobile phases were 0.1% formic acid in water (A) and 90% acetonitrile in water (B). The gradient program was: -3.6–1.0 min, 15% B; 1.0–25.0 min, linear increase to 95% B; 25.0–26.0 min, increase to 100% B; 26.0–27.0 min, hold at 100% B. The ESI source (HESI II, Thermo Fisher Scientific, Waltham, MA, USA) was operated in positive ion mode (ESI⁺) for alkaloid analysis and in negative ion mode (ESI⁻) for the phenolic profile, at 2.0 and 1.5 kV, respectively, using data-dependent acquisition (DDA).

Environmental impact assessment of the extraction process

The environmental impact of MAE- and UAE-assisted extraction processes was assessed using the innovative quantitative environmental metrics tool AGREEprep, proposed by Wojnowski *et al.* (2022), which is based on 10 principles of Green Analytical Chemistry, including criteria such as solvent type, sample size, and energy consumption.^{22,23} The open-access AGREEprep software was available at mostwiedzy.pl/AGREEprep or *via* the online emulator at the <https://www.agreeprep.anvil.app> website. The quantitative results of the tool are presented as pictograms, with scores from 0 to 1 represented by colors ranging from red to green. Red indicates a high negative impact, and green indicates a process that meets the green chemistry criteria.

The following criteria were considered in the CBS bioactive extraction process. Criterion 1: the CBS extraction process is realized prior to the analytical test; therefore, *ex situ* preparation was considered. Criterion 2: ethanol was used as a solvent, avoiding the use of methanol and hydrochloric acid, among others. Additionally, up to 75% of the solvent was recovered, enabling a sustainable process (criterion 3). Criterion 4: products derived from the separation of bioproducts after centrifugation are considered waste. Criterion 5: the amount of CBS needed to obtain one liter of extract in one hour is considered under optimal MAE and UAE conditions. Criterion 6: the

maximum capacity of the equipment used in an extraction batch (e.g., mL per batch or units per batch). Criterion 7: this consisted of the number of steps in the extraction process, as well as the level of instrumentation from sample preparation to extraction. Criterion 8: the maximum energy output required to obtain a batch of CBS extract was considered. Criterion 9: the analytical technique and instrumentation were defined to validate the presence of metabolites in the extract. Criterion 10: operating conditions based on moderate/high temperatures for extraction, but below the boiling point of the solvents.

Statistical assay

Stat-Ease® software from Stat-Ease, Inc. (MN, USA) was used for statistical analysis. A matrix was generated using optimal response surface methodology (RSM). Optimal designs are recommended when categorical and numeric factors are modeled using a custom model or a cubic or higher-order model.²⁴

The optimization of the UAE extraction was achieved using an optimal response surface design with three continuous numerical factors: extraction time (min), temperature ($^{\circ}\text{C}$), and *S-L* ratio (mL g^{-1}), and a nominal categorical factor, granulometry of the CBS fraction, with two levels: A (75–150 μm) and B (180–250 μm) (Table 1). On the other hand, for microwave-assisted extraction (MAE), the ratios of the sample to solvent (mL g^{-1}), power (W), time (min), and granulometry with two levels: A (75–150 μm) and B (180–250 μm), were selected (Table 2).

The optimized extraction conditions (UAE and MAE) were validated for maximum TPC, TFC, and *in vitro* antioxidant activities (DPPH, FRAP, and ABTS) using RSM-derived values. All responses were again determined under the optimized extraction conditions. The experimental values were compared with the model-predicted values to assess the model's validity.

Results and discussion

ANOVA analysis

The experimental design consisted of 24 runs, as shown in Table 3, with the responses TPC, DPPH, ABTS, and FRAP presented. Response values varied widely across all experiments: TPC ranged from 94 to 191 mg GAE per g, DPPH from 35 to 76%, ABTS from 111 to 274 mg TE per g, and FRAP from 124 to 253 mg TE per g, indicating a strong influence of extraction parameters. The influence of the parameters on the response was evaluated using analysis of variance (ANOVA), with the corresponding sum of squares (SS), degrees of freedom (df), and mean squares (MS). A *p*-value lower than 0.05 means that the model term is significant, whereas values higher than 0.100 are considered not significant. Furthermore, hierarchical ordering is used to assess model terms with *p*-values > 0.05.

It can be observed that the operating conditions of UAE and MAE influence the responses of phenol, flavonoid, and FRAP content. On the other hand, variations in DPPH and ABTS responses do not significantly affect the operating conditions. ANOVA evaluation of phenols, flavonoids, and FRAP is presented in Tables 4–11.



Table 3 Design of experiment matrix for extraction of TPC (mg GAE per g extract bs) = total phenolic content; scavenging activity, DPPH (%) = 2,2-diphenyl-1-picryl-hydrazyl-hydrate, and ABTS (mg ET per g extract bs) = 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid); antioxidant capacity: FRAP (mg ET per g extract bs) = ferric reducing antioxidant power

Run order	UAE – factors				Response				MAE – factors				Response			
	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	TPC	DPPH	ABTS	FRAP	<i>E</i>	<i>F</i>	<i>G</i>	<i>H</i>	TPC	DPPH	ABTS	FRAP
	<i>T</i> (°C)	<i>t</i> (min)	Ratio	Gr (µm)					Ratio	Power	<i>t</i> (min)	Gr (µm)				
1	60	90	60	B	132	61	179	158	69	665	3	B	142	41	158	168
2	76	80	71	B	158	61	247	194	60	665	4	A	147	64	188	186
3	80	45	72	B	165	61	260	208	65	950	4	B	128	59	153	161
4	80	45	60	A	160	62	221	192	69	665	3	B	129	47	130	160
5	60	59	66	B	121	52	170	152	80	950	2	B	126	45	134	166
6	72	45	80	A	143	53	211	158	66	570	2	A	149	55	187	194
7	70	68	70	A	166	63	241	210	60	95	3	A	122	51	141	165
8	80	90	80	A	134	55	210	148	77	95	2	A	140	42	154	185
9	70	90	80	B	143	52	219	166	80	570	3	A	146	53	202	163
10	70	45	60	B	144	76	211	195	80	570	3	A	170	54	200	181
11	80	90	60	B	115	52	179	135	60	950	2	B	115	53	134	124
12	80	64	80	B	100	38	130	127	70	950	3	A	150	60	196	163
13	60	83	72	B	102	47	132	132	77	760	2	A	158	47	134	186
14	60	90	80	A	148	51	206	187	70	950	3	A	153	72	194	168
15	80	73	62	A	123	60	193	169	72	190	4	A	131	49	115	144
16	68	90	60	A	153	53	223	211	77	950	3	B	124	52	111	146
17	62	46	72	A	157	51	240	210	69	665	3	B	94	47	147	151
18	60	45	80	B	138	57	203	179	61	190	2	A	105	45	145	146
19	80	90	60	B	115	57	171	156	80	950	4	B	101	52	161	170
20	80	90	80	A	139	41	201	174	80	190	3	B	147	35	127	150
21	80	45	72	B	148	48	209	190	65	190	2	B	105	40	136	136
22	60	45	80	B	169	45	238	214	75	475	4	B	151	40	131	141
23	60	90	80	A	191	47	274	252	61	190	4	B	124	38	129	133
24	60	45	60	A	178	61	262	253	72	190	4	A	182	39	164	160

An equation in terms of coded factors was generated correlating reaction parameters (independent variables) with the concentration response (dependent variable). In all cases, the final equation in default-coded terms assigns the highest value as 1 and the lowest value as -1 . The influence of each factor will be compared according to its coefficient in the equation.

Comparison of TPC extraction methods

Table 3 presents the TPC per gram of dry weight of CBS extracts obtained using UAE and MAE. The TPC values of the CBS obtained by UAE ranged from 100 to 191 mg GAE per g extract d.w., while those obtained by MAE ranged from 94 to 182 mg GAE per g extract d.w.

Table 4 ANOVA of extraction parameters in TPC UAE extraction

Source	SS	df	MS	<i>F</i> -value	<i>p</i> -value	
Model	4688.56	3	1563	4.15	0.0194	Significant
<i>B</i> - <i>t</i> (min)	1241.06	1	1241	3.29	0.0846	
<i>D</i> -Gr (µm)	2199.24	1	2199	5.84	0.0254	
<i>B</i> ²	1181.16	1	1181	3.13	0.0919	
Residual	7537.40	20	377			
Lack of fit	5995.69	15	400	1.30	0.4151	Not significant
Pure error	1541.72	5	308			
Cor total	12 225.97	23				

The TPC model equations are expressed in coded factor form for UAE (2) and MAE (3). In the UAE equation, time (*B*) shows a negative linear effect and a positive quadratic term (*B*²), indicating an optimal time. Granulometry (*D*) has a negative coefficient because *D* is coded (-1 for fine and $+1$ for coarse), so finer particles increase TPC. In the MAE equation, the ratio (*E*), power (*F*), and time (*G*) exhibit positive linear effects, while granulometry (*H*) is negative (finer particles are beneficial). The interaction terms *EF* and *FG* are negative, suggesting antagonistic effects when these factor pairs are increased simultaneously.

Table 5 ANOVA of extraction parameters in TPC MAE extraction

Source	SS	df	MS	<i>F</i> -value	<i>p</i> -value	
Model	6872.76	6	1145.46	4.64	0.0058	Significant
<i>E</i> -ratio	1311.57	1	1311.57	5.31	0.0341	
<i>F</i> -power	11.63	1	11.63	0.0471	0.8308	
<i>G</i> - <i>t</i> (min)	560.98	1	560.98	2.27	0.1502	
<i>H</i> -Gr (µm)	3071.26	1	3071.26	12.43	0.0026	
<i>EF</i>	1118.46	1	1118.46	4.53	0.0483	
<i>FG</i>	901.01	1	901.01	3.65	0.0732	
Residual	4200.88	17	247.11			
Lack of fit	1377.89	12	114.82	0.2034	0.9889	Not significant
Pure error	2822.99	5	564.6			
Cor total	11 073.64	23				



Table 6 ANOVA of extraction parameters in FRAPS UAE extraction

Source	SS	df	MS	F-value	p-value	
Model	11 042	3	3680.66	4.73	0.0119	Significant
A-T (°C)	3926.82	1	3926.82	5.05	0.0361	
D-Gr (µm)	4531.66	1	4531.66	5.82	0.0255	
AD	3711.13	1	3711.13	4.77	0.0411	
	15 566	20	778.32			
	12 155	15	810.35	1.19	0.4593	
Residual	3411.18	5	682.24			
Lack of fit	26 608	23				Not significant
Pure error	11 042	3	3680.66	4.73	0.0119	
Cor total	3926.82	1	3926.82	5.05	0.0361	

$$\text{TPC (UAE)} = 129.46 - 8.14B - 9.66D + 19.10B^2 \quad (2)$$

$$\text{TPC (MAE)} = 135.32 + 10.63E + 0.9897F + 6.38G - 11.81H - 12.60EF - 10.33FG \quad (3)$$

Both UAE and MAE showed that finer particles significantly increased TPC, but while UAE exhibited a curvilinear effect of time, indicating an optimum, MAE presented positive linear effects of the ratio, power, and time, along with two negative interactions (*EF* and *FG*). These differences reflect the distinct physical principles of each technique, conventional thermal diffusion in UAE *versus* microwave-induced heating in MAE, underscoring that optimization strategies must be tailored to the extraction method. All factor interpretations were based on statistical significance ($p < 0.05$) derived from the ANOVA.

Comparison of antioxidant capacity extraction methods

In this study, the FRAP, DPPH, and ABTS methods were employed to assess the antioxidant capacity of CBS extracts obtained *via* UAE and MAE. The FRAP values for the MAE and UAE extracts ranged from 124–194 and 127–253 mg TE per g extract d.w., respectively. Similarly, the DPPH scavenging effects of MAE and UAE extracts ranged from 35 to 72% and 38 to 76%, respectively. In addition, the ABTS values for MAE and UAE extracts ranged from 111 to 202 and 130 to 274 mg TE per g extract d.w., respectively (Table 3). Each antioxidant model equation (FRAP, DPPH, and ABTS) is expressed in the coded factor form for UAE (4, 6, 8) and MAE (5, 7, 9).

Table 7 ANOVA of extraction parameters in FRAPS MAE extraction

Source	SS	df	MS	F-value	p-value	
Model	5524.89	7	789.27	5.71	0.0019	Significant
E-ratio	279.86	1	279.86	2.02	0.174	
F-power	689.5	1	689.5	4.99	0.0402	
G-t (min)	98.74	1	98.74	0.7142	0.4105	
H-Gr (µm)	2215.38	1	2215.38	16.02	0.001	
EG	1008.24	1	1008.24	7.29	0.0158	
FG	892.06	1	892.06	6.45	0.0218	
F ²	490.39	1	490.39	3.55	0.078	
Residual	2212.12	16	138.26			
Lack of fit	1761.06	11	160.1	1.77	0.2735	Not significant
Pure error	451.06	5	90.21			
Cor total	7737.01	23				

Table 8 ANOVA of extraction parameters in ABTS UAE extraction

Source	SS	df	MS	F-value	p-value	
Model	5190.87	1	5190.87	4.33	0.0492	Significant
D-Gr (µm)	5190.87	1	5190.87	4.33	0.0492	
Residual	26 349.5	22	1197.7			
Lack of fit	22 015.73	17	1295.04	1.49	0.348	Not significant
Pure error	4333.77	5	866.75			
Cor total	31 540.37	23				

For the FRAP equation obtained by UAE, it is evident that temperature (*A*) is the most significant factor negatively influencing concentration. Also, the effect of particle size (*D*) produced by the AD interaction is suppressed mainly by the *D* term. If the AD interaction is negative (*i.e.*, -1 smaller particle size), the *D* term will be positive, and *vice versa*. In the case of MAE, the main factors, power (*F*) and ratio (*E*), have the most significant positive effects, and the power (*F*²) quadratic term has the most significant negative effect on FRAP extraction.

$$\text{FRAP (UAE)} = 183.45 - 14.91A - 13.80D + 14.49AD \quad (4)$$

$$\text{FRAP (MAE)} = 165.83 + 4.93E + 7.72F - 2.70G - 10.12H - 11.83EG + 10.46FG - 11.63F^2 \quad (5)$$

Regarding the UAE-derivate DPPH equation, the quadratic temperature term (*A*) is the most negative in the extraction, independent of particle size. On the other hand, MAE extraction is primarily influenced by power (*F*) and, to a lesser extent, by granulometry (*H*).

$$\text{DPPH (UAE)} = 59.83 - 0.3654A - 5.60C - 6.75A^2 \quad (6)$$

$$\text{DPPH (MAE)} = 48.47 - 3.56E + 9.61F + 1.94G - 5.31H + 2.80FG \quad (7)$$

Finally, the ABTS equation for UAE and MAE is represented by the *a* plane without any quadratic term (curvature). Granulometry has a positive effect on the extraction of smaller size (*D* and *H*, -1). Power (*F*) and time (*G*) in MAE have a positive effect. In contrast, granulometry (*H*) negatively affects extraction.

$$\text{ABTS (UAE)} = 210.75 - 14.76D \quad (8)$$

Table 9 ANOVA of extraction parameters in ABTS MAE extraction

Source	SS	df	MS	F-value	p-value	
Model	11 124.54	5	2224.91	6.06	0.0019	Significant
F-power	2993.18	1	2993.18	8.15	0.0105	
G-t (min)	230.16	1	230.16	0.6267	0.4389	
H-Gr (µm)	6732.67	1	6732.67	18.33	0.0004	
FG	1246.66	1	1246.66	3.39	0.0819	
FH	1395.37	1	1395.37	3.8	0.067	
Residual	6610.1	18	367.23			
Lack of fit	5005.58	13	385.04	1.2	0.4513	Not significant
Pure error	1604.52	5	320.9			
Cor total	17 734.64	23				



Table 10 ANOVA of extraction parameters in DPPH UAE extraction

Source	SS	df	MS	F-value	p-value	
Model	724.1	3	241.37	6.2	0.0038	Significant
A-T (°C)	2.38	1	2.38	0.0611	0.8073	
C-ratio	521.51	1	521.51	13.39	0.0016	
A ²	178.64	1	178.64	4.59	0.0447	
Residual	778.98	20	38.95			
Lack of fit	528.24	15	35.22	0.7022	0.7281	Not significant
Pure error	250.74	5	50.15			
Cor total	1503.08	23				

$$\text{ABTS (MAE)} = 153.51 + 15.66F + 4.08G - 17.65H + 12.18FG - 10.69FH \quad (9)$$

Across the three antioxidant assays, granulometry (particle size) consistently influenced extraction in both methods, with finer particles (*D* or *H* coded as -1) favouring higher activity in UAE for FRAP and ABTS, and in MAE for DPPH, FRAP, and ABTS. Other key factors differed between techniques, with UAE showing a dominant role in temperature (*A*) and its quadratic term in FRAP and DPPH, indicating an optimal temperature, whereas ABTS was affected solely by particle size. In contrast, MAE exhibited a more complex behaviour, with microwave power (*F*) significant across all three assays, often accompanied by quadratic (FRAP) or interaction terms (FRAP, ABTS, and DPPH), while the solvent-to-sample ratio (*E*) and time (*G*) contributed selectively depending on the assay. These differences reflect the distinct energy transfer mechanisms of each method and highlight that the optimal conditions for antioxidant recovery are assay-dependent. The interpretation of model terms was based on their *p*-values from the ANOVA.

The antioxidant activity values obtained for CBS extracts were higher than those reported for CBS flours (ABTS = 3.39–11.55 and FRAP = 3.84–7.62 mg TE per g sample).²⁵ However, in both studies, particle size influences antioxidant activity, with antioxidant activity inversely proportional to particle size. The smaller the particle size, the greater the recovery of bioactive compounds and the higher the antioxidant activity. Particle size favors mass transfer and solvent accessibility to bioactive compounds, facilitating their permeation and diffusion through the plant matrix.²⁶

Table 11 ANOVA of extraction parameters in DPPH MAE extraction

Source	SS	df	MS	F-value	p-value	
Model	1489.05	5	297.81	18.83	< 0.0001	Significant
E-ratio	148.08	1	148.08	9.36	0.0067	
F-power	1096.56	1	1096.56	69.34	< 0.0001	
G-t (min)	51.73	1	51.73	3.27	0.0872	
H-Gr (µm)	622.05	1	622.05	39.34	< 0.0001	
FG	66.32	1	66.32	4.19	0.0555	
Residual	284.64	18	15.81			
Lack of fit	135.9	13	10.45	0.3514	0.9403	Not significant
Pure error	148.74	5	29.75			
Cor total	1773.69	23				

Surface response analysis

Graphs represent a compound increase in a colour scale. In the case of TPC, as measured by UAE and MAE, the increase shifts from blue to red, ranging from approximately 120 to 180 mg GAE per g extract d.w. The use of a graphical contrast was proposed to highlight higher and lower extraction concentrations of each compound.

The TPC graphs show that MAE-assisted extraction yields higher extraction at higher ratios, with a significant effect of particle size (Fig. 2C). Granulometry is a significant factor influencing extraction concentration in the UAE (Fig. 2A and B). Additionally, increased time reduces the extraction of TPC from samples with larger particles.

Higher TPC concentrations by MAE can be attributed to microwave heating, which facilitates the direct transfer of electromagnetic energy to the matrix-solvent system, enabling rapid and uniform heating at the molecular level.^{27,28} This accelerates extraction kinetics, resulting in shorter extraction times than with UAE.

The FRAP graph shows an increase from blue to red, ranging from approximately 120 to 253 mg TE per g extract d.w. As observed in the FRAP graph (Fig. 3), a higher concentration is achieved with MAE at a 60 ratio, high power, and small particle size (Fig. 3C). On the other hand, using UAE, low temperatures, and small granulometry sizes increase FRAP concentration (Fig. 3A). In both cases, a significant reduction in concentration of the compounds is observed with higher particle size.


These FRAP trends closely parallel the TPC results discussed previously, where MAE also yielded higher phenolic extraction under similar conditions (high power and small particle size). This positive correlation suggests that phenolic compounds are major contributors to the antioxidant activity measured by FRAP. However, FRAP values were higher (up to 253 mg TE per g with UAE at *A*: 60, *B*: 45, *C*: 60, and *D*: *A*) than those with MAE, indicating that other factors may be influencing the reducing capacity.

Synergistic effects or interference from non-phenolic constituents with reducing power can enhance FRAP values, even when they lack biological relevance as antioxidants. Indeed, sugars and citric acid are recognized as common interferences,¹⁸ while organic acids (*e.g.*, tartaric acid) and minerals have also been shown to significantly influence FRAP results.²⁹ In the present study, gluconic acid (a reducing organic acid) was identified by UHPLC, and its presence likely contributed to the elevated FRAP values. Furthermore, the pronounced influence of particle size on both TPC and FRAP supports this, as smaller particles facilitate the release of a broader range of reducing species beyond phenolics.³⁰


The DPPH concentration with MAE increases at high power levels and with smaller particles. On the other hand, in the UAE, DPPH values increase at lower ratios, as indicated by ANOVA analysis (Fig. 4C). In MAE, there is a linear increase with power reduction and time that generates a planar surface. On the other hand, the UAE reaches higher values in the middle of the experimental window (Fig. 4A and B). The UAE exhibits a slight increase with a lower ratio and shows no significant influence of particle size. Slightly higher DPPH values were observed in the MAE extracts than in the UAE (72% vs. 65%). Phenolic



A: TPC




TPC (UAE):
Ratio (C) = 70
Granulometry (D) = A
C: TPC




TPC (MAE):
Time (G) = 4
Granulometry (H) = A

B: TPC



TPC (UAE):
Ratio (C) = 70
Granulometry (D) = B
D: TPC



TPC (MAE):
Time (G) = 4
Granulometry (H) = B

Fig. 2 Three-dimensional graphs of UAE and MAE under different conditions for phenols with UAE (A and B) and MAE (C and D) with different particle sizes.

compounds, which donate hydrogen from their hydroxyl groups, are potent DPPH radical scavengers, and a positive correlation between DPPH activity and total phenolic content is generally expected.³¹

However, as with FRAP, DPPH results must be interpreted with caution. The DPPH radical is sterically hindered, making the reaction rate more dependent on the antioxidant's ability to access the radical center than on its intrinsic chemical properties.¹⁹ Consequently, large polyphenols may react slowly and be underestimated, whereas small molecules with good accessibility can react quickly regardless of their true antioxidant capacity.¹⁹ Moreover, the DPPH assay is susceptible to interference from reducing agents, which can cause decolorization independently of

genuine radical-scavenging activity.^{18,20} In the present study, gluconic acid (identified by UHPLC) likely contributed to the measured radical scavenging activity, leading to an overestimation of the actual antioxidant capacity, a factor often overlooked in conventional DPPH-based analyses.

In ABTS, the UAE values were higher than those for MAE (274 > 202 mg ET per g CBS extract d.w.), as shown in Table 3 (Fig. 5C). Higher extraction was observed in MAE at smaller particle sizes, higher power, and a 4 minute time (Fig. 5C). ABTS extraction shows a linear increase in both cases, with no quadratic terms in their equations.

The discordance among FRAP, DPPH, and ABTS results highlights that electron-transfer-based antioxidant assays






Fig. 3 Three-dimensional graphs of UAE and MAE under different conditions for FRAP with UAE (A and B) and MAE (C and D) with different particle sizes.


measure the combined redox activity of multiple chemical families rather than a single class of compounds.²⁰ The Folin-Ciocalteu reagent is reduced by both phenolic and non-phenolic compounds, while ABTS and FRAP respond to the overall reducing capacity of a sample, often leading to poor correlation when total phenolic content is low or when non-phenolic reductants are abundant.²⁰ In the present study, FRAP and DPPH values were notably influenced by the presence of gluconic acid (identified by UHPLC), a reducing organic acid that can overestimate antioxidant capacity independently of phenolic content. This explains why UAE extracts showed higher FRAP values despite lower TPC, whereas MAE yielded higher phenolic extraction. Since theobromine and caffeine (both alkaloids) were found to be prominent compounds in CBS

extracts, their presence accounts for the higher ABTS values observed with UAE, despite the lower phenolic content. Taken together, these findings indicate that MAE favors phenolic extraction (as reflected in higher TPC), whereas UAE may enhance the recovery of alkaloids, organic acids, or other reducing species that interfere with FRAP, DPPH, and ABTS assays. Consequently, antioxidant capacity is not an absolute property but rather depends on both the extraction method and the assay employed, highlighting the need for chemical characterization and multiple assays when evaluating bioactive extracts.


Optimal conditions were established to achieve the highest component concentration in the extract (Tables 12 and 13). Desirability criteria were used to guide the optimization, aiming



A: DPPH




DPPH (UAE):
Ratio (C) = 60
Granulometry (D) = Average
C: DPPH




DPPH (MAE):
Ratio (E) = 60
Granulometry (H) = A

B: DPPH



DPPH (UAE):
Ratio (C) = 80
Granulometry (D) = Average
D: DPPH



DPPH (MAE):
Ratio (E) = 60
Granulometry (H) = B

Fig. 4 Three-dimensional graphs of UAE and MAE under different conditions for DPPH with UAE (A and B) and MAE (C and D).

to maximize the concentration obtained for each component simultaneously. Desirability is an objective function that ranges from ($d = 0$) to the unit ($d = 1$).²⁴

For CBS-UAE, a desirability value of 0.67 was obtained under optimal extraction conditions of 45 min, 65 °C, a ratio of 70 mL g⁻¹, and a particle size of 75–150 μm (Table 12). On the other hand, the optimal conditions for CBS-MAE yielded a desirability of 0.85 with higher power (950 watts), a ratio of 60 mL g⁻¹, and a particle size similar to that of UAE (Table 13).


Ultrasound-assisted extraction (UAE) demonstrated advantages over microwave-assisted extraction (MAE), yielding higher phenol concentrations and greater antioxidant capacity (as measured by ABTS and FRAP).

Graphical optimization plots are shown in Fig. 6 and 7. They indicate the regions where the desired criteria are met. The bright yellow (default) indicates where the entire range of all intervals meets the specified criteria (maximum), and the dark gold shows where the requirements are partially met.


The experimental verification values obtained under optimal extraction conditions, all within the expected range, are shown in Table 14. For the UAE, the experimental values were 160.01 mg GAE per g CBS extract (TPC), 31.99% antioxidant activity (DPPH), and 265.80 and 212.86 mg ET per g CBS extract for ABTS and FRAP, respectively. For MAE, the values were 166.74 mg GAE per g CBS extract (TPC), 42.12% (DPPH), and 253.59 and 183.83 mg ET per g CBS extract for ABTS and FRAP.



A: ABTS




ABTS (UAE):
Ratio (C) = 70
Granulometry (D) = A
C: ABTS




ABTS (MAE):
Ratio (E) = 70
Granulometry (H) = A

B: ABTS



ABTS (UAE):
Ratio (C) = 70
Granulometry (D) = B
D: ABTS



ABTS (MAE)
Ratio (E) = 70
Granulometry (H) = B

Fig. 5 Three-dimensional graphs of UAE and MAE under different conditions for ABTS with UAE (A and B) and MAE (C and D) with different particle sizes.

When comparing the two methods, UAE exhibited higher antioxidant capacity in the ABTS (266 > 254 mg ET per g) and FRAP (213 > 184 mg ET per g) assays, whereas MAE showed

higher DPPH radical-scavenging activity (42% vs. 32%). The confidence intervals for TPC overlapped, indicating no statistically significant difference between the methods. Both

Table 12 Optimal conditions in UAE maximized extraction of all components: ratio = 70 and granulometry = A

T (°C)	t (min)	Ratio	Gr (μm)	TPC	DPPH	ABTS	FRAP	Desirability
65	45	70	A	166	58	226	213	0.67

Table 13 Optimal condition for MAE maximized extraction of all components: time = 4 min and granulometry = A

Ratio	Power	t (min)	Gr (μm)	TPC	DPPH	ABTS	FRAP	Desirability
60	950	4	A	146	72	214	187	0.85






Fig. 6 Optimization graph of the UAE for all compounds with a granulometry of A.

extraction techniques outperformed traditional systems that rely on mechanical agitation.³²

Table 15 compares the extraction of bioactive compounds from cocoa bean shells using different techniques. Among the reported UAE and MAE methods, our approach achieved competitive results while using ethanol as a simpler, conventional solvent. Under optimal MAE conditions (95 W, 2 min, 80 mL g⁻¹), we obtained 42.12% DPPH and 166.74 mg g⁻¹ TPC, with UAE yielding 31.99% DPPH and 160.01 mg g⁻¹ TPC. Notably, our TPC values substantially outperformed water-based extractions (22.1–35.9 mg g⁻¹),³² demonstrating ethanol's superior efficiency for phenolic compound recovery. In comparison, deep eutectic solvents have shown high DPPH scavenging (74.81%),³³ though they require complex solvent preparation, whereas UAE with water has been reported to be efficient for isolating bioactive compounds from CBSs.³⁴

It can be assumed that there is a more efficient mass transfer mechanism in MAE than in UAE, which, unlike localized acoustic cavitation in UAE, activates a mechanism known as temperature-induced diffusion (TID), in which thermal

gradients alter the internal chemical potential of the biomass, promoting greater water absorption in the cells and generating high internal pressures that weaken or break the cell walls.^{36,37} This effect facilitates the release of intracellular bioactive compounds, thereby increasing extraction yield and significantly reducing processing time compared to the UAE. Furthermore, microwave-induced heating produces localized temperature increases within the plant matrix that exceed the average temperature of the extraction system, thereby accelerating the solubilization of compounds in the solvent and improving overall extraction efficiency.³⁷

In addition, MAE efficiency is linked to the dielectric properties of the system (solvent-plant matrix), especially the dielectric constant (ϵ') and the dielectric loss factor (ϵ''), which determines the material's ability to absorb energy and convert it into heat.³⁸ In the case of CBS, its heterogeneous composition, moisture, minerals, and fibers significantly influence its dielectric response. The higher the humidity and ions, the greater the dielectric loss (ϵ''), which translates into greater microwave energy absorption and an increase in the heating rate.^{38,39}

The use of polar solvents, such as ethanol–water (50 : 50), improves microwave energy absorption by adjusting the medium's dielectric constant, increasing internal temperature, and facilitating cell disruption.⁴⁰ Factors such as humidity, volumetric density, and the presence of CBS ions can increase the dielectric loss factor, intensify the heating rate, and release bioactive compounds, including polyphenols and methylxanthines, present in this residue.^{38,41} While this temperature increase is beneficial for cell disruption and extraction yield, it also introduces a critical process control consideration: if dielectric and thermal parameters are not precisely regulated, localized overheating may occur, leading to partial degradation of thermolabile antioxidant compounds. For this reason, beyond this inherent risk, the primary technical restriction to the use of MAE in industry is process scalability, as limitations in heat and mass transfer at large volumes must be addressed.

Chromatography identification of metabolites in optimized CBS extracts

Although the chemical profile obtained by UHPLC-Q-Orbitrap-MS/MS reveals no significant differences between both extraction methods, the analyzed extracts contain a complex combination of primary and secondary metabolites resulting from fermentation and drying. Metabolites were identified using deuterated standards; others were putatively identified based on matching retention times, exact masses, and MS2 spectra (Table 16). Retention times ranged from 1.50 to 6.00 min, with a predominance of highly polar molecules characteristic of lignocellulosic matrices, such as CBSs, and oxidized carbohydrates, amino acids, and alkaloids as a significant bioactive fraction. Table 16 summarizes the identified compounds, ordered based on retention time (RT).

On the one hand, two of the most intense signals found, in positive mode, for the CBS extracts assisted by MAE/UAE were theobromine (RT 3.38 min; m/z 181.0720) and caffeine (RT




Fig. 7 Optimization graph of MAE for all compounds with a granulometry of A.



Table 14 Verification of the extraction of bioactives by UAE and MAE

Method	95% CI	TPC	DPPH	ABTS	FRAP
UAE	Low for mean	134	48	204	199
	High for mean	165	58	247	254
	SD	19.4	6.24	34.6	27.8
	Experimental	160.01 ± 0.27	31.99 ± 0.77 ^a	265.8 ± 5.1	212.86 ± 4.95
MAE	Low for mean	138	51	159	166
	High for mean	157	56	183	186
	SD	15.7	3.97	19.1	11.7
	Experimental	166.74 ± 6.47	42.12 ± 1.53	253.59 ± 4.51	183.83 ± 15.03

^a Note: DPPH was diluted 1/4 for the DOE and 1/20 for validation due to the high initial concentration, which explains the lower DPPH values observed during validation.

5.86 min; m/z 195.0876), suggesting their high tolerance to energy-assisted extraction treatments, consistent with previous reports highlighting their thermal resistance and their high solubility in hydroalcoholic matrices.⁴² Their stability during MAE and UAE treatments suggests that both processes are suitable for obtaining alkaloid-enriched extracts, as reported by other authors who used MAE and UAE to extract alkaloids from cocoa husk.^{27,34} The presence of these methylxanthines reinforces the extract's potential for neurostimulant, antioxidant, and nutraceutical applications. It serves as a reliable chemical marker for cocoa crop authenticity, since the theobromine/caffeine ratio varies according to the cocoa genotype.⁴³

On the other hand, polyalcohols and organic acids, such as mannitol (RT 2.11 min; m/z 181.0713) and gluconic acid (RT

2.13 min; m/z 195.0507), were identified in negative mode. These are typical products of carbohydrate transformation during cocoa fermentation, so their presence in the extract suggests disruption of the cell wall and release of partially or totally oxidized sugars during dielectric heating (MAE) and cavitation (UAE). On the one hand, mannitol is produced primarily by yeasts and lactic acid bacteria (LAB), especially *Lactobacillus fermentum*, which uses fructose as an alternative electron acceptor, allowing its reduction to mannitol during anaerobic fermentation.^{44–46} Mannitol is associated with bitterness, astringency, and fruity and floral descriptors in chocolate.⁴⁶ Gluconic acid, on the other hand, is an undesirable product of cocoa fermentation, associated with a vinegary or unpleasant flavor in chocolate.⁴⁷ Its production is attributed to

Table 15 Optimal conditions and components obtained by different techniques^a

Method	Optimal conditions	Obtained compounds (d.b.)	Reference
UAE	45 min, 64 °C, ratio 70 mL g ⁻¹ , and a particle size of 75–150 μm	TPC: 160.01 % DPPH: 31.99 ABTS: 265.80 FRAP: 212.86	This study
MAE	95 W, 2 min, 80 mL g ⁻¹ , and a particle size of 75–150 μm	TPC: 166.74 % DPPH: 42.12 ABTS: 253.59 FRAP: 183.83	This study
UAE	69.45 °C; S/L 50 mL g ⁻¹ ; 37 kHz; UAE power (70%), and extraction time, 44.26 min	TPC: 118.38 % DPPH: 81.846	N. Pavlović <i>et al.</i> , 2021 (ref. 34)
UAE	40 kHz; 296 W; 50 mL g ⁻¹ ; 80% v/v ethanol; 55 °C; 45 min; extracto liofilizado	% DPPH: 30 FRAP: 319.20 μM Fe II per g extract	Md Yusof <i>et al.</i> , 2019 (ref. 35)
MAE	5 min; pH 12, 97 °C; 25 mL g ⁻¹ ; water solvent; extracto seco a 40 °C	TPC: 35.9 FRAP: 35.5 ABTS: 33.6	Mellinas <i>et al.</i> , 2020 (ref. 32)
Conventional extraction	22.2 mL g ⁻¹ ; water solvent; 100 °C; 90 min; extracto seco a 40 °C	TPC 22.1 FRAP 16.0 ABTS 22.2	
MAE	Stirring; 63% v/v ethanol; 399.88 W; 217.42 s; 69.71 mL g ⁻¹ ; extracto diluido a 100 mL	TPC: 45.68 FRAP*: 41.29	Rincón Soledad, 2023. ²⁷

^a TPC (mg GAE per g extract bs) = total phenolic content; scavenging activity, DPPH (%) = 2,2-diphenyl-1-picryl-hydrazyl-hydrate, and ABTS (mg ET per g extract bs) = 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid); antioxidant capacity: FRAP (mg ET per g extract bs) = ferric reducing antioxidant power (*).



Table 16 Identification of metabolites of CBS extract by UHPLC-Q-Orbitrap-MS/MS

Component name	Formula	Reference ion	Retention time (min)	Theoretical mass	Observed m/z	ppm ^a	Fragments
Arginine	C ₆ H ₁₄ N ₄ O ₂	[M + H] ⁺	1.88	175.1190	175.1191	0.85	60.0562, 68.05019, 70.06577, 71.04976
Adenine	C ₅ H ₅ N ₅	[M + H] ⁺	2.08	136.0617	136.0619	0.95	55.0297, 65.01408, 67.02972, 77.01400, 82.0404, 92.02483, 94.04040, 119.035949
Mannitol	C ₆ H ₁₄ O ₆	[M-H] ⁻	2.11	181.0718	181.0713	2.54	59.0128, 71.01301
Gluconic acid	C ₆ H ₁₂ O ₇	[M-H] ⁻	2.13	195.0510	195.0507	1.69	59.0128, 75.00780, 71.01285, 72.99220
Aminoundecanoic acid	C ₁₁ H ₂₃ NO ₂	[M + H] ⁺	2.25	202.1802	202.1807	2.70	55.05489, 57.0705, 58.06578, 65.0391, 67.05479, 79.05473, 81.07034, 84.08130, 85.06528
Theobromine ^b	C ₇ H ₈ N ₄ O ₂	[M + H] ⁺	3.38	181.0720	181.0720	0.00	54.0345, 56.0501, 67.02970, 69.04534, 81.0452, 83.06092, 94.0404, 108.05600, 110.0715
Phenylethanamine	C ₈ H ₉ N	[M + H] ⁺	3.46	120.0807	120.0810	1.92	51.02362, 53.03927, 63.0235, 65.03922, 67.0548, 75.02357, 77.03918, 91.05474, 103.05458
Phenylalanine	C ₉ H ₁₁ NO ₂	[M + H] ⁺	3.46	166.0863	166.0863	0.30	95.04965, 103.05461
Caffeine ^b	C ₈ H ₁₀ N ₄ O ₂	[M + H] ⁺	5.86	195.0877	195.0876	0.26	69.04535, 83.06095, 109.0400, 110.07171, 138.06639

^a Parts per million, unit of mass precision of the mass spectrometer. ^b Available commercial standard (STD) for validation.

Gluconobacter and *Acetobacter pasteurianus*, with *Gluconobacter* preferentially oxidizing glucose to gluconic acid when glucose has not been depleted by acetic acid bacteria (AAB), whereas *Acetobacter* acts at the end of fermentation, after ethanol depletion.^{47,48} Finally, nitrogen-rich and Maillard reaction metabolites were detected, such as arginine (RT 1.88 min, m/z 175.1191), phenylalanine (RT 3.46 min, m/z 166.0863), phenylethanamine (RT 3.46 min, m/z 120.0810), and aminoundecanoic acid (RT 2.25 min, m/z 202.1807). The presence of these compounds may indicate early Maillard reactions during roasting and the release of free amino acids during metabolite extraction.⁴⁹ Notably, the roasting conditions employed (a gradual ramp from 90 to 120 °C over 50 min) are relatively mild, which is consistent with the detection of early-stage Maillard products rather than advanced browning compounds.

Arginine was also identified, mainly due to proteolysis during grain fermentation, where this amino acid contributes to the slightly bitter taste.⁵⁰ Phenylalanine plays a structural role in proteins and is among the predominant amino acids following cocoa fermentation.^{51–53} It is also a key precursor in Maillard reactions, contributing to the aromas and color of cocoa, as well as to the formation of biogenic amines such as phenylethanamine, a product of its decarboxylation.^{54,55} The co-detection of phenylethanamine with phenylalanine in the extract suggests active degradation and conversion pathways

during post-harvest processing of cocoa beans. Amino-undecanoic acid is associated with lipid derivatives formed during processing through the modification of fatty acids.^{56,57} Finally, adenine was identified (RT 2.08 min; m/z 136.0619), suggesting possible changes in another nitrogen-rich metabolic pathway.⁵⁸

Environmental impact assessment

AGREEprep has been used to select processes with lower environmental impact or to identify problem points in terms of “greenness” during the design and development of new extractive processes. AGREEprep uses a scale of 0 to 1, with scores above 0.5 indicating an acceptable method or above 0.6, indicating an environmentally friendly process.⁵⁹ The results of the AGREEprep assessments are shown in Fig. 8, and the weight criterion is shown in Table 17. The extraction procedures using UAE and MAE yielded overall scores of 0.58/1.00 and 0.65/1.00, respectively. Although both extraction processes exceed the threshold for ecological or eco-friendly treatments,⁵⁹ MAE would be more sustainable. Likewise, the UAE score technique compared to MAE shows notable advantages in performance (criterion 6) and energy consumption (criterion 8). At the same time, both techniques shared weaknesses in the use of hazardous substances in the bioactive recovery process and in






Fig. 8 Comparison of the results of the AGREEprep evaluation of CBS bioactive extraction techniques.

Table 17 Criteria and weighting of criteria for evaluating CBS bioactive extraction techniques using the AGREEprep method

Assessment criteria	Weight	Score	
		UAE	MAE
1 Sample preparation: <i>ex situ</i>	1	0.00	0.00
2 Volume [mL] of problematic substances	2	0.00	0.00
3 Use of sustainable or renewable materials	2	0.75	0.75
4 Mass [g] of waste generated	4	0.82	0.90
5 Mass [g] of the sample	4	0.77	0.79
6 Number of samples per hour	3	0.42	0.76
7 Integrate steps and automation	2	0.00	0.00
8 Energy consumption	4	0.87	1.00
9 Post-sample preparation configuration for analysis	2	0.75	0.75
10 Number of identified hazards	3	0.50	0.50
	Total	0.58	0.65

the number of steps required (criteria 2 and 7). Nevertheless, ethanol is considered one of the most environmentally friendly solvents, enabling more sustainable, scalable extraction of bioactives on an industrial scale.

To distinguish between the two processes using AGREEprep criteria, greater weight (4) was assigned to criteria 4, 5, and 8, as these directly reflect critical differences in operational efficiency and resource utilization.

In criterion 4 (waste generation), MAE achieved a higher score (0.90) compared to UAE (0.82), reflecting its more efficient use of materials and lower waste output per batch. For criterion 5 (sample mass), both methods performed similarly (UAE: 0.77; MAE: 0.79), indicating that they handle sample sizes efficiently and show no meaningful difference. Criterion 8 (energy consumption) showed the most pronounced disparity, with the value for MAE at 1.00 compared to that of UAE at 0.76, confirming that the shorter extraction time of MAE translates into substantially lower energy demand.

For criterion 1 (*ex situ* sample preparation), both UAE and MAE were assigned a score of 0.00, as sample preparation is performed prior to extraction, as shown in Fig. 1 (grinding and sieving of the dried sample). Regarding criterion 3 (use of sustainable or renewable materials), both methods received a score of 0.75, reflecting a good commitment to renewable sources. In criterion 6

(number of samples per hour), MAE (0.76) substantially outperformed UAE (0.42). For UAE, each extraction batch required 45 minutes (including setup and cooling), allowing approximately 6 batches per hour, which yielded a score of 0.42. For MAE, the extraction time was 4 minutes per batch, enabling up to 25 batches per hour and resulting in a score of 0.76. These scores reflect the tool's normalization based on throughput efficiency. For criterion 9 (post-sample preparation configuration for analysis), both methods received a score of 0.75, indicating a moderate scope for direct coupling or reduced handling. Finally, criteria 2 (volume of problematic samples) and 10 (number of identified hazards) yielded a score of 0.50 for both UAE and MAE, primarily due to the use of ethanol, which, despite being one of the most environmentally friendly solvents for scalable extractions, remains flammable and requires careful safety measures.

In summary, the higher overall AGREEprep score for MAE (0.65 vs. 0.58) reflects its better performance in throughput (criterion 6), waste generation (criterion 4), and energy consumption (criterion 8). The results suggest that future efforts should focus primarily on reducing waste generation (criterion 4), improving extraction yield per unit mass (criterion 5), and optimizing energy consumption (criterion 8), as these factors were most decisive in distinguishing between the two techniques.



Conclusions

Both MAE and UAE demonstrated high extraction efficiency for bioactive compounds from cocoa husk at fine particle sizes (75–150 μm), with similar results for total phenolic content (TPC) and DPPH radical-scavenging activity. The selection of the most suitable method depends on the primary objective: if the goal is to maximize total phenolic content (TPC), MAE is preferred, as it yields higher phenolic extraction. Conversely, if the objective is to obtain extracts with high overall reducing capacity (as measured by FRAP and ABTS), UAE may be more suitable, though these assays are susceptible to interference from non-phenolic compounds such as gluconic acid and alkaloids, identified by HPLC. The UAE yielded significantly higher ABTS and FRAP values (indicating antioxidant activity), supporting its viability for industrial-scale applications. MAE is better suited when processing speed is the main driver, offering a comparable phytochemical profile while requiring careful management of its inherent risks. The temperature increase inherent to MAE is beneficial for cellular disruption and yield, but if not precisely regulated, localized overheating due to uneven dielectric absorption can degrade thermolabile antioxidant compounds. Furthermore, the primary technical limitation for MAE remains its scalability, given the complexities of achieving uniform microwave distribution in large-volume systems.

UHPLC–HRMS analysis showed that UAE and MAE produce extracts with highly comparable chemical profiles, characterized by intense signals for key methylxanthines, including theobromine and caffeine, as well as polyalcohols, organic acids, amino acids, and nitrogen-rich metabolites derived from cocoa fermentation and post-harvest processing. These findings confirm that both green technologies effectively release diverse and stable bioactive compounds from cocoa bean shells. Moreover, the similarity of the identified metabolites, predominantly alkaloids such as theobromine and caffeine, highlights the potential of these extracts for applications in nutraceutical formulations, antioxidant systems, functional beverages, and the development of micro- and nano-encapsulated products, given their compatibility with polymeric matrices.

Finally, the environmental assessment using the AGREEprep method validated both technologies as clean for the recovery of high-value metabolites, thereby supporting agro-industrial waste utilization and the cocoa value chain. However, MAE achieved a higher overall score (0.65 vs. 0.58) due to better throughput, lower waste generation, and reduced energy consumption, the key differentiating factors (criteria 4, 5, and 8). Therefore, future efforts should prioritize minimizing waste, improving extraction yield per unit mass, and optimizing energy use.

Author contributions

Erick Alvarez-Yanamango: conceptualization, methodology, investigation, resources, writing – original draft, writing – review & editing, visualization, supervision, project administration, funding acquisition. Milagros Huaytalla-Ramirez,

Nereyda Sarmiento-Perez, Gerald Chumpitaz-Huanqui, and Luis Napan-Tacca: investigation, formal analysis. Daniel Obregon-Valencia: methodology, formal analysis, validation, writing – review & editing, visualization. Alfredo Ibañez: investigation, supervision, formal analysis, writing – review & editing.

Conflicts of interest

The authors declare that no conflicts of interest or known personal connections could have influenced the work presented in this article.

Data availability

No primary research results, software or code have been included and no new data were generated or analysed as part of this study.

Acknowledgements

The authors acknowledge the financial support of the National Program for Scientific Research and Advanced Studies (PROCIENCIA) of the National Council of Science, Technology and Technological Innovation (CONCYTEC) through the project with contract PE501086489-2024-PROCIENCIA, and the Vice-Rectorate of Research of the Pontifical Catholic University of Peru for its financial support in the CAP 2024 - PI1115 project. Erick Alvarez-Yanamango acknowledges the Doctoral Scholarship in Engineering from Pontificia Universidad Católica del Perú (PUCP).

Notes and references

- 1 J. C. Hashimoto, J. C. Lima, R. M. S. Celeghini, A. B. Nogueira, P. Efraim, R. J. Poppi and J. A. L. Pallone, Quality Control of Commercial Cocoa Beans (*Theobroma cacao* L.) by Near-infrared Spectroscopy, *Food Anal. Methods*, 2018, **11**, 1510–1517.
- 2 M. Sánchez, A. Laca, A. Laca and M. Díaz, Cocoa Bean Shell: A By-Product with High Potential for Nutritional and Biotechnological Applications, *Antioxidants*, 2023, **12**(5), 1028, DOI: [10.3390/antiox12051028](https://doi.org/10.3390/antiox12051028).
- 3 ICCO, *Quarterly Bulletin of Cocoa Statistics - Production*, 2024.
- 4 O. Rojo-Poveda, L. Barbosa-Pereira, G. Zeppa and C. Stévigny, Cocoa bean shell—a by-product with nutritional properties and biofunctional potential, *Nutrients*, 2020, **12**(4), 1123, DOI: [10.3390/nu12041123](https://doi.org/10.3390/nu12041123).
- 5 Z. Ö. Cinar, M. Atanassova, T. B. Tumer, G. Caruso, G. Antika, S. Sharma, J. Sharifi-Rad and R. Pezzani, Cocoa and cocoa bean shells role in human health: An updated review, *J. Food Compos. Anal.*, 2021, **103**, 104115, DOI: [10.1016/j.jfca.2021.104115](https://doi.org/10.1016/j.jfca.2021.104115).
- 6 M. C. Castro, H. Villagarcía, A. Nazar, L. G. Arbeláez, M. L. Massa, H. Del Zotto, J. L. Ríos, G. R. Schinella and F. Francini, Cacao extract enriched in polyphenols prevents endocrine-metabolic disturbances in a rat model of prediabetes triggered by a sucrose rich diet, *J.*



- Ethnopharmacol.*, 2020, 247, 112263, DOI: [10.1016/j.jep.2019.112263](https://doi.org/10.1016/j.jep.2019.112263).
- 7 J. Li, H. D. Sesso, E. Kim, J. E. Manson, G. Friedenberg, A. Clar, T. Copeland, A. H. Shadyab, J. Wactawski-Wende, L. Tinker and S. Liu, Cocoa Extract Supplementation and Risk of Type 2 Diabetes: The Cocoa Supplement and Multivitamin Outcomes Study (COSMOS) Randomized Clinical Trial, *Diabetes Care*, 2023, 46, 2278–2284.
- 8 L. Goya, J. E. Kongor and S. de Pascual-Teresa, From Cocoa to Chocolate: Effect of Processing on Flavanols and Methylxanthines and Their Mechanisms of Action, *Int. J. Mol. Sci.*, 2022, 23(22), 14365, DOI: [10.3390/ijms232214365](https://doi.org/10.3390/ijms232214365).
- 9 D. Janitschke, A. A. Lauer, C. M. Bachmann, J. Winkler, L. V. Griebisch, S. M. Pilz, E. L. Theiss, H. S. Grimm, T. Hartmann and M. O. W. Grimm, Methylxanthines Induce a Change in the AD/Neurodegeneration-Linked Lipid Profile in Neuroblastoma Cells, *Int. J. Mol. Sci.*, 2022, 23(4), 2295, DOI: [10.3390/ijms23042295](https://doi.org/10.3390/ijms23042295).
- 10 M. Sánchez, A. Laca, A. Laca and M. Díaz, Cocoa Bean Shell: A By-Product with High Potential for Nutritional and Biotechnological Applications, *Antioxidants*, 2023, 12(5), 1028, DOI: [10.3390/antiox12051028](https://doi.org/10.3390/antiox12051028).
- 11 E. Alvarez-Yanamango, D. Obregon and A. Ibañez, A Comprehensive Review of the Development of Green Extraction Methods and Encapsulation of Theobromine from Cocoa Bean Shells for Nutraceutical Applications, *Food Eng. Rev.*, 2025, 17, 1083–1104.
- 12 V. L. Singleton and J. A. Rossi, Colorimetry Of Total Phenolics With Phosphomolybdic-Phosphotungstic Acid Reagents, *Am. J. Enol. Vitic.*, 1965, 16, 144–158, DOI: [10.5344/ajev.1965.16.3.144](https://doi.org/10.5344/ajev.1965.16.3.144).
- 13 W. Brand-Williams, M. E. Cuvelier and C. Berset, Use of a Free Radical Method to Evaluate Antioxidant Activity, *LWT – Food Sci. Technol.*, 1995, 28(1), 25–30, DOI: [10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5).
- 14 R. Re, N. Pellegrini, A. Proteggente, A. Pannala, M. Yang and C. Rice-Evans, Original Contribution Antioxidant Activity Applying An Improved ABTS Radical Cation Decolorization Assay, *Free Radical Biol. Med.*, 1999, 26(9–10), 1231–1237, DOI: [10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3).
- 15 I. F. F. Benzie and J. J. Strain, The Ferric Reducing Ability of Plasma (FRAP) as a Measure of “Antioxidant Power”: The FRAP Assay, *Anal. Biochem.*, 1996, 239(1), 70–76, DOI: [10.1006/abio.1996.0292](https://doi.org/10.1006/abio.1996.0292).
- 16 N. B. Sadeer, D. Montesano, S. Albrizio, G. Zengin and M. F. Mahomoodally, The versatility of antioxidant assays in food science and safety—chemistry, applications, strengths, and limitations, *Antioxidants*, 2020, 9(8), 709, DOI: [10.3390/antiox9080709](https://doi.org/10.3390/antiox9080709).
- 17 M. Minarti, N. Ariani, M. Megawati, A. Hidayat, M. Hendra, G. Primahana and A. Darmawan, in *E3S Web of Conferences*, EDP Sciences, 2024, vol. 503.
- 18 R. L. Prior, X. Wu and K. Schaich, Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements, *J. Agric. Food Chem.*, 2005, 53(10), 4290–4302, DOI: [10.1021/jf0502698](https://doi.org/10.1021/jf0502698).
- 19 K. M. Schaich, X. Tian and J. Xie, Hurdles and pitfalls in measuring antioxidant efficacy: A critical evaluation of ABTS, DPPH, and ORAC assays, *J. Funct. Foods*, 2015, 14, 111–125, DOI: [10.1016/j.jff.2015.01.043](https://doi.org/10.1016/j.jff.2015.01.043).
- 20 F. Shahidi and A. Samarasinghe, How to assess antioxidant activity? Advances, limitations, and applications of in vitro, in vivo, and ex vivo approaches, *Food Prod., Process. Nutr.*, 2025, 7(1), 1, DOI: [10.1186/s43014-025-00326-z](https://doi.org/10.1186/s43014-025-00326-z).
- 21 Y. Rodríguez-Carrasco, A. Gaspari, G. Graziani, A. Santini and A. Ritieni, Fast analysis of polyphenols and alkaloids in cocoa-based products by ultra-high performance liquid chromatography and Orbitrap high resolution mass spectrometry (UHPLC-Q-Orbitrap-MS/MS), *Food Res. Int.*, 2018, 111, 229–236.
- 22 W. Wojnowski, M. Tobiszewski, F. Pena-Pereira and E. Psillakis, AGREEprep – Analytical greenness metric for sample preparation, *TrAC, Trends Anal. Chem.*, 2022, 149, 116553, DOI: [10.1016/j.trac.2022.116553](https://doi.org/10.1016/j.trac.2022.116553).
- 23 F. Pena-Pereira, M. Tobiszewski, W. Wojnowski and E. Psillakis, A Tutorial on AGREEprep an Analytical Greenness Metric for Sample Preparation, *Adv. Sample Prep.*, 2022, 149, 116553, DOI: [10.1016/j.sampre.2022.100025](https://doi.org/10.1016/j.sampre.2022.100025).
- 24 M. J. Anderson and P. J. Whitcomb, *Optimizing Processes Using Response Surface Methods for Design of Experiments Second Edition RSM Simplified*, Boca Raton, FL, 2016.
- 25 C. Botella-Martínez, R. Lucas-Gonzalez, C. Ballester-Costa, J. Á. Pérez-álvarez, J. Fernández-López, J. Delgado-Ospina, C. Chaves-López and M. Viuda-Martos, Ghanaian cocoa (*Theobroma cacao* l.) bean shells coproducts: Effect of particle size on chemical composition, bioactive compound content and antioxidant activity, *Agronomy*, 2021, 11(2), 401, DOI: [10.3390/agronomy11020401](https://doi.org/10.3390/agronomy11020401).
- 26 A. Meullemiestre, E. Petitcolas, Z. Maache-Rezzoug, F. Chemat and S. A. Rezzoug, Impact of ultrasound on solid-liquid extraction of phenolic compounds from maritime pine sawdust waste. Kinetics, optimization and large scale experiments, *Ultrason. Sonochem.*, 2016, 28, 230–239.
- 27 M. C. Rincón Soledad, *Extracción asistida por microondas de compuestos fenólicos de los subproductos del beneficio del cacao (Theobroma cacao L.)*, 2023.
- 28 S. R. Dewi, L. Stevens, D. Irvine, R. Ferrari and E. Binner, *Microwave-Assisted Extraction Of Phenolic From Cacao Pod Husks-An Alternative For Valorisation*, Zenodo, 2021.
- 29 I. R. N. de Oliveira, R. F. Teófilo, E. B. de Oliveira, A. M. Ramos, F. A. R. de Barros, M. P. Maia and P. C. Stringheta, Evaluation of potential interfering agents on in vitro methods for the determination of the antioxidant capacity in anthocyanin extracts, *Int. J. Food Sci. Technol.*, 2017, 52, 511–518.
- 30 V. Lavelli, P. S. C. Sri Harsha, L. Torri and G. Zeppa, Use of winemaking by-products as an ingredient for tomato puree: The effect of particle size on product quality, *Food Chem.*, 2014, 152, 162–168.
- 31 A. Itam, M. S. Wati, V. Agustin, N. Sabri, R. A. Jumanah and M. Efdi, Comparative Study of Phytochemical, Antioxidant,



- and Cytotoxic Activities and Phenolic Content of *Syzygium aqueum* (Burm. f. Alston f.) Extracts Growing in West Sumatera Indonesia, *Sci. World J.*, 2021, DOI: [10.1155/2021/5537597](https://doi.org/10.1155/2021/5537597).
- 32 A. C. Mellinas, A. Jiménez and M. C. Garrigós, Optimization of microwave-assisted extraction of cocoa bean shell waste and evaluation of its antioxidant, physicochemical and functional properties, *LWT*, 2020, **127**, 109361, DOI: [10.1016/j.lwt.2020.109361](https://doi.org/10.1016/j.lwt.2020.109361).
- 33 N. Pavlović, S. Jokić, M. Jakovljević, M. Blažić and M. Molnar, Green extraction methods for active compounds from food waste - Cocoa bean shell, *Foods*, 2020, **9**(2), 140, DOI: [10.3390/foods9020140](https://doi.org/10.3390/foods9020140).
- 34 N. Pavlović, M. Jakovljević, M. Molnar and S. Jokić, Ultrasound-Assisted Extraction Of Active Compounds From Cocoa Bean Shell, *Hrana u zdravlju i bolesti*, 2021, **10**(2), 77–88.
- 35 A. H. Md Yusof, S. S. Abd Gani, U. H. Zaidan, M. I. E. Halmi and B. H. Zainudin, Optimization of an Ultrasound-Assisted Extraction Condition for Flavonoid Compounds from Cocoa Shells (*Theobroma cacao*) Using Response Surface Methodology, *Molecules*, 2019, **24**(4), 711, DOI: [10.3390/molecules24040711](https://doi.org/10.3390/molecules24040711).
- 36 A. Taqi, E. Farcot, J. P. Robinson and E. R. Binner, Understanding microwave heating in biomass-solvent systems, *Chem. Eng. J.*, 2020, **393**, 124741, DOI: [10.1016/j.cej.2020.124741](https://doi.org/10.1016/j.cej.2020.124741).
- 37 Y. Mao, J. Robinson and E. Binner, Understanding heat and mass transfer processes during microwave-assisted and conventional solvent extraction, *Chem. Eng. Sci.*, 2021, **233**, 116418, DOI: [10.1016/j.ces.2020.116418](https://doi.org/10.1016/j.ces.2020.116418).
- 38 X. Zhu, W. Guo, X. Wu and S. Wang, Dielectric properties of chestnut flour relevant to drying with radio-frequency and microwave energy, *J. Food Eng.*, 2012, **113**, 143–150.
- 39 D. Wang, X. Li, X. Hao, J. Lv and X. Chen, The Effects of Moisture and Temperature on the Microwave Absorption Power of Poplar Wood, *Forests*, 2022, **13**(2), 309, DOI: [10.3390/f13020309](https://doi.org/10.3390/f13020309).
- 40 M. D. Pavlović, A. V. Buntić, S. S. Šiler-Marinković and S. I. Dimitrijević-Branković, Ethanol influenced fast microwave-assisted extraction for natural antioxidants obtaining from spent filter coffee, *Sep. Purif. Technol.*, 2013, **118**, 503–510.
- 41 Q. Hu, Y. He, F. Wang, J. Wu, Z. Ci, L. Chen, R. Xu, M. Yang, J. Lin, L. Han and D. Zhang, Microwave technology: a novel approach to the transformation of natural metabolites, *Chin. Med.*, 2021, **16**(1), 87, DOI: [10.1186/s13020-021-00500-8](https://doi.org/10.1186/s13020-021-00500-8).
- 42 J. Zhong, N. Tang, B. Asadzadeh and W. Yan, Measurement and Correlation of Solubility of Theobromine, Theophylline, and Caffeine in Water and Organic Solvents at Various Temperatures, *J. Chem. Eng. Data*, 2017, **62**, 2570–2577.
- 43 A. C. Aprotosoaie, S. V. Luca and A. Miron, Flavor Chemistry of Cocoa and Cocoa Products-An Overview, *Compr. Rev. Food Sci. Food Saf.*, 2016, **15**, 73–91.
- 44 T. Lefeber, M. Janssens, F. Moens, W. Gobert and L. De Vuyst, Interesting starter culture strains for controlled cocoa bean fermentation revealed by simulated cocoa pulp fermentations of cocoa-specific lactic acid bacteria, *Appl. Environ. Microbiol.*, 2011, **77**, 6694–6698.
- 45 H. Nishimura, Y. Shiwa, S. Tomita and A. Endo, Microbial composition and metabolic profiles during machine-controlled intra-factory fermentation of cocoa beans harvested in semitropical area of Japan, *Biosci. Microbiota, Food Health*, 2024, **43**, 29–42.
- 46 Y. Kitani, S. P. Putri and E. Fukusaki, Investigation of the effect of processing on the component changes of single-origin chocolate during the bean-to-bar process, *J. Biosci. Bioeng.*, 2022, **134**, 138–143.
- 47 C. Xin Hui, N. Lindley, Y. Kanagasundaram and Y. Chow, in *Theobroma Cacao - Past, Present and Future Insights [Working Title]*, IntechOpen, 2025.
- 48 F. Moens, T. Lefeber and L. De Vuyst, Oxidation of metabolites highlights the microbial interactions and role of *Acetobacter pasteurianus* during cocoa bean fermentation, *Appl. Environ. Microbiol.*, 2014, **80**, 1848–1857.
- 49 R. F. Peña-Correa, B. A. Mogol, C. Fryganas and V. Fogliano, Fluidized-Bed-Roasted Cocoa Has Different Chemical Characteristics than Conventionally Roasted Cocoa, *J. Agric. Food Chem.*, 2023, **71**, 10203–10211.
- 50 C. R. Balcázar-Zumaeta, E. Fernández-Romero, A. S. Lopes, N. R. Ferreira, G. C. A. Chagas-Júnior, I. Yoplac, H. A. López-Trigoso, M. L. Tuesta-Occ, I. Maldonado-Ramirez, J. L. Maicelo-Quintana, I. S. Cayo-Colca and E. M. Castro-Alayo, Amino acid profile behavior during the fermentation of Criollo cocoa beans, *Food Chem.: X*, 2024, **22**, 101486, DOI: [10.1016/j.fochx.2024.101486](https://doi.org/10.1016/j.fochx.2024.101486).
- 51 V. Barišić, M. Kopjar, A. Jozinović, I. Flanjak, Đ. Ačkar, B. Miličević, D. Šubarić, S. Jokić and J. Babić, The chemistry behind chocolate production, *Molecules*, 2019, **24**(17), 3163, DOI: [10.3390/molecules24173163](https://doi.org/10.3390/molecules24173163).
- 52 M. Granvogl, S. Bugan and P. Schieberle, Formation of amines and aldehydes from parent amino acids during thermal processing of cocoa and model systems: New insights into pathways of the strecker reaction, *J. Agric. Food Chem.*, 2006, **54**, 1730–1739.
- 53 C. Hue, Z. Gunata, A. Breyse, F. Davrieux, R. Boulanger and F. X. Sauvage, Impact of fermentation on nitrogenous compounds of cocoa beans (*Theobroma cacao* L.) from various origins, *Food Chem.*, 2016, **192**, 958–964.
- 54 K. Purbaningrum, C. Hidayat, L. D. Witasari and T. Utami, Flavor Precursors and Volatile Compounds Improvement of Unfermented Cocoa Beans by Hydrolysis Using Bromelain, *Foods*, 2023, **12**(4), 820, DOI: [10.3390/foods12040820](https://doi.org/10.3390/foods12040820).
- 55 C. Q. Chen and E. Robbins, Chocolate flavor via the Maillard reaction, *ACS Symp. Ser.*, 2000, **754**, 286–292.
- 56 T. F. Soares and M. B. P. P. Oliveira, Cocoa By-Products: Characterization of Bioactive Compounds and Beneficial Health Effects, *Molecules*, 2022, **27**(5), 1625, DOI: [10.3390/molecules27051625](https://doi.org/10.3390/molecules27051625).
- 57 X.-Q. Zheng, Y. Koyama, C. Nagai and H. Ashihara, Biosynthesis, accumulation and degradation of theobromine in developing *Theobroma cacao* fruits, *J.*



- Plant Physiol.*, 2004, **161**(4), 363–369, DOI: [10.1078/0176-1617-01253](https://doi.org/10.1078/0176-1617-01253).
- 58 P. H. Pour, F. M. Suzaei and S. M. Daryanavard, Greenness assessment of microextraction techniques in therapeutic drug monitoring, *Bioanalysis*, 2024, **16**, 249–278.
- 59 S. S. Ferreira, T. A. Brito, A. P. R. Santana, T. G. S. Guimarães, R. S. Lamarca, K. C. Ferreira, P. C. F. L. Gomes, A. Oliveira, C. D. B. Amaral and M. H. Gonzalez, Greenness of procedures using NADES in the preparation of vegetal samples: Comparison of five green metrics, *Talanta Open*, 2022, **6**, 100131, DOI: [10.1016/j.talo.2022.100131](https://doi.org/10.1016/j.talo.2022.100131).

