



Cite this: DOI: 10.1039/d6fb00080k

# Fully unlocking the potential of Chenpi (*Citri Reticulatae Pericarpium*) as a functional ingredient in food development

Guoqiang Zhang,<sup>a</sup> HioTong Mak,<sup>c</sup> Zheng Zhang,<sup>b</sup> Ting Zhang,<sup>b</sup> Yuanhui Wang,<sup>de</sup> Run-Yang Zhang,<sup>f</sup> Hua-Min Liu,<sup>f</sup> Shaobo Zhou<sup>g</sup> and Bin Wu<sup>b</sup>

A shift to a sustainable and healthy diet is key to the establishment of a sustainable food system and the achievement of the sustainable development agenda. To enable this dietary shift, the exploration and development of sustainable, accessible, and affordable ingredients that can be integrated into the daily diet and the protection and promotion of traditional healthy diets are important actions. In the traditional Chinese diet, Chenpi (*Citri Reticulatae Pericarpium*) is a medicinal and food homology ingredient with extensive health-promoting effects and is widely used for culinary applications and food development. This makes Chenpi a valuable ingredient that has high potential to contribute to the achievement of a widespread, sustainable and healthy diet. Therefore, this review provides an overview of the major constituents of Chenpi and current advances in its development and food applications, aiming to maximise its potential in daily diets and contribute to the achievement of sustainable and healthy diets and food systems.

Received 17th March 2026

Accepted 16th April 2026

DOI: 10.1039/d6fb00080k

rsc.li/susfoodtech

## Sustainability spotlight

Chenpi, a medicine and food homology ingredient in China, contains abundant bioactive compounds with various health-promoting benefits. According to the EAT-Lancet Commission, a healthy and sustainable diet is key to the transition to a sustainable food system and to achieving the sustainable development agenda. In this context, the concept of medicine and food homology and the development of Chenpi could offer novel insights and opportunities to ensure global nutrition security by bridging traditional dietary wisdom with the national and global initiatives (e.g. the planetary health diet and the Sustainable Development Goals), thereby promoting the transition towards healthy and sustainable diets and food systems, which aligns well with SDG 2, SDG 3, and SDG 12.

## 1. Introduction

The rapid increase in global population and food demand, along with the food waste issue in the current food system, exerts significant pressure on the environment and strongly threatens food security and human survival.<sup>1,2</sup> Therefore, the EAT-Lancet Commission proposed a healthy and sustainable reference diet (referred to as the planetary health diet) to promote a global shift in food systems and diets, thereby

reducing environmental impact, promoting food system sustainability and human health, and achieving win-win outcomes between humans and the environment.<sup>3,4</sup> To achieve this, a series of actions are required, such as education, food environment interventions and the protection and promotion of healthy traditional diets.<sup>4,5</sup> In this context, the concept of medicine and food homology has received much attention, which emphasises the close connection and mutual transformation between food and medicine.<sup>6,7</sup> In other words, food not only serves various functions to meet human needs, but also contains many pharmacological properties that can promote human health.<sup>8</sup> This dietary concept has encouraged the continued exploration of the potential of all ingredients and their complete utilisation and development in daily diets to meet the demand for natural and healthy foods.<sup>9</sup>

Chenpi, also known as *Citri Reticulatae Pericarpium*, originates from the dried or aged peel of *Citrus reticulata* Blanco or its cultivars and has been widely used as both a medicine and food ingredient based on the concept of medicine and food homology in China.<sup>10,11</sup> In addition, in traditional Chinese medicine, Chenpi has many health-promoting benefits, particularly in maintaining

<sup>a</sup>School of Science, Coventry University, Coventry, UK. E-mail: ae4214@coventry.ac.uk<sup>b</sup>Institute of Agro-products Storage and Processing, Xinjiang Academy of Agricultural Science, Urumqi 830091, China<sup>c</sup>Faculty of Health and Life Sciences, Bristol Medical School, University of Bristol, UK<sup>d</sup>College of Food Science and Engineering, Henan University of Technology, Zhengzhou, 450001, China<sup>e</sup>Henan Province Wheat-flour Staple Food Engineering Technology Research Centre, Henan University of Technology, Zhengzhou, 450001, China<sup>f</sup>College of Food Science and Engineering, Institute of Special Oilseed Processing and Technology, Henan University of Technology, Zhengzhou 450001, China<sup>g</sup>School of Science, Faculty of Engineering and Science, University of Greenwich, Kent, UK

gastrointestinal health, such as relieving indigestion and loss of appetite.<sup>12,13</sup> Besides, many studies have reported and demonstrated the various biological activities of Chenpi, such as antioxidant, anti-inflammatory, and the modulation of gut microbiota.<sup>14–17</sup> This makes Chenpi a valuable ingredient in China and its total market size exceeded 10 billion Chinese yuan in 2021.<sup>18</sup> However, citrus peel is regarded as a major by-product in the citrus fruit processing line, and is scarcely utilised and often considered waste that is discarded in landfills.<sup>19,20</sup> Importantly, citrus peel has high potential value and should be considered a valuable ingredient rather than waste.<sup>21</sup> In this regard, utilisation of citrus peel as a value ingredient not only reduces food waste generation, but also offers opportunities to produce healthy and sustainable food for human consumption.<sup>22</sup> Nevertheless, reintroducing citrus peel into the food supply chain requires the development of appropriate valorisation strategies and the establishment of a new value chain.<sup>23,24</sup> In this regard, the traditional dietary wisdom of Chenpi and its development could present a novel utilisation pathway that transforms citrus peel into healthy, affordable, and sustainable ingredients in daily diets, thereby reducing citrus waste generation and promoting human health. Therefore, a better understanding of Chenpi and its development and applications can not only further unlock the potential of Chenpi in sustainable and healthy diets, but also promote the sustainable utilisation of citrus resources.<sup>25–27</sup>

To this end, this review provides an overview of the major constituents of Chenpi and summarises the recent advances in Chenpi development, particularly in the extraction of bioactive compounds and food applications, aiming to promote a better understanding of Chenpi and its further development in food products and to offer new insights for the value innovation of citrus peel to produce a healthier and more sustainable food supply, thereby promoting global human well-being and contributing to healthier and sustainable food systems and diets.

## 2. Overview of bioactive compounds in Chenpi

Chenpi (*Citri Reticulatae Pericarpium*, Fig. 1) is a traditional medicine and food homologous ingredient in China, where 4 varieties are commercially cultivated and consumed, namely *C. reticulata* “Chachi”, *C. reticulata* “Dahongpao”, *C. reticulata* “Unshiu”, and *C. reticulata* “Tangerina”.<sup>28,29</sup> Chenpi contains abundant bioactive constituents such as polysaccharides, alkaloids, carotenoids, terpenoids, and phenolic compounds, which are associated with Chenpi's extensive health benefits,



Fig. 1 *Citrus reticulata* fruit and its dried pericarp (Chenpi).

such as antioxidant, anti-inflammatory, anti-hyperglycaemic and anti-hyperlipidaemic activities, as well as the modulation of gut microbiota.<sup>16,17,30–33</sup> Notably, the content of bioactive compounds in Chenpi varies depending on many factors, such as variety, harvesting conditions, drying methods, and storage time (aging).<sup>28,34–37</sup> Zhu *et al.*<sup>37</sup> reported that the contents of ascorbic acid and  $\beta$ -carotene decrease with increasing storage (aging) time. In contrast, Chen *et al.*<sup>36</sup> reported that the total flavonoid content of Chenpi (*Citrus reticulata* ‘Chachi’) increased during storage (aging), which could be attributed to bacterial biotransformation. During the aging process of Chenpi, the main fungal species, including *Penicillium* and *Aspergillus*, actively participate in flavonoid metabolism through their enzymatic activities.<sup>38,39</sup> Tang *et al.*<sup>40</sup> demonstrated that *Aspergillus tubingensis* converted nobiletin into 3'-demethylnobiletin during the Chenpi aging process. In addition, *Aspergillus niger* has been reported to be involved in the transformation of hesperidin into hesperetin in the aging of Chenpi, thereby enhancing antioxidant capacity.<sup>41,42</sup> On the other hand, a variety of flavonoids exist in plant cells and are not easily released into the compact cell wall structures; however, the action of microbial enzymes (*e.g.* cellulase and  $\beta$ -glucosidase) can disrupt the plant cell wall structure and then promote flavonoid release.<sup>43,44</sup>

Regarding bioactive compounds in Chenpi, approximately 150 phenolic compounds have been identified in Chenpi, among which flavonoids are relatively abundant.<sup>45–48</sup> In the study by Zhang *et al.*,<sup>47</sup> 95 phenolic compounds were identified in Chenpi, including 92 flavonoids, 2 coumarins, and 1 phenolic acid. In contrast, Bian *et al.*<sup>48</sup> identified 65 flavonoids and 51 phenolic acids in Chenpi using ultra-high performance liquid chromatography-quadrupole time-of-flight mass spectrometry (UPLC-Q-TOF/MS) analysis. In addition, in the study by Chai *et al.*,<sup>46</sup> the authors identified and quantified 35 phenolic compounds in Chenpi using an ultra-performance liquid chromatography (UHPLC) system coupled with a Q-orbitrap mass spectrometer, and subsequently 10 phenolic compounds were identified as major antioxidant contributors through an *in vitro* chemical antioxidant capacity test (ABTS assay), including ferulic acid (0.057–0.164 mg g<sup>-1</sup>), *cis*-4-hydroxycinnamic acid (0.009–0.037 mg g<sup>-1</sup>), caffeic acid (0.004–0.014 mg g<sup>-1</sup>), protocatechuic acid (0.004–0.009 mg g<sup>-1</sup>), hesperidin (0.280–0.506 mg g<sup>-1</sup>), orientin (0.008–0.096 mg g<sup>-1</sup>), rutin (0.016–0.104 mg g<sup>-1</sup>), diosmin (0.030–0.227 mg g<sup>-1</sup>), isoorientin (0.004–0.030 mg g<sup>-1</sup>), and chrysoeriol (0–0.015 mg g<sup>-1</sup>). In addition, Shi *et al.*<sup>28</sup> used UPLC to identify and quantify 9 phenolic compounds (3 phenolic acids and 5 flavonoids) in Chenpi, including protocatechuic acid (0.043–0.075 mg g<sup>-1</sup>), vanillic acid (0.085–0.114 mg g<sup>-1</sup>), ferulic acid (0.186–0.437 mg g<sup>-1</sup>), narirutin (0.452–1.159 mg g<sup>-1</sup>), hesperidin (45.863–80.389 mg g<sup>-1</sup>), sinensetin (0.483–0.583 mg g<sup>-1</sup>), nobiletin (4.062–5.358 mg g<sup>-1</sup>), and tangeretin (2.870–3.833 mg g<sup>-1</sup>). In terms of dietary fibre, the dietary fibre content in Chenpi reaches 62.87% (w/w), whereas the contents of insoluble dietary fibre (IDF) and soluble dietary fibre (SDF) are 49.61% (w/w) and 13.02% (w/w), respectively.<sup>49</sup> Furthermore, Wang *et al.*<sup>49</sup> detected uronic acid (129.48 mg galacturonic acid equivalents per g) in Chenpi's SDF, indicating



that complex pectic polysaccharides are present in Chenpi. Notably, pectin has many biological activities and is widely used in the food industry as a gelling agent, emulsifier, and thickener.<sup>50,51</sup> This has driven increasing interest in the extraction of polysaccharides from Chenpi as functional or health-promoting ingredients, which will be introduced in Section 4.

### 3. Key volatile aroma compounds in Chenpi

Flavour is an important parameter of food quality and consumer acceptance, since it determines the overall palatability of food products and the sensory experience of consumers.<sup>52–54</sup> As demand and interest continue to grow for high sensory quality, healthier, and more sustainable food products, this drives the food industry to seek natural functional flavour ingredients in food innovation, aiming to enhance food nutritional quality and provide more attractive flavours to meet consumer demand.<sup>55,56</sup> In this regard, Chenpi, as a highly medicinal and food ingredient, presents a promising option for food development to enhance food flavour and provide additional health benefits.<sup>57</sup> Therefore, the identification of key volatile compounds in Chenpi could help better understand its overall taste and flavour perception and further facilitate its incorporation into suitable food systems.<sup>58,59</sup> Table 1 presents the identified volatile aroma active compounds in Chenpi that contribute to shaping its overall flavour profile.

It can be seen from Table 1 that  $\alpha$ -pinene (fresh, woody, piney, and herbal),  $\beta$ -myrcene (spicy and woody),  $\beta$ -limonene (citrus, fruity, and spicy), linalool (aniseed, citrus, floral, pleasant scent, and sweet), dimethyl anthranilate (floral, pepper, spicy, and citrusy),  $\gamma$ -terpinene (pine-like smell, woody, lemon-flavoured, sweet, and floral), and 2-methoxy-4-vinylphenol (pungent and floral) are considered to be important contributors to Chenpi flavour, providing fresh, woody, piney, herbal, citrus, fruity, spicy, floral, and sweet notes. In addition, other high-content compounds, such as  $\alpha$ -terpineol (floral and woody), octanal (orange flavour), thymol (herbal and medicine), and  $\beta$ -pinene (woody, pine-like smell, herbal, and pungent) enhance the pine-like, floral, citrus, woody, sweet, and herbal notes in Chenpi flavour. Besides, Li *et al.*<sup>57</sup> reported that the Chenpi aroma (contributing citrus and herbal notes) can enhance the flavour perception of sweetness and saltiness and has high application potential as a functional flavour agent in sugar- and salt-reduced foods.

Notably, the volatile profile of Chenpi is dependent on many factors, including geographical origin, cultivated methods, seasonality, processing methods and conditions, and storage time.<sup>28,60–63</sup> Cao *et al.*<sup>64</sup> explored the effect of tree age on volatile compounds of Chenpi and reported that Chenpi obtained from trees aged 5 years exhibited a fresher, more citrus aroma, whereas Chenpi obtained from trees aged 20 years exhibited more pronounced caramel and green leaf notes.

Jiang *et al.*<sup>65</sup> coupled headspace solid-phase microextraction gas chromatography-mass spectrometry (HS-SPME-GC-MS)

with gas chromatography-olfactometry-mass spectrometry (GC-O-MS) to evaluate the changes in aroma compounds of Chenpi (obtained from Xinhui, Guangdong, China) during storage, ranging from 2 to 25 years. The contents of citrus and fruity aroma compounds (*i.e.*  $\beta$ -limonene and  $\gamma$ -terpinene) decreased during storage, while those with spicy, woody, and herbal aroma notes (*i.e.*  $\alpha$ -farnesene) increased, resulting in an overall rise in spicy, woody, and herbal notes and a reduction in citrus and fruity notes. Similarly, Liu *et al.*<sup>66</sup> analysed the effect of aging stages on the flavour characteristics of Chenpi (obtained from Xinhui, Guangdong, China), supporting the finding of Liu *et al.*<sup>66</sup> The authors employed 5 different aging stages of Chenpi, including 5, 10, 20, 30, and 40 years and reported that the aroma attributes of citrus decreased with the prolongation of aging time, while the fresh, floral, grassy, and woody attributes increased. This change could be attributed to the synergistic effects of multiple reactions, such as oxidative, hydrolytic, and enzymatic reactions, which increase the complexity of volatile compounds and alter the aroma balance rather than only the degradation or transformation of monoterpenes, *i.e.*  $\beta$ -limonene and  $\gamma$ -terpinene.<sup>60</sup>

Zhang *et al.*<sup>41</sup> found that the lemon, sweet and musk aromas of Chenpi changed to apple, coffee, and pineapple aromas through fermentation using *Aspergillus niger*. In another study, Shen *et al.*<sup>67</sup> fermented Chenpi using *Monascus anka* and *Saccharomyces cerevisiae* for 14 days and discovered that the citrus, sweet, herbal, woody, and pine-like aromas of Chenpi were enhanced, while the pungent odours (*e.g.* grass, green, and sour) were decreased. In addition, Chen *et al.*<sup>61</sup> investigated the effect of steaming treatment on the volatile profile of Chenpi and the results showed that short steaming treatment (20 s) could reduce the overall pungency level, thereby improving the sensory quality. In a recent study, Wu *et al.*<sup>60</sup> reported that enzymatic hydrolysis (using pectinase and papain) can promote the production of various key aroma compounds (associated with fruity, floral, and oily attributes) and alter the overall aroma of Chenpi.

The above studies indicate that processing is a viable approach to modulate the aroma profile of Chenpi, thereby improving flavour quality and consumer acceptance. Nevertheless, more studies are needed to investigate the effects of different processing methods and conditions on the flavour characteristics of Chenpi and discover the mechanisms of flavour formation, thereby providing more information to enhance Chenpi flavour quality and better expand the applications of Chenpi and its derivatives in food innovation. Besides, during the aging process, the overall flavour and sensory properties of Chenpi can be changed due to the degradation and/or transformation of volatile compounds. According to traditional Chinese concepts, the flavour of Chenpi develops uniquely during the aging process; however, such changes may influence the sensory quality of the final Chenpi-based products and consumer acceptance when considering different consumer groups.<sup>43,60</sup> In this regard, the identification of flavour differences and consumer acceptance for future research would guide the subsequent development of Chenpi products.



Table 1 Most common volatile compounds in Chenpi, including content, the aroma description, and the odour activity value (OAV)

Number	Volatile compounds	Content (mg kg <sup>-1</sup> )	Odour activity value (OAV)	Aroma description	References
1	$\alpha$ -Pinene	43.78–1550	273–3127.014	Fresh, camphor, earthy, woody odour, pine-like, and herbal	28, 135 and 136
2	$\beta$ -Pinene	170–1160	26–36	Woody-green, pine-like smell, herbal, and pungent	28 and 136
3	$\beta$ -Myrcene	31.782–1930	3427–21188.2	Ethereal, resinous, soapy, spicy odour, woody, and pungent	28 and 135
4	$\alpha$ -Terpinene	3.47–250	1.45–2.38	Woody, lemon-flavored, and orange	28, 135 and 137
5	<i>D</i> -Limonene	3291.64–78360	11720–24026.58	Citrus, ethereal, fruity odour, and spicy	28 and 135–137
6	$\gamma$ -Terpinene	130.53–13280	6.54–130.53	Pine-like smell, woody, lemon-flavoured, sweet, and floral	28 and 135–137
7	Terpinolene	110–700	—	Citrus flavor	28
8	Linalool	20–561.39	368–20049.61	Aniseed, citrus, floral, terpene (pleasant scent), and sweet	28, 136 and 137
9	Citronellal	20–80	—	Japanese pepper tree, floral, and lemon scents	28
10	(+)-4-Terpineol	40–370.81	57.94	Camphoraceous, earthy, musty odour (pleasant), and woody	28 and 137
11	(S)-(-)-Perillaldehyde	20–70	—	Minty and herbal	28
12	Thymol	20–70	381.80	Herbal, pleasant aromatic odour, and medicine	28 and 137
13	Carvacrol	30–40	3–5	Spicy and citrusy	28 and 136
14	Dimethyl anthranilate	30–740	23917–28563	Floral, pepper, spicy, and citrus	28 and 136
15	(-)- $\beta$ -Caryophyllene	10–110	—	Spicy	28
16	2-Methoxy-4-vinylphenol	253.67	13351.10	Pungent and floral	137
17	Geraniol	16.99	1699.27	Rose	137
18	Octanal	78.71	342.23	Orange flavour	137
19	$\alpha$ -Terpineol	4.92–250.54	15.66–291.32	Floral and woody	137
20	$\beta$ -Ionone	1.94	231.49	Woody	137
21	Cymenol	29.62	164.56	Pungent and refreshing	137
22	1-Octanol	3.32	144.30	Oily and fruity	137
23	(-)-Carvone	6.18	92.26	Mint and spicy	137
24	Furfural	274.80	91.60	Nut	137
25	Nonanal	6.45	64.50–364	Oily, sweet, and orange	136 and 137
26	<i>n</i> -Decanoic acid	7.17	55.19	Floral	137
27	1-Nonanol	10.10	10.10	Orange scent	137
28	Copaene	3.36	1.61	—	135
29	Decanal	—	573–669	Fresh and spicy	136
30	$\alpha$ -Phellandrene	—	7–8	Cream and fruity	136



Table 2 Extraction of polysaccharides from Chenpi

Extraction method	Environment impact	Key findings	References
Hot water extraction	Low-medium	A yield of 8.9%	80
Hot water extraction	Low-medium	Molecular weight of 55.4 kDa The yield ranges from 17% to 18.6%, and the yield decreased with increasing storage time (from 5–20 years)	81
Hot water extraction	Low-medium	A yield of 4.28% The monosaccharide composition (mol%): arabinose (49.68%), galacturonic acid (26.02%), galactose (10.70%), rhamnose (5.25%), glucose (4.34%), mannose (3.26%), and xylose (0.73%) In an animal model (male Balb/c mice), treatment with extracted pectin (50 mg kg <sup>-1</sup> and 250 mg kg <sup>-1</sup> ) alleviated gastric lesions, decreased oxidative stress, and suppressed inflammatory responses in alcohol-induced mice	83
Hot water extraction	Low-medium	The monosaccharide composition (mol%): galacturonic acid (64.7%), arabinose (16.8%), galactose (13.7%), rhamnose (2.9%), and glucose (1.9%) In a high-fat diet-induced casper zebrafish model of non-alcoholic fatty liver disease, incorporation of extracted polysaccharides into zebrafish feed (10 µg mL <sup>-1</sup> and 40 µg mL <sup>-1</sup> ) ameliorated hepatic steatosis, and this effect was enhanced with increasing treatment time and dosage	82
Acid extraction (using HCl)	High	The yield reached 25.77% under the optimised extraction conditions, namely, a liquid-to-solid ratio of 25 : 1 (v/w), a temperature of 80 °C, a pH of 0.5, and an extraction time of 90 min The molecular weight of extracted polysaccharides obtained using different storage times (0, 1, 5, 10, and 15 years) ranged from 10–16 kDa, whereas the longest-aged Chenpi resulted in the lowest molecular weight of polysaccharide (10.57 kDa) The degree of esterification ranged from 54.68%–72.40%	38
Fermentation ( <i>Bacillus licheniformis</i> )	Low	A molecular weight of 3.72 kDa A solubility of 189.85 mg mL <sup>-1</sup> The monosaccharide molar ratio of arabinose (31.75%), galactose (24.77%), galacturonic acid (20.94%), fucose (7.83%), rhamnose (7.53%), glucose (4.08%), mannose (1.93%), and glucuronic acid (1.15%) <i>In vitro</i> fermentation, the extracted polysaccharides increased the relative abundances of <i>Bacteroides</i> , <i>Parabacteroides</i> , <i>Phascolarctobacterium</i> , and <i>Lachnospira</i> and decreased the relative abundances of <i>Shigella</i> , <i>Megamonas</i> , and <i>Haemophilus</i>	93
Ultrasound-assisted extraction	Low	The yield of 7.0% was under optimised extraction conditions, namely a liquid-to-solid ratio of 30 : 1 (v/w), a pH of 4.4, a temperature of 90 °C, a power of 250 W, and an extraction time of 20 min A molecular weight of 122.0 kDa The monosaccharide ratio (mol%): galacturonic acid (51.17%), arabinose (25.63%), galactose (12.25%), and glucose (5.46%) In an animal model, the addition of extracted polysaccharides (60 mg kg <sup>-1</sup> d <sup>-1</sup> ) to the feed of male mice (high fat diet, 45% fat, 16.8 kcal per g diet) for 4 weeks reduced body weight and adipose tissue accumulation The extracted polysaccharides stimulated the growth of <i>Lactobacillus johnsonii</i>	88



## 4. Extraction of Chenpi polysaccharides

Polysaccharides are natural bioactive compounds that are widely applied as functional ingredients in the food, pharmaceutical, and cosmetics industries, owing to their favourable physicochemical properties (e.g. gelling and emulsifying properties) and biological activities (e.g. antioxidant, anti-inflammatory, antibacterial, and anti-diabetes activities).<sup>68–71</sup> With changes in dietary patterns, there is increasing interest in food products with potential health benefits.<sup>72,73</sup> This makes Chenpi polysaccharides valuable and high potential ingredients in functional food development. Notably, many studies have demonstrated that extraction methods determine the physicochemical properties and biological activities of polysaccharides and consequently affect their value and further applications.<sup>74–77</sup> As mentioned earlier, polysaccharides are important bioactive compounds in Chenpi; therefore, appropriate extraction methods are important for developing high-quality Chenpi polysaccharides and are key to their commercial applications.<sup>78</sup> Table 2 summarises current advances in the extraction of Chenpi polysaccharides.

Hot water extraction is the most commonly used method for the extraction of polysaccharides, with the advantages of low cost, simple operation, and low equipment requirements.<sup>79</sup> In the study by Yang *et al.*,<sup>80</sup> the yield of Chenpi polysaccharides reached 8.9% at a liquid-to-solid ratio of 10 : 1 (w/w) at 65 °C for 120 min. After purification using DEAE Sepharose anion exchange chromatography, the authors reported that the molecular weight of Chenpi polysaccharides was 55.4 kDa, and they mainly consisted of arabinose and galacturonic acid (with a molar ratio of 1 : 2.3). In another study by Zhou *et al.*,<sup>81</sup> the yield reached 18.6% under extraction conditions of 80 °C for 120 min. Besides, the extracted Chenpi polysaccharides increased the viability of RAW264.7 cells at a concentration of 0.2 mg mL<sup>-1</sup>. In addition, Wang *et al.*<sup>82</sup> extracted Chenpi polysaccharides using the hot water method and subsequently investigated their monosaccharide composition and anti-inflammatory effects on a high-cholesterol-fed (HCF) casper zebrafish model. The results showed that the molecular weight (MW) of the extracted Chenpi polysaccharide was 9.8 kDa, and the monosaccharide composition consisted of rhamnose (2.9 mol%), arabinose (16.8 mol%), galactose (13.7 mol%), glucose (1.9 mol%), and galacturonic acid (64.7 mol%). Afterwards, in the HCF casper zebrafish model, dietary treatment with 0.5 wt% of Chenpi polysaccharides significantly reduced ( $p < 0.05$ ) hepatic macrophage infiltration, while treatment at 1 wt% significantly reverted hepatic macrophage infiltration to normalcy. In a recent study, Liang *et al.*<sup>83</sup> extracted the pectic polysaccharide fraction from Chenpi in a liquid-to-solid ratio of 15 : 1, v/w at 90 °C for 90 min for 3 extraction cycles, and the yield was 4.28%. Furthermore, the monosaccharide composition analysis results showed that the extracted pectic Chenpi polysaccharide was composed of arabinose (49.26 mol%), galacturonic acid (26.02 mol%), rhamnose (5.25 mol%), glucose (4.34 mol%), mannose (3.26 mol%), and xylose (0.73 mol%).

In addition to the hot water extraction method, acid extraction has also been applied in the extraction of Chenpi polysaccharides. Yue *et al.*<sup>88</sup> employed acid extraction (using HCl to adjust the pH) to extract Chenpi polysaccharides and optimised the extraction conditions in terms of pH, temperature, and liquid-to-solid ratio. The yield reached 25.77% under the optimised extraction conditions, namely, a liquid-to-solid ratio of 25 : 1 (v/w), pH of 0.5, and temperature of 80 °C. Notably, HCl (highly corrosive and hazardous solvent) use and its subsequent neutralisation steps can increase the risk of accidents and generate waste, which have impact on human health and the environment.<sup>84,85</sup> From an environmental perspective and the principles of green chemistry, avoiding toxic and hazardous chemicals in the extraction process is important to prevent waste generation, reduce the risk of accidents, and support sustainability.<sup>86,87</sup> Therefore, while acid extraction in the above study offers high yield for the extraction of Chenpi polysaccharides, its practical application should carefully balance extraction efficiency with environmental and safety considerations.

In another study, Li *et al.*<sup>88</sup> used the ultrasound-assisted extraction method to extract polysaccharides from Chenpi and optimised the extraction conditions using a Box-Behnken design, namely, a liquid-to-material ratio of 30 : 1 (v/w), a pH of 4.4, a temperature of 90.0 °C, an ultrasonic power of 250.0 W, and an extraction time of 20 min. Under these conditions, the polysaccharide yield reached 7.0%. After purification, the molecular weight of the extracted Chenpi polysaccharide was 122.0 kDa, and the monosaccharide composition was galacturonic acid (51.17%), arabinose (25.63%), galactose (12.25%), and glucose (5.46%). In addition, the authors observed that the extracted Chenpi polysaccharide exhibited antidiabetic and anti-obesogenic properties and modulated gut microbiota, which stimulated the growth of *Lactobacillus johnsonii* and normalised the levels of plasma total triacylglycerol (TG), total cholesterol (CHO), low-density lipoprotein-cholesterol (LDL-C), alanine aminotransferase (ALT), and aspartate aminotransferase (AST) in a high-fat-diet mice model following an oral dosage of 60 mg kg<sup>-1</sup> for 4 weeks. Notably, although UAE can significantly reduce extraction time and solvent usage compared to conventional hot water extraction, its scalability remains an important consideration for industrial application.<sup>89,90</sup> In large volumes, it is difficult to achieve ultrasound wave uniformity, resulting in uneven energy distribution and uneven extraction efficiency.<sup>91</sup> Therefore, the results may give an overly positive view of its performance in practical applications.<sup>92</sup> To this end, further research is needed to explore pilot-scale or industry scale production to provide more comparable results for practical applications.

In a recent study by Liu *et al.*,<sup>93</sup> the authors established a liquid fermentation method (using *Bacillus licheniformis* at 5%, v/v for 12 h) to obtain Chenpi polysaccharides. The obtained Chenpi polysaccharide (MW of 3.72 kDa) was composed of arabinose (31.75%), galactose (24.77%), galacturonic acid (20.94%), fucose (7.83%), rhamnose (7.53%), glucose (4.08%), mannose (1.93%), and glucuronic acid (1.15%). Besides, the authors used an *in vitro* fermentation method to evaluate the



prebiotic effect of the obtained Chenpi polysaccharide, and the results showed that the obtained polysaccharide promoted the growth of *Bacteroides*, *Parabacteroides*, and *Collinsella*, while reducing the relative abundance of *Shigella* and *Klebsiella*.

Overall, the above studies make promising progress in the extraction of polysaccharides from Chenpi and exhibit their promoting health effects. However, the existing studies mainly focus on hot water extraction, while the applications of other methods are relatively limited. Therefore, further studies could explore more extraction methods for the extraction of polysaccharides from Chenpi to provide more comparable results to increase extraction efficiency and obtain high quality products for further applications.<sup>94,95</sup> Besides, the biological activities of polysaccharides are closely associated with their structures, such as molecular weight (MW) and monosaccharide composition.<sup>96,97</sup> Liang *et al.*<sup>98</sup> evaluated the structure–activity relationship of black garlic polysaccharides and found that the higher uronic acid content and lower MW were associated with higher antioxidant capacity, with MW exhibiting a greater influence. In another study, Liang *et al.*<sup>99</sup> analysed the relationship between the structural features of goji berry polysaccharides and antioxidant and anti-aging activities. The results showed that the antioxidant capacity was positively correlated with galacturonic acid (GalA) content, while the neutral multi-branched chains may contribute to anti-aging activity. To date, research on the structure–activity relationship of Chenpi polysaccharides remains limited. Therefore, future studies could focus on establishing the relationship between the structural features (*e.g.*, monosaccharide composition and molecular weight) and the different biological activities of Chenpi polysaccharides, which would facilitate the production of Chenpi polysaccharides with enhanced biological activity and expand their applicable scenarios.<sup>100–102</sup>

## 5. Food applications

Chenpi is a traditional medicine and food homology ingredient in China and has been widely processed into various food products (Fig. 2).<sup>103,104</sup> The famous Chenpi tea beverage is Ganpu tea, which is commonly made from Chenpi and Pu-erh tea.<sup>62</sup> In addition, the traditional Chenpi snack, nine-processed Chenpi, is produced using the “nine-processed” processing technique, which includes multiple steps, such as selection (CP1), soaking (CP2), keeping fresh, peeling, pickling (CP3),

draining, seasoning, repeated drying, and finally storage (aging, CP4).<sup>39,105</sup> Notably, the “nine-processed” method could take longer time and have high energy consumption, leading to a substantial environmental footprint and not meeting the concept of energy-saving and sustainable development.<sup>106</sup> Therefore, it is necessary to explore new processing techniques to quickly improve the quality of Chenpi’s product and to meet the concept of sustainable development. Considering environmental burden and sustainable food systems, emerging technologies could be investigated to replace and/or minimise traditional “nine-processed” method steps in the production of traditional Chenpi snacks. Fermentation, as an eco-friendly, technologically flexible method, has demonstrated its potential in the production of Chenpi products, which can not only be used for minimising the impact on the environment but also for improving the quality of Chenpi.<sup>40,93,107</sup> Therefore, continuously exploring the fermentation method in Chenpi product production and comparing it with the traditional processing method (nine-processed technique) would contribute to the sustainability and diversification of Chenpi product production.<sup>67</sup>

In addition to the above, Peng *et al.*<sup>108</sup> developed a functional jelly product that was made of Chenpi, orange juice, and pectin and reported that the jelly exhibited excellent antioxidant capacity as determined by *in vitro* chemical assay [ABTS of 88.87  $\mu\text{mol}$  Trolox equivalents (TE) per g, DPPH of 12.44  $\mu\text{mol}$  TE per g, and FRAP of 28.24  $\mu\text{mol}$  TE per g]. In another study, Zhang *et al.*<sup>109</sup> employed Chenpi in soy yogurt development. In the formulation, Chenpi partly replaced soybean in the yogurt formulation at 1%, 2%, 3%, and 4%, w/w. The results showed that Chenpi addition (2–4%) increased the total acid value of yoghurt. Furthermore, Chenpi addition (1–3%) increased the adhesiveness and chewiness of yoghurt. In terms of rheological properties, 1% Chenpi addition resulted in the highest apparent viscosity, while 4% Chenpi addition resulted in lowest apparent viscosity. This difference could be attributed to the Chenpi pectin polysaccharide; an appropriate level of Chenpi pectin addition as a filler could fill the spaces of protein aggregates to enhance the gel strength, while excessive Chenpi pectin addition may interfere with the cross-linking of protein aggregates and then hinder formation of the gel network, resulting in a weak gel network.<sup>109,110</sup> In the sensory evaluation, the soybean yoghurt that contained 2% Chenpi had the highest sensory scores among all products in terms of overall acceptance, colour, texture, odour, and flavour.

In a recent study, Zhang *et al.*<sup>111</sup> developed a cold brew Chenpi beverage and explored the effects of different assisted extraction methods, namely, high pressure processing (HPP), ultrasound-assisted (UAE), and combination of HPP and UAE, on the flavour quality of the final Chenpi beverage. The results showed that the combination of HPP and UAE significantly promoted the release of volatile flavor substances (fatty and woody aroma-related compounds) from Chenpi and endowed the Chenpi cold brew with a richer woody aroma.

Overall, the above studies exhibit the potential of Chenpi in food development and point to a novel avenue for further food innovation. Nevertheless, outside of Asia, Chenpi is an



Fig. 2 Chenpi-based products.



uncommon ingredient in most countries and is scarcely utilised and introduced into the daily diets, which could affect the acceptance of Chenpi-based food products and their further promotion.<sup>112,113</sup> Food selection is a complex process and varies depending on multiple factors, such as cultural contexts, eating habits, and personal taste experience; therefore, it is necessary to explore the consumer performance and perception of Chenpi-based food products in different groups and countries, which can in turn guide further formulation optimisation and innovation and promote their widespread acceptance.<sup>114–116</sup> Besides, the above studies report promising antioxidant activity of Chenpi-based food in *in vitro* chemical assays (*e.g.* DPPH and ABTS); these results may not directly translate into pharmacologically relevant concentrations in the human body.<sup>117</sup> There is increasing evidence that phenolic compounds in different food matrices affect their bioavailability and consequently biological potential in the human body.<sup>118–121</sup> Ribnicky *et al.*<sup>122</sup> reported that anthocyanins from blueberry bound to a protein-rich matrix were more bioaccessible. In the study by Mandalari *et al.*,<sup>123</sup> the biscuit matrix decreased bioaccessibility of flavonols from almond skin. Therefore, future studies could conduct *in vivo* approaches to evaluate the interaction between phenolic compounds from Chenpi and different food matrices and its consequent phenolic bioavailability to better understand its derived health benefits.

## 6. Industrial scalability and techno-economic considerations

According to the above, Chenpi shows potential as a valuable functional ingredient for sustainable diets, which can provide a promising insight into utilisation of citrus processing by-products as high value-added ingredients.<sup>124</sup> However, the large-scale industrial production of Chenpi from citrus waste requires careful consideration of scalability and techno-economic feasibility. First, the quality of citrus peel varies widely depending on citrus variety, cultivation conditions, and processing methods.<sup>125</sup> Therefore, establishing quality standardisation and standardised processing protocols is important to ensure raw materials with maximum retention of bioactive compounds and pharmacological efficacy and to support their reproducibility in industrial applications.<sup>126,127</sup> Additionally, in terms of extraction of high value-added ingredients, advanced extraction technologies (*e.g.* UAE) show high efficiency and environmental benefits, but may face barriers to industrial scale-up, including high energy costs, equipment limitations, and maintenance requirements.<sup>128,129</sup> In this regard, a comprehensive techno-economic analysis is necessary; in addition, optimising the process and combination of different techniques could overcome the mentioned limitations, thereby enhancing selectively, reducing cost, and then fostering industry applications.<sup>130,131</sup> On the other hand, consumer acceptance is important for the commercial success of Chenpi-based products.<sup>132</sup> Development of a wider range of Chenpi-based products and systematic consumer studies focusing on consumer perception and willingness are important to meet the diversified needs of different

consumer groups, contributing to their market success, and be widely promoted and incorporated into daily diets.<sup>133,134</sup>

## 7. Conclusions

Chenpi, one of the widely recognised and used medicinal and edible ingredients in the traditional Chinese diet, contains abundant bioactive compounds with extensive pharmacological benefits, such as antioxidant and anti-inflammatory properties and the modulation of gut microbiota. This makes Chenpi a valuable ingredient, and its wider utilisation in the daily diet could contribute to a shift towards a sustainable and healthy diet. However, efforts to develop Chenpi-based health products and widely integrate them into daily diet continue to address some challenges. First, Chenpi is an uncommon ingredient in most countries, and most customers lack an understanding of Chenpi; therefore, strengthening science communication and promoting the utilisation method are important directions for future development. In terms of bioactive compound extraction, the extraction method determines the quality and functions of target extracted compounds; therefore, more studies are needed in the extraction field to enhance the product quality, which could further expand the applications. In terms of functional food development, further studies could focus on the development of a wider range of Chenpi-based products to meet the diversified needs of different consumer groups.

## Author contributions

Guoqiang Zhang: conceptualization, investigation, visualization, writing – original draft, writing – review & editing. HioTong Mak: resources, visualization. Zheng Zhang: writing – review & editing. Ting Zhang: writing – review & editing. Yuanhui Wang: writing – review & editing, resources. Run-Yang Zhang: writing – review & editing. Hua-Min Liu: writing – review & editing. Shaobo Zhou: writing – review & editing. Bin Wu: writing – review & editing, funding acquisition.

## Conflicts of interest

The authors declare that they have no competing interests.

## Data availability

No primary research results, software or code have been included and no new data were generated or analysed as part of this review.

## Acknowledgements

The authors acknowledge the financial support from the ‘Xinjiang Tianshan Talent’ Cultivation Program (No. 2022TSYCLJ0040) and the Key R&D Special Project of Xinjiang Autonomous Region (No. 2022B02026). The graphical abstract image was created using some icons from <https://www.Biorender.com>.



## References

- 1 X. Li, X. Wang, B. A. Engel, E. Hua, S. Sun, P. Wu, *et al.*, Integrating environmental footprints and planetary boundaries for assessing environmental sustainability in food production, *Sustainable Prod. Consumption*, 2024, **48**, 205–218, DOI: [10.1016/j.spc.2024.05.018](https://doi.org/10.1016/j.spc.2024.05.018).
- 2 G. Zhang, Y. Wang and B. Wu, Advancing sustainable food innovation with house cricket (*Acheta domesticus*) and mealworm (*Tenebrio molitor*): nutrition, applications, and product quality, *J. Agric. Food Res.*, 2026, **25**, 102565, DOI: [10.1016/j.jafr.2025.102565](https://doi.org/10.1016/j.jafr.2025.102565).
- 3 Y. Li, P. He, Y. Shan, Y. Li, Y. Hang, S. Shao, *et al.*, Reducing climate change impacts from the global food system through diet shifts, *Nat. Clim. Change*, 2024, **14**, 943–953, DOI: [10.1038/s41558-024-02084-1](https://doi.org/10.1038/s41558-024-02084-1).
- 4 J. Rockström, S. H. Thilsted, W. C. Willett, L. J. Gordon, M. Herrero, C. C. Hicks, *et al.*, The EAT–Lancet Commission on healthy, sustainable, and just food systems, *Lancet*, 2025, **406**, 1625–1700, DOI: [10.1016/S0140-6736\(25\)01201-2](https://doi.org/10.1016/S0140-6736(25)01201-2).
- 5 Y. Guo, P. He, T. D. Searchinger, Y. Chen, M. Springmann, M. Zhou, *et al.*, Environmental and human health trade-offs in potential Chinese dietary shifts, *One Earth*, 2022, **5**, 268–282, DOI: [10.1016/j.oneear.2022.02.002](https://doi.org/10.1016/j.oneear.2022.02.002).
- 6 L. Wang, X. Zhu, H. Liu and B. Sun, Medicine and food homology substances: a review of bioactive ingredients, pharmacological effects and applications, *Food Chem.*, 2025, **463**, 141111, DOI: [10.1016/j.foodchem.2024.141111](https://doi.org/10.1016/j.foodchem.2024.141111).
- 7 A. Ma, F. Zou, R. Zhang and X. Zhao, The effects and underlying mechanisms of medicine and food homologous flowers on the prevention and treatment of related diseases, *J. Food Biochem.*, 2022, **46**, e14430, DOI: [10.1111/jfbc.14430](https://doi.org/10.1111/jfbc.14430).
- 8 Z. Liu, S. Sun, Y. Zhang, H. Zhao, C. Xu, Y. Liu, *et al.*, Food as Medicine: Integrating Food-Medicine Homology into Foods for Special Medical Purposes for Targeted Symptom Management and Nutritional Support, *Food Rev. Int.*, 2026, 1–35, DOI: [10.1080/87559129.2026.2639506](https://doi.org/10.1080/87559129.2026.2639506).
- 9 S. Zhou, Y. Huang and W. Chu, Medicine-food homology substances: ecological engineers of gut microbiota for precision nutrition, *Food Humanit.*, 2026, **6**, 101007, DOI: [10.1016/j.foohum.2026.101007](https://doi.org/10.1016/j.foohum.2026.101007).
- 10 C. Zheng, H. Zheng, Z. Ye and Q. Li, Chenpi years identification based on terahertz spectral data and time series deep learning, *Appl. Food Res.*, 2025, **5**, 101463, DOI: [10.1016/j.afres.2025.101463](https://doi.org/10.1016/j.afres.2025.101463).
- 11 X. K. Li, L. J. Tang, Z. Y. Li, D. Qiu, Z. L. Yang, X. Y. Zhang, *et al.*, Geographical origin discrimination of Chenpi using machine learning and enhanced mid-level data fusion, *NPJ Sci. Food*, 2025, **9**, 17, DOI: [10.1038/s41538-025-00376-0](https://doi.org/10.1038/s41538-025-00376-0).
- 12 F. Li, Y. Lu, C. Li, R. Huang, E. Tian, E. Tan, *et al.*, trnL-trnF copy number is inversely correlated with storage time of Guang Chenpi, the aged sun-dried peels of *Citrus reticulata* ‘Chachi’, *J. Stored Prod. Res.*, 2022, **97**, 101982, DOI: [10.1016/j.jspr.2022.101982](https://doi.org/10.1016/j.jspr.2022.101982).
- 13 X. Sun, H. Deng, B. Shan, Y. Shan, J. Huang, X. Feng, *et al.*, Flavonoids contribute most to discriminating aged Guang Chenpi (*Citrus reticulata* ‘Chachi’) by spectrum-effect relationship analysis between LC-Q-Orbitrap/MS fingerprint and ameliorating spleen deficiency activity, *Food Sci. Nutr.*, 2023, **11**, 7039–7060, DOI: [10.1002/fsn3.3629](https://doi.org/10.1002/fsn3.3629).
- 14 M. Falduto, F. Smedile, M. Zhang, T. Zheng, J. Zhu, Q. Huang, *et al.*, Anti-obesity effects of Chenpi: an artificial gastrointestinal system study, *Microb. Biotechnol.*, 2022, **15**, 874–885, DOI: [10.1111/1751-7915.14005](https://doi.org/10.1111/1751-7915.14005).
- 15 A. Li, N. Wang, N. Li, B. Li, F. Yan, Y. Song, *et al.*, Modulation effect of *chenpi* extract on gut microbiota in high-fat diet-induced obese C57BL/6 mice, *J. Food Biochem.*, 2021, **45**, e13541, DOI: [10.1111/jfbc.13541](https://doi.org/10.1111/jfbc.13541).
- 16 Z. Cen, Z. Chen, D. Wang, Y. Zuo, J. Peng, J. Chen, *et al.*, Guang Chenpi Extract as a Multifunctional Phytotherapeutic: Enhanced Effects with Ergothioneine and Polydeoxyribonucleotide on Redox Homeostasis and Tissue Resilience, *Pharmaceuticals*, 2026, **19**, 293, DOI: [10.3390/ph19020293](https://doi.org/10.3390/ph19020293).
- 17 L. Xu, W. Liu, X. Wu, T. Chen, Y. Gao, J. Huang, *et al.*, *Citrus Reticulatae Pericarpium* improves obesity in OVX mice by regulating the gut microbiota and suppressing appetite, *Food Funct.*, 2025, **16**, 9545–9561, DOI: [10.1039/D5FO03974F](https://doi.org/10.1039/D5FO03974F).
- 18 E. Tan, F. Li, X. Lin, S. Ma, G. Zhang, H. Zhou, *et al.*, Comparative study on comprehensive quality of Xinhui chenpi by two main plant propagation techniques, *Food Sci. Nutr.*, 2023, **11**, 1104–1112, DOI: [10.1002/fsn3.3148](https://doi.org/10.1002/fsn3.3148).
- 19 P. Sharma, R. Vishvakarma, K. Gautam, A. Vimal, V. Kumar Gaur, A. Farooqui, *et al.*, Valorization of citrus peel waste for the sustainable production of value-added products, *Bioresour. Technol.*, 2022, **351**, 127064, DOI: [10.1016/j.biortech.2022.127064](https://doi.org/10.1016/j.biortech.2022.127064).
- 20 Q. Peng, Y. Zhang, M. Zhu, F. Bao, J. Deng and W. Li, Polymethoxyflavones from citrus peel: advances in extraction methods, biological properties, and potential applications, *Crit. Rev. Food Sci. Nutr.*, 2024, **64**, 5618–5630, DOI: [10.1080/10408398.2022.2156476](https://doi.org/10.1080/10408398.2022.2156476).
- 21 M. A. Augustin, L. Sanguansri, E. M. Fox, L. Cobiac and M. B. Cole, Recovery of wasted fruit and vegetables for improving sustainable diets, *Trends Food Sci. Technol.*, 2020, **95**, 75–85, DOI: [10.1016/j.tifs.2019.11.010](https://doi.org/10.1016/j.tifs.2019.11.010).
- 22 G. Zhang, Z. Li, A. Chatzifragkou and D. Charalampopoulos, Effect of processing methods on the phytochemical content of melon seeds (*Cucumis melo* L.), *Future Foods*, 2024, **10**, 100399, DOI: [10.1016/j.fufo.2024.100399](https://doi.org/10.1016/j.fufo.2024.100399).
- 23 S. Manzoor, U. Fayaz, A. H. Dar, K. K. Dash, R. Shams, I. Bashir, *et al.*, Sustainable development goals through reducing food loss and food waste: a comprehensive review, *Future Foods*, 2024, **9**, 100362, DOI: [10.1016/j.fufo.2024.100362](https://doi.org/10.1016/j.fufo.2024.100362).
- 24 G. Zhang, J. Guo and J. Guo, A sustainable approach in pumpkin seed oil processing line: recent advances in pumpkin seed oil and oil processing by-products, *Food*



- Chem.:X*, 2025, 26, 102259, DOI: [10.1016/j.fochx.2025.102259](https://doi.org/10.1016/j.fochx.2025.102259).
- 25 D. Granato, I. Zabetakis and A. Koidis, Sustainability, nutrition, and scientific advances of functional foods under the new EU and global legislation initiatives, *J. Funct. Foods*, 2023, 109, 105793, DOI: [10.1016/j.jff.2023.105793](https://doi.org/10.1016/j.jff.2023.105793).
- 26 A. Kumar, S. Kumar, R. N. Komal and P. Singh, Role of Traditional Ethnobotanical Knowledge and Indigenous Communities in Achieving Sustainable Development Goals, *Sustainability*, 2021, 13, 3062, DOI: [10.3390/su13063062](https://doi.org/10.3390/su13063062).
- 27 D.-X. Sun-Waterhouse, X.-Y. Chen, Z.-H. Liu, I. N. Waterhouse G and W.-Y. Kang, Transformation from traditional medicine-food homology to modern food-medicine homology, *Food Med. Homol.*, 2024, 1, 9420014, DOI: [10.26599/FMH.2024.9420014](https://doi.org/10.26599/FMH.2024.9420014).
- 28 J. Shi, L. Peng, W. Chen, W. Qiao, K. Wang, Y. Xu, *et al.*, Evaluation of chemical components and quality in Xinhui Chenpi ( *Citrus reticulata* 'Chachi') with two different storage times by GC-MS and UPLC, *Food Sci. Nutr.*, 2024, 12, 5036–5051, DOI: [10.1002/fsn3.4154](https://doi.org/10.1002/fsn3.4154).
- 29 X. Yu, S. Sun, Y. Guo, Y. Liu, D. Yang, G. Li, *et al.*, Citri Reticulatae Pericarpium (Chenpi): botany, ethnopharmacology, phytochemistry, and pharmacology of a frequently used traditional Chinese medicine, *J. Ethnopharmacol.*, 2018, 220, 265–282, DOI: [10.1016/j.jep.2018.03.031](https://doi.org/10.1016/j.jep.2018.03.031).
- 30 Y. Wang, X. Zhang, C. Zhou, H. Khan, M. Fu and W. S. Cheang, Citri Reticulatae Pericarpium (Chenpi) Protects against Endothelial Dysfunction and Vascular Inflammation in Diabetic Rats, *Nutrients*, 2022, 14, 5221, DOI: [10.3390/nu14245221](https://doi.org/10.3390/nu14245221).
- 31 Y. Qian, Z. Gao, C. Wang, J. Ma, G. Li, F. Fu, *et al.*, Effects of Different Treatment Methods of Dried Citrus Peel (Chenpi) on Intestinal Microflora and Short-Chain Fatty Acids in Healthy Mice, *Front. Nutr.*, 2021, 8, 702559, DOI: [10.3389/fnut.2021.702559](https://doi.org/10.3389/fnut.2021.702559).
- 32 M. Zhang, J. Zhu, X. Zhang, D. Zhao, Y. Ma, D. Li, *et al.*, Aged citrus peel ( *chenpi* ) extract causes dynamic alteration of colonic microbiota in high-fat diet induced obese mice, *Food Funct.*, 2020, 11, 2667–2678, DOI: [10.1039/C9FO02907A](https://doi.org/10.1039/C9FO02907A).
- 33 Q. Wang, Z. Qiu, Y. Chen, Y. Song, A. Zhou, Y. Cao, *et al.*, Review of recent advances on health benefits, microbial transformations, and authenticity identification of *Citri reticulatae* Pericarpium bioactive compounds, *Crit. Rev. Food Sci. Nutr.*, 2024, 64, 10332–10360, DOI: [10.1080/10408398.2023.2222834](https://doi.org/10.1080/10408398.2023.2222834).
- 34 G. Xu, J. Zhao, J. Yao, Y. Xu, X. Yuan and S. Pan, Effects of aging on the fine structure, chain conformation, and morphology of Chenpi polysaccharides, *Carbohydr. Polym.*, 2025, 349, 122970, DOI: [10.1016/j.carbpol.2024.122970](https://doi.org/10.1016/j.carbpol.2024.122970).
- 35 C. Tian, H. Xu, J. Li and Z. Han, Characteristics and intestinal immunomodulating activities of water-soluble pectic polysaccharides from Chenpi with different storage periods, *J. Sci. Food Agric.*, 2018, 98, 3752–3757, DOI: [10.1002/jsfa.8888](https://doi.org/10.1002/jsfa.8888).
- 36 L. Chen, G. Xiao, Y. Xu, W. S. Cheang, J. Wu, Y. Yu, *et al.*, Effects of aging on the bioactive flavonoids and fungal diversity of Pericarpium Citri Reticulatae (Citrus reticulata 'Chachi'), *LWT-Food Sci. Technol.*, 2024, 200, 116135, DOI: [10.1016/j.lwt.2024.116135](https://doi.org/10.1016/j.lwt.2024.116135).
- 37 Y. Zhu, Y. Zhu, X. Liu and J. Qi, Change in nutrients of Citri Reticulatae 'Chachi' in relation to browning during aging, *J. Food Compos. Anal.*, 2022, 107, 104351, DOI: [10.1016/j.jfca.2021.104351](https://doi.org/10.1016/j.jfca.2021.104351).
- 38 F. Yue, F. Zhang, Q. Qu, C. Wang, Y. Qin, L. Ma, *et al.*, Effects of ageing time on the properties of polysaccharide in tangerine peel and its bacterial community, *Food Chem.*, 2023, 417, 135812, DOI: [10.1016/j.foodchem.2023.135812](https://doi.org/10.1016/j.foodchem.2023.135812).
- 39 M. Fu, Y. Wang, X. Zhu, M. Li, Y. Yu, J. Wu, *et al.*, Metabolomics reveal changes of flavonoids during processing of "nine-processed" tangerine peel (Jiuzhi Chenpi), *LWT-Food Sci. Technol.*, 2024, 214, 117132, DOI: [10.1016/j.lwt.2024.117132](https://doi.org/10.1016/j.lwt.2024.117132).
- 40 Z. Tang, Y. Miao, H. Lu, C. Chen, S. Li and W. Zhang, Aging of Citri Reticulatae pericarpium: flavonoid metabolite dynamics, fungal community succession, and functional analysis of key transformation strains, *Food Res. Int.*, 2026, 227, 118220, DOI: [10.1016/j.foodres.2025.118220](https://doi.org/10.1016/j.foodres.2025.118220).
- 41 Y. Zhang, X. Tong, B. Chen, S. Wu, X. Wang, Q. Zheng, *et al.*, Novel application of HS-GC-IMS for characteristic fingerprints and flavor compound variations in citrus reticulatae pericarpium during storage with different *Aspergillus niger* fermentation, *Food Chem.:X*, 2023, 18, 100653, DOI: [10.1016/j.fochx.2023.100653](https://doi.org/10.1016/j.fochx.2023.100653).
- 42 R. Liu, Y. Tao, T. Xu, T. Wu, Q. Yu, S. Pan, *et al.*, Antioxidant activity increased due to dynamic changes of flavonoids in orange peel during *Aspergillus niger* fermentation, *Int. J. Food Sci. Technol.*, 2023, 58, 3329–3336, DOI: [10.1111/ijfs.16239](https://doi.org/10.1111/ijfs.16239).
- 43 Y. Fu, C. Wang, Z. Gao, Y. Liao, M. Peng, F. Fu, *et al.*, Microbes: drivers of Chenpi manufacturing, biotransformation, and physiological effects, *Food Chem.*, 2025, 464, 141631, DOI: [10.1016/j.foodchem.2024.141631](https://doi.org/10.1016/j.foodchem.2024.141631).
- 44 J. Xie, F. Wei, L. Luo, L. Dai, L. Zeng and X. Wang, Effect of cell wall-disruption processes on wall disruption, antioxidant activity and nutrients in tea pollen, *Int. J. Food Sci. Technol.*, 2022, 57, 3361–3374, DOI: [10.1111/ijfs.15612](https://doi.org/10.1111/ijfs.15612).
- 45 J. Shi, L. Peng, W. Chen, W. Qiao, K. Wang, Y. Xu, *et al.*, Evaluation of chemical components and quality in Xinhui Chenpi ( *Citrus reticulata* 'Chachi') with two different storage times by GC-MS and UPLC, *Food Sci. Nutr.*, 2024, 12, 5036–5051, DOI: [10.1002/fsn3.4154](https://doi.org/10.1002/fsn3.4154).
- 46 H. Chai, X. Li, S. Chen, D. Lin, R. Cai, A. Yang, *et al.*, Comparative Study Between Orah Mandarin Peel and Traditional Mandarin Peels (Chenpi) Based on Data-Repository-Enabled Ultra-High-Performance Liquid Chromatography-Quadrupole-Orbitrap-Mass Spectrometry



- Analysis, *Chem. Biodiversity*, 2025, 22(12), e01661, DOI: [10.1002/cbdv.202501661](https://doi.org/10.1002/cbdv.202501661).
- 47 J. Zhang, X. Wu, J. Qiu, L. Zhang, Y. Zhang, X. Qiu, *et al.*, Comprehensive Comparison on the Chemical Profile of Guang Chen Pi at Different Ripeness Stages Using Untargeted and Pseudotargeted Metabolomics, *J. Agric. Food Chem.*, 2020, 68, 8483–8495, DOI: [10.1021/acs.jafc.0c02904](https://doi.org/10.1021/acs.jafc.0c02904).
- 48 X. Bian, X. Xie, J. Cai, Y. Zhao, W. Miao, X. Chen, *et al.*, Dynamic changes of phenolic acids and antioxidant activity of Citri Reticulatae Pericarpium during aging processes, *Food Chem.*, 2022, 373, 131399, DOI: [10.1016/j.foodchem.2021.131399](https://doi.org/10.1016/j.foodchem.2021.131399).
- 49 L. Wang, H. Xu, F. Yuan, Q. Pan, R. Fan and Y. Gao, Physicochemical characterization of five types of citrus dietary fibers, *Biocatal. Agric. Biotechnol.*, 2015, 4, 250–258, DOI: [10.1016/j.bcab.2015.02.003](https://doi.org/10.1016/j.bcab.2015.02.003).
- 50 T. Xiang, R. Yang, L. Li, H. Lin and G. Kai, Research progress and application of pectin: a review, *J. Food Sci.*, 2024, 89, 6985–7007, DOI: [10.1111/1750-3841.17438](https://doi.org/10.1111/1750-3841.17438).
- 51 S. Roy, R. Priyadarshi, Ł. Łopusiewicz, D. Biswas, V. Chandel and J.-W. Rhim, Recent progress in pectin extraction, characterization, and pectin-based films for active food packaging applications: a review, *Int. J. Biol. Macromol.*, 2023, 239, 124248, DOI: [10.1016/j.ijbiomac.2023.124248](https://doi.org/10.1016/j.ijbiomac.2023.124248).
- 52 R. K. R., S. Gopi and P. Balakrishnan, *Introduction to Flavor and Fragrance in Food Processing*, 2022, pp. 1–19, DOI: [10.1021/bk-2022-1433.ch001](https://doi.org/10.1021/bk-2022-1433.ch001).
- 53 H. Jiang, S. Yang, H. Tian and B. Sun, Research progress in the use of liquid-liquid extraction for food flavour analysis, *Trends Food Sci. Technol.*, 2023, 132, 138–149, DOI: [10.1016/j.tifs.2023.01.005](https://doi.org/10.1016/j.tifs.2023.01.005).
- 54 Y. Yang, Y. Wang, B. Li, C. Chen, Y. Zhao, L. Pan, *et al.*, Analysis of microbial diversity of traditional Chinese starters and screening of special flavor yeast for Chinese baked flatbread, *Front. Nutr.*, 2026, 12, 1755466, DOI: [10.3389/fnut.2025.1755466](https://doi.org/10.3389/fnut.2025.1755466).
- 55 F. B. Siewe, C. W. Makebe, W. Claudette Bakisu Muala, A. Laya, N. Raymond Nkongho, M. O. Meliko, *et al.*, Advances in processing, reaction pathways, stabilisation and food applications of natural seafood flavourings, *Food Biosci.*, 2024, 58, 103627, DOI: [10.1016/j.fbio.2024.103627](https://doi.org/10.1016/j.fbio.2024.103627).
- 56 C. I. Kothe, C. Carøe, F. Mazel, D. Zilber, P. Cruz-Morales, N. Mohellibi, *et al.*, Novel misos shape distinct microbial ecologies: opportunities for flavourful sustainable food innovation, *Food Res. Int.*, 2024, 189, 114490, DOI: [10.1016/j.foodres.2024.114490](https://doi.org/10.1016/j.foodres.2024.114490).
- 57 H. Li, L. Lin, Y. Feng and M. Zhao, Exploration of optimal preparation strategy of Chenpi (pericarps of Citrus reticulata Blanco) flavouring essence with great application potential in sugar and salt-reduced foods, *Food Res. Int.*, 2024, 175, 113669, DOI: [10.1016/j.foodres.2023.113669](https://doi.org/10.1016/j.foodres.2023.113669).
- 58 Y.-H. Wang, Y.-Y. Yang, H.-Q. Li, Q.-D. Zhang, F. Xu and Z.-J. Li, Characterization of aroma-active compounds in steamed breads fermented with Chinese traditional sourdough, *LWT–Food Sci. Technol.*, 2021, 152, 112347, DOI: [10.1016/j.lwt.2021.112347](https://doi.org/10.1016/j.lwt.2021.112347).
- 59 S. Urlass, Y. Wu, T. T. L. Nguyen, P. Winberg, M. S. Turner and H. Smyth, Unravelling the aroma and flavour of algae for future food applications, *Trends Food Sci. Technol.*, 2023, 138, 370–381, DOI: [10.1016/j.tifs.2023.06.018](https://doi.org/10.1016/j.tifs.2023.06.018).
- 60 S. Wu, X. Song, G. Huo, L. Li, M. Huo, M. Zhao, *et al.*, Elucidation of the diverse molecular signatures and transformation pathways of flavor metabolites in age-stratified citri reticulatae pericarpium (ChenPi) and their enzymatic hydrolysates: a perspective from the flavor space, *Food Res. Int.*, 2026, 225, 118054, DOI: [10.1016/j.foodres.2025.118054](https://doi.org/10.1016/j.foodres.2025.118054).
- 61 S. Chen, Z. Chen, D. O'Neill Rothenberg, Y. Long, H. Li, X. Zeng, *et al.*, Short-term steaming during processing impacts the quality of Citri Reticulatae 'Chachi' peel, *Food Chem.*, 2024, 447, 138964, DOI: [10.1016/j.foodchem.2024.138964](https://doi.org/10.1016/j.foodchem.2024.138964).
- 62 X. Deng, S. He, Y. Han and Y. Chen, Metabolites profiling reveals the dynamic changes of non-volatiles in Pu-erh during Ganpu tea processing, *Food Chem.:X*, 2023, 19, 100774, DOI: [10.1016/j.fochx.2023.100774](https://doi.org/10.1016/j.fochx.2023.100774).
- 63 T.-X. Li, F.-Y. Xiong, W. Cai, Y.-N. Fang, X. Yang, J.-X. Li, *et al.*, Chemical compositions and bioactivity of essential oils of Citri Reticulata Pericarpium Viride from three provinces in China, *Food Sci. Biotechnol.*, 2026, 35, 1147–1157, DOI: [10.1007/s10068-026-02112-0](https://doi.org/10.1007/s10068-026-02112-0).
- 64 K. Cao, C. Yao, J. Xiao, R. Huang, Z. Qin, Y. Shi, *et al.*, Elucidation of dynamic evolution patterns of non-volatile and volatile compounds in peel of Citrus reticulata cv. Chachiensis with different tree ages: a metabolomics-based analysis, *Food Res. Int.*, 2026, 227, 118254, DOI: [10.1016/j.foodres.2025.118254](https://doi.org/10.1016/j.foodres.2025.118254).
- 65 K. Jiang, K. Xu, J. Wang, F. Meng and B. Wang, Based on HS-SPME-GC-MS combined with GC-O-MS to analyze the changes of aroma compounds in the aging process of Citri Reticulatae Pericarpium, *Food Biosci.*, 2023, 54, 102798, DOI: [10.1016/j.fbio.2023.102798](https://doi.org/10.1016/j.fbio.2023.102798).
- 66 H. Liu, M. Fu, J. Wu, Y. Yu, W. Si, Y. Xu, *et al.*, Flavor characterization of aged Citri Reticulatae Pericarpium from core regions: an integrative approach utilizing GC-IMS, GC-MS, E-nose, E-tongue, and chemometrics, *Food Chem.*, 2025, 490, 144995, DOI: [10.1016/j.foodchem.2025.144995](https://doi.org/10.1016/j.foodchem.2025.144995).
- 67 F. Shen, T. Wang, R. Zhang, B. Zhong and Z. Wu, Metabolism and release of characteristic components and their enzymatic mechanisms in Pericarpium Citri Reticulatae co-fermentation, *Food Chem.*, 2024, 432, 137227, DOI: [10.1016/j.foodchem.2023.137227](https://doi.org/10.1016/j.foodchem.2023.137227).
- 68 T. Liu, M. Zhao, Y. Zhang, R. Xu, Z. Fu, T. Jin, *et al.*, Polysaccharides from Phellinus linteus attenuate type 2 diabetes mellitus in rats via modulation of gut microbiota and bile acid metabolism, *Int. J. Biol. Macromol.*, 2024, 262, 130062, DOI: [10.1016/j.ijbiomac.2024.130062](https://doi.org/10.1016/j.ijbiomac.2024.130062).
- 69 G. Chen, N. Jiang, J. Zheng, H. Hu, H. Yang, A. Lin, *et al.*, Structural characterization and anti-inflammatory activity of polysaccharides from Astragalus membranaceus, *Int. J.*



- Biol. Macromol.*, 2023, **241**, 124386, DOI: [10.1016/j.ijbiomac.2023.124386](https://doi.org/10.1016/j.ijbiomac.2023.124386).
- 70 S. Wang, G. Li, X. Zhang, Y. Wang, Y. Qiang, B. Wang, *et al.*, Structural characterization and antioxidant activity of Polygonatum sibiricum polysaccharides, *Carbohydr. Polym.*, 2022, **291**, 119524, DOI: [10.1016/j.carbpol.2022.119524](https://doi.org/10.1016/j.carbpol.2022.119524).
- 71 Y. Qiao, Y. Shen, H. Jiang, D. Li and B. Li, Structural characterization, antioxidant and antibacterial activity of three pectin polysaccharides from blueberry, *Int. J. Biol. Macromol.*, 2024, **262**, 129707, DOI: [10.1016/j.ijbiomac.2024.129707](https://doi.org/10.1016/j.ijbiomac.2024.129707).
- 72 G. Zhang and L. Liu, Insights into potential value and sustainable utilisation of pumpkin peel: a review, *Future Foods*, 2025, **11**, 100575, DOI: [10.1016/j.fufo.2025.100575](https://doi.org/10.1016/j.fufo.2025.100575).
- 73 V. Dixit, S. W. Joseph Kamal, P. Bajrang Chole, D. Dayal, K. K. Chaubey, A. K. Pal, *et al.*, Functional Foods: Exploring the Health Benefits of Bioactive Compounds from Plant and Animal Sources, *J. Food Qual.*, 2023, **2023**, 1–22, DOI: [10.1155/2023/5546753](https://doi.org/10.1155/2023/5546753).
- 74 X. Wang, Y. Su, J. Su, J. Xue, R. Zhang, X. Li, *et al.*, Optimization of Enzyme-Assisted Aqueous Extraction of Polysaccharide from Acanthopanax senticosus and Comparison of Physicochemical Properties and Bioactivities of Polysaccharides with Different Molecular Weights, *Molecules*, 2023, **28**, 6585, DOI: [10.3390/molecules28186585](https://doi.org/10.3390/molecules28186585).
- 75 Z. Zhang, G. Lv, T. Song, Z. Xu and M. Wang, Effects of different extraction methods on the structural and biological properties of Hericium coralloides polysaccharides, *Food Chem.*, 2024, **445**, 138752, DOI: [10.1016/j.foodchem.2024.138752](https://doi.org/10.1016/j.foodchem.2024.138752).
- 76 Q. Wei and Y. Zhang, Ultrasound-assisted polysaccharide extraction from Cercis chinensis and properites, antioxidant activity of polysaccharide, *Ultrason. Sonochem.*, 2023, **96**, 106422, DOI: [10.1016/j.ultsonch.2023.106422](https://doi.org/10.1016/j.ultsonch.2023.106422).
- 77 A. D. Premarathna, T. A. E. Ahmed, G. Kulshreshtha, S. Humayun, C. N. Shormeh Darko, V. Rjabovs, *et al.*, Polysaccharides from red seaweeds: effect of extraction methods on physicochemical characteristics and antioxidant activities, *Food Hydrocolloids*, 2024, **147**, 109307, DOI: [10.1016/j.foodhyd.2023.109307](https://doi.org/10.1016/j.foodhyd.2023.109307).
- 78 G. Zhang, Z. Li, L. Liu and Q. Xiang, Road to valorisation of melon seeds (*Cucumis melo* L.): a comprehensive review of nutritional profiles, biological activities, and food applications, *Sustainable Food Technol.*, 2024, **2**, 1166–1182, DOI: [10.1039/D4FB00119B](https://doi.org/10.1039/D4FB00119B).
- 79 G. Huang, F. Chen, W. Yang and H. Huang, Preparation, deproteinization and comparison of bioactive polysaccharides, *Trends Food Sci. Technol.*, 2021, **109**, 564–568, DOI: [10.1016/j.tifs.2021.01.038](https://doi.org/10.1016/j.tifs.2021.01.038).
- 80 J. Yang, L. Wen, Y. Zhao, Y. Jiang, M. Tian, H. Liu, *et al.*, Structure identification of an arabinogalacturonan in Citrus reticulata Blanco ‘Chachiensis’ peel, *Food Hydrocolloids*, 2018, **84**, 481–488, DOI: [10.1016/j.foodhyd.2018.06.022](https://doi.org/10.1016/j.foodhyd.2018.06.022).
- 81 T. Zhou, Y. Jiang, L. Wen and B. Yang, Characterization of polysaccharide structure in Citrus reticulata ‘Chachi’ peel during storage and their bioactivity, *Carbohydr. Res.*, 2021, **508**, 108398, DOI: [10.1016/j.carres.2021.108398](https://doi.org/10.1016/j.carres.2021.108398).
- 82 L. Wang, W. Li, Y. Li, G. Chen, L. Zhao, W. Li, *et al.*, Dried tangerine peel polysaccharide (DTPP) alleviates hepatic steatosis by suppressing TLR4/MD-2-mediated inflammation and endoplasmic reticulum stress, *Bioorg. Chem.*, 2024, **147**, 107369, DOI: [10.1016/j.bioorg.2024.107369](https://doi.org/10.1016/j.bioorg.2024.107369).
- 83 H. Liang, Y. Liang, L. Wu, L. Lin, Y. Yao, J. Deng, *et al.*, Pretreatment with Citrus reticulata ‘Chachi’ Polysaccharide Alleviates Alcohol-Induced Gastric Ulcer by Inhibiting NLRP3/ASC/Caspase-1 and Nrf2/HO-1 Signaling Pathways, *Nutrients*, 2025, **17**, 2062, DOI: [10.3390/nu17132062](https://doi.org/10.3390/nu17132062).
- 84 C. Castiello, P. Junghanns, A. Mergel, C. Jacob, C. Ducho, S. Valente, *et al.*, GreenMedChem: the challenge in the next decade toward eco-friendly compounds and processes in drug design, *Green Chem.*, 2023, **25**, 2109–2169, DOI: [10.1039/D2GC03772F](https://doi.org/10.1039/D2GC03772F).
- 85 V. Venugopal, A. Sasidharan and T. Rustad, Green Chemistry to Valorize Seafood Side Streams: An Ecofriendly Roadmap toward Sustainability, *J. Agric. Food Chem.*, 2023, **71**, 17494–17509, DOI: [10.1021/acs.jafc.3c03126](https://doi.org/10.1021/acs.jafc.3c03126).
- 86 Á. I. López-Lorente, F. Pena-Pereira, S. Pedersen-Bjergaard, V. G. Zuin, S. A. Ozkan and E. Psillakis, The ten principles of green sample preparation, *TrAC, Trends Anal. Chem.*, 2022, **148**, 116530, DOI: [10.1016/j.trac.2022.116530](https://doi.org/10.1016/j.trac.2022.116530).
- 87 F. Kurul, B. Doruk and S. N. Topkaya, Principles of green chemistry: building a sustainable future, *Discov. Chem.*, 2025, **2**, 68, DOI: [10.1007/s44371-025-00152-9](https://doi.org/10.1007/s44371-025-00152-9).
- 88 Y. Li, Z. Li, B. Chen, Y. Hou, Y. Wen, L. Gan, *et al.*, Ultrasonic assisted extraction, characterization and gut microbiota-dependent anti-obesity effect of polysaccharide from Pericarpium Citri Reticulatae ‘Chachiensis’, *Ultrason. Sonochem.*, 2023, **95**, 106383, DOI: [10.1016/j.ultsonch.2023.106383](https://doi.org/10.1016/j.ultsonch.2023.106383).
- 89 Y. Deng, W. Wang, S. Zhao, X. Yang, W. Xu, M. Guo, *et al.*, Ultrasound-assisted extraction of lipids as food components: mechanism, solvent, feedstock, quality evaluation and coupled technologies – a review, *Trends Food Sci. Technol.*, 2022, **122**, 83–96, DOI: [10.1016/j.tifs.2022.01.034](https://doi.org/10.1016/j.tifs.2022.01.034).
- 90 S. Sethi and V. K. Rathod, Recent advances in ultrasound-assisted extraction of natural products using novel solvents: a mini-review, *Curr. Opin. Chem. Eng.*, 2025, **48**, 101132, DOI: [10.1016/j.coche.2025.101132](https://doi.org/10.1016/j.coche.2025.101132).
- 91 U. Roobab, R. M. Aadil, S. S. Kurup and S. Maqsood, Comparative evaluation of ultrasound-assisted extraction with other green extraction methods for sustainable recycling and processing of date palm bioresources and by-products: a review of recent research, *Ultrason. Sonochem.*, 2025, **114**, 107252, DOI: [10.1016/j.ultsonch.2025.107252](https://doi.org/10.1016/j.ultsonch.2025.107252).



- 92 J. Z. Lian, T. Liu, J. Yang, H. Zhu, N. Chalermchotiwong, Y. Grain, *et al.*, Sustainable extraction of bioactive compounds: a life cycle perspective on technologies, solvents, and process scale-up, *RSC Sustainability*, 2026, **4**, 1180–1202, DOI: [10.1039/D5SU00832H](https://doi.org/10.1039/D5SU00832H).
- 93 Y. Liu, Y. Huang, H. Hu, H. Yang, L. Xiong, J. Zheng, *et al.*, Structural characterization and probiotic activity of Chenpi polysaccharides prepared by fermentation with *Bacillus licheniformis*, *Food Biosci.*, 2025, **74**, 107789, DOI: [10.1016/j.fbio.2025.107789](https://doi.org/10.1016/j.fbio.2025.107789).
- 94 W. Hu, A. Yu, H. Bi, Y. Gong, H. Wang, H. Kuang, *et al.*, Recent advances in *Artemisia argyi* Lev. et Vant. polysaccharides: extractions, purifications, structural characteristics, pharmacological activities, and existing and potential applications, *Int. J. Biol. Macromol.*, 2024, **279**, 135250, DOI: [10.1016/j.ijbiomac.2024.135250](https://doi.org/10.1016/j.ijbiomac.2024.135250).
- 95 P. Otero, M. Carpena, P. Garcia-Oliveira, J. Echave, A. Soria-Lopez, P. Garcia-Perez, *et al.*, Seaweed polysaccharides: emerging extraction technologies, chemical modifications and bioactive properties, *Crit. Rev. Food Sci. Nutr.*, 2023, **63**, 1901–1929, DOI: [10.1080/10408398.2021.1969534](https://doi.org/10.1080/10408398.2021.1969534).
- 96 B. Wang, L. Yan, S. Guo, L. Wen, M. Yu, L. Feng, *et al.*, Structural Elucidation, Modification, and Structure-Activity Relationship of Polysaccharides in Chinese Herbs: A Review, *Front. Nutr.*, 2022, **9**, 908175, DOI: [10.3389/fnut.2022.908175](https://doi.org/10.3389/fnut.2022.908175).
- 97 J. Wang, L. Wang, H. Zhou, X. Liang, M. Zhang, Y. Tang, *et al.*, The isolation, structural features and biological activities of polysaccharide from *Ligusticum chuanxiong*: a review, *Carbohydr. Polym.*, 2022, **285**, 118971, DOI: [10.1016/j.carbpol.2021.118971](https://doi.org/10.1016/j.carbpol.2021.118971).
- 98 J. Liang, Y. Zhao, F. Yang, L. Zheng, Y. Ma, Q. Liu, *et al.*, Preparation and structure-activity relationship of highly active black garlic polysaccharides, *Int. J. Biol. Macromol.*, 2022, **220**, 601–612, DOI: [10.1016/j.ijbiomac.2022.08.115](https://doi.org/10.1016/j.ijbiomac.2022.08.115).
- 99 X. Liang, M. Liu, Y. Wei, L. Tong, S. Guo, H. Kang, *et al.*, Structural characteristics and structure-activity relationship of four polysaccharides from *Lycii fructus*, *Int. J. Biol. Macromol.*, 2023, **253**, 127256, DOI: [10.1016/j.ijbiomac.2023.127256](https://doi.org/10.1016/j.ijbiomac.2023.127256).
- 100 L. Shi, Q. He, J. Li, Y. Liu, Y. Cao, Y. Liu, *et al.*, Polysaccharides in fruits: biological activities, structures, and structure-activity relationships and influencing factors—a review, *Food Chem.*, 2024, **451**, 139408, DOI: [10.1016/j.foodchem.2024.139408](https://doi.org/10.1016/j.foodchem.2024.139408).
- 101 B. Hu, S. Zhang, Z. Wang, Q. Han, D. Zhang, Y. Zheng, *et al.*, Degradation Method, Structural Characteristics, Biological Activity and Structure-Activity Relationship of Degraded Polysaccharides, *Food Rev. Int.*, 2024, **40**, 2378–2407, DOI: [10.1080/87559129.2023.2273933](https://doi.org/10.1080/87559129.2023.2273933).
- 102 X. Ji, J. Guo, T. Cao, T. Zhang, Y. Liu and Y. Yan, Review on mechanisms and structure-activity relationship of hypoglycemic effects of polysaccharides from natural resources, *Food Sci. Hum. Wellness*, 2023, **12**, 1969–1980, DOI: [10.1016/j.fshw.2023.03.017](https://doi.org/10.1016/j.fshw.2023.03.017).
- 103 Y. Li, Y. Chen, Y. Zhou, J. He, Q. Zhou and M. Wang, Unveiling the potentials and action mechanisms of *Citri reticulatae* Pericarpium as an anti-inflammatory food, *Food Front.*, 2025, **6**, 163–184, DOI: [10.1002/fft2.506](https://doi.org/10.1002/fft2.506).
- 104 X. Lu, C. Zhao, H. Shi, Y. Liao, F. Xu, H. Du, *et al.*, Nutrients and bioactives in citrus fruits: different citrus varieties, fruit parts, and growth stages, *Crit. Rev. Food Sci. Nutr.*, 2023, **63**, 2018–2041, DOI: [10.1080/10408398.2021.1969891](https://doi.org/10.1080/10408398.2021.1969891).
- 105 M. Fu, Y. Wang, Y. Yu, J. Wen, M. S. Cheong, W. S. Cheang, *et al.*, Changes of volatile substance composition during processing of nine-processed tangerine peel (Jiuzhi Chenpi) determined by gas chromatography-ion mobility spectrometry, *Front. Nutr.*, 2022, **9**, 963655, DOI: [10.3389/fnut.2022.963655](https://doi.org/10.3389/fnut.2022.963655).
- 106 R. Wang, P. Li, S. Chen, J. Guo, Y. Liu, H. Shen, *et al.*, Elucidating the effects of nine-steaming-nine-drying on the morphology, metabolomic profile, and active ingredients of *Polygonatum sibiricum* Redouté, *Front. Food Sci. Technol.*, 2025, **5**, 1666149, DOI: [10.3389/frfst.2025.1666149](https://doi.org/10.3389/frfst.2025.1666149).
- 107 J. Sun and X. Mao, An environmental friendly process for Antarctic krill (*Euphausia superba*) utilization using fermentation technology, *J. Cleaner Prod.*, 2016, **127**, 618–623, DOI: [10.1016/j.jclepro.2016.04.020](https://doi.org/10.1016/j.jclepro.2016.04.020).
- 108 M. Peng, Z. Gao, Y. Liao, J. Guo and Y. Shan, Development of Citrus-Based Functional Jelly and an Investigation of Its Anti-Obesity and Antioxidant Properties, *Antioxidants*, 2022, **11**, 2418, DOI: [10.3390/antiox11122418](https://doi.org/10.3390/antiox11122418).
- 109 R. Zhang, J. Zhang, Y. Zhang, H. Zhang, Y. Cao and W. Xu, Effects of *Citri Reticulatae* Pericarpium on the physicochemical, rheological, and sensory properties of soy yogurt, *LWT—Food Sci. Technol.*, 2025, **234**, 118569, DOI: [10.1016/j.lwt.2025.118569](https://doi.org/10.1016/j.lwt.2025.118569).
- 110 P. R. Mary, S. Mutturi and M. Kapoor, Non-enzymatically hydrolyzed guar gum and orange peel fibre together stabilize the low-fat, set-type yogurt: a techno-functional study, *Food Hydrocolloids*, 2022, **122**, 107100, DOI: [10.1016/j.foodhyd.2021.107100](https://doi.org/10.1016/j.foodhyd.2021.107100).
- 111 J. Zhang, K. Shi, B. Li, C. Chen, S. Jia and Y. Liu, Effects of High Pressure Processing and Ultrasound-Assisted Extraction on Physicochemical Properties, Antioxidant Activity and Flavor Compounds of Cold Brew *Citri Reticulatae* Pericarpium Beverage, *Foods*, 2026, **15**, 757, DOI: [10.3390/foods15040757](https://doi.org/10.3390/foods15040757).
- 112 G. Zhang, Y. Wang, Z. Liang, J. Guo and B. Wu, Nutritional value and advances of mealworm (*Tenebrio molitor*) as a sustainable functional ingredient in bakery products: a key step toward sustainable diets, *Future Foods*, 2025, **12**, 100716, DOI: [10.1016/j.fufo.2025.100716](https://doi.org/10.1016/j.fufo.2025.100716).
- 113 A. A. Faria and J. Kang, It's not just about the food: motivators of food patterns and their link with sustainable food neophobia, *Appetite*, 2022, **174**, 106008, DOI: [10.1016/j.appet.2022.106008](https://doi.org/10.1016/j.appet.2022.106008).
- 114 S. E. García-Barón, R. P. Carmona-Escutia, E. J. Herrera-López, D. A. Leyva-Trinidad and A. Gschaedler-Mathis, Consumers' Drivers of Perception and Preference of Fermented Food Products and Beverages: A Systematic Review, *Foods*, 2025, **14**, 713, DOI: [10.3390/foods14050713](https://doi.org/10.3390/foods14050713).



- 115 K. Estay, C. Proserpio, C. Cattaneo and M. Laureati, Children's food neophobia across different socioeconomic backgrounds in Chile: Exploring acceptance and willingness to try unfamiliar vegetables, *Food Qual. Prefer.*, 2025, **129**, 105511, DOI: [10.1016/j.foodqual.2025.105511](https://doi.org/10.1016/j.foodqual.2025.105511).
- 116 C. Günden, P. Atakan, M. Yercan, K. Mattas and M. Knez, Consumer Response to Novel Foods: A Review of Behavioral Barriers and Drivers, *Foods*, 2024, **13**, 2051, DOI: [10.3390/foods13132051](https://doi.org/10.3390/foods13132051).
- 117 D. Granato, A. Mocan and J. S. Câmara, Is a higher ingestion of phenolic compounds the best dietary strategy? A scientific opinion on the deleterious effects of polyphenols in vivo, *Trends Food Sci. Technol.*, 2020, **98**, 162–166, DOI: [10.1016/j.tifs.2020.01.010](https://doi.org/10.1016/j.tifs.2020.01.010).
- 118 A. Arce-Reynoso, R. Mateos, E. J. Mendivil, V. M. Zamora-Gasga and S. G. Sáyago-Ayerdi, Bioavailability of bioactive compounds in Hibiscus sabdariffa beverage as a potential anti-inflammatory, *Food Res. Int.*, 2023, **174**, 113581, DOI: [10.1016/j.foodres.2023.113581](https://doi.org/10.1016/j.foodres.2023.113581).
- 119 H. Wu, Z. Liu, P. Lu, C. Barrow, F. R. Dunshea and H. A. R. Suleria, Bioaccessibility and bioactivities of phenolic compounds from roasted coffee beans during in vitro digestion and colonic fermentation, *Food Chem.*, 2022, **386**, 132794, DOI: [10.1016/j.foodchem.2022.132794](https://doi.org/10.1016/j.foodchem.2022.132794).
- 120 D. Mechi, B. Baccouri, D. Martín-Vertedor and L. Abaza, Bioavailability of Phenolic Compounds in Californian-Style Table Olives with Tunisian Aqueous Olive Leaf Extracts, *Molecules*, 2023, **28**, 707, DOI: [10.3390/molecules28020707](https://doi.org/10.3390/molecules28020707).
- 121 C. X. Li, F. R. Wang, B. Zhang, Z. Y. Deng and H. Y. Li, Stability and antioxidant activity of phenolic compounds during in vitro digestion, *J. Food Sci.*, 2023, **88**, 696–716, DOI: [10.1111/1750-3841.16440](https://doi.org/10.1111/1750-3841.16440).
- 122 D. M. Ribnicky, D. E. Roopchand, A. Oren, M. Grace, A. Poulev, M. A. Lila, *et al.*, Effects of a high fat meal matrix and protein complexation on the bioaccessibility of blueberry anthocyanins using the TNO gastrointestinal model (TIM-1), *Food Chem.*, 2014, **142**, 349–357, DOI: [10.1016/j.foodchem.2013.07.073](https://doi.org/10.1016/j.foodchem.2013.07.073).
- 123 G. Mandalari, M. Vardakou, R. Faulks, C. Bisignano, M. Martorana, A. Smeriglio, *et al.*, Food Matrix Effects of Polyphenol Bioaccessibility from Almond Skin during Simulated Human Digestion, *Nutrients*, 2016, **8**, 568, DOI: [10.3390/nu8090568](https://doi.org/10.3390/nu8090568).
- 124 H. Liu, Y. Wang, Y. Wang, J. Wang, H. Hu, X. Zhong, *et al.*, Effective Identification of Variety and Origin of Chenpi Using Hyperspectral Imaging Assisted with Chemometric Models, *Foods*, 2025, **14**, 1979, DOI: [10.3390/foods14111979](https://doi.org/10.3390/foods14111979).
- 125 F. Cirrincione, P. Ferranti, A. Ferrara and A. Romano, A critical evaluation on the valorization strategies to reduce and reuse orange waste in bakery industry, *Food Res. Int.*, 2024, **187**, 114422, DOI: [10.1016/j.foodres.2024.114422](https://doi.org/10.1016/j.foodres.2024.114422).
- 126 M. F. S. Mohamed, Y.-L. Yu, L.-H. Wang, M.-K. Lin and J. Chao, Advances in phytochemistry, pharmacology, and sustainable utilization of orange peels, *Int. J. Food Prop.*, 2026, **29**(1), 2637277, DOI: [10.1080/10942912.2026.2637277](https://doi.org/10.1080/10942912.2026.2637277).
- 127 C.-A. Danciu, A.-G. Mag, C. Stanciu, L. Vidu and M. Stanciu, Valorization of Plant-Based Food By-Products Through Green Extraction of Bioactive Compounds for Functional Food, *Molecules*, 2026, **31**, 646, DOI: [10.3390/molecules31040646](https://doi.org/10.3390/molecules31040646).
- 128 S. Yadav, K. Malik, J. M. Moore, B. R. Kamboj, S. Malik, V. K. Malik, *et al.*, Valorisation of Agri-Food Waste for Bioactive Compounds: Recent Trends and Future Sustainable Challenges, *Molecules*, 2024, **29**, 2055, DOI: [10.3390/molecules29092055](https://doi.org/10.3390/molecules29092055).
- 129 S. Zheng, Z. Huang, L. Dong, D. Li, X. Hu, F. Chen, *et al.*, Sustainable Extraction Technology of Fruit and Vegetable Residues as Novel Food Ingredients, *Foods*, 2025, **14**, 331, DOI: [10.3390/foods14020331](https://doi.org/10.3390/foods14020331).
- 130 Y. A. Bhadange, J. Carpenter and V. K. Saharan, A Comprehensive Review on Advanced Extraction Techniques for Retrieving Bioactive Components from Natural Sources, *ACS Omega*, 2024, **9**, 31274–31297, DOI: [10.1021/acsomega.4c02718](https://doi.org/10.1021/acsomega.4c02718).
- 131 B. L. Angioletti Decker, T. V. Fonteles, F. A. N. Fernandes and S. Rodrigues, Green extraction technologies for valorising Brazilian agri-food waste, *Sustainable Food Technol.*, 2026, **4**, 62–88, DOI: [10.1039/D5FB00323G](https://doi.org/10.1039/D5FB00323G).
- 132 M. Laureati, A. De Boni, A. Saba, E. Lamy, F. Minervini, A. M. Delgado, *et al.*, Determinants of Consumers' Acceptance and Adoption of Novel Food in View of More Resilient and Sustainable Food Systems in the EU: A Systematic Literature Review, *Foods*, 2024, **13**, 1534, DOI: [10.3390/foods13101534](https://doi.org/10.3390/foods13101534).
- 133 L. G. Lima Nascimento, D. Odelli, A. Fernandes de Carvalho, E. Martins, G. Delaplace, P. Peres de sa Peixoto Júnior, *et al.*, Combination of Milk and Plant Proteins to Develop Novel Food Systems: What Are the Limits?, *Foods*, 2023, **12**, 2385, DOI: [10.3390/foods12122385](https://doi.org/10.3390/foods12122385).
- 134 J. Núñez-Pérez, J. L. Burbano-García, R. Espín-Valladares, M. V. Lara-Fiallos, J. C. DelaVega-Quintero, M. Cevallos-Vallejos, *et al.*, Cascade Valorisation of Lemon Processing Residues (Part II): Integrated Biorefinery Design, Circular Economy, and Techno-Economic Feasibility, *Foods*, 2026, **15**, 1041, DOI: [10.3390/foods15061041](https://doi.org/10.3390/foods15061041).
- 135 Z. Lan, X. Zhang, X. Cai, W. Pan, W. Li, Y. Sun, *et al.*, Metabolite profiling and 'dryness'-like effect analysis of 'Chenpi' plant tea (*Citri Reticulatae Pericarpium peel*) with and without steaming peocessing, *Food Res. Int.*, 2025, **206**, 116089, DOI: [10.1016/j.foodres.2025.116089](https://doi.org/10.1016/j.foodres.2025.116089).
- 136 Y. Liu, H. Wen, J. Kong, Z. Hu, Y. Hu, J. Zeng, *et al.*, Flavor characterization of *Citri Reticulatae Pericarpium* (*Citrus reticulata* 'Chachiensis') with different aging years via sensory and metabolomic approaches, *Food Chem.*, 2024, **443**, 138616, DOI: [10.1016/j.foodchem.2024.138616](https://doi.org/10.1016/j.foodchem.2024.138616).
- 137 D. Yang, X. Wu, H. Shi, J. Zhang and C. Wang, Essential aroma substances and release pattern of Xinhui Chenpi, *Beverage Plant Res.*, 2022, **2**, 1–8, DOI: [10.48130/BPR-2022-0022](https://doi.org/10.48130/BPR-2022-0022).

