

Faraday Discussions

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

This article can be cited before page numbers have been issued, to do this please use: D. Turculet, S. Miao, K. Agg and S. Perkin, *Faraday Discuss.*, 2026, DOI: 10.1039/D5FD00145E.

Speciation and hydration forces in sodium carbonate/bicarbonate aqueous solutions nanoconfined between mica sheets[†]

Daria Turculet, Shurui Miao, Kieran J. Agg and Susan Perkin*

Received 00th January 20xx, Accepted 00th January 20xx DOI: 10.1039/x0xx00000x

Abstract

The equilibrium between hydrated and hydrolysed forms of CO₂ in water is central to a multitude of processes in geology, oceanography and biology. Chemistry of the carbonate system is well understood in bulk solution, however processes such as mineral weathering and biomineralisation frequently occur in nano-confined spaces where carbonate chemistry is less explored. For confined systems, the speciation equilibria are expected to tilt due to surface reactivity, electric fields and reduced configurational entropy. In this discussion paper we provide measurements of interaction force between negatively charged aluminosilicate (mica) sheets across aqueous carbonate/bicarbonate solutions confined to nanoscale films in equilibrium with a reservoir of the solution. By fitting the measurements to a Poisson-Boltzmann equation modified to account for charge regulation at the bounding walls, we discuss features of the bicarbonate speciation in confinement. We find that (i) the presence of bicarbonate in the bulk reservoir causes a repulsive excess pressure in the slit compared to pH-neutral salt solutions at the same concentration, arising from a higher (negative) effective charge on the mica surfaces; (ii) the electrostatic screening length is lower for solutions of Na₂CO₃ compared to NaHCO₃ at the same bulk concentration, due to a shift in the speciation equilibria with pH and in accordance with Debye-Hückel theory; (iii) hydration forces are observed at distances below 2 nm with features of size 0.1 nm and 0.3 nm; this was reproducible across the various bicarbonate electrolytes studied, and contrasts with hydration forces of uniform step size measured in pH-neutral electrolytes.

Introduction

Carbonates play a key role in regulating Earth's carbon cycle and often serves as reliable record of the co-evolution of Earth and life. The geological formation of carbonate minerals is highly sensitive to variations in calcium and magnesium concentrations. Even subtle changes in the relative abundance of these ions can shift the delicate balance of electrostatic interactions, hydration forces, and mineral formation and dissolution kinetics, favouring the precipitation of specific polymorphs such as calcite, aragonite, or dolomite.^{1,2} Likewise, life on Earth has evolved intricate and diverse mechanisms to control carbonate mineralization. Organisms such as corals, mollusks, and plankton actively mediate carbonate mineral formation by regulating local ion concentrations and pH, enabling the construction of shells and skeletons.³ In addition to the structural role of biomineralisation, the carbonate-bicarbonate system is also one of the most fundamental acid-base regulation systems in biology. It is widespread among life forms ranging from vertebrates and microbes, where it serves as a buffer system to maintain pH balance⁴; to algae and plants, where it is used to produce dissolved CO₂ as an alternative carbon source for photosynthesis⁵.

In aqueous environments, sodium carbonate and bicarbonate solutions constitute an acid-base buffer system. Dissolved inorganic carbon (DIC) does not exist as a single species but as a dynamic equilibrium among carbonic acid (H₂CO₃), bicarbonate (HCO₃⁻), and carbonate (CO₃²⁻), whose relative abundances are governed by the bulk pH through the following equilibria (at *T* = 298 K):



The equilibrium constants⁶ (or acid dissociation constants) determine the speciation of carbon: carbonic acid dominates under acidic conditions, bicarbonate prevails in the near-neutral range, and carbonate ions become significant at high pH. The fractional abundances of each species as a function of pH are illustrated in Figure 1(a), which highlights the transition from H₂CO₃ to HCO₃⁻ and finally to CO₃²⁻ with increasing pH.

Many geological and biological processes involving carbonates, such as weathering and biomineralisation, occur within confined environments. The interfaces and confining geometry imposed by mineral pore spaces or biomolecular matrices such liposomes or extracellular fluid compartments direct reactivity towards distinct pathways³. Such confinement appears to strongly influence the nucleation, growth, and dissolution of carbonate⁷⁻⁹, however the mechanisms by which this occurs are complex and not fully known^{10,11}. Electric fields arising from surface charges, specific chemical interactions with the confining surfaces, and restricted configurations and limited ion mobility in confined regions are all likely to play a role. Understanding how charged solid-liquid interfaces, and degrees of confinement, shape the thermodynamic and kinetic behaviour of carbonate-bicarbonate systems is therefore essential for linking molecular processes to macroscopic geochemical and biological outcomes.






Figure 1 (a) Fractional concentrations of the three dissolved inorganic carbon (DIC) species present in bicarbonate-containing solutions as a function of electrolyte pH. Calculated using the equilibrium constants as in equation 1 as shown in the SI. (b) Schematic diagram of the SFB used to measure interactions between mica sheets across electrolyte solution. The mica-coated lens setup is immersed in bulk solution, so that confined system is in equilibrium with a reservoir.

In this work we present model experiments representing an idealised confined geometry: we study aqueous solutions of the carbonate/bicarbonate system confined to nanoscale films between two atomically smooth muscovite mica sheets using a surface force balance (SFB)¹². Muscovite mica is well suited to high-resolution experimental studies of this sort due to its ideal cleavage along the basal plane revealing an atomically-smooth crystal surface over macroscopic ($\approx \text{cm}^2$) areas. Muscovite mica is a phyllosilicate clay with overall composition $\text{KSi}_3\text{Al}_3\text{O}_{10}(\text{OH})_2$ consisting of tetrahedral-octahedral-tetrahedral (TOT) silicate layers. 1 out of 4 of the Si atoms in the tetrahedral layers is substituted for Al, leading to a net negative charge on the TOT lattice which is neutralised by K^+ ions lying between the aluminosilicate sheets¹³. When immersed in electrolyte solutions, the surface K^+ ions can exchange with cations from the bulk electrolyte. The adsorbed cations are either directly associated with the aluminosilicate lattice or retain their primary hydration layer; the latter form the so-called Stern layer¹⁴. However, ions adsorbed (hydrated or otherwise) to mica in aqueous solution usually do not fully neutralise the surface charge; instead, the surface retains an effective charge which is neutralised by dissolved ions in nearby solution extending over a distance characterised by the electrostatic screening length (κ_D^{-1})¹⁵. According to mean-field descriptions, two particles or surfaces approaching to distances such that the electrical double layers overlap will experience an interaction force, always repulsive for two identical surfaces, arising from the excess osmotic pressure in the confined region. Measurements of this force/pressure can be fitted (in the far-field) using the Poisson-Boltzmann equation with appropriate boundary conditions; the result is fitted empirical values for the effective surface potential or surface charge, screening length, dielectric permittivity, etc. under different solution conditions.

At very strong confinement (to surface separations below a few nm), an additional *hydration force* is often measured between two mica sheets across electrolyte solutions¹⁶. In previous measurements, it was found that the nature of the hydration forces is strongly connected to the ion exchange process. For example, in acidic solutions there were no hydration forces at all; face-to-face mica sheets experience an attraction into direct contact as predicted by DLVO theory. On the other hand, in solutions containing alkali halide salts (LiCl, NaCl, KCl, CsCl) above a critical concentration an additional short-range repulsion was measured. These additional short-range forces, or hydration forces, were attributed to the adsorption of hydrated ions at the mica surface which are not de-hydrated under the attractive van der Waals pressure between the mica sheets^{17–19}. Detailed investigation of the hydration forces revealed that the repulsion is not monotonic, but instead consists of oscillations with a wavelength close to the size of a water molecule^{16,20,21}. The observation of oscillatory hydration force has been accompanied by a heuristic interpretation of ‘squeezing out’ water layers, while theoretical models more rigorously indicate that oscillations arise from a resonance of the bulk structure in the cavity between the surfaces^{22,23}.

Measurements of interaction forces between mica sheets across a wide range of electrolyte solutions have been made in the past^{17,18,24}, revealing trends in the surface adsorption properties and hydration forces. However, since mica is not a simple protic surface characterised by a single pK value, the relation between solution conditions and the adsorption equilibria are ion-specific and quite complex²⁵. Here, we report surface force measurements between mica sheets across aqueous solutions of NaHCO_3 and Na_2CO_3 at different concentrations, and compare these to literature measurements for KCl. We consider the fitted effective surface charge, screening length, and hydration forces observed. Each of these appears to be distinctly different in the bicarbonate-containing solutions compared to KCl; we discuss the role of speciation on surface and colloidal interactions, and make connections to recent grand-canonical Monte-Carlo simulations of surface forces in a similar charge regulating system.

Methods

The interaction force, F , as a function of separation distance, D , between atomically smooth mica sheets in crossed-cylinder configuration and immersed in aqueous solutions was measured using a Surface Force Balance (SFB)¹². When the radius of curvature of the mica



cylinders, R , is much greater than the separation between them, *i.e.* $R \gg D$, we can write that $F/2\pi R = W^{\parallel}$ where W^{\parallel} is the free energy of interaction (potential of mean force) between parallel sheets, per unit area, at separation D (the Derjaguin approximation). This is essentially exact in our case where $R \approx 1$ cm, $D \approx 1 - 100$ nm. For comparison to theory and simulations, W^{\parallel} is the excess grand potential since the electrolyte between the mica sheets is in (electro-)chemical and thermal equilibrium with the reservoir²⁶.

White light interferometry is used to determine D (equivalent to the electrolyte film thickness) to 0.1 nm precision and 0.2-0.5 nm accuracy relative to mica-mica contact; in the present work we present measurements of distance relative to the closest approach within an experiment which we denote D_0 . Forces are measured via deflection of a spring in the direction perpendicular to the crossed cylinders' axes. A schematic of the SFB is provided in Figure 1(b). Interactions were measured from ≈ 200 nm down to contact, however we present only the region at shorter distances where non-zero forces were detected. The detailed design and operating principles of the apparatus have been described extensively elsewhere¹². Details specific to the present study are as follows. Ruby muscovite mica sheets (S&J Trading Inc.) were cleaved in a particle-free environment to ensure atomically smooth surfaces. The mica pieces used had thicknesses between 2-5 μm and were back-silvered with a 45 nm thick silver layer prior to being mounted on cylindrical glass lenses using EPON glue. Electrolyte solutions consisted of sodium bicarbonate (NaHCO_3) and sodium carbonate (Na_2CO_3) salts obtained in anhydrous form from Fisher Scientific UK Ltd. with a purity of 99.99% (Puratronic[®]) and potassium chloride (KCl) obtained from Thermo Scientific with a purity of 99.997% (Puratronic[®]). The water used was generated from a Milli-Q high-purity system, with the total organic carbon (TOC) < 4 ppb and the resistivity of 18.2 $\text{M}\Omega\text{cm}$. Solutions were prepared at concentrations of 1 mM for KCl, 1 mM and 10 mM for NaHCO_3 , and 10 mM for Na_2CO_3 . The pH of each electrolyte solution was measured using a calibrated pH meter (HI5221, HANNA[®] Instruments)

Results and Discussion

General features of the force-distance profiles

The interaction force (F) as a function of separation distance (D), was measured between atomically smooth mica surfaces immersed in aqueous solutions of sodium bicarbonate and sodium carbonate for various bulk concentrations in the range 1-100 mM. Forces were measured on both approach and retraction of the surfaces, for distances in the range ≈ 200 nm down to 0 nm. A representative force-distance profile is shown in Figure 2, for 10 mM Na_2CO_3 . The mica surfaces experience a repulsive force, measurable from approximately 10-15 nm, which increases exponentially with decreasing separation down to ≈ 2 nm. Below 2 nm, small jumps inwards are observed, of size 0.1 nm then 0.3 nm. On retraction of the surfaces a hysteresis is observed in the region of the jumps (spring instabilities) but the force profile becomes reversible again beyond 1-2 nm. At contact or near-contact separations, $0 < D < 0.2$ nm, the interaction force remains repulsive at all pressures applied; there is no adhesion. Qualitatively similar features were observed for all bicarbonate electrolytes studied, and can be interpreted in terms of (reversible) mean-field DLVO forces at longer range and structural forces, or *hydration forces*, at short range arising from the discontinuous squeeze-out of hydrated ions from the film between the mica sheets. In the following we discuss quantitative differences observed between bicarbonate solutions at varying concentration and pH, and their comparison to simple salt solutions.




Figure 2 Interaction free energy per unit area, $W^{\parallel} = F/2\pi R$, between mica sheets as a function of their separation distance (relative to closest separation), D_0 , measured across 10 mM Na_2CO_3 aqueous solution at 294 K. Interactions measured on approach (blue curve) and retraction (red curve) are shown. The force profiles are reversible in the range down to 1 nm, then small steps and a hysteresis are seen (enlarged in the inset). At the smallest separations the force is again reversible, showing absence of adhesion.



Bicarbonate in the reservoir adjusts the effective surface charge

Figure 3(a) compares the measured interaction force across 1 mM NaHCO₃ to the interaction force across 1 mM KCl. The range of the interaction, captured by the exponential decay length, is similar for the two electrolytes. However, the magnitude of the force is substantially larger in the case of NaHCO₃ despite the same 1 mM bulk concentration. To interpret these observations we consider a simple model for the mean-field interaction force between symmetric charged surfaces across electrolyte, involving a sum of van der Waals and electrostatic forces. Electrostatic forces are modelled using the Poisson-Boltzmann equation, expected to be suitable at this low bulk ion concentration, with allowance for charge regulation as the surfaces approach. To account for the charge regulation we use a constant regulation approximation^{15,27}, involving a regulation parameter, p , taking values between 0 and 1 to interpolate between the limits of constant potential (CP; $p = 0$) and constant charge (CC; $p = 1$)^{28,29}. The total interaction force including the van der Waals term is:

$$\frac{F}{2\pi R} = W^{\parallel} = -\frac{A}{12\pi D^2} + 2\varepsilon_0\varepsilon_r\kappa\psi_{\text{eff}}^2 \frac{e^{-\kappa D}}{1 + (1-2p)e^{-\kappa D}} \quad (2)$$

where A is the Hamaker constant, ε_0 is the permittivity of free space, ε_r is the relative permittivity, ψ_{eff} is the mica effective surface potential (at large surface separation), κ^{-1} is the screening length. F , R and D are measured in the experiment, A and $\varepsilon_0\varepsilon_r$ are known for the system, and ψ_{eff} , κ and p are treated as fitting parameters. κ^{-1} may also be compared to the theoretical Debye-Hückel screening length, κ_D^{-1} (see below). The fitted values for these force profiles are shown in Table 1. The fitted screening length κ^{-1} is the same within error for the two electrolytes and is in accordance with the calculated Debye-Hückel screening length for a 1:1 electrolyte at 1 mM concentration. However, there are two clear differences between the measured force profiles for 1 mM NaHCO₃ and 1 mM KCl: (i) 1 mM NaHCO₃ has a larger fitted ψ_{eff} , and (ii) the fitted p -values imply almost constant charge boundary condition for 1 mM NaHCO₃ in contrast to KCl which facilitates charge regulation. We will consider these two observations in turn.




Figure 3 Interaction energy per unit area measured between mica sheets as a function of separation distance. (a) 1 mM NaHCO₃ compared to 1 mM KCl²¹. (b) 1 mM NaHCO₃ compared to 10 mM NaHCO₃ and 10 mM Na₂CO₃. In all cases the dashed lines are fits to equation 2, with fitting parameters in Table 1. The decay length was treated as a free fitting parameter and obtained by nonlinear least-squares fitting of the force–distance data for distances down to 5 nm (1 mM NaHCO₃), 3.2 nm (10 mM NaHCO₃), 3.5 nm (10 mM Na₂CO₃), and 3 nm (1 mM KCl).

Table 1 Fitting parameters for the datasets shown in Figure 3. These fitting parameters and associated errors are relevant for the individual profiles; in the SI we provide fitting parameters for the multiple force-runs carried out in each solution including across multiple experiments (different mica sheets).

Parameter	1 mM NaHCO ₃	10 mM NaHCO ₃	10 mM Na ₂ CO ₃	1 mM KCl
κ^{-1} / nm	10.2 ± 0.2	3.2 ± 0.05	2.2 ± 0.1	10.0 ± 0.2
p	0.85 ± 0.10	0.74 ± 0.04	0.77 ± 0.09	0.75 ± 0.04
ψ_{eff} / mV	87 ± 10	64 ± 3	35 ± 3	58 ± 2


The difference in ψ_{eff} between 1 mM KCl and 1 mM NaHCO₃ is unlikely to arise from the K⁺/Na⁺ cations¹⁸. Instead, the origin of the striking difference in ψ_{eff} is likely attributed to the higher reservoir pH of 1 mM NaHCO₃; the bicarbonate speciation equilibria, equation 1,



act as a sink for protons and cations, driving the mica surface adsorption equilibria in the direction of more negative surface charge. To illustrate this, a simplistic model for the mass-action equilibria involving mica and its counterions is shown in equation 3. The equilibrium involves mica neutralised by adsorbed ions (here M^+ implies metal, e.g. Na^+ or K^+), mica with hydrated ions adsorbed (often called "Stern layer"), and mica holding a negative charge due to dissolution of counterions into the diffuse part of the electrical double layer. The mica equilibrium is distinct from that of a protic surface such as silica, and mica is not normally considered to have a simple isoelectric point or surface acidity constant. Nonetheless adsorption and surface charge respond to the bulk electrolyte conditions (concentration, pH) through multiple coupled processes.

We are not aware of calculations or simulations of the relevant mica equilibria in the presence of bicarbonate or other pH-buffering solution, although suitable methods have recently been devised and applied to similar scenarios^{30,31} as discussed below. Complicating factors include the variation in H^+ concentration in the electrical double layer vs. bulk solution due to the requirement for constant electrochemical potential between the reservoir and confined/surface regions³² and solvent dielectric variations between the Stern layer and bulk.

Nonetheless, it is clear that the mica surface equilibria are coupled to the bicarbonate bulk speciation equilibria, equation 1, through the hydronium ion: an increase in pH due to the presence of bicarbonate in the reservoir (away from the surfaces) drives the mica equilibrium in the direction of more negative charge and higher negative surface potential.



While the mica-charging and ψ_{eff} effects discussed above apply to mica-solution interfaces in general (no confinement; $D = \infty$), charge regulation is intrinsically a confinement effect ($D \rightarrow 0$). The fitted p -values for 1 mM NaHCO_3 and 1 mM KCl reveal differences in charge regulation, with 1 mM NaHCO_3 closer to the constant-charge limit and 1 mM KCl displaying more pronounced charge regulation, albeit with only a modest but systematic difference rather than a sharp transition between boundary conditions. This is directly apparent on inspection of the force profiles: the curvature of $\ln(W)$ vs. D below about 10 nm (*i.e.* $D < \kappa^{-1}$) across 1 mM NaHCO_3 is convex, deviating from the plain exponential decay at larger distances where the boundary condition has little effect on the functional form. As D decreases in the range $D < \kappa^{-1}$, the boundary condition (CC or CR) has an increasingly strong influence on the interaction. The molecular origin of CR, for 1 mM KCl , involves adsorption of diffuse-layer ions to the surfaces as D decreases to lower the overall free energy of interaction. In 1 mM NaHCO_3 , on the other hand, there is little charge regulation implying that ions are not driven to adsorb on the surfaces during compression of the film; the surface charge remains constant (and high), giving rise to a steeper increase in excess pressure. Looking again at the simple mass-action expressions in equation 3, we can interpret that, for KCl , the position of equilibrium shifts to the left as two mica surfaces approach whereas in NaHCO_3 this is not the case. It may be that, for KCl solutions, adsorption of H_3O^+ onto the mica or incorporation of H^+ within it provide the mechanism for CR whilst this mechanism is not available at high pH and low ionic strength. A supporting counterexample is the observation that acidic solutions displayed even stronger reduction of the mica surface charge during approach of two surfaces leading to attraction and adhesion¹⁷.

The effect of bulk sodium bicarbonate concentration is illustrated by comparing the 1 mM NaHCO_3 measurement to 10 mM NaHCO_3 in Figure 3(b) and corresponding fitting parameters in Table 1. With increased NaHCO_3 concentration the surface potential is lower, the range (screening length) is shorter, and the boundary condition best fitting the measurement is with substantial charge regulation. It appears that, with a higher concentration of Na^+ and HCO_3^- in solution, charge regulation is now favoured during surface approach; the 10-fold increase in solution Na^+ concentration appears sufficient to shift the equilibrium fraction of charged mica sites. Although ψ_{eff} is lower in 10 mM compared to 1 mM NaHCO_3 , it remains substantially higher than in KCl or other 1:1 salt solutions at similar concentration. Therefore the influence of high pH on effective charge and ψ_{eff} remain at this higher concentration. The decrease in screening length is anticipated within the Debye-Hückel theory; quantitative effects of the speciation involved are discussed in the next section.

Screening length depends on carbonate speciation

In Figure 3(b) we make a further comparison between the mica-mica interaction free energy across solutions of sodium bicarbonate (NaHCO_3) and sodium carbonate (Na_2CO_3). As anticipated by the speciation equilibria for bicarbonate solutions in equation 1, addition of sodium bicarbonate tips the equilibria to the right and the resulting solution has higher pH; the measured pH and absolute concentrations of each species in the equilibrium are provided for our experimental scenarios in Table 2. The impact of this shift in the speciation equilibria on surface forces is clear: we see that the absolute magnitude of the repulsion is lower, and the screening length κ^{-1} is also shorter, for Na_2CO_3 compared to NaHCO_3 at the same 10 mM reservoir concentration. To rationalise the lower κ^{-1} in Na_2CO_3 , we calculate the theoretical Debye screening length:

$$\kappa_D^{-1} = \left(\frac{\epsilon_r \epsilon_0 k_B T}{\frac{1}{2} \sum_i c_i e^2 z_i^2} \right)^{1/2} \quad (4)$$

with e the electron charge, c_i the concentration of ionic species i and z_i its valence. Taking the sum over all ionic species present: Na^+ , HCO_3^- , CO_3^{2-} (with H^+ and OH^- having negligible concentration), and calculating their concentrations using the equilibrium constants for speciation as in equation 1, we calculate theoretical screening lengths as shown in Table 2. The total ionic strength, $I = \frac{1}{2} \sum c_i z_i^2$, is about three times higher in Na_2CO_3 due to almost 100 times higher concentration of CO_3^{2-} and the strong influence of divalent ions on



effective ionic strength of an electrolyte. The result of this is a small decrease in the predicted (theoretical) screening length, κ_D^{-1} from 3.0 nm to 1.7 nm, which is closely similar to the decrease seen in the experimental measurements in Figure 3(b). This effect of speciation on the observed screening length, well anticipated in classic electrolyte theory, was clear and reproducible for the single example electrolyte and concentrations reported here. However we have not yet extended our studies to higher concentrations, where non-ideality (not taken into account in equation 4 or our calculation) is likely to be more significant, or to other buffering species.

Table 2 Theoretical screening length, κ_D^{-1} , and speciation corresponding to the measured pH of the solutions measured and shown in Figure 3.

Solution	pH	$f_{\text{H}_2\text{CO}_3}$ (%)	$f_{\text{HCO}_3^-}$ (%)	$f_{\text{CO}_3^{2-}}$ (%)	Total DIC (M)	$c_{\text{H}_2\text{CO}_3}$ (M)	$c_{\text{HCO}_3^-}$ (M)	$c_{\text{CO}_3^{2-}}$ (M)	Ionic strength (M)	κ_D^{-1} (nm)
1 mM NaHCO ₃	8.03	2.11	97.39	0.5	1×10^{-3}	2.11×10^{-5}	9.74×10^{-4}	5.02×10^{-6}	9.98×10^{-4}	9.6
10 mM NaHCO ₃	8.20	0.00	97.82	0.75	1×10^{-2}	1.43×10^{-4}	9.78×10^{-3}	7.46×10^{-5}	1.00×10^{-2}	3.0
10 mM Na ₂ CO ₃	11.6	0.00	4.96	95.04	1×10^{-2}	2.89×10^{-9}	4.96×10^{-4}	9.50×10^{-3}	3.12×10^{-2}	1.7

Short-range hydration forces in carbonate/bicarbonate systems

At separations below ≈ 2 -3 nm, the force-distance profiles deviate substantially from the mean-field behaviour described in equation 2 and instead displayed discrete, reproducible steps in the measured separation indicative of structural layering transitions in the confined liquid. Two characteristic step sizes were identified: one of approximately 0.1 nm and another of 0.3 nm. These are shown in Figure 4, with one example from each of the bicarbonate solutions studied. The smaller 0.1 nm step consistently appeared at slightly larger separations, preceding the 0.3 nm transition during approach. The 0.3 nm step corresponds closely to the molecular diameter of a water molecule and can be attributed to the expulsion of a single hydration layer between the approaching mica surfaces, a phenomenon commonly observed in SFB and AFM studies of aqueous systems and attributed to structural hydration forces. The smaller 0.1 nm step, however, is unusual. Its reproducibility across multiple measurements and solution conditions, always when bicarbonate-containing electrolytes are present, suggests a genuine interfacial event rather than a mechanical or instrumental artefact. The origin of this 0.1 nm step remains uncertain, one possible explanation is the partial displacement or rearrangement of hydrated protons or tightly bound ion-water complexes near the charged mica surface. The fact that this step always precedes the water-layer squeeze-out supports the idea that subtle protonic or hydration-shell reorganizations occur before complete dehydration of the confined region. Further experimental and theoretical work will be required to uncover its origin.




Figure 4 High resolution measurement of the interaction as a function of separation below ≈ 2 nm, in (a)-(c) for each of the solutions discussed above, showing discontinuities attributed to structural changes in the film. In each case a step of 0.1 nm precedes a step of 0.3 nm.

Conclusions

We have presented initial results for the interaction free energy, derived from direct force measurements, between muscovite mica sheets across aqueous solutions of carbonate and bicarbonate salts and in equilibrium with bulk solution. The results are explained using classical theories for colloidal interactions incorporating electrostatic/entropic and van der Waals terms, except for the hydration forces observed at very short distances. Quantitative comparison to simple salt solutions reveals a strong influence of pH on the surface effective potential and, at low concentrations, the near absence of charge-regulation during approach of the surfaces. At 1-2 nm range, the hydration forces also displayed different features.

Recently, grand-canonical Monte-Carlo (GCMC) simulation methods have been developed to handle electrolytes confined between charged surfaces comprised of individual ionisable groups and in equilibrium with a bulk reservoir³⁰⁻³². There it was shown that the force between the surfaces increases more rapidly with decreasing distance for solutions at higher pH, and furthermore that surfaces held at constant charge boundary condition led to steeper repulsion than charge regulating surfaces³¹. The GCMC findings are in accordance with our initial measurements presented here for bicarbonate solutions between mica surfaces, although we note that the simulated surfaces involved a simpler proton-exchange equilibrium compared to the complex system of physisorption normal for aprotic mineral surfaces such as mica. The experiments presented here represent an initial data set and further measurements are required, in particular to approach higher



concentrations of both bicarbonate and background salt.

Our initial findings highlight how speciation of bicarbonates in bulk solution influences interfacial forces and molecular organization in confined electrolytes. Even though the negatively charge carbonate and bicarbonate ions are themselves present at only negligible concentration in the confined region due repulsion from the negatively charged bounding surfaces, they act strongly to influence the pressure and structural forces via the electrochemical potential equilibria between reservoir and film. These findings, and future experiments exploring the bicarbonate system in more detail, may be relevant towards understanding carbonate speciation effects on ion transport and stability in natural and technological aqueous environments¹¹.

Data availability

All raw data is available from the Oxford University Research Archive. See DOI: <https://doi.org/10.5287/ora-9ogqgd0dq>.

Author Contributions

S.P., S.M. and D.T. conceived and planned the experiments. D.T., S.M., and K.A. performed the experiments and fitting. S.P. D.T., S.M., K.J.A. interpreted the data and wrote the paper.

Conflicts of interest

There are no conflicts to declare.

Acknowledgments

The authors gratefully acknowledge funding from the European Research Council under grant 101001346 ELECTROLYTE. S.M. is supported by a Career Development Research Fellowship from St. John's College, Oxford. K.J.A. would like to acknowledge support from The Oxford-The Queen's College Graduate Scholarship in partnership with the Clarendon Fund, University of Oxford.

References

- [1] R. J. P. Williams and R. Rickaby, *Evolutions Destiny*, The Royal Society of Chemistry, 2012.
- [2] J. Kim, Y. Kimura, B. Puchala, T. Yamazaki, U. Becker and W. Sun, *Science*, 2023, **382**, 915–920.
- [3] H. A. Lowenstam and S. Weiner, *On Biomineralization*, Oxford University Press, 1989.
- [4] D. Voet, J. G. Voet and C. W. Pratt, *Fundamentals of biochemistry : life at the molecular level*, John Wiley & Sons, Hoboken, NJ, Fifth edition. edn, 2016.
- [5] M. Giordano, J. Beardall and J. A. Raven, *Annual Review of Plant Biology*, 2005, **56**, 99–131.
- [6] *CRC Handbook of Chemistry and Physics*, ed. W. M. Haynes, CRC Press, Boca Raton, FL, 100th edn, 2019.
- [7] C. J. Stephens, S. F. Ladden, F. C. Meldrum and H. K. Christenson, *Advanced Functional Materials*, 2010, **20**, 2108–2115.
- [8] F. Kohler, O. Pierre-Louis and D. K. Dysthe, *Nature Communications*, 2022, **13**, 6990.
- [9] J. Dziadkowiec, G. Linga, L. Kalchgruber, S. Kavunga, H.-W. Cheng, O. Nilsen, C. Campsteijn, B. Pokroy and M. Valtiner, *Crystal Growth & Design*, 2024, **24**, 4930–4943.
- [10] B. Jin, Y. Chen, H. Pyles, M. D. Baer, B. A. Legg, Z. Wang, N. M. Washton, K. T. Mueller, D. Baker, G. K. Schenter, C. J. Mundy and J. J. De Yoreo, *Nature Materials* 2024, 2024.
- [11] Y. Xu, J. M. Galloway, L. J. Hasselt and F. C. Meldrum, *Chemical Reviews*, 2025, **0**, null.
- [12] H. J. Hayler, T. S. Groves, A. Guerrini, A. Southam, W. Zheng and S. Perkin, *Reports on Progress in Physics*, 2024, **87**, 046601.
- [13] H. K. Christenson and N. H. Thomson, *Surface Science Reports*, 2016, **71**, 367–390.
- [14] S. S. Lee, P. Fenter, K. L. Nagy and N. C. Sturchio, *Langmuir*, 2012, **28**, 8637–8650.
- [15] A. M. Smith, M. Borkovec and G. Trefalt, *Advances in Colloid and Interface Science*, 2020, **275**, 102078.
- [16] J. N. Israelachvili and R. M. Pashley, *Nature*, 1983, **306**, 249–250.
- [17] R. Pashley, *Journal of Colloid and Interface Science*, 1981, **80**, 153–162.
- [18] R. Pashley, *Journal of Colloid and Interface Science*, 1981, **83**, 531–546.
- [19] S. R. van Lin, K. K. Grotz, I. Siretanu, N. Schwierz and F. Mugele, *Langmuir*, 2019, **35**, 5737–5745.

⁰Physical and Theoretical Chemistry Laboratory, Department of Chemistry, University of Oxford, Oxford OX1 3QZ, UK. E-mail: susan.perkin@chem.ox.ac.uk

[†] Supplementary Information available: additional measurement runs of force profiles and derivation of species fractions for the carbonate system. See DOI: 00.0000/00000000.



- [20] J. E. Hallett, K. J. Agg and S. Perkin, *Proceedings of the National Academy of Sciences*, 2023, **120**, e2215585120.
- [21] K. J. Agg, J. E. Hallett and S. Perkin, *Biophysical Journal*, 2025.
- [22] J. G. Hedley, H. Berthoumieux and A. A. Kornyshev, *The Journal of Physical Chemistry C*, 2023, **127**, 8429–8447.
- [23] T. S. Groves and S. Perkin, *Faraday Discuss.*, 2024, **253**, 193–211.
- [24] T. Baimpos, B. R. Shrestha, S. Raman and M. Valtiner, *Langmuir*, 2014, **30**, 4322–4332.
- [25] B. A. Legg, M. D. Baer, J. Chun, G. K. Schenter, S. Huang, Y. Zhang, Y. Min, C. J. Mundy and J. J. De Yoreo, *Journal of the American Chemical Society*, 2020, **142**, 6093–6102.
- [26] R. Evans and U. Marini Bettolo Marconi, *The Journal of Chemical Physics*, 1987, **86**, 7138–7148.
- [27] G. Trefalt, S. H. Behrens and M. Borkovec, *Langmuir*, 2016, **32**, 380–400.
- [28] S. L. Carnie and D. Y. Chan, *Journal of Colloid and Interface Science*, 1993, **161**, 260–264.
- [29] R. Pericet-Camara, G. Papastavrou, S. H. Behrens and M. Borkovec, *Journal of Physical Chemistry B*, 2004, **108**, 19467–19475.
- [30] Y. Levin and A. Bakhshandeh, *The Journal of Chemical Physics*, 2023, **159**, 111101.
- [31] D. Izzo, A. Bakhshandeh and Y. Levin, *The Journal of Chemical Physics*, 2025, **163**, 064703.
- [32] A. Bakhshandeh, D. Frydel and Y. Levin, *Langmuir*, 2022, **38**, 13963–13971.
- ...



The supporting data has been provided as part of the Supplementary information. Supplementary information: Figures S1-S20 contain the data collected and interpreted in the work. All raw data points have also been uploaded to the Oxford Research Archive, currently awaiting DOI assignment.

[View Article Online](#)

DOI: 10.1039/D5FD00145E

