



Cite this: DOI: 10.1039/d5fo03328d

# Dietary *N*-acylethanolamines are bioaccessible in the small intestine and modulate postprandial hormonal responses: a randomized crossover trial in subjects with ileostomy

Silvia Tagliamonte,<sup>a</sup> Holly R. Neill,<sup>b</sup> Brian Óg Murphy,<sup>b</sup> Kirsty L. Pourshahidi,<sup>b</sup> Francesca De Filippis,<sup>a,c</sup> Danilo Ercolini,<sup>a,c</sup> Chris I. R. Gill,<sup>b</sup> Koos Natalia,<sup>b</sup> Brendan Curran,<sup>b</sup> Murray Nicole,<sup>b</sup> Slevin Mary,<sup>b</sup> Sara Dobani,<sup>b</sup> Massimiliano Fontana<sup>b</sup> and Paola Vitaglione<sup>id \*a,c</sup>

*N*-Acylethanolamines (NAEs) are bioactive lipid mediators involved in the regulation of appetite, inflammation, and gut–brain signaling. This study investigated the metabolic fate of dietary NAEs following the consumption of two test meals with differing NAE contents in subjects with ileostomy and evaluated their effects on gastrointestinal hormones, glycaemia, and appetite regulation. An acute, double-blind, randomized, crossover postprandial study was conducted in ileostomy patients who consumed either a high-NAE meal (HNM) or a low-NAE meal (LNM) on two separate occasions. Ileal fluid and plasma samples were collected over an 8-hour postprandial period for analysis of NAEs and endocannabinoids (ECs). Baseline ileal microbiota composition was assessed. At the end of the 8-hour period, participants completed a buffet meal test to evaluate *ad libitum* energy intake. Dietary NAEs were significantly recovered in ileal fluids after HNM intake, with concentrations approximately 3-fold higher than those after LNM, suggesting partial digestion and release from the food matrix. No significant differences in postprandial plasma NAE concentrations were observed between meals. HNM consumption led to higher postprandial levels of plasma insulin, C-peptide, and glucose-dependent insulinotropic polypeptide, despite no differences in glycemic response or subsequent *ad libitum* energy intake. Metagenomic analysis identified clusters of ileal microbial taxa associated with circulating lipid profiles, suggesting a role of the small intestinal microbiota in the metabolism of NAEs and ECs. Dietary NAEs reach the small intestine at active concentrations and may influence local signaling via GPR119, with microbiota composition influencing their release from food.

Received 5th August 2025,  
 Accepted 29th April 2026

DOI: 10.1039/d5fo03328d

rsc.li/food-function

## 1. Introduction

*N*-acylethanolamines (NAEs) and endocannabinoids (ECs) are a group of endogenous lipid mediators that play important roles in a wide variety of biological processes such as appetite cues, food intake, blood pressure, inflammation, glycaemia, cognition and immunity.<sup>1</sup> NAEs are the congeners of ECs, and both classes of compounds, along with their receptors and enzymes involved in their synthesis and breakdown, constitute the “endocannabinoid system” that is active in several tissues, including the gastrointestinal (GI) tract.<sup>1</sup>

The ECs consist of *N*-arachidonylethanolamide (AEA) and 2-arachidonoylglycerol (2-AG). They may have agonist activity on cannabinoid receptors CB1 and CB2, which are located in the central nervous system (CNS) and in peripheral tissues such as the enteric nervous system (ENS), the liver and adipose tissue.<sup>2</sup> On the other hand, NAEs are known as “endocannabinoid-like” molecules and include oleoylethanolamine (OEA), linoleoylethanolamine (LEA), palmitoylethanolamine (PEA) and stearoylethanolamide (SEA). AEA and NAEs are synthesized from low-abundance phospholipids, *N*-acyl-phosphatidylethanolamines (NAPES).<sup>1,3</sup> In contrast to ECs, they can bind G-protein coupled receptor 119 (GPR119) and peroxisome proliferator-activated receptors (PPARs) located not only in the brain but also in the GI tract mucosa.<sup>4–6</sup> The synthesis of ECs and NAEs is closely interconnected, both among themselves and with other biological systems.<sup>3</sup> Emerging evidence also suggests that OEA mitigates obesity-associated dysmetabolism such as dyslipidemia in chronic diseases, by inducing the

<sup>a</sup>Department of Agricultural Sciences, University of Naples Federico II, 80055 Portici, Italy. E-mail: paola.vitaglione@unina.it

<sup>b</sup>Nutrition Innovation Centre for Food and Health (NICHE), School of Biomedical Sciences, Ulster University, Coleraine, UK

<sup>c</sup>Task Force on Microbiome Studies, University of Naples Federico II, Naples, Italy



expression of genes involved in energy homeostasis, feeding behavior, and lipid metabolism regulation, including GPR119 and PPAR- $\alpha$ .<sup>1,3</sup> As a PPAR- $\alpha$  ligand, it stimulates fatty acid oxidation and lipolysis in white adipose tissue<sup>7,8</sup> and slows gastric emptying.<sup>9</sup> By activating GPR119, OEA promotes the release of anorexigenic hormones such as GLP-1, thereby reducing food intake and enhancing satiety.<sup>10</sup>

Human studies have shown that daily supplementation with capsules containing *N*-oleyl-phosphatidylethanolamine (NOPE; 170 mg) and epigallocatechin-3-gallate (EGCG; 240 mg) can induce weight loss and increase feelings of fullness in overweight and obese individuals.<sup>11,12</sup>

Additionally, an 8-week supplementation with 2  $\times$  125 mg OEA was reported to increase PPAR- $\alpha$  gene expression while reducing appetite, body weight, and fat mass in obese subjects.<sup>13</sup> However, since the supplements used in these studies contained 85% OEA along with LEA and PEA, both known to influence food intake,<sup>14</sup> a combined effect of the three NAEs cannot be ruled out.

Diet composition has been shown to influence fasting and postprandial plasma levels of ECs, NAPEs, and NAEs by affecting the availability of their precursors, such as fatty acids and amines.<sup>1</sup> For instance, consuming an oleic acid-rich breakfast increased postprandial plasma OEA levels and reduced subsequent meal energy intake in humans.<sup>15</sup> More recently, NAPEs, NAEs, and ECs have been identified and quantified in various foods<sup>16</sup> and in the intestinal fluids of fasting ileostomy subjects.<sup>17</sup> Remarkably, ileal concentrations were 1.5 to 13 times higher than the EC<sub>50</sub> values of OEA, PEA, and LEA for GPR119 activation, while OEA concentrations exceeded by approximately 10-fold the levels needed to trigger a physiological response through PPAR- $\alpha$ .<sup>5,10,18</sup> Furthermore, NAEs and NAPEs were shown to be more effective in eliciting a leptogenic and anorexigenic effect from the intestinal lumen than in other tissues.<sup>19,20</sup>

We hypothesize that the levels of NAPEs, NAEs, and ECs in a meal may influence the concentrations of NAEs and ECs in the intestine, subsequently affecting postprandial hormonal response and appetite through interactions with receptors on the intestinal mucosa. To test this hypothesis, we conducted an acute double-blind, randomized, crossover postprandial study in a cohort of ileostomy patients who consumed, on two separate occasions, a high-NAE meal and a low-NAE meal and provided ileal fluids and plasma samples over time postprandially.

## 2. Materials and methods

### 2.1. Study design

This acute double-blind randomized crossover study in a cohort of ileostomy subjects was conducted at the Human Intervention Studies Unit (HISU) at Ulster University (Coleraine, UK). The study was approved by the Office for Research Ethics Committees Northern Ireland (ORECNI) and the University of Ulster Ethical Committee in accordance with the Declaration of Helsinki. All participants signed an informed consent form before enrollment.

The trial was registered at <https://www.clinicaltrials.gov> (NCT05845229). Participants (male and non-pregnant/non-lactating females) recruited to the study were aged 18–70 years, had previously undergone an ileostomy and were  $\geq 2$  years post-operative at the time of recruitment. Participants were excluded if they were lactose intolerant, smokers, or had their surgical procedure  $< 2$  years prior. The postprandial changes in NAEs and ECs in plasma and ileal fluids were registered as the primary outcomes of the study. Secondary outcomes included postprandial changes in plasma glucose, plasma gastrointestinal hormones (*i.e.* insulin, GLP-1, glucose-dependent insulinotropic polypeptide (GIP), glucagon, C-peptide, ghrelin and leptin), composition of the gut microbiome and energy intake at the subsequent meal during a buffet meal test.

### 2.2. Dietary intervention

Fourteen participants were randomized by permuted block randomization to receive either a high-NAE meal (HNM) or a low-NAE meal (LNM) according to the food NAE database.<sup>16</sup> The allocation ratio was 1 : 1. The HNM consisted of cow's milk (150 mL), white bread (46 g), jam (10 g), cocoa powder (15 g), and whole-grain cereal (30 g). The LNM consisted of cow's milk (150 mL), whole-grain bread (80 g), jam (10 g), butter (5 g), instant coffee powder (2 g), and dried apples (30 g). Food items were purchased from the local store – Tesco shop. The nutritional composition of both breakfasts is reported in the Table 1.

The two breakfast meals were designed to be isocaloric ( $\sim 360$ – $344$  kcal) and to provide a significantly different amount of NAEs and NAPEs. The HNM provided  $\sim 13$ -fold more total NAEs (13- and 34-fold more LEA and OEA, respectively) and 5-fold more NAPEs than the LNM (SI Table S1). Total NAPEs were expressed as *N*-arachidonoylphosphatidylethanolamine equivalents according to the food NAE database.<sup>16</sup> Although species-specific concentrations for individual foods were not determined, all NAPEs identified included the major *N*-acyl chains (palmi-

**Table 1** Nutritional composition of the high-NAE meal (HNM) and the low-NAE meal (LNM)

	HNM	LNM
Total weight (g)	251	277
Energy (kcal)	360	344
Carbohydrates (g) (% kcal)	59.2 (66)	59.2 (69)
Of which sugars	20.6	33.7
Fats (g) (%)	5.4 (14)	5.4 (14)
Of which saturated fats	2.5	3.0
Of which monounsaturated fats	2.1	1.9
Of which polyunsaturated fats	0.8	0.5
Proteins (g) (% kcal)	14.8 (16)	9.4 (11)
Dietary fiber (g) (% kcal)	7.9 (4)	10.3 (6)
NAEs ( $\mu\text{g}$ ) <sup>a</sup>	124.0	9.5
Of which LEA <sup>b</sup> ( $\mu\text{g}$ )	69.8	5.4
Of which PEA <sup>c</sup> ( $\mu\text{g}$ )	7.0	2.7
Of which OEA <sup>d</sup> ( $\mu\text{g}$ )	47.1	1.4
NAPEs <sup>e</sup> (mg)	57.8	12.1

<sup>a</sup> *N*-Acylethanolamines. <sup>b</sup> Linoleoylethanolamide. <sup>c</sup> Palmitoylethanolamide. <sup>d</sup> Oleoylethanolamide. <sup>e</sup> *N*-Acylphosphatidylethanolamides.






Fig. 1 Study design.

toyl, linoleoyl and oleoyl), representing the precursors of the monitored NAEs: PEA, LEA, and OEA, respectively. Fig. 1 shows the study design. On the day before the study visit, participants followed a fixed evening meal consisting of plain chicken and boiled white rice to maintain low dietary levels of NAEs, NAs, and ECs.<sup>16</sup> In addition, after this evening meal, participants attached a new stoma bag for the overnight ileal fluid collection. Following an overnight fast, the participants visited the HISU, Ulster University (Coleraine, UK) and provided the overnight ileal fluid samples and baseline blood samples. Anthropometric measures were collected at each visit. Weight (kg) was measured using electronic SECA alpha flat scales (Seca Ltd, Hamburg, Germany) and height (m) was measured using a SECA 213 portable stadiometer (Seca Ltd, Hamburg, Germany). Dietary behavior data were collected through the Three-Factor Eating Questionnaire (TFEQ)<sup>21</sup> and the power of food scale.<sup>22</sup> Individual psychological profiles were ascertained through the Depression, Anxiety and Stress Scale (DASS),<sup>23</sup> the Snaith–Hamilton Pleasure Scale,<sup>24</sup> and the health-related quality-of-life questionnaire (SF-12).<sup>25</sup> Participants also rated their baseline appetite sensations on visual analogue scale (VAS) questionnaires<sup>26</sup> and subsequently consumed the test meal: either a high-NAE meal (HNM) or a low-NAE meal (LNM). Thereafter, blood samples were collected at 2, 4, 6 and 8 hours after breakfast into an EDTA tube, and EDTA tubes were added with protease inhibitors. Ileal effluent samples were collected at 0–2, 2–4, 4–6, and 6–8 hours. At each time point, the participant removed the stoma bag and replaced it with a new one. Prior to glycaemia measurement, subjects rated their hunger, fullness and satiety by VAS questionnaires. After the final sample collection at 8 hours, participants underwent a buffet meal test. They were provided with a tray containing a variety of foods (SI Table S2) to assess *ad libitum* energy intake. The second and third (final) clinic visits took place at least one week later, during which the same procedures were repeated for the alternate treatment. Ileal fluid samples were processed as previously described.<sup>17</sup>

### 2.3. Analysis of EC and NAE contents in ileal and plasma samples

Ileal fluid and plasma samples, collected in EDTA tubes, were processed and analyzed for ECs and NAEs as previously described.<sup>17,27</sup> The chromatographic separation was performed

using an HPLC apparatus provided with two micropumps – PerkinElmer Series 200 (Norwalk, CT, USA). The compounds were separated on a Synergi Max RP 80 column (50 × 2.1 mm) (Phenomenex, USA) with a setting temperature of 30 °C and a flow rate of 0.2 mL min<sup>-1</sup>. The injection volume was 10 μL. The monitored compounds were eluted by a linear gradient of H<sub>2</sub>O and 0.2% formic acid (solvent A) and CH<sub>3</sub>CN (solvent B). According to Tagliamonte and colleagues (2021),<sup>27</sup> the eluting gradient was set as follows: 50–79% B (10 min), 79–95% B (1 min), constant at 95% B (2 min) and finally returning to the initial conditions within 2 min. The acquisition was performed in positive ion mode using an API 3000 triple quadrupole mass spectrometer (Applied Biosystems, Canada) in MRM (Multiple Reaction Monitoring). All the acquisition parameters are summarized in SI Table S3. ECs (2-arachidonoylglycerol, 2-AG; anandamide, AEA) and NAEs (oleoylethanolamide, OEA; linoleoylethanolamide, LEA; palmitoylethanolamide, PEA; SEA, stearoylethanolamide) standards were purchased from Cayman (Cayman Chemical, Ann Arbor, MI).

### 2.4. Ileal gut microbiome composition

The ileal microbiome was analysed at baseline (before breakfast meal consumption) by shotgun metagenomics as previously described.<sup>28</sup> Briefly, microbial DNA was extracted following the SOP 07 developed by the International Human Microbiome Standard Consortium (<https://www.microbiome-standards.org>). DNA libraries were sequenced on an Illumina NovaSeq platform, leading to 2 × 150 bp, paired-end reads. One sample failed library preparation due to poor DNA quality and was removed from the analysis. Therefore, microbiome analysis was carried out on 13 subjects. Human reads were removed using the Best Match Tagger (BMtagger; [https://hmpdacc.org/hmp/doc/HumanSequenceRemoval\\_SOP.pdf](https://hmpdacc.org/hmp/doc/HumanSequenceRemoval_SOP.pdf)). Non-human reads were quality-filtered using PRINSEQ 0.20.4 (trimming when a base with a Phred score <15 was found and discarding reads <75 bp).<sup>29</sup> High-quality reads were imported into MetaPhlan4<sup>30</sup> to obtain species-level, quantitative taxonomic profiles.

### 2.5. Blood glucose and gastrointestinal hormones

Glycaemia was measured at baseline and at postprandial time points (2 h, 4 h, 6 h and 8 h) by finger pricking and using a bedside glucometer (OneTouch SureStep; Life Scan Inc.). Accuracy of the glucometer was evaluated by using least squares linear regression analysis and was found to be 97% “clinically accurate” compared with reference (YSI2700) results. Blood samples were collected into EDTA tubes added with protease inhibitors as previously reported.<sup>31</sup> The samples were centrifuged at 1800g per 15 min at 4 °C, and the supernatants were stored at –80 °C before analysis. The simultaneous quantification of insulin, C-peptide, ghrelin, glucose-dependent insulinotropic peptide (GIP), glucagon-like peptide 1 (GLP-1), leptin and glucagon in plasma samples was achieved using Bio-Plex Pro immunoassay kits as previously described.<sup>31</sup>



## 2.6. Statistical analysis

The sample size was calculated by power analysis, considering the response of NAEs and ECs as endpoints. A sample size of 14 participants was calculated to identify significant changes in the response in accordance with Monteleone *et al.* (2012) and Gatta-Cherifi *et al.* (2012) with an 80% power and an  $\alpha = 0.05$ .<sup>32,33</sup> All statistical analyses were performed using R version 4.2.3. After checking for normality, variables showing a significant positive skewness were log-transformed [ $\ln(x + k)$ , with  $k$  values zeroing the skewness]. Postprandial differences over time within and between the groups for normally distributed variables were evaluated with a one-way repeated-measures ANOVA and Bonferroni adjustment for multiple comparisons; non-normally distributed variables were assessed with Friedman's tests and pairwise Wilcoxon's *post-hoc* tests. The postprandial incremental areas under the curve (iAUCs) were calculated as previously reported<sup>34</sup> and differences in iAUCs between the groups were assessed by an independent sample *t*-test. In order to explore the different responses to HNM *vs.* LNM consumption of gastrointestinal hormones, a multilevel partial least squares discriminant analysis for cross-over design (PLS-DA, *plsda* function) was performed on GI hormone iAUCs (library *mixOmics*). Within-subject correlations among the baseline microbial compositions of the ileal fluids and the postprandial response of ileal concentrations of NAEs were performed by using the *rmcorr* function. The resulting significant correlations were visualized as a network by using the *igraph* package illustrating the significant co-occurring microbial guilds and their interactions with postprandial ileal NAE concentrations.

## 3. Results

### 3.1. Study participants

The participant flow is shown in Fig. 2. Fourteen ileostomates completed the study. One participant was unable to provide a blood sample; therefore, their data were excluded from the NAE analysis. The general characteristics of the participants at baseline are reported in Table 2. Psychometric variables are reported in SI Table S4.

### 3.2. Postprandial concentrations of *N*-acylethanolamines and endocannabinoids in ileal fluids

No difference was found at baseline in the amount of NAEs and ECs quantified in the ileal fluids before the consumption of the LNM or HNM. Among the NAEs, LEA was the most abundant with a quantity on average 50 times greater than OEA, followed by PEA and SEA. The ECs were less abundant than NAEs, with 2-AG being found at an average amount 5-fold greater than AEA (Fig. 3), whereas significantly higher amounts of LEA and OEA were present in ileal fluids (4–6 h and 6–8 h intervals) following HNM compared to LNM consumption (Fig. 4). Specifically, LEA concentrations were approximately 5-fold higher during the 4–6 h interval and 3-fold higher during the 6–8 h interval following the HNM compared to the LNM. Specifically, LEA concen-




Fig. 2 CONSORT diagram of the study.

Table 2 General and anthropometric characteristics of participants ( $n = 14$ ) at baseline

Sex ( $n$ , F/M)	6F, 8M
Age (years)	49.5 ± 12.6
Height (cm)	170 ± 0.01
Weight (kg)	83.4 ± 26.8
BMI (kg m <sup>-2</sup> )	28.6 ± 7.8

Data are expressed as means ± SD.  $n$ , number of participants; F, female; M, male; y, years.





Fig. 3 Ileal fluid content of ECs and NAEs at baseline in the 13 participants before consumption of the LNM (low-NAE meal, in blue) and the HNM (high-NAE meal, in green). 2-AG, 2-arachidonoylglycerol; AEA, arachidonylethanolamide; LEA, linoleylethanolamide; OEA, oleylethanolamide; PEA, palmitoylethanolamide; SEA, stearoylethanolamide.

trations were approximately 5-fold higher during the 4–6 h interval and 3-fold higher during the 6–8 h interval following the HNM compared to the LNM. Similarly, OEA concentrations were around 6-fold higher at 4–6 h and 4-fold higher at 6–8 h post-HNM intake.

No difference over time was shown for the other monitored compounds. Overall, the cumulative 8-hour postprandial recovery of OEA and LEA in ileal fluids was significantly higher after





**Fig. 4** Post-prandial ileal fluid content of ECs and NAEs at each collection interval post-consumption of the LNM (low-NAE meal, in blue) and the HNM (high-NAE meal, in green). Data are shown as mean  $\pm$  SEM. \**P*-value <0.05, between-meal difference versus the HNM vs. the LNM assessed by one-way repeated-measures ANOVA and Bonferroni adjustment. 2-AG, 2-arachidonoylglycerol; AEA, arachidonoylethanolamide; LEA, linoleoylethanolamide; OEA, oleoylethanolamide; PEA, palmitoylethanolamide; SEA, stearoylethanolamide.


HNM intake, with total amounts reaching  $34 \pm 6.7$   $\mu$ g (OEA) and  $1.5 \pm 0.3$  mg (LEA), compared to  $10.3 \pm 1.7$   $\mu$ g and  $0.5 \pm 0.1$  mg, respectively, following the LNM (Fig. 5).

### 3.3. Ileal microbiome and its association with the postprandial lipid mediators in the ileum

The composition of the ileal microbiome in 13 participants at the phylum and species levels is reported in Fig. 6A and B.

One sample failed library preparation due to poor DNA quality and was removed from the analysis.

At the phylum level, the microbiome was mainly dominated by Bacillota (formerly Firmicutes; mean abundance: 74.9%) and



**Fig. 5** Post-prandial ileal fluid total content of ECs and NAEs upon consumption of the LNM (low-NAE meal, in blue) and the HNM (high-NAE meal, in green). \**P*-value <0.05, between-meal difference versus the HNM vs. the LNM assessed by an independent sample *t*-test. 2-AG, 2-arachidonoylglycerol; AEA, arachidonoylethanolamide; LEA, linoleoylethanolamide; OEA, oleoylethanolamide; PEA, palmitoylethanolamide; SEA, stearoylethanolamide.

Pseudomonadota (formerly Proteobacteria; mean abundance: 20.6%). In contrast, Bacteroidota (formerly Bacteroidetes) was detected at >1% abundance in only four subjects, with relative abundances ranging from 0 to 8.8%. At the species level, high variability among the subjects was observed (Fig. 6B).

For subjects where Pseudomonadota was the dominant phylum, *Escherichia coli* and *Haemophilus parainfluenzae* prevailed at the species level. Within Bacillota, *Streptococcus* (*S. salivarius*, *S. parasanguinis*, *S. lutetiensis*, *S. galloyticus* and *S. mitis*), *Romboutsia timonensis*, *Veillonella* spp., *Clostridium perfringens*, *C. disporicum* and *Ligilactobacillus salivarius* were the main species. To explore the associations between the ileal microbiome and the lipid mediators formed in the ileal fluids following the consumption of the HNM and LNM, a network analysis was conducted, and the results are shown in Fig. 6C.

It revealed that each lipid mediator displayed a unique interaction pattern with specific microbial taxa, highlighting their diverse roles within the intestinal ecosystem. OEA and LEA were positively correlated with a wide variety of taxa, including *Clostridium* and *Enterococcus* species and the *Fusobacteriota* phylum, while they were negatively associated with the *Bacillota* phylum. Similarly, PEA, SEA and AEA exhibited overlapping clusters, sharing microbial associations such as *Bifidobacterium* and *Enterococcus* species. In contrast, 2-AG formed a separate and well-defined cluster, primarily involving the *Lactobacillus* genus. A negative correlation was shown with the *Lactobacillus* genus and the pro-inflammatory *R. gnavus* and a positive association with the probiotic *Bifidobacterium bifidum* and the *Actinobacteriota* phylum.

### 3.4. Postprandial plasma *N*-acylethanolamines and endocannabinoids

Following the consumption of the HNM and LNM, the postprandial plasma concentrations of AEA, LEA, OEA and PEA decreased significantly at 2 hours compared to baseline (Fig. 7). No between-meal differences were observed for any of the monitored compounds.


### 3.5. Gastrointestinal hormone response and glycaemia

The postprandial gastrointestinal hormone response to the LNM and HNM interventions is reported in Fig. 8A.


Data showed that the C-peptide and insulin response increased 2 hours after both LNM and HNM intake compared to baseline. Conversely, GIP significantly increased, while ghrelin decreased, 2 hours after HNM intake compared to baseline. The C-peptide and GIP responses were significantly higher 2 hours and 4 hours after HNM than LNM consumption, respectively. The concentration of GLP-1 was below the limit of detection in all the samples analysed. The postprandial incremental areas under the curve (iAUCs) for C-peptide, GIP, and insulin were significantly higher after HNM intake compared to LNM intake (Fig. 8B), determining a distinct response between the meals when analyzed using PLS-DA (Fig. 8C).

No between- and within-meal differences were observed in glycaemic response (Fig. 8D).





**Fig. 6** Composition of the ileal microbiome in 13 participants at the phylum (A) and species (B) levels; (C): network illustrating significant ( $p$ -value  $< 0.05$ ) co-occurring microbial guilds in the intestinal lumen at baseline and their within-subject interaction with the post-prandial ileal total content of ECs and NAEs following consumption of the LNM (low-NAE meal) and the HNM (high-NAE meal).  $R$  coefficients were calculated by performing a repeated-measures correlation. Blue lines indicate positive correlations and red lines indicate negative correlations between variables ( $R > 0$  and  $< 0$ , respectively). Only moderate-to-strong associations ( $r > 0.4$  and  $r < -0.4$ ) were included; weak correlations were excluded from the analysis. The names of the taxon levels are abbreviated as p-phyllum, g-genus and s-species. 2-AG, 2-arachidonoylglycerol; AEA, arachidonylethanolamide; LEA, linoleylethanolamide; OEA, oleoylethanolamide; PEA, palmitoylethanolamide; SEA, stearoylethanolamide.



**Fig. 7** Post-prandial plasma circulating ECs and NAEs upon consumption of the LNM (low-NAE meal, in blue) and the HNM (high-NAE meal, in green). Data are shown as mean  $\pm$  SEM. \* $P$ -value  $< 0.05$ , within-group difference versus baseline assessed by one-way repeated-measures ANOVA and Bonferroni adjustment. 2-AG, 2-arachidonoylglycerol; AEA, arachidonylethanolamide; LEA, linoleylethanolamide; OEA, oleoylethanolamide; PEA, palmitoylethanolamide; SEA, stearoylethanolamide.

### 3.6. Energy intake at subsequent lunch

After 8 hours, participants underwent a buffet meal test to measure *ad libitum* energy intake. No significant effect was


observed for either breakfast type on subsequent energy intake or the macronutrient composition of the consumed meal (SI Fig. S1).

## 4. Discussion

### 4.1. NAEs and ECs increase in the ileal fluids postprandially

Results of this study clearly show a causal relationship between the contents of NAEs, NAEs, and ECs in the meal and the levels of NAEs and ECs in the intestinal lumen postprandially. Indeed, baseline levels of the monitored lipids in ileal fluids were homogeneous across participants, reflecting adherence to a standardized 24-hour diet prior to each test day. In contrast, the levels of NAEs in the ileal fluids increased postprandially when the participants consumed a meal richer in NAEs and in the phosphorylated precursors (NAPes). Specifically, the amount of OEA significantly doubled and LEA tripled the basal values between 4 and 8 hours after consuming the HNM. This finding confirms that food digestion in the upper GI tract releases NAEs and NAPes from the food matrix,





**Fig. 8** (A) Post-prandial concentrations of plasma gastrointestinal hormones upon consumption of the LNM (low-NAE meal, in blue) and the HNM (high-NAE meal, in green). \**P*-value < 0.05, within-meal difference versus the HNM vs. the LNM assessed by one-way repeated-measures ANOVA and Bonferroni adjustment or the non-parametric Friedman test depending on data normal distribution. GIP, glucose-dependent insulinotropic polypeptide. (B) Post-prandial incremental areas under the curve (iAUCs) of the gastrointestinal hormones upon consumption of the LNM (low-NAE meal, in blue) and the HNM (high-NAE meal, in green). \**P*-value < 0.05, between-meal difference versus the HNM vs. the LNM assessed by an independent sample *t*-test. Red arrows indicate individual increases in iAUCs, while black arrows represent individual decreases. (C) The multilevel Partial Least Squares Discriminant Analysis (PLS-DA) for cross-over design by integrating gastrointestinal hormone iAUCs after the HNM (high-NAE meal, in violet) and LNM (low-NAE meal, in yellow). The AUC was 0.82 (*p*-value = 0.0052) when comparing the hormone response upon HNM and LNM intake. (D) Post-prandial plasma glucose upon consumption of the LNM (low-NAE meal, in blue) and the HNM (high-NAE meal, in green). Data are shown as mean  $\pm$  SEM.

continuing the process initiated by mastication in the mouth.<sup>35–37</sup>

In a previous study, the intestinal bioaccessibility of OEA, assessed through *in vitro* digestion, was found to be higher from a liquid than a semisolid food,<sup>38</sup> confirming that the food matrix substantially influenced the ability of digestive enzymes to gain access to it, limiting the rate and extent of the release in the lumen (for a review, see ref. 39). To reduce the impact of the physical form and structure of the two meals on chemical digestion, in the present study, the two breakfast

meals were designed to include foods with different contents of NAEs and NAPEs but similar matrices and macronutrient composition. Interestingly, the ratio of cumulative OEA to LEA recovery in ileal fluids did not mirror the relative proportions of these compounds in the meals. This observation may reflect selective conversion from their respective dietary NAPE precursors or cross-regulation among NAE species within the intestinal environment.<sup>40,41</sup> Differences in the abundance and hydrolysis efficiency of NAPE oleoyl (18:1) versus linoleoyl (18:2) NAPE acyl chains,<sup>40</sup> as well as luminal processes such as re-esterification or microbial metabolism, may have contributed to modulating the observed NAE profile.<sup>41</sup> Moreover, we observed considerable interindividual variability in postprandial NAE levels, which is likely explained by physiological differences among participants, including the lipid digestion efficiency, intestinal motility and uptake, and differences in the intestinal microbiota. Such variability is well recognized in studies of lipid-derived mediators and is consistent with the heterogeneous physiology of the small intestine.<sup>42–44</sup>

#### 4.2. Ileal microbiome may influence dietary NAE metabolism

Results of the current study also suggest *in vivo* that the ileal microbiome may influence the intestinal levels of ECs and NAEs, due to microbial metabolism.

Although less densely populated than the colon, the small intestine harbors a metabolically active microbiota<sup>43</sup> that may contribute to the early formation and transformation of luminal NAEs from dietary precursors, including NAPEs, prior to the extensive microbial processing of dietary lipids and the subsequent formation of metabolites, including NAEs, in the colon.<sup>41,45,46</sup>

The ileal microbiome is a topic still underexplored in scientific literature, and due to the difficulties in sampling and ethical reasons, the knowledge is limited to subjects suffering from some intestinal diseases.<sup>47</sup> Compared with the colon microbiome, the small intestine is usually dominated by facultative anaerobes (*e.g.*, Bacillota, such as *Streptococcaceae*, *Lactobacillaceae* and *Veillonellaceae*, and Pseudomonadota), due to higher oxygen concentration, while Bacteroidota abundance is usually limited. The findings of this study confirmed *Streptococcus*, *Haemophilus* and *Veillonella* as members of a core ileal microbiome, present with varying abundances in most of the subjects.<sup>47</sup> However, high inter-personal variability in composition was also observed. Indeed, the ileum microbiome is subject to higher fluctuations in composition and abundance compared with the colon, and dynamic changes of the microbiome are reported during fasting and fed states.<sup>43,48</sup>

Interestingly, some of the participants of this study were characterized by a particularly high abundance of *Escherichia coli* and *Klebsiella*. The overgrowth of these taxa has been previously associated with inflammation, an impaired intestinal barrier and malabsorption.<sup>49</sup>

Recent *in vitro* studies have demonstrated a clear role of the colonic microbiota in the formation and transformation of ECs and NAEs in the intestinal lumen. Specifically, changes in EC and NAE profiles were observed in the simulated chyme



undergoing fecal fermentation, with certain metabolites decreasing and others emerging over time.<sup>45</sup> Similarly, Roussel *et al.* (2024) showed that human fecal microbiota can both generate and hydrolyze specific NAEs (such as *N*-stearidonoyl ethanolamine, LEA, and PEA) from dietary polyunsaturated NAPE substrates in the Mucosal Simulator of the Human Intestinal Microbial Ecosystem (M-SHIME) model.<sup>41</sup> These findings align with the known enzymatic capacities of various intestinal microorganisms, including phospholipase, amidase, and acyltransferase activities, which can hydrolyse or remodel NAPEs and thereby influence the profile of NAEs derived from dietary precursors.

*In vivo* evidence further supports a microbiota-dependent contribution to luminal NAE and broader endocannabinoidome dynamics. In mice, OEA was undetectable in the cecal contents of microbiota-depleted animals, but was readily produced in wild-type mice receiving an intra-cecal oleic acid infusion.<sup>46</sup> Moreover, intraperitoneal OEA administration has been shown to modulate gut microbial composition while increasing hepatic OEA levels and exerting anti-inflammatory effects,<sup>50</sup> suggesting bidirectional interactions between circulating NAEs and the gut microbiota mediated through host metabolic pathways.

In humans, a cross-sectional and dietary-intervention study demonstrated that the fecal endocannabinoidome profile is shaped predominantly by gut microbiota composition and dietary intakes, rather than by circulating mediator levels.<sup>51</sup> Reinforcing this causal link between diet, microbial ecology, and endocannabinoidome signaling, findings from an 8-week Mediterranean diet intervention showed that a personalized dietary shift elicited coordinated changes in circulating NAE profiles alongside increased abundance of key microbial taxa such as *Akkermansia muciniphila*, independent of weight loss.<sup>27</sup>

In the present study, the analysis of the microbiome in the basal ileal fluids and the postprandial levels of ECs and NAEs in the ileal fluids, allowed the identification of clusters of microbial taxa associated with the monitored lipids, thus suggesting that the microbial communities present in the small intestine may influence the metabolism of ECs and NAEs *in vivo*. Interestingly, OEA, PEA, LEA, and AEA formed interconnected clusters, indicating overlap in their microbial associations. For instance, *Lactobacillus* and *Streptococcus* spp. interact with several lipid mediators, predominantly showing negative associations with the postprandial formation of OEA and LEA in the ileum. Indeed, these genera, together with *Escherichia coli* (also abundant in the study participants), have been reported to be involved in lipid metabolism in the small intestine.<sup>52,53</sup> OEA and LEA belong to the same cluster and are primarily associated with a broad range of taxa from the Fusobacteriota phylum, while negatively correlating with Bacillota. Bacillota, including certain species of *Enterococcus*, *Streptococcus*, and *Clostridium*, harbor ethanolamine utilization genes, which may facilitate the metabolism of dietary NAEs as a carbon and/or nitrogen source.<sup>54,55</sup> Additionally, several strains within *Lactobacillus*, *Bifidobacteria*, *Streptococcus*, *Clostridium* and *Enterococcus* are known for their capacity to metabolize ethanolamine.<sup>54</sup>

However, further studies integrating metagenomic functional profiling with targeted metabolomics are needed to better elucidate the microbial pathways involved in NAE and EC metabolism.

#### 4.3. NAE levels in the intestine are sufficient to elicit GPR119

It is worth noting that the postprandial concentrations of NAEs in the ileal fluids were found in quantities exceeding their reported EC<sub>50</sub> for GPR119.<sup>5,10,18</sup> GPR119 is a receptor for monoacylglycerols, lysophosphatidylcholine, and fatty acid ethanolamides.<sup>56</sup> It is expressed on enteroendocrine cells and, by sensing lipids, produces incretins, such as GLP-1 and GIP after a meal.<sup>1,57</sup> Therefore, results of this study suggest that OEA and LEA, delivered from the food, by activating GPR119 likely contributed to the increase in C-peptide, insulin, and particularly GIP after the HNM compared to the LNM.

On the other hand, GLP-1 was undetectable in the present study. In a previous study involving ileostomy subjects, the postprandial GLP-1 response was stimulated by the presence of lipids in the duodenum, occurring approximately 150 min after ingestion of a high-fat meal (comprising 78% of energy from fats). This response was significantly lower in ileostomates compared to healthy subjects.<sup>58</sup> The discrepancy between GIP and GLP-1 responses is likely attributable to differences in the distribution of enteroendocrine cells. K cells, which secrete GIP, are most densely located in the duodenum, whereas L cells, responsible for GLP-1 secretion, show a gradient of increasing density along the GI tract, with the highest concentrations found in the distal ileum and colon.<sup>59</sup> This anatomical distribution supports the interpretation that GLP-1 release predominantly occurs in the distal gut, explaining both the diminished GLP-1 response in ileostomy subjects and the undetectable levels observed in the present study. The lack of observable differences in glycemic responses between meals suggests that the hormonal fluctuations measured were insufficient to affect glucose regulations at the monitored time points. Moreover, the postprandial increase in GIP and the simultaneous reduction in ghrelin following HNM consumption did not appear to influence subsequent energy intake during the buffet meal. The 8-hour fasting period imposed after breakfast, prior to the buffet, may have contributed to this result, as behavioral and cognitive factors could potentially override hormonal cues.<sup>60</sup>

#### 4.4. NAE levels in the bloodstream do not reflect intestinal levels

An additional finding of this study was the lack of significant differences in the postprandial plasma concentrations of NAEs and ECs between the two meals, as well as the absence of correlation between ileal and plasma concentrations. These observations suggest two possible hypotheses. The first hypothesis is that dietary NAEs may not exert a direct effect on systemic circulation under the conditions of this study, possibly due to predominant intracellular physiological mechanisms (*e.g.*, metabolic processing, receptor binding, or enzymatic degradation) that limit their release into the bloodstream.<sup>38,39,52,61</sup>



The second hypothesis is that absorbed NAEs are rapidly utilized by peripheral tissues, which could account for the observed postprandial decline in plasma concentrations 2 hours after breakfast. This interpretation is consistent with previous findings<sup>1</sup> and suggests that this rapid clearance may mask other ongoing physiological phenomena. In both scenarios, future feeding studies using isotopically labeled (e.g., <sup>13</sup>C or <sup>2</sup>H) NAEs would provide a more detailed understanding of their metabolic fate, from ingestion through intestinal absorption to systemic distribution.

However, the lack of correlation between ileal and plasma endocannabinoid mediators observed in our study is consistent with previous findings showing a similar disconnect between fecal or ileal tissue and circulating endocannabinoid profiles in humans.<sup>51,62</sup>

This study has four main limitations. First, due to the 2-hour intervals used for glycemia monitoring, potential differences in postprandial glycemic responses between the two breakfasts, especially within the initial 2 hours after consumption, may have been missed. However, this design allowed for synchronized collection of biological samples, minimizing participant burden and improving data consistency. Second, although total carbohydrate content was comparable between the two meals, the proportion of sugars differed. However, postprandial glycemia was not significantly different following the two meals. Therefore, while we cannot entirely exclude the possibility that differences in sugar composition may have slightly influenced gastrointestinal hormone responses, any such effect is likely minimal. Third, the 8-hour interval between breakfast and buffet may have been too long to effectively evaluate the impact of breakfast composition on subsequent energy intake. After such an extended fasting period, behavioral and cognitive drivers of hunger may override hormonal or metabolic signals. Nevertheless, this extended interval enabled a more accurate assessment of the kinetics of dietary NAE bioaccessibility in ileal fluids, without interference from an additional meal. Finally, the ileostomy-based study may limit the generalizability of the findings to individuals with an intact gastrointestinal tract, although it allows direct assessment of early, diet-driven NAE bioaccessibility in the small intestine; consequently, subsequent colonic microbial transformations potentially contributing to fecal and systemic NAE pools are not captured.

## 5. Conclusions

In conclusion, the present study demonstrates that dietary NAEs are released into the small intestinal lumen and recovered at higher levels following a high-NAE meal, confirming their bioaccessibility under physiological conditions. However, no corresponding increase in plasma NAEs was observed, indicating that circulating levels do not reflect local intestinal concentrations. Notably, the higher ileal recovery of OEA and LEA was accompanied by increased postprandial GIP, insulin, and C-peptide responses, supporting a link between luminal NAEs

and gut-derived hormonal signaling. Furthermore, baseline microbial composition was associated with variability in postprandial NAE responses, suggesting a modulatory role of the ileal microbiome. Overall, these findings highlight the importance of the small intestinal environment in shaping the local metabolism and physiological effects of dietary NAEs.

## Author contributions

Paola Vitaglione: conceptualization, project administration, and writing – review & editing; Chris R Gill: supervision, resources, and writing – review & editing; Silvia Tagliamonte: methodology, investigation, formal analysis, writing – original draft, and data curation; Danilo Ercolini: writing – review & editing; Francesca De Filippis: formal analysis and writing – review & editing; Holly R Neill: investigation and data curation; L. Kirsty Pourshahidi: supervision; Brian Óg Murphy: investigation and data curation; Koos Natalia: data curation; Brendan Curran: investigation and data curation; Murray Nicole: investigation and data curation; Slevin Mary: supervision; Sara Dobani: data curation; and Massimiliano Fontana: data curation.

## Conflicts of interest

There are no conflicts to declare.

## Data availability

The raw sequence reads generated in this study have been deposited in the Sequence Read Archive (SRA) of the NCBI under accession number PRJNA1238284.

Supplementary information (SI) is available. See DOI: <https://doi.org/10.1039/d5fo03328d>.

## References

- 1 R. F. Witkamp, The role of fatty acids and their endocannabinoid-like derivatives in the molecular regulation of appetite, *Mol. Aspects Med.*, 2018, **64**, 45–67.
- 2 Y. Wang, M. Balvers, H. Hendriks, T. Wilpshaar, T. van Heek, R. Witkamp and J. Meijerink, Docosahexaenoyl serotonin emerges as most potent inhibitor of IL-17 and CCL-20 released by blood mononuclear cells from a series of N-acyl serotonins identified in human intestinal tissue, *Biochim. Biophys. Acta, Mol. Cell Biol. Lipids*, 2017, **1862**, 823–831.
- 3 C. J. Hillard, Circulating endocannabinoids: from whence do they come and where are they going?, *Neuropsychopharmacology*, 2018, **43**, 155–172.
- 4 H. S. Hansen and T. A. Diep, N-acylethanolamines, anandamide and food intake, *Biochem. Pharmacol.*, 2009, **78**, 553–560.



- 5 J. Fu, S. Gaetani, F. Oveisi, J. L. Verme, A. Serrano, F. R. de Fonseca, A. Rosengarth, H. Luecke, B. Di Giacomo, G. Tarzia and D. Piomelli, Oleylethanolamide regulates feeding and body weight through activation of the nuclear receptor PPAR- $\alpha$ , *Nature*, 2003, **425**, 90.
- 6 J. Lo Verme, S. Gaetani, J. Fu, F. Oveisi, K. Burton and D. Piomelli, Regulation of food intake by oleoylethanolamide, *Cell. Mol. Life Sci.*, 2005, **62**, 708.
- 7 A. V. Contreras, N. Torres and A. R. Tovar, PPAR- $\alpha$  as a key nutritional and environmental sensor for metabolic adaptation, *Adv. Nutr.*, 2013, **4**, 439–452.
- 8 A. Ostadrahimi, Y. Khajebishak, F. Moradi and L. Payahoo, The effect of oleoylethanolamide supplementation on lipid profile, fasting blood sugar and dietary habits in obese people: a randomized double-blind placebo-control trial, *BMC Endocr. Disord.*, 2024, **24**, 1–9.
- 9 A. Sarro-Ramírez, D. Sánchez-López, A. Tejeda-Padrón, C. Frías, J. Zaldívar-Rae and E. Murillo-Rodríguez, Brain molecules and appetite: the case of oleoylethanolamide, *Cent. Nerv. Syst. Agents Med. Chem.*, 2013, **13**, 88–91.
- 10 K. B. Hansen, M. M. Rosenkilde, F. K. Knop, N. Wellner, T. A. Diep, J. F. Rehfeld, U. B. Andersen, J. J. Holst and H. S. Hansen, 2-Oleoyl glycerol is a GPR119 agonist and signals GLP-1 release in humans, *J. Clin. Endocrinol. Metab.*, 2011, **96**, E1409–E1417.
- 11 D. Barbaro, A. Menasci, B. Baldini, C. Pasquini and P. Lapi, The dietary supplement EGCG: NOPE (N-oleyl-phosphatidylethanolamine and epigallocatechin-3-gallate formula) helps patients to follow a flexible dietary regimen and induces loss of weight in patients who had previously experienced no response to other weight loss intervention, *J. Obes. Weight Loss Ther.*, 2011, **1**, 105.
- 12 M. Rondanelli, A. Opizzi, S. B. Solerte, R. Trotti, C. Klersy and R. Cazzola, Administration of a dietary supplement (N-oleyl-phosphatidylethanolamine and epigallocatechin-3-gallate formula) enhances compliance with diet in healthy overweight subjects: a randomized controlled trial, *Br. J. Nutr.*, 2008, **101**, 457–464.
- 13 L. Payahoo, Y. Khajebishak, M. R. Alivand, H. Soleimanzade, S. Alipour, A. Barzegari and A. Ostadrahimi, Investigation the effect of oleoylethanolamide supplementation on the abundance of Akkermansia muciniphila bacterium and the dietary intakes in people with obesity: a randomized clinical trial, *Appetite*, 2019, **141**, 104301.
- 14 F. R. De Fonseca, M. Navarro, R. Gomez, L. Escuredo, F. Nava, J. Fu, E. Murillo-Rodríguez, A. Giuffrida, J. LoVerme, S. Gaetani, S. Kathuria, C. Gall and D. Piomelli, An anorexic lipid mediator regulated by feeding, *Nature*, 2001, **414**, 209.
- 15 I. Mennella, M. Savarese, R. Ferracane, R. Sacchi and P. Vitaglione, Oleic acid content of a meal promotes oleoylethanolamide response and reduces subsequent energy intake in humans, *Food Funct.*, 2015, **6**, 203–209.
- 16 L. De Luca, R. Ferracane, N. C. Ramírez and P. Vitaglione, N-acylphosphatidylethanolamines and N-acylethanolamines increase in saliva upon food mastication: the influence of the individual nutritional status and fat type in food, *Food Funct.*, 2020, **11**, 3382–3392.
- 17 S. Tagliamonte, C. I. Gill, L. K. Pourshahidi, M. M. Slevin, R. K. Price, R. Ferracane, R. Lawther, G. O'Connor and P. Vitaglione, Endocannabinoids, endocannabinoid-like molecules and their precursors in human small intestinal lumen and plasma: does diet affect them?, *Eur. J. Nutr.*, 2021, **60**, 2203–2215.
- 18 H. A. Overton, A. J. Babbs, S. M. Doel, M. C. Fyfe, L. S. Gardner, G. Griffin, H. C. Jackson, M. J. Procter, C. M. Rasamison, M. Tang-Christensen, P. S. Widdowson, G. M. Williams and C. Reynet, Deorphanization of a G protein-coupled receptor for oleoylethanolamide and its use in the discovery of small-molecule hypophagic agents, *Cell Metab.*, 2006, **3**, 167–175.
- 19 Z. Chen, L. Guo, Y. Zhang, R. L. Walzem, J. S. Pendergast, R. L. Printz, L. C. Morris, E. Matafonova, X. Stien, L. Kang, D. Coulon, O. P. McGuinness, K. D. Niswender and S. S. Davies, Incorporation of therapeutically modified bacteria into gut microbiota inhibits obesity, *J. Clin. Invest.*, 2014, **124**, 3391–3406.
- 20 Z. Chen, Y. Zhang, L. Guo, N. Dosoky, L. de Ferra, S. Peters, K. D. Niswender and S. S. Davies, Leptogenic effects of NAPE require activity of NAPE-hydrolyzing phospholipase D, *J. Lipid Res.*, 2017, **58**, 1624–1635.
- 21 A. J. Stunkard and S. Messick, The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger, *J. Psychosom. Res.*, 1985, **29**, 71–83.
- 22 E. R. Didie, *The Power of Food Scale (PFS): development and theoretical evaluation of a self-report measure of the perceived influence of food*, Drexel University, 2003.
- 23 P. F. Lovibond and S. H. Lovibond, The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories, *Behav. Res. Ther.*, 1995, **33**, 335–343.
- 24 I. H. Franken, E. Rassin and P. Muris, The assessment of anhedonia in clinical and non-clinical populations: further validation of the Snaith–Hamilton Pleasure Scale (SHAPS), *J. Affective Disord.*, 2007, **99**, 83–89.
- 25 J. E. Ware, M. Kosinski and S. D. Keller, A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity, *Med. Care*, 1996, **34**, 220–233.
- 26 S. M. Green, H. J. Delargy, D. Joanes and J. E. Blundell, A satiety quotient: a formulation to assess the satiating effect of food, *Appetite*, 1997, **29**, 291–304.
- 27 S. Tagliamonte, M. Laiola, R. Ferracane, M. Vitale, M. A. Gallo, V. Meslier, N. Pons, D. Ercolini and P. Vitaglione, Mediterranean diet consumption affects the endocannabinoid system in overweight and obese subjects: possible links with gut microbiome, insulin resistance and inflammation, *Eur. J. Nutr.*, 2021, **60**, 3703–3716.
- 28 F. De Filippis, L. Paparo, R. Nocerino, G. Della Gatta, L. Carucci, R. Russo, E. Pasolli, D. Ercolini and R. Berni Canani, Specific gut microbiome signatures and the associ-



- ated pro-inflammatory functions are linked to pediatric allergy and acquisition of immune tolerance, *Nat. Commun.*, 2021, **12**, 5958.
- 29 R. Schmieder and R. Edwards, Quality control and preprocessing of metagenomic datasets, *Bioinformatics*, 2011, **27**, 863–864.
- 30 A. Blanco-Míguez, F. Beghini, F. Cumbo, L. J. McIver, K. N. Thompson, M. Zolfo, P. Manghi, L. Dubois, K. D. Huang, A. M. Thomas, W. A. Nickols, G. Piccinno, E. Piperni, M. Punčochář, M. Valles-Colomer, A. Tett, F. Giordano, R. Davies, J. Wolf, S. E. Berry, T. D. Spector, E. A. Franzosa, E. Pasolli, F. Asnicar, C. Huttenhower and N. Segata, Extending and improving metagenomic taxonomic profiling with uncharacterized species using MetaPhlan 4, *Nat. Biotechnol.*, 2023, **41**, 1633–1644.
- 31 R. B. Lumaga, S. Tagliamonte, T. De Rosa, V. Valentino, D. Ercolini and P. Vitaglione, Consumption of a sourdough-leavened croissant enriched with a blend of fibers influences fasting blood glucose in a randomized controlled trial in healthy subjects, *J. Nutr.*, 2024, **154**, 2976–2987.
- 32 P. Monteleone, F. Piscitelli, P. Scognamiglio, A. M. Monteleone, B. Canestrelli, V. Di Marzo and M. Maj, Hedonic eating is associated with increased peripheral levels of ghrelin and the endocannabinoid 2-arachidonoylglycerol in healthy humans: a pilot study, *J. Clin. Endocrinol. Metab.*, 2012, **97**, E917–E924.
- 33 B. Gatta-Cherifi, I. Matias, M. Vallee, A. Tabarin, G. Marsicano, P. V. Piazza and D. Cota, Simultaneous postprandial deregulation of the orexigenic endocannabinoid anandamide and the anorexigenic peptide YY in obesity, *Int. J. Obes.*, 2012, **36**, 880.
- 34 F. Brouns, I. Bjorck, K. N. Frayn, A. L. Gibbs, V. Lang, G. Slama and T. M. S. Wolever, Glycaemic index methodology, *Nutr. Res. Rev.*, 2005, **18**, 145–171.
- 35 L. De Luca, R. Ferracane, N. C. Ramírez and P. Vitaglione, N-acylphosphatidylethanolamines and N-acylethanolamines increase in saliva upon food mastication: the influence of the individual nutritional status and fat type in food, *Food Funct.*, 2020, **11**, 3382–3392.
- 36 X. Kong, R. Ferracane, L. De Luca and P. Vitaglione, Salivary concentration of N-acylethanolamines upon food mastication and after meal consumption: influence of food dietary fiber, *Food Res. Int.*, 2016, **89**, 186–193.
- 37 I. Mennella, R. Di Monaco, A. Balazy, R. Ferracane, N. A. Miele, S. Cavella and P. Vitaglione, Salivary endocannabinoids and N-acylethanolamines upon mastication of a semisolid food: implications in fat taste, appetite and food liking, *Food Funct.*, 2018, **9**, 476–484.
- 38 G. Boudry, I. Mennella, O. Menard, R. Janvier, I. Nogret, A. Madadlou, A. Cahu, L. Le Normand, E. Bobillier-Chaumont, R. Ferracane, P. Vitaglione, D. Dupont and D. Val-Laillet, Development of a functional dairy snack containing oleoylethanolamide that reduces food intake in normal-weight and obese minipigs, *J. Funct. Foods*, 2023, **111**, 105916.
- 39 C. M. Weaver and D. I. Givens, Overview: the food matrix and its role in the diet, *Crit. Rev. Food Sci. Nutr.*, 2025, **65**(1), 1–18.
- 40 E. Leishman, K. Mackie, S. Luquet and H. B. Bradshaw, Lipidomics profile of a NAPE-PLD KO mouse provides evidence of a broader role of this enzyme in lipid metabolism in the brain, *Biochim. Biophys. Acta, Mol. Cell Biol. Lipids*, 2016, **1861**, 491–500.
- 41 C. Roussel, M. Sola, J. Lessard-Lord, N. Nallabelli, P. Génereux, C. Cavestri, O. A. Wallen, R. Villano, F. Raymond, N. Flamand, C. Silvestri and V. Di Marzo, Human gut microbiota and their production of endocannabinoid-like mediators are directly affected by a dietary oil, *Gut Microbes*, 2024, **16**, 2335879.
- 42 I. Mennella, M. Savarese, R. Ferracane, R. Sacchi and P. Vitaglione, Oleic acid content of a meal promotes oleoylethanolamide response and reduces subsequent energy intake in humans, *Food Funct.*, 2015, **6**, 203–209.
- 43 B. Yilmaz, T. Fuhrer, D. Morgenthaler, N. Krupka, D. Wang, D. Spari, D. Candidnas, B. Misselwitz, G. Beldi, U. Sauer and A. J. Macpherson, Plasticity of the adult human small intestinal stoma microbiota, *Cell Host Microbe*, 2022, **30**, 1773–1787.
- 44 K. J. Bowen, P. M. Kris-Etherton and S. G. West, Dietary fatty acids and postprandial lipid metabolism: variability and clinical relevance, *Curr. Opin. Lipidol.*, 2018, **29**, 46–52.
- 45 M. Kråkström, A. M. Dickens, M. A. Alves, S. D. Forssten, A. C. Ouwehand, T. Hyötyläinen, M. Orešič and S. Lamichhane, Dynamics of the lipidome in a colon simulator, *Metabolites*, 2023, **13**, 355.
- 46 M. Igarashi, T. Hayakawa, H. Tanabe, K. Watanabe, A. Nishida and I. Kimura, Intestinal GPR119 activation by microbiota-derived metabolites impacts feeding behavior and energy metabolism, *Mol. Metab.*, 2023, **67**, 101649.
- 47 S. Yersin and P. Vonaesch, Small intestinal microbiota: from taxonomic composition to metabolism, *Trends Microbiol.*, 2024, **32**, p970–p983.
- 48 A. C. D. Machado, S. D. Brown, A. Lingaraju, V. Sivaganesh, C. Martino, A. Chaix, P. Zhao, A. F. M. Pinto, M. W. Chang, R. A. Richter, A. Saghatelian, A. R. Saltiel, R. Knight, S. Panda and A. Zarrinpar, Diet and feeding pattern modulate diurnal dynamics of the ileal microbiome and transcriptome, *Cell Rep.*, 2022, **40**, 111111.
- 49 J. M. Collard, L. Andrianonimiadana, A. Habib, M. Rakotondrainipiana, P. Andriantsalama, R. Randriamparany, M. A. N. Rabenandrasana, F. X. Weill, N. Sauvonnet, R. V. Randremanana, V. Guillemot, P. Vonaesch and P. J. Sansonetti, High prevalence of small intestine bacteria overgrowth and asymptomatic carriage of enteric pathogens in stunted children in Antananarivo, Madagascar, *PLoS Neglected Trop. Dis.*, 2022, **16**, e0009849.
- 50 F. Cimmino, C. Silvestri, G. Trinchese, L. Petrella, G. Cavaliere, C. Fogliano, F. Piscitelli, L. Cristino, B. Avallone, S. Banni, J. Sihang, V. Di Marzo and M. P. Mollica, Anti-obesity effects of Oleoylethanolamide:



- Modulation of mitochondrial bioenergetics, endocannabinoidome and gut microbiome, *Biomed. Pharmacother.*, 2025, **188**, 118201.
- 51 S. Castonguay-Paradis, S. Lacroix, G. Rochefort, L. Parent, J. Perron, C. Martin, B. Lamarque, F. Raymond, N. Flamand, V. Di Marzo and A. Veilleux, Dietary fatty acid intake and gut microbiota determine circulating endocannabinoidome signaling beyond the effect of body fat, *Sci. Rep.*, 2020, **10**, 15975.
- 52 J. R. Araújo, A. Tazi, O. Burlen-Defranoux, S. Vichier-Guerre, G. Nigro, H. Licandro, S. Demignot and P. J. Sansonetti, Fermentation products of commensal bacteria alter enterocyte lipid metabolism, *Cell Host Microbe*, 2020, **27**, 358–372.
- 53 A. Tazi, J. R. Araújo, C. Mulet, E. T. Arena, G. Nigro, T. Pedron and P. J. Sansonetti, Disentangling host–microbiota regulation of lipid secretion by enterocytes: insights from commensals *Lactobacillus paracasei* and *Escherichia coli*, *mBio*, 2018, **9**, e01493-18.
- 54 K. G. Kaval and D. A. Garsin, Ethanolamine utilization in bacteria, *mBio*, 2018, **9**, e00066-18.
- 55 N. Fornelos, E. A. Franzosa, J. Bishai, J. W. Annand, A. Oka, J. Lloyd-Price, T. D. Arthur, A. Garner, J. Avila-Pacheco, H. J. Haiser, A. C. Tolonen, J. A. Porter, C. B. Clish, R. B. Sartor, C. Huttenhower, H. Vlamakis and R. J. Xavier, Growth effects of N-acyl ethanolamines on gut bacteria reflect altered bacterial abundances in inflammatory bowel disease, *Nat. Microbiol.*, 2020, **5**, 486–497.
- 56 H. S. Hansen, M. M. Rosenkilde, J. J. Holst and T. W. Schwartz, GPR119 as a fat sensor, *Trends Pharmacol. Sci.*, 2012, **33**, 374–381.
- 57 A. Psichas, F. Reimann and F. M. Gribble, Gut chemosensing mechanisms, *J. Clin. Invest.*, 2015, **125**, 908–917.
- 58 M. D. Robertson, G. Livesey, L. M. Morgan, S. M. Hampton and J. C. Mathers, The influence of the colon on postprandial glucagon-like peptide 1 (7–36) amide concentration in man, *J. Endocrinol.*, 1999, **161**, 25–32.
- 59 G. Tolhurst, F. Reimann and F. M. Gribble, Nutritional regulation of glucagon-like peptide-1 secretion, *J. Physiol.*, 2009, **587**, 27–32.
- 60 A. P. Goldstone, C. G. Precht de Hernandez, J. D. Beaver, K. Muhammed, C. Croese, G. Bell, G. Durighel, E. Hughes, A. D. Waldman, G. Frost and J. D. Bell, Fasting biases brain reward systems towards high-calorie foods, *Eur. J. Neurosci.*, 2009, **30**, 1625–1635.
- 61 F. Fanelli, M. Mezzullo, A. Repaci, I. Belluomo, D. I. Gasparini, G. Di Dalmazi, M. Mastroberto, V. Vicennati, A. Gambineri, A. M. Morselli-Labate, R. Pasquali and U. Pagotto, Profiling plasma N-acyl ethanolamine levels and their ratios as a biomarker of obesity and dysmetabolism, *Mol. Metab.*, 2018, **14**, 82–94.
- 62 V. Rakotoarivelo, B. Allam-Ndoul, C. Martin, L. Biertho, V. Di Marzo, N. Flamand and A. Veilleux, Investigating the alterations of endocannabinoidome signaling in the human small intestine in the context of obesity and type 2 diabetes, *Heliyon*, 2024, **10**, e26968.

