

Cite this: *Food Funct.*, 2026, **17**, 2430

Antioxidant pine nut protein hydrolysate as a therapeutic oral immunotherapy agent in a murine model of pine nut allergy

 Nourelimane Benzitoune,^{a,b,c,d} David Menchén-Martínez,^a Daniel Lozano-Ojalvo,^a Nabil Kadri,^c Rosina López-Fandiño,^b ^a Elena Molina^a and Sara Benedé ^{*a}

Oral immunotherapy (OIT) is one of the most promising treatments for food allergy, although no protocols have yet been described for pine nut allergy. We aim to produce a pine nut protein hydrolysate to be used as a treatment agent in OIT. Hydrolysates were produced using three proteases (pepsin, trypsin, and Alcalase) alone or in combination and their antioxidant activity, antigenicity, and immunomodulatory properties were determined *in vitro*. Pine nut-sensitized C57BL/6 mice were subjected to OIT with intact and hydrolysed pine nut protein. The trypsin hydrolysate showed higher antioxidant activity and lower IgE binding capacity than the intact protein, and it was able to desensitize mice in a similar way to the unhydrolyzed protein. Therefore, it could be regarded as a good candidate as a therapeutic agent in OIT on the grounds of its effectiveness in desensitizing allergic mice and its improved antioxidant and immunoreactive properties with respect to intact pine nut protein. Oral immunotherapy is one of the most promising treatments for food allergy, although no protocols have yet been described for pine nut allergy. To identify pine nut protein hydrolysates with potential for OIT, we applied enzymatic hydrolysis using food-grade enzymes and a two-step selection strategy. Hydrolysates were first screened for *in vitro* antioxidant activity and then assessed for their ability to modulate allergen-specific T-cell responses *ex vivo*. The trypsin-derived hydrolysate showed the most promising profile, leading to further evaluation of its IgE-binding properties and therapeutic potential in pine nut-sensitized C57BL/6 mice subjected to OIT with intact and hydrolysed pine nut protein. Mice treated with the trypsin hydrolysate showed reduced specific IgE and IgG1 levels and were protected against oral pine nut-induced anaphylaxis, similar to those treated with intact pine nut protein. Due to its ability to desensitize allergic mice, combined with lower IgE binding and higher antioxidant activity, the trypsin-hydrolysed pine nut protein appears to be a promising candidate for OIT.

Received 4th November 2025,
Accepted 1st February 2026

DOI: 10.1039/d5fo04650e

rsc.li/food-function

Introduction

Food allergy has emerged as a pressing global public health concern, displaying a steadily increasing prevalence over the past few decades. It encompasses a range of immune-mediated adverse responses triggered by specific food proteins, affecting up to 10% of certain populations, with a predilection for industrialized regions and emerging economies.¹

Nine primary allergenic foods have been identified, including tree nuts, which are known for causing potentially fatal anaphylactic reactions, often triggered by IgE-mediated responses. Tree nut allergies are closely linked to specific protein families, such as 2S albumins, vicilins, legumins, and nonspecific lipid transfer proteins.^{2,3} Allergic reactions to pine nuts are relatively less common than to other nuts, such as walnuts, cashews, hazelnuts or almonds, although they are particularly alarming due to the significant frequency of severe anaphylactic responses (approximately 80%) triggered by even small quantities of pine nuts.^{4,5} A high proportion (70%) of pine nut allergic patients are monosensitized^{5–7} and there is limited cross-reactivity between pine nuts and more commonly consumed nuts, primarily due to the evolutionary differences between gymnosperms (pine nuts) and angiosperms (other nuts), which contribute to the distinct characteristics of pine nut allergy.⁴

Pine nut allergy management is still limited to strict dietary avoidance, basically because allergen-specific immunotherapy using the whole allergenic food is often accompanied by

^aInstituto de Investigación en Ciencias de la Alimentación (CIAL), CSIC-UAM, Nicolás Cabrera, 9, 28049 Madrid, Spain. E-mail: s.benede@csic.es

^bLaboratoire de Gestion et Valorisation des Ressources Naturelles et Assurance Qualité (LGVRNAQ), Faculté des Sciences de la Nature et de la Vie et des Sciences de la Terre, Université de Bouira, 10000 Bouira, Algérie

^cDépartement des Sciences Biologiques, Faculté des Sciences de la Nature et de la Vie et des Sciences de la Terre, Université de Bouira, 10000 Bouira, Algérie

^dDépartement de Biologie, Faculté des Sciences de la Nature et de la Vie et des Sciences de la Terre, Université de Khemis Miliana, 44000 Ain defla, Algérie



severe side effects.⁴ Oral immunotherapy (OIT) has been implemented mainly against peanut, egg, and milk allergies to induce immunomodulation and protect against accidental allergen exposure. However, concerns about its safety and ability to establish permanent tolerance persist.^{8,9} Alternative food allergy treatments have emerged, such as hydrolysed protein-based OIT, which offers the advantage of a reduced risk of adverse reactions compared to intact allergen-based OIT. In addition, the peptides resulting from the hydrolysis of allergens may have immunomodulatory properties maintaining and reinforcing barrier function and promoting immune homeostasis through their action on the cells of the innate and adaptive immune system.^{10–12} Peptides derived from food proteins may also exhibit antioxidant properties,¹³ which could have an additional beneficial effect during immunotherapy treatments as there is ample evidence that high levels of oxidative stress may play an important role in the pathogenesis of allergic disorders.^{14,15} An imbalance between reactive oxygen species (ROS) production and antioxidant defence during allergen encounter may favour a T helper 2 cell immune response.¹⁶ In fact, certain clinical trials suggest that specific antioxidants from diet or vitamin supplements might improve asthma control or lung function in asthmatic children and adults.^{17,18}

To the best of our knowledge, there are no previous reports on the use of pine nut protein hydrolysates in OIT. To identify hydrolysates with optimal properties, after performing enzymatic hydrolysis of pine nut proteins using food-grade enzymes, either individually or sequentially, a two-step selection strategy was applied: first, hydrolysates were evaluated for antioxidant activity using multiple *in vitro* methods as the primary criterion; second, their ability to down-regulate *ex vivo* T-cell responses induced by the pine nut allergen was considered as a secondary criterion. Among the tested hydrolysates, the trypsin-derived hydrolysate exhibited the most favourable profile. After assessing its IgE-binding properties with sera from pine nut-allergic patients, we further investigated its therapeutic potential *in vivo* using the C57BL/6 mouse model.

Materials and methods

Preparation of protein isolate and protein hydrolysates

Pine nuts (*Pinus pinea* L), skinned and shelled, were bought from a local market (Blida, Algeria). To prepare pine nut protein isolate (PI), clean and dried pine nuts were ground and defatted by Soxhlet extraction using *n*-hexane. Then, a solution of defatted flour in MilliQ water at 125 mg mL⁻¹ was prepared and centrifuged at 3000g for 10 min and the supernatant was freeze-dried.

Pine nut protein hydrolysates were obtained using pepsin (P; Sigma-Aldrich, St. Louis, MO, USA) at an enzyme-to-substrate (E/S) ratio of 128.034 U mg⁻¹ protein in 200 mM KCl/HCl buffer, pH 2.0, at 37 °C for 4 h; trypsin (T; Sigma-Aldrich) at an E/S ratio of 0.3377 U mg⁻¹ protein in 50 mM phosphate

buffer, pH 8.0, at 37 °C for 4 h; and Alcalase (A; Novozyme, Bagsvaerd, Denmark) at an E/S ratio of 62.205 U mg⁻¹ protein in 50 mM phosphate buffer, pH 8.0, at 55 °C for 3 h. Hydrolysis was also carried out using combinations of pepsin/trypsin (PT), pepsin/Alcalase (PA), trypsin/Alcalase (TA), and pepsin/trypsin/Alcalase (PTA). Sequential hydrolysis was initiated by adding the first enzyme and allowing it to hydrolyse the protein under the conditions mentioned above. Then, the pH was adjusted with 6 M NaOH, and the next enzyme was added and incubated for an additional 2 h at the indicated temperature. Upon completion of the hydrolysis, the reaction mixtures were heated at 85 °C for 15 min to inactivate enzymes and centrifuged at 8000g for 15 min. Supernatants were collected and freeze-dried for further analysis.

The protein content of PI and its hydrolysates was quantified by determining their nitrogen percentage by elemental analysis. The lipopolysaccharide present in the samples was removed by using Pierce™ high capacity endotoxin removal spin columns (Thermo Fisher Scientific, Waltham, MA, USA) and its absence was confirmed by using the transfected cell line THP1-XBlue™, stably expressing an NF-κB/AP-1-inducible secreted alkaline phosphatase reporter assessed by the QUANTI-Blue™ assay (InvitroGen, Carlsbad, CA, USA), following the manufacturer's instructions.

SDS-PAGE analyses

The SDS-PAGE analysis of the samples was performed as previously described.¹⁹ Concisely, the samples were dissolved in sample buffer containing 2-mercaptoethanol, heated at 95 °C for 5 min and loaded (30 µg protein per well) onto 4–12% Bis-Tris Criterion XT precast gel (Bio-Rad, Hercules, CA, USA). The gel was stained with Bio-safe Coomassie blue R-250 (Bio-Rad). As a molecular weight marker, Precision Plus Protein™ Dual Xtra Prestained Protein Standard (Bio-Rad) was used.

Determination of antioxidant activity

The ABTS, DPPH, and OH radical scavenging activities of the samples were assessed, respectively, according to the methods described by Zhang *et al.*,²⁰ Farvin *et al.*,²¹ and Mintah *et al.*²² The oxygen radical absorbance capacity (ORAC) of the samples was measured following the method described by Vasquez-Rojas *et al.*²³ The ability of samples to chelate ferrous iron was evaluated based on the procedure described by Xia *et al.*,²⁴ and the method described by He *et al.*²⁵ was followed for the determination of the ferric reducing power (FRAP) of the samples.

In vitro gastrointestinal digestion

The *in vitro* gastrointestinal digestion of PI and T was performed according to the INFOGEST 2.0 protocol.²⁶ For the gastric phase, PI and T solutions (15 mg mL⁻¹ in the final mixture), mixed with simulated gastric fluid, were adjusted to a pH of 3.0 with 1 M HCl. Pepsin from porcine gastric mucosa was added to achieve a concentration of 2000 U mL⁻¹ in the final mixture, followed by a 2-hour incubation at 37 °C. Pepsin activity was halted by raising the pH to 7.0 using 1 M NaHCO₃. Subsequently, gastric samples were mixed with simulated



intestinal fluid. The pH was adjusted to 7.0 with 1 M NaOH, and CaCl₂ (0.075 mM in the final digestion mixture) was introduced. To simulate the intestinal phase, pancreatin from porcine pancreas (100 U mL⁻¹ in the final digestion mixture based on its trypsin activity) and bovine bile extract (10 mM in the final digestion mixture) were added. The mixtures were subjected to an additional 2-hour incubation and the process was halted with a Bowman–Birk protease inhibitor. Samples were immediately frozen and stored at –80 °C for subsequent analyses.

Immunodetection assays

The human IgE binding capacity of samples was assessed by inhibition ELISA and western blotting as previously reported,²⁷ using a pool of 5 sera from pine nut-allergic patients diagnosed through a compatible clinical history and positive skin prick test. Serum samples were obtained from the Institute of Food Science Research (CIAL) serum library in compliance with relevant institutional and national guidelines and regulations and in accordance with the Declaration of Helsinki. The study protocol was approved by the CIAL Bioethics Committee. Written informed consent was obtained from all participants prior to inclusion. Human samples were processed in a coded manner to protect participant confidentiality and stored at –80 °C until analysis.

Ex vivo experiments in spleen cells from sensitized mice

The immunomodulatory properties of the hydrolysates were assessed on splenocytes from mice sensitized to pine nut protein. Six-week-old female C57BL/6 mice were purchased from Charles River Laboratories (Saint Germain sur l'Arbresle, France). Mice ($n = 5$) were sensitized to PI by the oral administration of 5 mg of PI protein per mouse plus 10 µg of cholera toxin (CT, List Biologicals, Campbell, CA) in PBS, during 3 consecutive days on the first week and once per week during the following 5 weeks (SI Fig. 1A). Mice were euthanized by CO₂ inhalation and spleens were collected and processed under sterile conditions. Splenocytes from individual animals were cultured in 96-well plates (4×10^6 cells per mL) and stimulated in triplicate with PI alone or in combination with its hydrolysates (200 µg mL⁻¹). Stimulation with concanavalin A (2.5 µg mL⁻¹) was used as a positive control, and PBS as a negative control. Inactivated enzymes were also tested at a concentration equivalent to that present in 200 µg mL⁻¹ hydrolysates. Supernatants were collected after 72 h at 37 °C in 5% CO₂ and detection of IL-4 and IFN-γ was performed by ELISA, using commercial kits (Invitrogen).

In vivo experiments in sensitized mice

For the *in vivo* evaluation of the therapeutic capacity of pine nut hydrolysate with trypsin (T), three groups of six-week-old female C57BL/6 mice ($n = 5$) were sensitized to PI as described above, while mice from a fourth group ($n = 5$) just received 10 µg of CT (sham-sensitized mice). One week apart, mice from two of the PI-sensitized groups were orally treated with 5 mg of PI or T protein per mouse, three times per week for 3

weeks. Mice from the other PI-sensitized group and the sham-sensitized group received PBS during the therapy. Three days after the last treatment dose, all mice were orally challenged with 50 mg of PI (SI Fig. 1B). Anaphylactic responses were evaluated by measuring the body temperature with a rectal thermometer and scoring clinical signs 30 min after challenge with the scale described in Pablos-Tanarro *et al.*²⁸ Blood samples were collected for pine nut specific IgE and IgG1 analyses by indirect ELISA as previously described.²⁸ Mouse mast-cell protease 1 (MCP-1) was analysed using an ELISA Kit (Invitrogen).

All protocols involving animals followed the European Legislation (Directive 2010/63/EU) and were approved by the Comunidad de Madrid (Ref PROEX 286.8/20).

Statistical analyses

Statistical analyses were performed using GraphPad Prism version 7.0e (GraphPad Software Inc., San Diego, CA). For Fig. 1, 2, and 3C, differences among groups were assessed by one-way ANOVA followed by Tukey's multiple comparisons test. For Fig. 3, differences were assessed using a two-tailed Student's *t* test. For Fig. 4 and 5A, data were analyzed by two-way ANOVA followed by Šidák's multiple comparisons test. Clinical sign scores (Fig. 5b) were analyzed using the Mann-Whitney *U* test. Statistical significance was set at $P < 0.05$.

Results

Antioxidant and immunomodulant activity of pine nut hydrolysates

Under reducing conditions, PI showed protein bands from ~100 to ~10 kDa. Enzymatic hydrolysis removed most high-molecular-mass proteins, leaving only <10 kDa bands after Alcalase or combined enzyme treatments, while P and T retained bands up to ~15 and 20 kDa, respectively (SI Fig. 2). The antioxidant capacity of the hydrolysates was measured by different *in vitro* methods (Fig. 1). Overall, all hydrolysates showed better antioxidant capacity than PI. Radical scavenging assays revealed that the lowest IC₅₀ values were observed for PA in the ABTS assay, for T, PT, and PA in the DPPH assay, and for T, PT, PA, and TA in the OH radical assay. All hydrolysates showed ORAC antioxidant capacity values ranging from 439 to 615 µM trolox equivalents per g except for A which showed a higher capacity reaching up to 1090 µM trolox equivalents per g. T and TA were the hydrolysates that showed the highest ferrous iron chelating capacity with IC₅₀ values of 19.5 and 42.4 µg mL⁻¹, respectively. The hydrolysate that showed the best ferric reducing ability was P, but it was excluded from further consideration due to its comparatively low antioxidant capacity in the other assays performed. On the basis of the highest overall antioxidant capacity obtained with the 6 methods tested, we selected T, PT, PA, and TA for further evaluation of their immunomodulatory capacity.

In order to test whether the allergen-induced responses on immune cells from pine nut sensitized mice (SI Fig. 1A) could






Fig. 1 Antioxidant activity of pine nut protein isolate (PI) and its hydrolysates obtained using pepsin (P), trypsin (T), Alcalase (A), pepsin/trypsin (PT), pepsin/Alcalase (PA), trypsin/Alcalase (TA), and pepsin/trypsin/Alcalase (PTA) measured by ABTS, DPPH, and OH radical scavenging capacity (SC) assays, oxygen radical absorbance capacity (ORAC) assay, Fe²⁺ chelating activity (CA), and ferric reducing power (FRAP) assays. Data are represented as mean ± SEM (*n* = 5). Different letters indicate significant differences among groups (*p* < 0.05).




Fig. 2 Immunomodulatory effects of pine nut protein isolate (PI) and its hydrolysates obtained using trypsin (T), pepsin/trypsin (PT), pepsin/Alcalase (PA), and trypsin/Alcalase (TA) on the secretion of IL-4 and IFN-γ by spleenocytes from pine nut sensitized mice. Data are expressed as mean ± SEM (*n* = 5). Different letters indicate significant differences among groups (*p* < 0.05).

be modulated by the hydrolysates, splenocytes were stimulated with the PI used for sensitization in the presence of its hydrolysates at a concentration of 200 µg mL⁻¹ of each substance, and the impact on the secretion of Th2 (IL-4) and Th1 (IFN-γ) cytokines was determined (Fig. 2). PI-induced production of IL-4 was inhibited by co-incubation of spleen cells from PI-sensitized mice with the hydrolysates of PI with T and TA. On the other hand, these hydrolysates tended to promote the release of the Th1 cytokine IFN-γ, when used together with PI for the stimulation of the spleen cells from sensitized mice. However, PT and PA failed to reduce IL-4 secretion and increase the levels of IFN-γ induced by PI.

Characterisation of pine nut hydrolysate with trypsin

According to the results explained above, T and TA were the hydrolysates with the highest overall antioxidant capacity and were also able to modulate specific T-cell responses induced by the sensitizing allergen. Ultimately, we opted for T as a candidate for peptide immunotherapy because a single-step






Fig. 3 IgE binding to pine nut protein isolate (PI) and its hydrolysate obtained using trypsin (T), estimated by inhibition ELISA (A) and western blotting (B) with sera from pine nut allergic patients. Data are represented as mean \pm SEM ($n = 3$). Asterisks indicate statistically significant differences. $***P < 0.001$. Full, uncropped blots including all lanes and molecular mass markers are provided in the SI. All samples and controls used for comparative analysis were run on the same gel. Cropping was applied only for clarity, and lane boundaries are indicated.

hydrolysis simplifies production and ensures high reproducibility. In order to evaluate its potential allergenicity, IgE binding was determined by inhibition ELISA using a pool of sera from 5 pine nut allergic patients. Fig. 3 shows that the IgE

binding capacity of PI was significantly reduced after hydrolysis with trypsin, with EC₅₀ values (concentration for 50% of maximum IgE binding) a thousand times higher. Western blot (Fig. 3B) showed that most protein bands in PI were recognized by patient IgE. T still retained residual IgE binding capacity, mainly due to two bands of approximately 15 and 20 kDa that remained unhydrolyzed.

We next evaluated the influence of *in vitro* gastrointestinal digestion on the immunoreactive and antioxidant properties of the hydrolysate. The IgE binding capacity of both PI and T decreased significantly throughout the digestion process, with the percentage of IgE binding being the same for both samples after gastric digestion and remaining similar until the end of duodenal digestion (data not shown). Fig. 4 shows the results of the antioxidant activity determined by different methods after 2 h of simulated gastric digestion, followed by 2 h of duodenal digestion of PI and T. After undergoing pepsin action during gastric digestion, the antioxidant activity of T was significantly higher than that of PI. The same trend was observed after subsequent duodenal digestion when the antioxidant activity was measured by the ABTS and DPPH radical scavenging assays and the FRAP assay. However, when the antioxidant activity was measured by the OH radical scavenging, ferrous iron chelating, and ORAC methods, the values obtained for the gastrointestinal digests of PI and T were similar. Although *in vitro* digestion models do not fully replicate physiologically relevant conditions, these findings suggest




Fig. 4 Antioxidant activity of undigested (C) pine nut protein isolate (PI), its hydrolysate obtained using trypsin (T) and their gastric (G) and gastrointestinal (D) digests, measured by ABTS, DPPH, and OH radical scavenging capacity assays, oxygen radical absorbance capacity (ORAC) assay, Fe²⁺ chelating activity assay, and ferric reducing power (FRAP) assays. Data are represented as mean \pm SEM ($n = 3$). Asterisks indicate statistically significant differences. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$.






Fig. 5 (A) serum EW-specific IgE and IgG1 of mice after sensitization (orange bars) with pine nut protein isolate (PI) plus cholera toxin (CT) and subsequent oral immunotherapy (blue bars) with PI and its hydrolysate obtained using trypsin (T). Untreated mice were administered PBS. (B) Anaphylaxis in mice after oral challenge with PI assessed by the clinical signs, body temperature, and serum concentration of mouse mast cell protease-1, MCP-1. Values are means \pm SEM ($n = 5$), except for clinical signs, which are expressed as medians. Asterisks indicate statistically significant differences. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$.

that the hydrolysate may possess superior properties *in vivo* compared to the parent extract.

***In vivo* evaluation of the therapeutic capacity of pine nut hydrolysate with trypsin**

Pine nut allergy was induced in mice by using the oral sensitization and boosting protocol shown in SI Fig. 1B. Starting at week 6 of the protocol, mice were given 4 weekly oral treatments with PI or T for 5 weeks and the concentrations of specific antibodies were measured in the serum. The levels of specific IgE and IgG1 were significantly reduced in both groups of treated mice, while the levels in the non-treated group remained unchanged (Fig. 5A). Three days after the last treatment dose, mice were orally challenged with PI and no visual signs of anaphylaxis or decrease in body temperature were detected in the treated groups (Fig. 5B). In contrast, the untreated group experienced obvious clinical signs of anaphylaxis and a significant drop in body temperature. Moreover, serum levels of MCP-1, a protease released by activated intestinal mast cells, which is indicative of mast cell degranulation, were significantly lower in the treated mice than in the non-treated ones (Fig. 5B).

Discussion

We used enzymatic hydrolysis of pine nut protein isolate to produce protein hydrolysates with antioxidant and immunomodulatory activities. SDS-PAGE analysis showed protein bands (6–11 kDa) that resisted enzymatic cleavage and were present in all hydrolysates. Cabanillas *et al.* identified a band with a molecular mass of around 6 kDa under reducing conditions as a 2S albumin known as Pin p 1, the first characterized allergen in *Pinus pinea* L. nuts.⁵ This protein has been reported to be resistant to heat treatment and cleavage by digestive enzymes. However, a band with this molecular mass

was reactive to human IgE only in the non-hydrolysed extract (PI).

Our investigation delved into the antioxidant activity of pine nut protein hydrolysates using a diverse array of assays to cover the variety of mechanisms by which antioxidants inhibit oxidative routes, including inactivation of reactive oxygen species, scavenging of free radicals, chelation or sequestration of metal catalysts, or reduction of ferric ions, among others.²⁹ The results obtained with each of the methodologies used depend on the presence of amino acids in peptides that serve as electron or hydrogen donors,³⁰ with sequence position and hydrophobicity also playing crucial roles.³¹ The ABTS assay may be more useful than the DPPH assay for detecting the scavenging capacity of hydrophilic antioxidants in foods,³² while the presence of hydrophobic and aromatic amino acids could contribute to the ability of hydrolysates to neutralize DPPH radicals.³³ The hydroxyl radical (OH) is one of the most toxic reactive oxygen radicals known, which causes damage to all important biomolecules involving DNA, proteins, and membranes.³⁴ Notably, OH radical scavenging activity was observed in T, PT, and TA hydrolysates, likely attributed to the presence of amino acids with hydrogen atom donation capabilities that could mitigate hydrogen peroxide-induced oxidative stress,³⁵ such as Arg and Lys, which can become exposed after trypsin cleaves at their carboxyl side. In the ORAC assay, A exhibited a particularly high antioxidant activity, probably attributed to the ability of this enzyme to expose hydrophobic amino acids during hydrolysis,³⁶ potentially enhancing hydrophobic interactions between peptides and the oxidizing agent.³¹ Pine nut hydrolysates also displayed high Fe²⁺ chelating activity and substantial ferric reducing antioxidant power, which could be attributed to peptides containing sulfhydryl amino acids such as Cys and Met. These residues can bind heavy metals, reducing their pro-oxidant activity,³⁷ and donate electrons to radicals, thereby stabilizing them and preventing oxidative damage.³⁸



Assessment of the immunomodulatory capacity of pine nut protein hydrolysates on splenocytes from sensitized C57BL/6 mice revealed that, among the hydrolysates with the highest antioxidant activity, T and TA could be regarded as good candidates for peptide-based immunotherapy on the grounds of their potential to reduce *ex vivo* Th2 responses induced by pine nut protein, while also enhancing Th1 responses. Previous results from our group have shown that ovalbumin and egg white hydrolysates, selected for their low IgE binding and their ability to reduce Th2 responses in splenocytes from sensitized mice while enhancing Th1 responses, effectively promoted oral tolerance to the corresponding intact allergens in mouse models.^{39,40} This effect was associated with the induction of a tolerogenic environment in the intestine that promotes the expansion of regulatory T cells.⁴¹ Overall, these findings indicate that allergen-derived peptides capable of dampening inflammation and redirecting immune responses of specific cell populations can enhance the effectiveness of immunotherapy. A simple hydrolysis step with trypsin appears to be a cheaper and simpler method for the reproducible production of pine nut-derived antioxidant and immunomodulant peptides than combined hydrolysis with trypsin and Alcalase. Furthermore, there is evidence that bacterial enzymes, such as Alcalase, may induce sensitization in susceptible individuals,⁴² while the European Food Safety Authority considers that the risk of allergic sensitisation to trypsin from porcine pancreas is low.⁴³ Therefore, the hydrolysate obtained with trypsin was selected for further examination, including the assessment of its potential hypoallergenic effect using sera from pine nut allergic subjects. Numerous studies have highlighted the efficacy of enzymatic hydrolysis for reducing the allergenicity of various food proteins.^{39,44} In our study, the IgE binding capacity of PI was significantly reduced after hydrolysis with trypsin, similarly to what has been previously reported for other hydrolysates of different food substrates obtained with this enzyme.⁴⁵ During the digestion process, antioxidant peptides may be further hydrolysed by gastrointestinal enzymes, thereby altering their activities. In this study, the antioxidant activity of T was higher than that of PI after the action of digestive enzymes, as described in the case of other food peptides,^{46,47} although this effect was more pronounced at the end of the gastric phase.

To evaluate the use of the pine nut protein hydrolysate with trypsin as a therapeutic agent, once pine nut allergy was established in mice, T was administered for 3 weeks. Mice treated with the hydrolysate were protected against anaphylaxis induced by an oral pine nut challenge to the same extent as those treated with intact pine nut protein. Protection was accompanied by a reduction in serum pine nut-specific IgE and IgG1 levels, consistent with previous reports of successful desensitization in mouse models.³⁹

The clinical efficacy of OIT is largely attributed to the activation of T cells by specific epitopes, which initiates counter-regulatory immune mechanisms that reduce Th2 cell activity and expansion.⁴⁸ Fragmented allergens offer a distinct advantage in OIT because they preserve T cell-stimulating epitopes

while lacking the ability to cross-link IgE on effector cells, thereby minimizing the risk of adverse reactions. However, the selection of peptides that balance immunostimulatory potential with reduced allergenicity remains challenging, as longer peptides provide broader T cell epitope coverage but increase the likelihood of IgE cross-linking.⁴⁹ Since enzymatic hydrolysis typically generates short fragments with limited sequence coverage, potentially reducing the proportion of full-length T epitopes, the comparable desensitization achieved with PI and T suggests that the latter may exert superior immunomodulatory effects, possibly by promoting a shift from Th2 to Th1 responses and/or inducing regulatory T cells. This dual benefit of reduced allergenicity combined with preserved or enhanced immunomodulatory capacity positions the pine nut hydrolysate with trypsin as a promising candidate for safer and more effective OIT strategies. Such an approach could mitigate adverse reactions during therapy, which are a major barrier to patient adherence, although this aspect cannot be properly evaluated in murine models.

Conclusion

In conclusion, the hydrolysate of pine nuts with trypsin could be regarded as a good candidate for a therapeutic agent in OIT on the grounds of its effectiveness in desensitizing pine nut allergic mice with the added properties of having lower IgE binding capacity and higher antioxidant activity than intact pine nut protein.

Author contributions

N. B.: investigation, methodology, and writing – original draft; D. M-M.: investigation; D. L-O. and N. K.: review; R. L-F. and E. M.: funding acquisition, conceptualization, validation, and writing – review & editing; and S. B.: investigation, methodology, writing – original draft, conceptualization, validation, and writing – review & editing. We acknowledge the Bioanalytical Techniques Unit (BAT) at CIAL for their technical support and services. D. M-M., D. L-O., R. L-F., E. M., and S. B. are members of the CSIC Interdisciplinary Thematic Platform (PTI) “Food Allergy”.

Conflicts of interest

The authors declare that they have no competing interests.

Data availability

The data generated during the current study are available from the corresponding author on request.

Supplementary information (SI) is available. See DOI: <https://doi.org/10.1039/d5fo04650e>.



Acknowledgements

This work was supported by the project PID2021-123236OB-I00. D. Menchén-Martínez was the recipient of an FPI grant (all from the Ministerio de Ciencia, Innovación y Universidades).

References

- J. P. Lopes and S. Sicherer, Food allergy: epidemiology, pathogenesis, diagnosis, prevention, and treatment, *Curr. Opin. Immunol.*, 2020, **66**, 57–64.
- M. P. Borres, S. Sato and M. Ebisawa, Recent advances in diagnosing and managing nut allergies with focus on hazelnuts, walnuts, and cashew nuts, *World Allergy Organ. J.*, 2022, **15**, 100641.
- B. Elrahi, Z. Mehanna, S. Haidar, M. Serhan and H. F. Hassan, Food allergies and allergens in Lebanon: characterization and perceptions toward labeling, *World Allergy Organ. J.*, 2023, **16**, 100743.
- B. Cabanillas and N. Novak, Allergic reactions to pine nut: a review, *J. Invest. Allergol. Clin. Immunol.*, 2015, **25**, 329–333.
- B. Cabanillas, J. F. Crespo, S. J. Maleki, J. Rodriguez and N. Novak, Pin p 1 is a major allergen in pine nut and the first food allergen described in the plant group of gymnosperms, *Food Chem.*, 2016, **210**, 70–77.
- B. Cabanillas, H. Cheng, C. C. Grimm, B. K. Hurlburt, J. Rodríguez, J. F. Crespo and S. J. Maleki, Pine nut allergy: clinical features and major allergens characterization, *Mol. Nutr. Food Res.*, 2012, **56**, 1884–1893.
- J. F. Crespo, C. Bueno, M. Villalba, L. Monaci, C. Cuadrado, N. Novak and B. Cabanillas, Epitope mapping of the major allergen 2S albumin from pine nut, *Food Chem.*, 2021, **339**, 127895.
- M. Vazquez-Ortiz and P. J. Turner, Improving the safety of oral immunotherapy for food allergy, *Pediatr. Allergy Immunol.*, 2016, **27**, 117–125.
- G. Pouessel and G. Lezmi, Oral immunotherapy for food allergy: translation from studies to clinical practice?, *World Allergy Organ. J.*, 2023, **16**, 100747.
- D. Lozano-Ojalvo, E. Molina and R. López-Fandiño, Hydrolysates of egg white proteins modulate T- and B-cell responses in mitogen-stimulated murine cells, *Food Funct.*, 2016, **7**, 1048–1056.
- D. Lozano-Ojalvo and R. López-Fandiño, Immunomodulating peptides for food allergy prevention and treatment, *Crit. Rev. Food Sci. Nutr.*, 2018, **58**, 1629–1649.
- D. Lozano-Ojalvo, M. Martínez-Blanco, L. Pérez-Rodríguez, E. Molina and R. López-Fandiño, Oral immunotherapy with egg peptides induces innate and adaptive tolerogenic responses, *Mol. Nutr. Food Res.*, 2019, **63**, e1900144.
- S. Benedé and E. Molina, Chicken egg proteins and derived peptides with antioxidant properties, *Foods*, 2020, **9**, 735.
- R. P. Bowler and J. D. Crapo, Oxidative stress in allergic respiratory diseases, *J. Allergy Clin. Immunol.*, 2002, **110**, 349–356.
- M. Han, D. Lee, S. H. Lee and T. H. Kim, Oxidative stress and antioxidant pathway in allergic rhinitis, *Antioxidants*, 2021, **10**, 1266.
- L. Utsch, C. Folisi, J. H. Akkerdaas, A. Logiantara, M. A. van de Pol, J. S. van der Zee, E. J. Krop, R. Lutter, R. van Ree and L. S. van Rijt, Allergic sensitization is associated with inadequate antioxidant responses in mice and men, *Allergy*, 2015, **70**, 1246–1258.
- C. H. Guo, P. J. Liu, K. P. Lin and P. C. Chen, Nutritional supplement therapy improves oxidative stress, immune response, pulmonary function, and quality of life in allergic asthma patients: an open-label pilot study, *Altern. Med. Rev.*, 2012, **17**, 42–56.
- E. Pendezza, C. Nava, A. Leone, F. Riccaboni, A. Bosetti, G. V. Zuccotti, S. Bertoli and E. D'Auria, Antioxidant potential of the diet in Italian children with food allergies, *Front. Nutr.*, 2023, **10**, 1096288.
- S. Benedé, I. López-Expósito, R. López-Fandiño and E. Molina, Identification of IgE-binding peptides in hen egg ovalbumin digested in vitro with human and simulated gastroduodenal fluids, *J. Agric. Food Chem.*, 2014, **62**, 152–158.
- W. Zhang, N. Jia, Z. Zhu, Y. Wang, J. Wang and C. Xue, Changes of antioxidative activities and peptidomic patterns of *Auxenochlorella pyrenoidosa* protein hydrolysates: effects of enzymatic hydrolysis and decoloration processes, *LWT*, 2021, **152**, 112306.
- K. H. Farvin, L. L. Andersen, H. H. Nielsen, C. Jacobsen, G. Jakobsen, I. Johansson and F. Jessen, Antioxidant activity of cod (*Gadus morhua*) protein hydrolysates: in vitro assays and evaluation in 5% fish oil-in-water emulsion, *Food Chem.*, 2014, **149**, 326–334.
- B. K. Mintah, R. He, M. Dabbour, M. K. Golly, A. A. Agyekum and H. Ma, Effect of sonication pretreatment parameters and their optimization on the antioxidant activity of *Hermetia illucens* larvae meal protein hydrolysates, *J. Food Process. Preserv.*, 2019, **43**, e14093.
- W. V. Vasquez-Rojas, D. Martín, B. Miralles, I. Recio, T. Fornari and M. P. Cano, Composition of Brazil nut (*Bertholletia excelsa* HBK), its beverage and by-products: a healthy food and potential source of ingredients, *Foods*, 2021, **10**, 3007.
- Y. Xia, F. Bamdad, M. Gänzle and L. Chen, Fractionation and characterization of antioxidant peptides derived from barley glutelin by enzymatic hydrolysis, *Food Chem.*, 2012, **134**, 1509–1518.
- R. He, A. T. Girgih, S. A. Malomo, X. Ju and R. E. Aluko, Antioxidant activities of enzymatic rapeseed protein hydrolysates and the membrane ultrafiltration fractions, *J. Funct. Foods*, 2013, **5**, 219–227.
- A. Brodkorb, L. Egger, M. Alminger, P. Alvito, R. Assunção, S. Balance, T. Bohn, C. Bourlieu-Lacanal, R. Boutrou, F. Carrière, A. Clemente, M. Corredig, D. Dupont,



- C. Dufour, C. Edwards, M. Golding, S. Karakaya, B. Kirkhus, S. Le Feunteun, U. Lesmes, A. Macierzanka, A. R. Mackie, C. Martins, S. Marze, D. J. McClements, O. Ménard, M. Minekus, R. Portmann, C. N. Santos, I. Souchon, R. P. Singh, G. E. Vegarud, M. S. J. Wickham, W. Weitschies and I. Recio, INFOGEST static in vitro simulation of gastrointestinal food digestion, *Nat. Protoc.*, 2019, **14**, 991–1014.
- 27 L. Martín-Pedraza, C. Mayorga, F. Gomez, C. Bueno-Díaz, N. Blanca-Lopez, M. González, M. Martínez-Blanco, J. Cuesta-Herranz, E. Molina, M. Villalba and S. Benedé, IgE-reactivity pattern of tomato seed and peel nonspecific lipid-transfer proteins after in vitro gastrointestinal digestion, *J. Agric. Food Chem.*, 2021, **69**, 3511–3518.
- 28 A. Pablos-Tanarro, D. Lozano-Ojalvo, E. Molina and R. López-Fandiño, Assessment of the allergenic potential of the main egg white proteins in BALB/c mice, *J. Agric. Food Chem.*, 2018, **66**, 2970–2976.
- 29 K. Jomova, S. Y. Alomar, S. H. Alwasel, E. Nepovimova, K. Kuca and M. Valko, Several lines of antioxidant defense against oxidative stress: antioxidant enzymes, nano-materials with multiple enzyme-mimicking activities, and low-molecular-weight antioxidants, *Arch. Toxicol.*, 2024, **98**, 1323–1367.
- 30 A. Sila and A. Bougatef, Antioxidant peptides from marine by-products: isolation, identification and application in food systems. A review, *J. Funct. Foods*, 2016, **21**, 10–26.
- 31 A. G. Samaranyaka and E. C. Li-Chan, Food-derived peptidic antioxidants: a review of their production, assessment, and potential applications, *J. Funct. Foods*, 2011, **3**, 229–254.
- 32 A. Floegel, D.-O. Kim, S.-J. Chung, S. I. Koo and O. K. Chun, Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods, *J. Food Compos. Anal.*, 2011, **24**, 1043–1048.
- 33 S. Zhang, M. Zhang, J. Xing and S. Lin, A possible mechanism for enhancing the antioxidant activity by pulsed electric field on pine nut peptide glutamine–tryptophan–phenylalanine–histidine, *J. Food Biochem.*, 2019, **43**, e12714.
- 34 S. B. Nimse and D. Pal, Free radicals, natural antioxidants, and their reaction mechanisms, *RSC Adv.*, 2015, **5**, 27986–28006.
- 35 A. Phaniendra, D. B. Jestadi and L. Periyasamy, Free radicals: properties, sources, targets, and their implication in various diseases, *Indian J. Clin. Biochem.*, 2015, **30**, 11–26.
- 36 V. G. Tacias-Pascacio, R. Morellon-Sterling, E. H. Siar, O. Tavano, Á. Berenguer-Murcia and R. Fernandez-Lafuente, Use of Alcalase in the production of bioactive peptides: a review, *Int. J. Biol. Macromol.*, 2020, **165**, 2143–2196.
- 37 U. Urbizo-Reyes, M. F. San Martín-González, J. García-Bravo, A. L. M. Vigil and A. M. Liceaga, Physicochemical characteristics of chia seed (*Salvia hispanica*) protein hydrolysates produced using ultrasonication followed by microwave-assisted hydrolysis, *Food Hydrocolloids*, 2019, **97**, 105187.
- 38 M. M. Aondona, J. K. Ikya, M. T. Ukeyima, T. W. J. Gborigo, R. E. Aluko and A. T. Girgih, *In vitro* antioxidant and antihypertensive properties of sesame seed enzymatic protein hydrolysate and ultrafiltration peptide fractions, *J. Food Biochem.*, 2021, **45**, e13587.
- 39 D. Lozano-Ojalvo, E. Molina and R. López-Fandiño, Hypoallergenic hydrolysates of egg white proteins modulate allergen responses induced ex vivo on spleen cells from sensitized mice, *Food Res. Int.*, 2016, **89**, 661–669.
- 40 D. Lozano-Ojalvo, L. Pérez-Rodríguez, A. Pablos-Tanarro, E. Molina and R. López-Fandiño, Hydrolysed ovalbumin offers more effective preventive and therapeutic protection against egg allergy than the intact protein, *Clin. Exp. Allergy*, 2017, **47**, 1342–1354.
- 41 D. Lozano-Ojalvo, M. Martínez-Blanco, L. Pérez-Rodríguez, E. Molina, C. Peláez, T. Requena and R. López-Fandiño, Egg white peptide-based immunotherapy enhances vitamin A metabolism and induces ROR γ t regulatory T cells, *J. Funct. Foods*, 2019, **52**, 204–211.
- 42 D. I. Bernstein, I. L. Bernstein, W. G. Gaines Jr., T. Stauder and E. R. Wilson, Characterization of skin prick testing responses for detecting sensitization to detergent enzymes at extreme dilutions: inability of the RAST to detect lightly sensitized individuals, *J. Allergy Clin. Immunol.*, 1994, **94**, 498–507.
- 43 EFSA Panel on Food Contact Materials, Enzymes and Processing Aids (CEP), C. Lambré, J. M. Barat Baviera, C. Bolognesi, P. S. Cocconcelli, R. Crebelli, D. M. Gott, K. Grob, E. Lampi, M. Mengelers, A. Mortensen, G. Rivière, I. L. Steffensen, C. Tlustos, H. Van Loveren, L. Vernis, H. Zorn, M. Andryszkiewicz, Y. Liu, S. Rainieri and A. Chesson, Safety evaluation of the food enzyme trypsin from porcine pancreas, *EFSA J.*, 2022, **20**, e07008.
- 44 C. Cuadrado, C. Arribas, A. Sanchiz, M. M. Pedrosa, P. Gamboa, D. Betancor, C. Blanco, B. Cabanillas and R. Linacero, Effects of enzymatic hydrolysis combined with pressured heating on tree nut allergenicity, *Food Chem.*, 2024, **451**, 139433.
- 45 S. J. Laly, A. Kumar, T. V. Sankar and S. K. Panda, In vitro characterisation of tropomyosin from flower tail shrimp, *Metapenaeus dobsoni*, and effect of hydrolysis on allergenicity of tropomyosin, *Int. J. Food Sci. Technol.*, 2022, **57**, 7434–7444.
- 46 S. Ketnawa, O. Martínez-Alvarez, S. Benjakul and S. Rawdkuen, Gelatin hydrolysates from farmed giant catfish skin using alkaline proteases and its antioxidative function of simulated gastro-intestinal digestion, *Food Chem.*, 2016, **192**, 34–42.
- 47 H. Li, H. Fan, Z. Wang, Q. Zhua and J. Wu, Effect of simulated gastrointestinal digestion on antioxidant and anti-inflammatory activities of bioactive peptides generated in



- sausages fermented with *Staphylococcus simulans* QB7, *Food Sci. Hum. Wellness*, 2024, **13**, 1662–1671.
- 48 M. D. Kulis, S. U. Patil, E. Wambre and B. P. Vickery, Immune mechanisms of oral immunotherapy, *J. Allergy Clin. Immunol.*, 2018, **141**, 491–498.
- 49 S. R. Prickett, A. L. Voskamp, T. Phan, A. Dacumos-Hill, S. I. Mannering, J. M. Rolland and R. E. O’Hehir, Ara h 1 CD4⁺ T cell epitope-based peptides: candidates for a peanut allergy therapeutic, *Clin. Exp. Allergy*, 2013, **43**, 684–697.

