



Cite this: DOI: 10.1039/d6lc90032a

Correction: Cell docking inside microwells within reversibly sealed microfluidic channels for fabricating multiphenotype cell arrays

Ali Khademhosseini,^{ab} Judy Yeh,^c George Eng,^c Jeffrey Karp,^c Hirokazu Kaji,^d Jeffrey Borenstein,^e Omid C. Farokhzad^f and Robert Langer^{*ac}

DOI: 10.1039/d6lc90032a

Correction for ‘Cell docking inside microwells within reversibly sealed microfluidic channels for fabricating multiphenotype cell arrays’ by Ali Khademhosseini *et al.*, *Lab Chip*, 2005, 5, 1380–1386, <https://doi.org/10.1039/B508096G>.

rsc.li/loc

An issue was identified with the original Fig. 5B, where certain elements of the image appeared repetitive. As the original data were unavailable, and the cause of the issue could not be determined, the experiment was independently repeated. The independently reperfomed Fig. 5 corroborates the conclusions presented in the original study.




Fig. 5 Microfluidic arrays with upstream microfluidic mixers. To lower the number of independent inlets into the device, micromixers can be incorporated upstream from the microchannel arrays: (a) represents experiments in which a concentration gradient was generated in an array of channels, and (b) is a monolayer of NIH-3T3 cells immobilized in such a microfluidic array.

An independent expert has viewed the independently reperfomed Fig. 5 and the associated raw data, and has confirmed that it is consistent with the discussions and conclusions presented in the original article.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Harvard-MIT Division of Health Sciences and Technology, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA. E-mail: rlanger@mit.edu

^b Department of Medicine, Brigham and Women’s Hospital, Harvard Medical School, Boston, MA, 02115, USA

^c Department of Chemical Engineering, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA

^d Department of Bioengineering and Robotics, Graduate School of Engineering, Tohoku University, Aoba, Sendai, 980-8579, Japan

^e Draper Laboratory, 555 Technology Square, Cambridge, MA, 02139, USA

^f Department of Anesthesiology, Brigham and Women’s Hospital, Harvard Medical School, Boston, MA, 02115, USA

