

RESEARCH ARTICLE

[View Article Online](#)
[View Journal](#) | [View Issue](#)Cite this: *RSC Med. Chem.*, 2026, 17, 1513Investigating the role of cytochrome *bd* oxidases in the antibacterial activity of madecassic acid and derivatives thereofSamantha A. Henry,^a Geraud N. Sansom,^a  Thao Thi Phuong Tran,^b 
Ryan A. Boughton,^a Guy Joiner,^a Calum M. Webster,^a H. Ireshika C. de Silva,^c 
Michelle D. Garrett,^a  Christopher J. Serpell,^d 
Gary K. Robinson  and Mark Shepherd *

Natural products are valuable starting points for drug discovery, although individual modes of action are often difficult to pin down. Ursanes such as madecassic acid have been shown to have antibacterial properties, but a variety of mechanisms have been proposed. In this paper, we report previously uninvestigated activity against cytochrome *bd* oxidases which are only found in prokaryotes and are therefore promising new targets, using madecassic acid and a set of synthetically modified derivatives. Our work shows that madecassic acid and its derivatives can block activity of these enzymes, while phenotypic effects in membrane and whole organism assays are more complex, consistent with modulation of multiple pathways depending upon molecular structure. This provides a new route to ursane-based antibacterial action while highlighting the importance of chemical modifications in fine-tuning biological activity of natural products.

Received 11th December 2025,
Accepted 28th December 2025

DOI: 10.1039/d5md01116g

rsc.li/medchem

Introduction

Ursanes are pentacyclic terpenoids which have been shown to have antibacterial properties.^{1–4} Madecassic acid, an ursane produced by the common Asian herb *Centella asiatica*, has been shown to have good activity against *S. aureus* and moderate activity against other bacterial species, with evidence consistent with a mechanism including increasing membrane and wall permeability, interaction with DNA, and inhibition of protein synthesis.⁵ This is similar to evidence for ursolic acid, a related compound also found to be active against Gram-positive bacteria.⁶ Unpurified extracts of *C. asiatica* have shown to be active against Gram-positive and -negative bacteria.⁷ These trends have been demonstrated across a range of modified ursanes,⁸ and the potential of asiatic acid to inhibit bacterial growth had led to its incorporation into gel materials to aid *in vivo* wound healing.⁹

Cytochrome *bd* oxidases are attractive drug targets for discovery of new antibacterials, since they are found only in prokaryotic organisms,¹⁰ and therefore permit specific killing of bacteria. The proteins are found on the inner membrane, where their role is to

couple reduction of molecular oxygen with oxidation of quinols, resulting in a proton motive force used in the generation of ATP. Blocking cytochrome *bd* oxidase activity would therefore impede the capacity of bacteria to access energy required for growth.¹¹ *Escherichia coli*, a Gram-negative bacterium responsible for many urinary tract and bloodstream infections, has two *bd* oxidases: *bd*-I which is expressed in microaerobic environments, and *bd*-II which is predominant in anaerobic conditions.¹²

We have discovered effective steroidal ligands for *bd*-I which inhibit growth and provide lethality against bacteria, with the strongest measured affinities in *E. coli*.¹³ Since ursane-type structures resemble sterols, it is possible that binding of bacterial cytochrome oxidases is a key component in the antibacterial mechanism of action of the compound class. Since literature reports that a number of mechanisms are at play, it would also be useful to know if the balance of these processes can be modulated through structural refinement of the compounds. In this paper we test madecassic acid (**MA**) and three synthetic analogues for cytochrome oxidase inhibition through molecular docking, measurement of oxygen consumption in isolated membranes, and testing against live *E. coli* models, each engineered to have access to different oxidases.

Experimental procedures

Chemical synthesis

Madecassic acid (**MA**) was obtained by extraction of leaves grown in Quang Tho II Commune, Hue Province, VietNam,

^a School of Natural Sciences, University of Kent, Canterbury, CT2 7NH, UK.

E-mail: M.Shepherd@kent.ac.uk

^b Institute of Chemistry, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet Road, Nghia Do, Hanoi, Vietnam^c Department of Chemistry, University of Colombo, Colombo 03, Sri Lanka^d School of Pharmacy, University College London, 29-39 Brunswick Square, London, WC1N 1AX, UK. E-mail: chris.serpell@ucl.ac.uk

following our reported procedure.¹⁴ Deuterated solvents were purchased from Cambridge Isotope Laboratories Inc. Solvents and reagents were purchased from Thermo Fisher Scientific, Fluorochem, or Alfa Aesar.

A Bruker AVII 400 MHz spectrometer was used to record NMR spectra, and each spectrum was calibrated to the known chemical shift of the residual solvent peak of the deuterated solvent used. Chemical shifts were reported in part per million (ppm) and J coupling values were reported in Hz. Proton NMR spectra were obtained at 400 MHz and ¹³C spectra were obtained at 101 MHz. Spectral data was processed using MestReNova software. Electrospray mass spectrometry data was obtained using a Thermo MSQPlus instrument fitted with a Zorbax SB-C18 5 μm 3.0 × 150 mm column using H₂O + 0.1% formic acid and MeOH + 0.1% formic acid or H₂O + 0.1% TFA and MeCN + 0.1% TFA mobile phases. Data was analysed using Chromeleon™ Chromatography Data System (CDS) Software. A Bruker micrOTOF-Q LCMS system was used to obtain high-resolution mass spectrometry data, samples were dissolved in HPLC-grade methanol and injected using direct injection mode with a mobile phase system of 50:50 MeOH and H₂O. Data was processed using Bruker Compass Hystar software.

MA-2. **MA** (201.6 mg, 0.40 mmol) was dissolved in 6 mL of dry pyridine under nitrogen and acetic anhydride (0.13 mL, 1.40 mmol) was injected into the reaction slowly. After 24 hours, 10 mL pyridine was added, and additional acetic anhydride (4 mL, 42.3 mmol) was added to the reaction. After 48 hours, TLC (after staining with vanillin) showed complete consumption of starting material. The reaction was transferred to a separating funnel, EtOAc (25 mL) was added, and the reaction organic layer was washed with 1 M HCl_(aq) (3 × 50 mL). The organic layer was dried over anhydrous MgSO₄, filtered and the solvent removed under reduced pressure. Crude product was purified using gradient silica gel column chromatography (0–50%, EtOAc: CH₂Cl₂) giving **MA-2** as a white needle crystalline solid (183 mg 72.6% yield). ¹H NMR (400 MHz, CDCl₃) δ 5.27 (d, J = 3.6 Hz, 1H), 5.22 (td, J = 11.3, 4.8 Hz, 2H), 5.00 (d, J = 10.3 Hz, 1H), 4.33 (d, J = 3.8 Hz, 1H), 3.93 (d, J = 11.9 Hz, 1H), 3.70 (d, J = 12.0 Hz, 1H), 2.19 (d, J = 11.2 Hz, 1H), 2.15–0.80 (m, 46H). ¹³C NMR (101 MHz, CDCl₃) δ 183.9, 171.0, 170.6, 170.6, 137.3, 125.7, 75.0, 70.0, 68.0, 65.5, 52.5, 48.3, 48.0, 47.9, 45.9, 42.5, 40.8, 39.2, 38.9, 38.7, 37.4, 36.7, 30.7, 28.0, 24.2, 23.6, 23.4, 21.3, 21.2, 21.2, 21.0, 20.9, 18.8, 18.6, 17.1, 15.5, 14.3. **HREI-MS:** *m/z* calculated for C₃₆H₅₄O₉Na [M + Na⁺] 653.3660; observed 653.3672.

MA-3. The *tert*-butyloxycarbonyl-protected version of **MA-3** (**MA-3Boc**) was first synthesised. HATU (131.1 mg, 0.334 mmol) and **MA-2** (175.5 mg, 0.278 mmol) were dissolved in 5 mL dry CH₂Cl₂ and DIPEA (0.145 mL, 0.835 mmol) was slowly added to the reaction and it was stirred for 1 hour. *Tert*-butyl 10-aminodecylcarbamate (152.8 mg, 0.556 mmol) was dissolved in dry CH₂Cl₂ (5 mL) and slowly added to the reaction and it was left to stir overnight. After 22 hours, the reaction was transferred to a separating funnel and diluted

with CH₂Cl₂ (30 mL). It was then extracted with 1 M HCl_(aq) (3 × 50 mL) and brine (1 × 50 mL). Dried over anhydrous MgSO₄, filtered and reduced under vacuum. The crude product was purified by isocratic column chromatography (25%:75%, EtOAc:CHCl₃) to yield **MA-3Boc** as a white solid (236 mg, 95.8%). ¹H NMR (400 MHz, CDCl₃) δ 5.85 (t, J = 5.4 Hz, 1H), 5.34 (t, J = 3.5 Hz, 1H), 5.22 (td, J = 11.0, 4.7 Hz, 1H), 5.01 (d, J = 10.3 Hz, 1H), 4.57 (s, 1H), 4.34 (t, J = 3.8 Hz, 1H), 3.94 (d, J = 11.9 Hz, 1H), 3.71 (d, J = 12.0 Hz, 1H), 3.23 (dq, J = 13.3, 6.8 Hz, 1H), 3.14–2.93 (m, 3H), 2.17–0.77 (m, 72H). ¹³C NMR (101 MHz, CDCl₃) δ 177.9, 170.9, 170.5, 170.5, 139.4, 125.5, 79.1, 75.0, 70.1, 67.4, 65.4, 54.2, 48.1, 47.9, 47.8, 46.0, 43.1, 42.5, 40.6, 39.9, 39.6, 39.2, 38.8, 38.8, 37.4, 37.3, 31.0, 30.2, 29.5, 29.5, 29.3, 29.3, 29.3, 29.2, 28.5, 27.9, 27.2, 26.9, 24.9, 23.5, 23.3, 21.3, 21.2, 21.0, 20.9, 18.7, 18.3, 17.3, 15.4. **HREI-MS:** *m/z* calculated for C₅₁H₈₄N₂O₁₀Na [M + Na⁺] 907.6018; observed 907.6052.

MA-3Boc (131.2 mg, 0.148 mmol) was then dissolved in CH₂Cl₂ (3.7 mL), 4 M HCl in dioxane (3.7 mL, 14.8 mmol) was added to the solution at 0 °C. The reaction was stirred and left to warm to room temperature overnight. After 21 hours, solvent was removed under reduced pressure. The residue was washed into a separating funnel with a mixture of CH₂Cl₂ (50 mL) and sat. NaHCO_{3(aq)} (50 mL) and aqueous layer removed. It was then extracted with a further sat. NaHCO_{3(aq)} (2 × 50 mL) and brine (1 × 50 mL). Dried over anhydrous MgSO₄, filtered and reduced under vacuum. Extracted product was clean by NMR and required no further purification yielding **MA-3** (105.8 mg, 90.9%). ¹H NMR (400 MHz, CDCl₃) δ 5.92 (t, J = 5.4 Hz, 1H), 5.34 (d, J = 3.7 Hz, 1H), 5.21 (td, J = 10.9, 4.5 Hz, 1H), 5.00 (d, J = 10.3 Hz, 1H), 4.83 (s, 2H), 4.31 (d, J = 5.1 Hz, 1H), 3.94 (d, J = 11.7 Hz, 1H), 3.70 (dd, J = 12.5, 2.6 Hz, 1H), 3.28–2.94 (m, 3H), 2.85 (s, 2H), 2.25–0.73 (m, 62H). ¹³C NMR (101 MHz, CDCl₃) δ 178.1, 171.1, 170.7, 170.6, 139.4, 125.5, 75.0, 70.2, 67.2, 65.4, 54.1, 48.1, 47.9, 47.8, 46.0, 43.8, 43.1, 42.5, 40.4, 39.9, 39.7, 39.2, 38.8, 37.4, 37.3, 31.0, 29.5, 29.4, 29.3, 29.3, 29.2, 29.1, 27.8, 27.1, 26.7, 24.9, 23.5, 23.3, 21.3, 21.3, 21.0, 20.9, 18.7, 18.3, 17.3, 15.4.

MA-4. **MA-3** (52.0 mg, 0.0662 mmol) was dissolved in MeOH (5 mL). 4 M NaOH_(aq) solution (0.080 mL, 0.331 mmol) was injected into the reaction, and it was left to stir overnight. After 20 hours, reaction was transferred to a separating funnel, EtOAc (50 mL) was added. The organic layer was extracted with 1 M NaOH_(aq) (3 × 50 mL). Dried over anhydrous MgSO₄, filtered and reduced under vacuum. Extracted product was pure and required no further purification, yielding **MA-4** (43.6 mg, 99.9%). ¹H NMR (400 MHz, methanol-*d*₄) δ 5.39 (t, J = 3.6 Hz, 1H), 4.37 (t, J = 3.4 Hz, 1H), 3.74 (ddd, J = 11.3, 9.3, 4.3 Hz, 1H), 3.58 (d, J = 11.1 Hz, 1H), 3.43 (d, J = 11.1 Hz, 1H), 3.30 (d, J = 8.0 Hz, 1H), 3.21–3.01 (m, 2H), 2.75–2.66 (m, 2H), 2.20–0.79 (m, 54H). ¹³C NMR (101 MHz, methanol-*d*₄) δ 180.1, 139.7, 127.1, 78.0, 69.6, 68.3, 65.7, 54.3, 50.2, 49.1, 48.9, 44.8, 43.9, 42.0, 41.2, 40.9, 40.8, 40.3, 40.1, 38.8, 38.5, 32.2, 32.0, 30.7, 30.7, 30.7, 30.6, 30.5, 30.3, 28.9, 28.3, 27.9, 25.2, 24.5, 24.1, 21.6, 19.3,



19.3, 17.7, 15.3. **HREI-MS**: m/z calculated for $C_{40}H_{71}N_2O_5$ $[M + H^+]$ 659.5357; observed 659.5392.

Computation and molecular docking

Ligand docking to an AlphaFold2 model of *E. coli* cytochrome *bd-I* was performed essentially as described in detail previously.¹³ Briefly, preparation of *in silico* ligand and protein files was performed using AutoDockTools and PyMOL.¹⁵ The equation $K_d = \exp(\Delta G/(R \times T))$ with ΔG = binding energy (kcal mol⁻¹), R = gas constant = (1.986 cal mol⁻¹ K⁻¹) and T = temperature (298 K) was used to estimate the dissociation constants.

Clog P values were calculated using the Chemical Properties tool in Signals ChemDraw v23.1.2.7.

Bacterial strains

EC958 is a multidrug-resistant *E. coli* O25: H4-ST131 isolate.^{16,17} Generation of the EC958 *bd-I* only strain (genotype $\Delta cyoA$ *appCB::Cm*) is described elsewhere.¹³

Growth assays

Starter cultures were grown in 10 mL LB (*E. coli*) in sterile 50 mL conical flasks at 180 rpm and 37 °C until stationary phase was reached and were used to inoculate 50 mL of fresh growth medium in 250 mL conical flasks. M9 minimal medium was used for *E. coli* (16 g L⁻¹ Na₂HPO₄·2H₂O, 3 g L⁻¹ KH₂PO₄, 0.5 g L⁻¹ NaCl, 1 g L⁻¹ NH₄Cl, 0.24 g L⁻¹ MgSO₄, 0.01 g L⁻¹ CaCl₂, 0.1% casamino acids and 2% glycerol). Drug stocks were prepared in DMSO so that their final concentrations were 40× higher than the working concentrations. Greiner F-bottom sterile 96-well plates were prepared by adding 100 μL of a 2× concentrated growth medium, 61.7 μL sterile milliQ H₂O, 5 μL of the drug and 33.3 μL of cell culture (final OD₆₀₀ of ~0.1). Cells were grown in a FLUOstar Omega plate reader at double orbital pattern setting.

Oxygen consumption assays

Membrane preparation and oxygen consumption were performed as described previously.¹³ Briefly, cells were grown to exponential phase then harvested *via* centrifugation for 20 min at 4000 rpm and 4 °C. The cell pellet was resuspended in ice-cold sonication buffer (20 mM Tris/HCl at pH 7.4, 2 mM MgCl₂, and 1 mM EGTA). The resuspended cells were sonicated (6 × 30s on ice at 15 μM) before centrifugation at 44000 rpm for 1 h and 4 °C to isolate membranes. The membrane pellet was resuspended in 20 mM Tris/HCl (pH 7.4) at a final concentration of 100 mg mL⁻¹ and stored at -20 °C. For oxygen measurements a Rank Brothers oxygen electrode with a 4 mL closed chamber was used at 37 °C, which contained 50 mM HEPES pH 7.4, 0.5 mg mL⁻¹ membranes (based on wet membranes) and DMSO-solubilised drug (added from 40× final concentration). A final concentration of 8 mM succinate (pH 7.4) was added (from

160 mM stock) to initiate the reaction with a single run lasting 15–20 min.

Viability assays

Viability assays were performed essentially as described previously.¹³ Briefly, *E. coli* was grown overnight in 10 mL LB medium and was used to inoculate M9 medium. 20 μL of each drug was added to the wells of row A of a 96 well plate followed by 180 μL of cells (OD₆₀₀ of 0.1). Cells were exposed to drug for 3 hours at 37 °C. Following drug exposure, serial dilutions were performed in 1 × phosphate buffered saline (pH 7.4) before being spotted onto LB agar plates overnight to determine changes in cell survival. Six repeats were performed for each concentration of drug which included two biological repeats and three technical repeats of each.

Results

Both **MA** and ubiquinol-8 (**UQ-8**) the natural substrate of cytochrome *bd-I* were docked to the quinol site of the *E. coli* AlphaFold2 cytochrome *bd-I* structural model.¹³ The binding affinity values predicted by AutoDock Vina (Fig. 1) indicate that **MA** is substantially more tightly bound by cytochrome *bd-I* (1.14 μM) than **UQ-8** (108.9 μM). The carboxylic acid of **MA** was found to reside in the hydrophobic cleft (red, Fig. 1) close to where the aromatic unit of **UQ-8** sits, while the pentacyclic hydrocarbon occupies a similar space to that of the oligoisoprene tail of **UQ-8**. The polar units on **MA** point out of the protein cavity towards the aqueous environment.

These predictions suggest that **MA** would be able to competitively inhibit cytochrome *bd-I*. To test this prediction, we performed oxygen consumption assays to measure oxidoreductase activity using isolated *E. coli* EC958 cytochrome *bd-I* only membranes (Fig. 2A),¹³ and **MA** obtained by extraction of *C. asiatica*,¹⁴ resulting in an IC₅₀ of 34 ± 11 μM. This activity is consistent with the docking results in that **MA** would be expected to outcompete **UQ-8**. On this basis, we advanced to assessing the ability of **MA** to inhibit the growth of cytochrome *bd-I* only *E. coli* cells (Fig. 2B), giving an IC₅₀ of 9.4 ± 1.5 μM, with a maximum inhibition of 63%. We further conducted a viability assay (Fig. 2C) which showed that **MA** did not kill *bd-I* only *E. coli* cells. The observed lack of lethality may relate to permeability of the cell membrane and/or wall to **MA**, but is consistent with other tests on Gram-negative bacteria.⁵

To investigate the contribution of cytochrome *bd-I* to **MA**-mediated respiratory inhibition, growth experiments were also performed on the wild type EC958 strain, which encodes an additional respiratory oxidase cytochrome *bo'* that is expressed under the aerobic conditions tested (Fig. S1A). These data revealed an IC₅₀ of ~10 μM and maximal inhibition of 48%, which is consistent with the wild type responding to respiratory inhibition in a similar way to the *bd-I* only strain. In addition, the wild type strain was not susceptible to **MA**-mediated killing at concentrations below 1






Fig. 1 Docking of ubiquinol-8 (A) and MA (B) to the quinol-binding site of an AlphaFold 2 model for *E. coli* cytochrome *bd-I*. Haem b_{558} is a cytochrome *bd-I* cofactor that accepts electrons from the ubiquinol substrate, and is labelled on the figure. Predicted affinities from Autodock Vina are shown in red.




Fig. 2 Impact of madecassic acid upon cytochrome *bd-I* activity, *E. coli* growth, and *E. coli* survival. (A) Dose inhibition curve of madecassic acid against oxygen consumption activity of *E. coli* EC958 cytochrome *bd-I*-only membranes. (B) Growth assay of MA against *E. coli* cytochrome *bd-I* only cells. (C) Survival assay of MA against *E. coli* EC958 cytochrome *bd-I* only cells. Dose response data were fitted to three- or four-parameter logistic equations using nonlinear regression (Sigmaplot) to generate IC_{50} values with standard error. Error bars represent standard deviations for at least four technical repeats, including two biological repeats.

mM (Fig. S1B), confirming that MA is bacteriostatic but not bactericidal towards both strains.

We then decided to explore variations upon the MA skeleton which might alter access to proteins on the bacterial inner membrane. For this reason, we examined modified versions. These were based on transformations prominent in the literature for such compounds,^{18,19} enabling easy development in future, and operated on opposite ends of the molecule to provide initial independent evaluation of modification sites. Acetylating the alcohols at positions 2, 3, and 23 gave MA-2, lowering polarity from $ClogP$ of MA at 3.97 to 6.64. Additionally, conversion of the carboxylic acid into an amide using 1,10-diaminodecane to add a hydrophobic chain and introduce a cationic group (likely to interact with bacterial lipids) gave MA-3 ($ClogP = 8.63$). We also produced the 1,10-diaminodecane amide without acetylation (MA-4, $ClogP = 5.96$) to create a double-mutant

cycle. These compounds have been previously studied for anticancer activity, and were synthesised in the same manner, starting from MA.¹⁸ As before, these were docked into the quinol binding site of cytochrome *bd-I* (Fig. 3). Compared with the predicted K_d of 1.14 μM for MA, acetylation resulted in a slightly stronger affinity ($K_d = 0.96 \mu\text{M}$) for MA-2. However, when the long chain amine was also added to give MA-3, the affinity was substantially weakened ($K_d = 14.4 \mu\text{M}$). The chain appears to add too much bulk, and impedes the binding geometry. Removal of some of the molecule's steric bulk in the unacetylated MA-4 resulted in a moderate improvement in K_d (8.64 μM). Importantly, all of these are modelled to bind more strongly than UQ-8, and therefore we could expect inhibitory activity.

Oxygen consumption assays were then performed with *bd-I*-only membranes (Fig. 4a). The tighter predicted binding of MA-2 compared with MA correlated with elevated






Fig. 4 Impact of madecassic acid derivatives upon *E. coli* growth, cytochrome *bd-I* activity, and *E. coli* survival. (A, D and G) Oxygen consumption activities of *E. coli* EC958 cytochrome *bd-I* only membranes exposed to madecassic acid derivatives. (B, E and H) Growth assays *E. coli* cytochrome *bd-I* only cells exposed to madecassic acid derivatives. (C, F and I) Survival assays for *E. coli* EC958 cytochrome *bd-I* only cells exposed to madecassic acid derivatives. Where possible, data were fitted to three- or four-parameter logistic equations using nonlinear regression (Sigmaplot) to generate IC_{50} values (or LC_{50} for killing data) with standard error. Error bars represent standard deviations for at least four technical repeats, including two biological repeats.

increased lipidic interaction could therefore both increase the local concentration of MA derivatives, or conversely result in competitive sequestration within the lipid bilayer, away from the protein. Convolution of these effects with the other mechanisms at play have led to the overall result that MA-4 has the greatest bactericidal activity. The myriad ways in which natural products such as MA can interact with biological macromolecules leads to complications which are hard to avoid – in our case, the possibility that MA might interact with succinate dehydrogenase⁵ was a potential limitation for the isolated membrane components of this study, since the oxygen consumption assay relies upon this complex to maintain the supply of ubiquinol for cytochrome

bd-mediated oxygen consumption. It was therefore necessary to perform additional control experiments to exclude MA-mediated SDH inhibition and further demonstrate the inhibition of cytochrome *bd-I* activity where electron delivery is independent of SDH. To introduce an additional level of rigour to these investigations, an alternative *bd-I* only strain was engineered where an oxidase null mutant strain was complemented with a plasmid encoding the cytochrome *bd* operon (SI). Membranes from this *E. coli* EcoM4 pSU2718G-*cydABX*-his₆ strain were prepared and SDH activity measurements confirmed that 1 mM MA did not inhibit SDH in isolated membranes (Fig. S2). To verify that MA directly targets cytochrome *bd-I* directly, oxygen consumption



experiments were undertaken using the same membranes and duroquinol as the electron donor (delivers electrons directly to *bd-I*). These data (Fig. S3) confirmed that 1 mM MA completely abolishes *bd-I* activity, further supporting the direct binding of MA to *bd-I* in EC958 membranes.

Plants have evolved a certain set of natural products which have a particular role for that plant. Because they interact with the biological machinery in one organism, there is a good chance that they will do something in other organisms, and this has historically led to many successful medicines.²⁰ However, there is no reason to expect natural products to behave exactly as we would wish, and off-target effects are common, since the compounds were selected for a different role. Chemical modification can make a big difference,²¹ and we have shown here that modifications can alter the behaviour of MA in bacterial models, and this provides a route to focus the activity of a natural product into a particular mechanism. Through strategic chemical modification, it should therefore be possible to both better understand the different mechanisms of triterpenoid antibacterial activity, and create molecules which maximise their potency against specific targets. Cytochrome *bd* oxidases are particularly attractive in this regard since they are only found in prokaryotic organisms. Given the versatile modification chemistry open to madecassic acid,^{18,19,22,23} further work to maximise its activity against these proteins would be well justified.

Conclusion

Madecassic acid and three derivatives thereof have been modelled to, and shown to have, a high affinity for cytochrome *bd-I* oxidase, and experimental observations of enzymatic activity, bacterial growth, and bactericidal assays are consistent with this modelling. Alterations of the chemical structure have been shown to influence the biological activity, although the relationships between the different assay outcomes are non-linear. These findings show that inhibition of cytochrome *bd* oxidases is a further component of the antibacterial activity of madecassic acid, in addition to mechanisms already investigated by others. Further strategic chemical modification could result in new antimicrobial compounds based upon madecassic acid or other similar plant-derived compounds, which are specifically targeted towards the unique aspects of bacterial biochemistry.

Author contributions

All authors contributed to writing the manuscript. TTPT extracted and purified the MA from *C. asiatica*. GNS and HICDS performed organic transformations. SAH was involved in planning and performing the microbiological experiments and data analysis. RAB developed the duroquinol assay, CMW engineered the complementation vector, GJ performed the SDH assays and duroquinol experiments, GKR

contributed to the planning of the microbiological experiments and data analysis. CJS and MDG obtained funding and supervised the chemistry. MS also obtained funding and was responsible for all microbiological experimental design and data analysis.

Conflicts of interest

There are no conflicts of interest.

Data availability

Processed data for this paper is included in the main manuscript. Raw data for this paper (molecular characterisation, assay readout) is available from the authors upon request. Supplementary information (SI) is available. See DOI: <https://doi.org/10.1039/d5md01116g>.

Acknowledgements

We thank the UK's Engineering and Physical Science Research Council and Global Challenges Research Fund (EP/T020164/1), the UK's Biotechnology and Biological Sciences Research Council (BB/T008741/1; BB/Z517343/1), the University of Kent, and Vietnam Academy of Science and Technology (VAST) for financial support.

References

- 1 D. Wojnicz, D. Tichaczek-Goska, K. Korzekwa, M. Kicia and A. Hendrich, Anti-Enterococcal Activities of pentacyclic Triterpenes, *Adv. Clin. Exp. Med.*, 2017, **26**(3), 483–490, DOI: [10.17219/acem/62245](https://doi.org/10.17219/acem/62245).
- 2 Z. Sycz, D. Tichaczek-Goska, A. Jezierska-Domaradzka and D. Wojnicz, Are Uropathogenic Bacteria Living in Multispecies Biofilm Susceptible to Active Plant Ingredient—Asiatic Acid?, *Biomolecules*, 2021, **11**(12), 1754, DOI: [10.3390/biom11121754](https://doi.org/10.3390/biom11121754).
- 3 W.-H. Liu, T.-C. Liu and M.-C. Mong, Antibacterial Effects and Action Modes of Asiatic Acid, *Biomedicine*, 2015, **5**(3), 16, DOI: [10.7603/s40681-015-0016-7](https://doi.org/10.7603/s40681-015-0016-7).
- 4 D. Wojnicz, D. Tichaczek-Goska and M. Kicia, pentacyclic Triterpenes Combined with Ciprofloxacin Help to Eradicate the Biofilm Formed in Vitro by *Escherichia coli*, *Indian J. Med. Res.*, 2015, **141**(3), 343–353, DOI: [10.4103/0971-5916.156631](https://doi.org/10.4103/0971-5916.156631).
- 5 C. Wei, P. Cui and X. Liu, Antibacterial Activity and Mechanism of Madecassic Acid against *Staphylococcus aureus*, *Molecules*, 2023, **28**(4), 1895, DOI: [10.3390/molecules28041895](https://doi.org/10.3390/molecules28041895).
- 6 G. Liu, P. Qin, X. Cheng, L. Wu, W. Zhao and W. Gao, Evaluation of the Mechanistic Basis for the Antibacterial Activity of Ursolic Acid against *Staphylococcus aureus*, *Front. Microbiol.*, 2024, **15**, DOI: [10.3389/fmicb.2024.1389242](https://doi.org/10.3389/fmicb.2024.1389242).
- 7 P. Chonsut, C. Romyasamit, A. Konyanee, N. Niyomtham, L. Goodla and A. Mordmuang, Potential Activities of Centella Asiatica Leaf Extract against Pathogenic Bacteria-Associated Biofilms and Its Anti-Inflammatory Effects, *Adv. Pharmacol. Pharm. Sci.*, 2024, **2024**(1), 5959077, DOI: [10.1155/2024/5959077](https://doi.org/10.1155/2024/5959077).



- 8 Z. Xie, X. Dai and L. Wang, Novel 23-Hydroxybetulinic Acid Derivatives: Semi-Synthesis and Antibacterial Activity, *Russ. J. Gen. Chem.*, 2024, **94**(10), 2770–2788, DOI: [10.1134/S1070363224100219](https://doi.org/10.1134/S1070363224100219).
- 9 K. Kandaswamy, S. P. Panda, M. R. Shaik, S. A. Hussain, P. Deepak, N. Thiyagarajulu, D. Jain, A. P. M. Antonyraj, R. Subramanian, A. Guru and J. Arockiaraj, Formulation of Asiatic Acid-Loaded Polymeric Chitosan-Based Hydrogel for Effective MRSA Infection Control and Enhanced Wound Healing in Zebrafish Models, *Int. J. Biol. Macromol.*, 2025, **293**, 137425, DOI: [10.1016/j.ijbiomac.2024.137425](https://doi.org/10.1016/j.ijbiomac.2024.137425).
- 10 V. B. Borisov, R. B. Gennis, J. Hemp and M. I. Verkhovskiy, The Cytochrome *bd* Respiratory Oxygen Reductases, *Biochim. Biophys. Acta, Bioenerg.*, 2011, **1807**(11), 1398–1413, DOI: [10.1016/j.bbabi.2011.06.016](https://doi.org/10.1016/j.bbabi.2011.06.016).
- 11 L. N. Jeffreys, A. Ardrey, T. A. Hafiz, L.-A. Dyer, A. J. Warman, N. Mosallam, G. L. Nixon, N. E. Fisher, W. D. Hong, S. C. Leung, G. Aljayoussi, J. Bibby, D. V. Almeida, P. J. Converse, N. Fotouhi, N. G. Berry, E. L. Nuermberger, A. M. Upton, P. M. O'Neill, S. A. Ward and G. A. Biagini, Identification of 2-Aryl-Quinolone Inhibitors of Cytochrome *Bd* and Chemical Validation of Combination Strategies for Respiratory Inhibitors against *Mycobacterium tuberculosis*, *ACS Infect. Dis.*, 2023, **9**(2), 221–238, DOI: [10.1021/acsinfecdis.2c00283](https://doi.org/10.1021/acsinfecdis.2c00283).
- 12 A. Giuffrè, V. B. Borisov, M. Arese, P. Sarti and E. Forte, Cytochrome *bd* Oxidase and Bacterial Tolerance to Oxidative and Nitrosative Stress, *Biochim. Biophys. Acta, Bioenerg.*, 2014, **1837**(7), 1178–1187, DOI: [10.1016/j.bbabi.2014.01.016](https://doi.org/10.1016/j.bbabi.2014.01.016).
- 13 S. A. Henry, C. M. Webster, L. N. Shaw, N. J. Torres, M.-E. Jobson, B. C. Totzke, J. K. Jackson, J. E. McGreig, M. N. Wass, G. K. Robinson and M. Shepherd, Steroid Drugs Inhibit Bacterial Respiratory Oxidases and Are Lethal Toward Methicillin-Resistant *Staphylococcus aureus*, *J. Infect. Dis.*, 2024, **230**(1), e149–e158, DOI: [10.1093/infdis/jiad540](https://doi.org/10.1093/infdis/jiad540).
- 14 T. T. P. Tran, N. T. Pham, N. X. Nguyen, C. V. Tran, L. V. Tran, L. T. Nguyen, D. T. Nguyen, N. T. N. Huynh, T. L. Nguyen, Y. T. H. Be, M. D. Garrett, C. J. Serpell and T. Van Sung, A Comparative Study of Organic and Conventional Production Methods on Biomass Accumulation and Bioactive Compounds Content in *Centella asiatica* (L.) Urban in Vietnam, *Ind. Crops Prod.*, 2025, **237**, 122251, DOI: [10.1016/j.indcrop.2025.122251](https://doi.org/10.1016/j.indcrop.2025.122251).
- 15 D. Seeliger and B. L. de Groot, Ligand Docking and Binding Site Analysis with PyMOL and Autodock/Vina, *J. Comput.-Aided Mol. Des.*, 2010, **24**(5), 417–422, DOI: [10.1007/s10822-010-9352-6](https://doi.org/10.1007/s10822-010-9352-6).
- 16 M.-H. Nicolas-Chanoine, X. Bertrand and J.-Y. Madec, *Escherichia coli* ST131, an Intriguing Clonal Group, *Clin. Microbiol. Rev.*, 2014, **27**(3), 543–574, DOI: [10.1128/cmr.00125-13](https://doi.org/10.1128/cmr.00125-13).
- 17 B. M. Forde, N. L. B. Zakour, M. Stanton-Cook, M.-D. Phan, M. Totsika, K. M. Peters, K. G. Chan, M. A. Schembri, M. Upton and S. A. Beatson, The Complete Genome Sequence of *Escherichia coli* EC958: A High Quality Reference Sequence for the Globally Disseminated Multidrug Resistant E. Coli O25b:H4-ST131 Clone, *PLoS One*, 2014, **9**(8), e104400, DOI: [10.1371/journal.pone.0104400](https://doi.org/10.1371/journal.pone.0104400).
- 18 T. Van Loc, V. T. Q. Nhu, T. Van Chien, L. T. T. Ha, T. T. P. Thao and T. Van Sung, Synthesis of Madecassic Acid Derivatives and Their Cytotoxic Activity, *Z. Naturforsch., B: J. Chem. Sci.*, 2018, **73**(2), 91–98, DOI: [10.1515/znbs-2017-0172](https://doi.org/10.1515/znbs-2017-0172).
- 19 C. V. Tran, T. T. P. Tran, A. T. Nguyen, L. V. Tran, N. T. Pham, L. T. Nguyen, D. T. Nguyen, M. D. Garrett, N. T. Nguyen, T. T. Do, C. J. Serpell and S. V. Tran, Synthesis and Cytotoxic Activity of Madecassic Acid–Silybin Conjugate Compounds in Liver Cancer Cells, *RSC Med. Chem.*, 2024, **15**, 3418–3432, DOI: [10.1039/D4MD00170B](https://doi.org/10.1039/D4MD00170B).
- 20 D. J. Newman and G. M. Cragg, Natural Products as Sources of New Drugs from 1981 to 2014, *J. Nat. Prod.*, 2016, **79**(3), 629–661, DOI: [10.1021/acs.jnatprod.5b01055](https://doi.org/10.1021/acs.jnatprod.5b01055).
- 21 M. Decker, Hybrid Molecules Incorporating Natural Products: Applications in Cancer Therapy, Neurodegenerative Disorders and Beyond, *Curr. Med. Chem.*, 2011, **18**(10), 1464–1475, DOI: [10.2174/092986711795328355](https://doi.org/10.2174/092986711795328355).
- 22 A. S. C. Valdeira, D. A. Ritt, D. K. Morrison, J. B. McMahon, K. R. Gustafson and J. A. R. Salvador, Synthesis and Biological Evaluation of New Madecassic Acid Derivatives Targeting ERK Cascade Signaling, *Front. Chem.*, 2018, **6**, 434, DOI: [10.3389/fchem.2018.00434](https://doi.org/10.3389/fchem.2018.00434).
- 23 A. S. C. Valdeira, E. Darvishi, G. M. Woldemichael, J. A. Beutler, K. R. Gustafson and J. A. R. Salvador, Madecassic Acid Derivatives as Potential Anticancer Agents: Synthesis and Cytotoxic Evaluation, *J. Nat. Prod.*, 2019, **82**(8), 2094–2105, DOI: [10.1021/acs.jnatprod.8b00864](https://doi.org/10.1021/acs.jnatprod.8b00864).

