

Materials Horizons

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. Das, L. Seethapathy, R. M. C. K. Hazra, J. Samanta and S. Pramanik, *Mater. Horiz.*, 2026, DOI: 10.1039/D5MH02406D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

New Concepts

View Article Online
DOI: 10.1039/D5MH02406D

This study introduces a new conceptual framework in supramolecular chemistry by demonstrating that a traditionally rigid, planar ligand, 2,4,6-tri(4-pyridyl)-1,3,5-triazine (TPT), can undergo pronounced and reversible conformational deformation without metal coordination. Simple *N*-alkylation induces an unprecedented planar-to-bowl transformation of the free ligand, revealing chemically triggered flexibility in an otherwise static scaffold. Importantly, this deformation is reversible: selective guest recognition restores the planar geometry, enabling size-adaptable induced-fit host-guest interactions reminiscent of biological receptors. The conformational switching directly modulates the electronic structure of the system, leading to controlled repositioning of the HOMO–LUMO levels. This feature allows chemical information to be encoded and decoded through donor–acceptor charge-transfer interactions, establishing a platform for molecular-level encryption and decryption. Furthermore, the enhanced electron deficiency of trialkylated TPT enables efficient metal-free oxidation chemistry. Overall, this work establishes chemical stimulus-induced flexibility in a rigid free ligand as a powerful new concept, integrating adaptive recognition, information processing, and reactivity within a single organic system.



Data Availability Statement

View Article Online
DOI: 10.1039/D5MH02406D

The data supporting this article have been included as part of the Supplementary Information (SI).



One System, Three Functions: An Electroactive Reconfigurable Organic Receptor for Adaptive Binding, Information Encoding, and Metal-Free Oxidation

View Article Online
DOI: 10.1039/C6MH02406D

Logeshwari Seethapathy,^a Rohith M,^a Chinmoy K. Hazra,^b Jayanta Samanta,^a Susnata Pramanik^a and Rajorshi Das^{*a}

^aDepartment of Chemistry, College of Engineering and Technology, SRM Institute of Science and Technology, Kattankulathur, Chennai - 603203, India

^bDepartment of Chemistry, Indian Institute of Technology Delhi, Hauz Khas, South Delhi - 110016, India.

*Corresponding author. E-mail: rajorshd@srmist.edu.in, rajorika@gmail.com

Abstract

Living systems rely on biomolecular receptors capable of selectively responding to multiple chemical cues, enabling precise information processing and adaptive functionality. Inspired by this paradigm, we report a multifunctional synthetic receptor that functions in stimuli-responsive conformational switching, programmable information encoding, and metal-free oxidation within a single molecular system. The receptor features a rigid, planar core that undergoes an unexpected and extensive deformation upon chemical alkylation, adopting a bowl-shaped geometry that can be fully reverted to its original flat structure upon guest recognition. This reversible transformation enables adaptive, induced-fit binding of diverse guest molecules, allowing the host to modulate its geometry to optimize donor–acceptor interactions and maximize complex stability. Electrochemical studies show that the resulting cationic species, **1**(PF₆)₃, is highly electron-deficient and serves as an efficient oxidant for metal-free oxidative transformations. Furthermore, the programmable conformational and redox states of the receptor provide a basis for constructing tunable donor–acceptor systems capable of molecular-level information encoding and decoding.

Introduction

Biomolecular receptors are essential components of living systems, processing information through selective responses to chemical signals.¹ They can recognize multiple stimuli and





exhibit induced-fit binding, where the presence of one molecule changes the shape and size of parent biological host influencing the cooperative binding.¹⁻⁴ The working principles of these natural systems provide a foundation for the synthesis of programmable molecular architectures, which can be designed according to the demand. Such systems may generate distinct outputs based on specific input(s), enabling applications in molecular logic,^{5,6} sensing,^{7,8} and catalysis.⁹

One promising approach to achieving such functionality is the use of self-assembled architecture with size-adaptable guest binding. Raymond,¹⁰ Fujita,¹¹⁻¹⁴ Clever,¹⁵⁻¹⁷ Nitschke,¹⁸⁻²¹ Mukherjee^{22,23} and some other groups²⁴⁻²⁷ have reported several supramolecular systems that can alter their architecture in the presence of external guest molecules. For example, Raymond's M_2L_3 helicate cage undergoes a structural transformation to tetrahedron via guest-encapsulation.¹⁰ An $M_{18}L_6$ molecular capsule reported by Fujita can expand its inner core by reassembling its components to a larger capsule $M_{24}L_8$ to accommodate guest molecules inside its cavity.¹³ In a similar fashion, an M_2L_4 cage of Clever converts its structure to an interlocked $[3]_3$ -catenane triggered by excess amount of halide ions.¹⁵ Sun's M_4L_2 metallohelix can also potentially transform its secondary structure to a tertiary protein-like structure after anion binding.²⁷

All the examples described above elaborate on the reassembling of supramolecular structures in the presence of guest molecules, which follows a self-correction pathway, a routine procedure of metal-ligand dissociation and reassociation to generate thermodynamically more stable product(s). In these cases, the rigid ligands used for the construction of supramolecular structures remain rigid, only the reconstitution of metal-ligand self-assembly with a greater number of metal ions and ligands occurs, leading to the formation of larger supramolecular structures. The induction of structural flexibility in a rigid 'free' ligand (where the ligand is not bound to any metal ion forming a constrained supramolecular assembly) through external stimuli is extremely difficult, therefore, it has been unprecedented. However, if possible, it would lead to the generation of novel materials that can not only show induced fit guest-recognition ability, but it may also exhibit other functions including chemical transformations and information security.






Scheme 1. *N*-Alkylation-driven structural and conformational switching in a planar rigid ligand. The first step shows the bowl formation from a planar TPT ligand. The next step elaborates on the conversion of the bowl to planar conformation via guest recognition.

Herein, in an unprecedented reaction, we demonstrate that the rigidity of a ligand can significantly be altered by chemical stimuli without changing the core chemical structure of the rigid ligand. We found that the alkylation of a planar 2,4,6-tri(4-pyridyl)-1,3,5-triazine (TPT) yielded an unusual 'bowl-shaped' trialkylated tripyridiumtriazine derivative $1(\text{PF}_6)_3$, which can regain its planarity upon charge-complementary molecular recognition (Scheme 1). This trialkylation not only imparts remarkable structural flexibility to the TPT molecule but also generates a highly electron-deficient, high-valent cationic system. Notably, the strong electron-accepting nature of triethylated TPT, $1(\text{PF}_6)_3$, makes it a promising candidate for its applications in data encryption/decryption and metal-free oxidation reactions.

Results and Discussion

Induction of chemical flexibility into rigid TPT motif

TPT is a well-known tripodal ligand widely used in supramolecular chemistry for the construction of a variety of self-assembled structures.^{28,29} According to our literature survey, over 300 crystal structures of TPT-metal complexes have been reported to date. Notably, the majority of these complexes exhibit TPT in a planar conformation, consistent with its inherent rigid molecular geometry. However, there is only an isolated example in which the TPT unit




adopts a non-planar configuration. In this rare case, the triazine core displays a measurable deviation from planarity, typically attributed to the severe strain induced by the formation of a rigid supramolecular cage.³⁰

In our approach, structural flexibility was introduced into the otherwise rigid TPT molecule by modulating its electronic environment through chemical and charge-transfer processes, as illustrated in Scheme 1. The tricationic species **1**(PF₆)₃ was synthesized via a straightforward two-step procedure. Initially, TPT was reacted with an excess of ethyl bromide at 50 °C for 48 hours in a sealed tube, resulting in the formation of the tri-N-ethylpyridinium tribromide salt, **1**(Br)₃. Subsequent anion exchange with hexafluorophosphate gave the corresponding hexafluorophosphate salt, **1**(PF₆)₃, in 65% yield as a colorless solid. The product **1**(PF₆)₃ exhibits good solubility in polar solvents such as acetonitrile, dimethylformamide, and dimethylsulfoxide, whereas it remains insoluble in diethyl ether and other non-polar solvents.

The formation of **1**(PF₆)₃ was confirmed by multinuclear NMR spectroscopy, high-resolution electrospray ionization mass spectrometry (HR-ESI MS), single-crystal X-ray diffraction (SCXRD) and cyclic voltametric analysis. In the ¹H NMR spectrum, two peaks were obtained in the aromatic region at $\delta = 9.52$ and 9.48 ppm in DMSO-*d*₆ for pyridinium protons (Fig. S1 in the Supporting Information). The other CH₃ and CH₂ protons appeared at $\delta = 1.68$ and 4.84 ppm, respectively. The ¹³C {¹H} NMR spectrum shows the appearance of the aromatic carbon atoms at $\delta = 169.0$, 148.5 , 146.2 and 127.1 ppm and the alkyl carbon atoms at 57.1 and 16.5 ppm, as expected (Fig. S2). The molecular composition of **1**(PF₆)₃ is further verified by the HR-ESI mass-spectrometry (Fig. S12). The spectrum showed the highest intensity peak for the molecular ion [**1**(PF₆)₂]⁺ with the correct isotopic distribution patterns (signal was obtained at $m/z = 689.1559$ for [**1**(PF₆)₂]⁺ (calcd $m/z = 689.1580$ for C₂₄H₂₇F₁₂P₂N₆).

The formation of the trication **1**(PF₆)₃ is unequivocally confirmed by the SCXRD analysis. Suitable crystals were grown by the slow evaporation of solvent from the concentrated acetonitrile solution of **1**(PF₆)₃. The compound crystallized in the trigonal space group R3m with a three-fold symmetry. The structural analysis reveals an unusual bowl-shaped structure of [**1**]³⁺, unlike the conventional planar form of metal-coordinated TPT. The molecular solid-state structure of [**1**]³⁺ along with its packing motifs are displayed in Fig. 1. The angle between the ethylated pyridinium ring and the central triazine plane measures 22° as depicted in Fig. 1a. The packing structure shows that one tricationic [**1**]³⁺ is in short contact with six nearby



$[1]^{3+}$ (Fig. 1b-d) and fifteen hexafluorophosphate (PF_6^-) ions by $\text{C-H}\cdots\pi$ (2.861 Å) (Fig. 1b,c) and $\text{C-H}\cdots\text{F}$ (3.158 Å, Fig. 1e,f) non-classical hydrogen bonds (C-H \cdots F 2.463 – 2.625 Å).





Fig. 1 The molecular structure of $[1]^{3+}$ in $1(\text{PF}_6)_3$. Hexafluorophosphate (PF_6^-) anions are removed for clarity. The side view of trication $[1]^{3+}$ shows its bowl-shaped structure featuring the positioning of the peripheral pyridinium rings at an angle of 22° . **a)** The angle shows the deviation from planarity, **b** and **c)** side view of the packing structure in stick and space-filling models shows the intermolecular interactions with the neighboring tricationic $[1]^{3+}$, **d)** top view of hexagonal packing of $[1]^{3+}$. Multiple $\text{C-H}\cdots\pi$, $\text{C-H}\cdots\text{F}$, $\text{P-F}\cdots\pi$ and electrostatic interactions of $[1]^{3+}$ with the fifteen neighboring PF_6^- anions in $1(\text{PF}_6)_3$, **e)** side view and **f)** top view, **g)** Possible sites for reductions and **h)** the cyclic voltammogram of $1(\text{PF}_6)_3$. The electrochemical measurements of the complexes were performed at a standard scan rate of 100 mV/s using $^n\text{Bu}_4\text{NPF}_6$ (0.1 M) as supporting electrolyte, Pt wire as working electrode and Fc/Fc^+ as internal standard in dry dimethyl formamide (DMF).

In order to realize the solution-state structure of $1(\text{PF}_6)_3$ and examine the possibility of conformational interconversion in solution, we performed two complementary ^1H NMR experiments. First, concentration-dependent ^1H NMR spectra of $1(\text{PF}_6)_3$ were recorded in DMSO-d_6 (2.5, 10, and 20 mg in 0.5 mL). At lower concentration, two well-resolved pyridinium proton signals are observed. However, at higher concentration, these peaks progressively merge into two broad resonances (Fig. S3). This behaviour suggests



intermolecular interactions at elevated concentration, likely arising from proximity between pyridinium units of neighbouring molecules, which may induce partial distortion of the bowl-shaped conformation and dynamic averaging of the signals.

Second, donor-acceptor titration experiment was conducted by adding the increasing amounts (0.2–1.0 eq.) of pyrene as a donor to a constant concentration of **1**(PF₆)₃. Upon gradual addition (notably around 0.6 eq.), the initially resolved pyridinium signals broaden and eventually coalesce into a single broad peak, consistent with strong π – π /charge-transfer interactions and guest-induced structural reorganization from bowl to a more planar geometry (Fig. S11).


Next, we investigated the electrochemical behaviour of **1**(PF₆)₃ to realize the electron-deficiency caused by the trialkylation of tripyridyltriazine. It is worth mentioning that the triazine is an inherent electron-deficient moiety frequently used as hydrophobic unit to impose electron-deficiency in a system. After the trialkylation, the electron-deficient TPT becomes even more electron-deficient due to the loss of electrons from the three pyridyl units. As expected, the cyclic voltammogram of **1**(PF₆)₃ shows two redox waves in the reduction region. The first wave corresponds to the one-electron reduction of the pyridyl cation that is observed at $E_{1/2} = -790$ mV and the second wave found at $E_{1/2} = -1102$ mV can be attributed to the one-electron reduction of the triazine unit. The cyclic voltametric data along with the square-wave voltammogram are depicted in Fig. 1g,h and Fig. S16. The observed potentials match well with the previously reported alkylated pyridinium compounds.³¹

Charge-complementary molecular recognition and bowl-to-disc conversion

Considering the cationic nature and the strong electron-affinity of **1**(PF₆)₃, we next investigated its guest-recognition ability toward electron-rich guest molecules. The initial screening of guests covers a range of differently sized molecules ranging from two-ring system naphthalene to a seven-ring system coronene molecule listed in Fig. 2a,b.

For all cases, 1.5 equivalents of guests were added to the acetonitrile solution of **1**(PF₆)₃. While the individual solutions of **1**(PF₆)₃ and naphthalene, anthracene, phenanthrene, pyrene or coronene, are colourless or light-yellow, their mixtures gave yellow, green or deep red coloured solutions as shown in Fig. 2c-e. The change of colours from their parent solutions indicates the charge-transfer from the electron-rich polyaromatic hydrocarbon (PAH) guests to the electron-poor tricationic **1**(PF₆)₃.

The formation of host-guest complexes was confirmed by ¹H NMR, UV-vis, infrared (IR) spectroscopy and X-ray crystallographic analysis. Suitable quality single crystals were



obtained by the slow evaporation of a concentrated acetonitrile solution from their host-guest mixtures. Dissolving the co-crystals in DMSO- d_6 , the solutions were subjected to ^1H NMR studies, which show the presence of both donors (neutral guests) and acceptor (cationic host) units confirming the formation of charge-complementary adducts (Fig. 2f and Fig. S4-9 in SI). The NMR spectral analysis also indicated the upfield shifts of the chemical resonances of all the guest molecules (donors) compared to the strong acceptor $\mathbf{1}(\text{PF}_6)_3$. For example, while the aromatic proton resonances of perylene appear at $\delta = 7.55, 7.80$ and 8.37 ppm and the pyridinium protons appear at 9.49 and 9.52 ppm, their donor-acceptor adduct shows the resonances at $\delta = 7.50, 7.75$ and 8.29 (for perylene) as well as 9.48 and 9.50 ppm (for pyridinium protons) in dmsO- d_6 . Similar observation has also been found for pyrene and anthracene molecules (Fig. S4-7).

The observed upfield shifts of the perylene proton signals upon complexation with $\mathbf{1}(\text{PF}_6)_3$ arise from the shielding effects generated by the host molecules, for example, π - π stacking, ring-current effects, and recognition between the two $[\mathbf{1}]^{3+}$. These effects primarily influence the electronic environment of the 'guest' perylene protons, leading to the detectable chemical shift changes. In contrast, the proton resonances of the host $\mathbf{1}(\text{PF}_6)_3$ remain essentially unchanged due to the fast exchange on the NMR time scale. This mainly happens in case of dynamic host-guest equilibrium. The rapid exchange between free and bound states can lead to averaged signals for the host. However, in case of $\mathbf{1}(\text{PF}_6)_3 \cdot \text{pyrene}$, the ^1H NMR spectrum shows a significant shift of the pyridinium protons to the upfield region (Fig. S10,11).

The ^1H NMR spectra also indicate that all the donor molecules exhibit fast exchange kinetics with the acceptor $\mathbf{1}(\text{PF}_6)_3$ on the NMR timescale. To further elucidate the binding stoichiometries and determine the corresponding binding constants, photoluminescence titration experiments were carried out.^{32,33} The binding studies revealed that the binding isotherms of the guests, such as naphthalene, anthracene, and pyrene, were best fitted to a 1:2 stoichiometric model, indicating that one planar donor can recognize up to two molecules of acceptor or vice-versa (Table 1, Table S6–S8, and Fig. S17–S19). For perylene, the binding isotherm shows a 1:1 binding stoichiometry. Owing to the larger and more extended π -conjugated surface of perylene, it can form a stronger and more extensive π - π interaction with a single acceptor unit (Table S9 and Fig. S20).

Table 1: Association constants (M^{-1}) for the complexation of $\mathbf{1}(\text{PF}_6)_3$ with different aromatic guests.[‡]

Entry	Guests	K_{11} (M^{-1})	K_{12} (M^{-1})	NMR kinetics
1	Naphthalene	4476 ± 121	70234 ± 152	Fast
2	Anthracene	10774 ± 28	699 ± 11	Fast



3	Pyrene	299 ± 32	2792 ± 51	Fast	View Article Online
4	Perylene	451 ± 4	Nil	Fast	DOI: 10.1039/D5MH02406D

$\neq K_{11}$ and K_{12} values are by calculated by the global fit V0.5 analysis from fluorescence titrations, using a constant concentration of guest of 0.01 mM and gradual addition of **1**(PF₆)₃ of 5 mM in ACN at 298K.




Fig. 2 Recognition of electron-rich molecules by **1**(PF₆)₃. **a)** Polyaromatic hydrocarbon (PAH) molecules and **b)** electron-rich heteroatom-containing molecules selected for recognition studies. The change of colors demonstrates the charge-transfer from donors to acceptor. Acetonitrile solution of **c)** **1**(PF₆)₃, **d)** all donors (from left): naphthalene, anthracene, phenanthrene, pyrene, perylene, coronene, decamethylferrocene, dibenzothiophene and TTF. **e)** Acetonitrile solutions of the donor•acceptor complexes (from left): **1**(PF₆)₃•naphthalene, **1**(PF₆)₃•anthracene, **1**(PF₆)₃•pyrene, **1**(PF₆)₃•peryrene, **1**(PF₆)₃•coronene, **1**(PF₆)₃•decamethylferrocene,




1(PF₆)₃•dibenzothiophene, **1**(PF₆)₃•TTF. **f**) Monitoring the donor-acceptor interactions by ¹H NMR spectroscopy. The significant upfield shift is observed for perylene protons compared to pyridinium protons of **1**(PF₆)₃.

The recognition of electron-rich molecules by the acceptor **1**(PF₆)₃ was unequivocally confirmed by X-ray crystallographic studies. The diffraction analysis of **1**(PF₆)₃•anthracene shows the presence of one anthracene and one **1**(PF₆)₃ in the asymmetric unit. The anthracene molecule is trapped between the two molecules of triethyltripyrindiniumtriazine units by the charge-complementary electrostatic as well as π - π stacking interactions (Fig. 3a,b and Fig. S13). The distance between the donor and the acceptor is around 3.5 Å (Fig. 3c). The two anthracene molecules separated by one [**1**]³⁺ are lying in two different orientations with an angle of 45° (Fig. 3d). The distance between the two [**1**]³⁺ units measure 7.1 Å as displayed in Fig. 3a.

Interestingly, the ‘bowl’-shaped structure of [**1**]³⁺ got flattened after the recognition of the electron-rich polyaromatic hydrocarbon anthracene molecule. The molecular recognition was driven by the charge-complementary interactions allowing the charge separation from the donor (D) to acceptor (A) components to form an infinite •••D^{d+}-A^{d-}-D^{d+}-A^{d-}-D^{d+}-A^{d-}••• chain by the electrostatic interactions between the anthracene and the tricationic [**1**]³⁺. In principle, the π - π interactions between the donor and acceptor components become strongest while both are planar giving the maximum overlap between them rather than the interaction between a planar and a bowl-shaped species which only give a point interaction. Thus, the overall stability of the charge-transfer complex drives the flattening of the ‘bowl’ shaped **1**(PF₆)₃ to a planar structure in presence of planar guest molecules.

To extend the concept of flattening of [**1**]³⁺ in the presence of planar donors, we also attempted the co-crystallization with other polyaromatic hydrocarbon donor molecules with **1**(PF₆)₃ (Fig. 3). Gratifyingly, the single crystal X-ray diffraction studies with the crystals of **1**(PF₆)₃•phenanthrene and **1**(PF₆)₃•pyrene unequivocally confirm the presence of planar conformation of [**1**]³⁺ trapped between the two phenanthrene or two pyrene molecules (Fig. 3e,h). Unlike the anthracene in **1**(PF₆)₃•anthracene, the orientation of phenanthrene and pyrene molecules are found to be perfectly aligned on the top of each other as depicted in Fig. 3f,g and 3i,j. Intriguingly, all the hexafluorophosphate (PF₆⁻) counter anions in these donor-acceptor adducts are pushed away from the central triazine core due to the strong π - π stacking of PAH



molecules, further elaborating the fact that the noncovalent electrostatic interactions in these systems are much stronger than the halogen (F) $\cdots \pi$ interactions (Fig. S13-15).





Fig. 3 Recognition of electron-rich molecules by $1(\text{PF}_6)_3$. X-ray molecular structures of **a**, **b**, **c** and **d**) $1(\text{PF}_6)_3 \cdot \text{anthracene}$, **e**, **f** and **g**) $1(\text{PF}_6)_3 \cdot \text{phenanthrene}$, **h**, **i** and **j**) $1(\text{PF}_6)_3 \cdot \text{pyrene}$. Donor molecules are displayed in colored, **a-d**) anthracene: cyan and magenta, **e-g**) phenanthrene: purple and **h-j**) pyrene: green. Monitoring the recognition of electron-rich molecules by $1(\text{PF}_6)_3$ via CT complex formation via UV-vis spectroscopy. The spectra of CT complexes are plotted along with their parent donor and acceptor constituents. **k**) $1(\text{PF}_6)_3 \cdot \text{naphthalene}$, **l**) $1(\text{PF}_6)_3 \cdot \text{anthracene}$, **m**) $1(\text{PF}_6)_3 \cdot \text{phenanthrene}$, **n**) $1(\text{PF}_6)_3 \cdot \text{pyrene}$, **o**) $1(\text{PF}_6)_3 \cdot \text{perylene}$, **p**) $1(\text{PF}_6)_3 \cdot \text{coronene}$.



The formation of charge-complementary interaction-driven recognition of electron-rich donors by $\mathbf{1}(\text{PF}_6)_3$ is also evident from the UV-vis spectroscopy. The characteristic colour change associated with CT interaction becomes prominent only at relatively high concentrations. Under these conditions, the absorbance values rapidly approach 5–6 a.u., exceeding the reliable linear detection range of the spectrophotometer and compromising quantitative analysis. At lower concentrations suitable for UV-vis measurements, the CT band is too weak to be distinctly resolved in solution. To address this limitation, we performed solid-state UV-vis measurements, where the donor-acceptor interaction could be clearly detected through the emergence of a CT band in the visible region. Thus, while solution-phase detection is constrained by instrumental limits, the solid-state data provide clear evidence of charge-transfer interaction.

The solid-state UV-vis spectra of all neutral donors and $\mathbf{1}(\text{PF}_6)_3$ and of the D-A adducts from $\mathbf{1}(\text{PF}_6)_3$ •naphthalene to $\mathbf{1}(\text{PF}_6)_3$ •coronene were recorded in an integrating sphere to diminish intensity losses due to the scattering and reflection. As shown in Fig. 3k-p, each of the D-A complexes displays a new band in their UV-vis spectra in addition to their individual bands of the neutral donor and acceptor units. Such bands, often found in the visible-NIR region, are characteristic for the charge-transfer (CT) transition in their D-A adducts, confirming the ability of molecular recognition of electron-rich donors by the newly synthesized $\mathbf{1}(\text{PF}_6)_3$ acceptor.

For example, while the $\mathbf{1}(\text{PF}_6)_3$ and anthracene³⁴ display intense bands at 220 and 300 nm, and 250 and 390 nm, respectively, their charge-complementary adduct $\mathbf{1}(\text{PF}_6)_3$ •anthracene shows discernible broad bands at 380 and at ca. 560 nm. Similarly, phenanthrene³⁵ to coronene molecules can also be successfully recognized by the tripyridiniumtriazine cation $[\mathbf{1}]^{3+}$. Notably, the CT bands for more electron-rich anthracene, pyrene, perylene and coronene are observed at lower energy region from 500-700 nm compared to the lesser electron-rich naphthalene showing its CT band at the relatively higher energy region at $\lambda_{\text{max}} = 440$ nm.

Interestingly, the acceptor $\mathbf{1}(\text{PF}_6)_3$ can be separated easily from its CT adducts by dissolving the D-A crystals in acetonitrile and then running a column chromatography (Scheme 1). The single crystals obtained from the purified $\mathbf{1}(\text{PF}_6)_3$ were further mounted on X-ray diffractometer which showed identical cell parameters, confirming the reversal of bowl-structure of $\mathbf{1}(\text{PF}_6)_3$.



Data encryption/decryption with $\mathbf{1(PF_6)_3}$

View Article Online
DOI: 10.1039/D5MH02406D

The increasing demand for secure information storage and transfer has driven chemists to explore a new frontier at the interface of chemistry, material science and cryptography.^{36–38} Unlike traditional digital encryption, molecular-level approaches harness chemical reactivity, conformational switching, supramolecular recognition, and stimuli-responsive behavior to encode, store, and conceal information in physical matters.^{38,39} These strategies provide unique advantages in non-digital operation, low detectability, and environmental responsiveness, offering fundamentally new platforms for secure communication and anti-counterfeiting technologies.

Since the electron-deficient receptor $\mathbf{1(PF_6)_3}$ undergoes predictable transformations and exhibits distinct color changes in response to chemical stimuli, we explored its potential for molecular-level data encryption and decryption using donor-acceptor-based charge-transfer interactions (Fig. 4). To demonstrate this concept, a piece of paper was encoded with the word ‘CHEMIST’ by writing with an acetonitrile solution of $\mathbf{1(PF_6)_3}$ (Fig. 4b). Because $\mathbf{1(PF_6)_3}$ is colorless, the message became completely invisible after drying for a few minutes, resulting in an unrecognizable white sheet (Fig. 4c). Spraying the surface with an acetonitrile solution of pyrene, a suitable electron donor that rapidly forms a visible charge-transfer complex with $\mathbf{1(PF_6)_3}$, revealed the hidden message (Fig. 4d,e). The decoded information could then be erased by gently washing the sheet with acetonitrile, which removed the donor and restored the blank appearance (Fig. 4f,g). Notably, this gentle washing step leaves the acceptor embedded within the paper, enabling regeneration of the message upon reapplication of the pyrene spray (Fig. 4h,i). Because the donor-acceptor binding is relatively weak, gentle washing with acetonitrile preferentially removes the externally applied donor, which is more loosely associated and more readily solvated. In contrast, the pre-deposited and dried $\mathbf{1(PF_6)_3}$ remains largely retained on the paper substrate due to its stronger interactions with the paper material (cellulose matrix), enabling the acceptor to persist under mild washing conditions. For complete removal of the encrypted data, the sheet can be washed two to three times to remove the residual acceptor from the sheet of paper.






Fig. 4. Demonstration of molecular data encryption and decryption using the electron-deficient receptor $\mathbf{1}(\text{PF}_6)_3$ and pyrene. **a)** Plain paper before encoding, **b)** writing the message ‘CHEMIST’ using an acetonitrile solution of $\mathbf{1}(\text{PF}_6)_3$, **c)** the message becomes invisible after drying, yielding an apparently blank sheet, **d,e)** decryption of the hidden message upon spraying with a pyrene solution, producing a visible donor–acceptor charge-transfer color, **f,g)** erasure of the revealed message by washing with acetonitrile to remove the donor, regenerating a blank sheet, **h,i)** re-encryption/decryption cycle: the message can be revealed again by reapplying pyrene spray.

This reversible process elaborated above highlights the potential of $\mathbf{1}(\text{PF}_6)_3$ for recyclable, stimuli-responsive, molecular-level information security. Interestingly, this method can also be employed for its application in rewritable printing technology. Fabricating the paper containing similar $\mathbf{1}(\text{PF}_6)_3$ acceptor-like molecules may serve as rewritable page, where pyrene solution can be used as ink.

$\mathbf{1}(\text{PF}_6)_3$ as Organo-Oxidant

Inspired by the results of charge-complementary molecular recognition of planar electron-rich PAH molecules by the electron-deficient $\mathbf{1}(\text{PF}_6)_3$, we next attempted its ability to fully oxidize electron-rich metal complex like decamethylferrocene (DMFC) as shown in Fig 5a. In principle, for a successful oxidation, the oxidation potential of a compound must be either equal to, less than or at least close to the reduction potential of an oxidant. In case of DMFC, the $\text{Fe}^{2+/3+}$ oxidation process is found to be at -550 mV against ferrocene/ferrocenium standard in dichloromethane,^{40–42} which is lower than the first reduction potential of $\mathbf{1}(\text{PF}_6)_3$, observed at



$E_{1/2}^{3+/0} = -290$ mV in acetonitrile. Since the redox potential strictly depends on the solvents used for measurement, it can be assumed that the tricationic $\mathbf{1}(\text{PF}_6)_3$ can potentially oxidize DMFC in acetonitrile. As expected, while the colourless $\mathbf{1}(\text{PF}_6)_3$ was added into the orange acetonitrile solution of DMFC, an immediate formation of a green solution was observed (Fig. 2d,e). The color change confirms the formation of decamethylferrocenium cation (DMFC^+).⁴³ The *in-situ* oxidation of DMFC can also be monitored by UV-vis spectroscopy. The pristine DMFC shows a band at $\lambda = 222$ nm. Upon gradual addition of the acceptor $\mathbf{1}(\text{PF}_6)_3$ a new band near 252 nm appears along with the rise of 222 nm band (Fig. 5c). The oxidation of DMFC is also verified with the addition of silver hexafluorophosphate ($E_{\text{Ag}/\text{Ag}^+} = 40$ mV in acetonitrile vs Fc/Fc^+ redox couple), which shows the identical result to that of $\mathbf{1}(\text{PF}_6)_3$ addition (Fig. S22, S23).




Fig. 5. a) Oxidation of decamethylferrocene by $\mathbf{1}(\text{PF}_6)_3$. b) Molecular structure of oxidized product decamethylferrocenium hexafluorophosphate obtained from SCXRD study. c) The titration of decamethylferrocene with $\mathbf{1}(\text{PF}_6)_3$ monitored by UV-vis spectroscopy ($[\text{DMFC}] = 10^{-5}$ M, $[\mathbf{1}(\text{PF}_6)_3] = 10^{-3}$ M). The inset figure shows the colour change from light orange to green occurred due to the oxidation.

After slow evaporation at ambient temperature, the resulted solution yielded green rod-shaped crystals which after X-ray crystallographic study confirmed the formation of decamethylferrocenium cation $[\text{Me}_5\text{C}_5\text{Fe}^{\text{III}}\text{C}_5\text{Me}_5]^+$, the one-electron oxidized product of parent DMFC. The molecular structure of decamethylferrocenium hexafluorophosphate is displayed in Fig. 5b.

It is noteworthy to mention that such oxidation has previously been performed only with strong oxidants, for example, cerium(IV) ammonium nitrate (CAN), ferric chloride (FeCl_3) or silver cation (Ag^+).⁴⁰ However, in our case, the oxidation can be done purely in metal-free condition.



Theoretical Calculations

The structural deformation of the rigid-planar TPT induced by trialkylation was analyzed by density functional theory (DFT) calculations. We first calculated the effect of alkylation on the TPT moiety. The highest occupied molecular orbitals (HOMOs) and the lowest unoccupied molecular orbitals (LUMOs) of TPT and triethylated TPT $\mathbf{1}(\text{PF}_6)_3$ were illustrated in Fig. 6a. The calculation shows a significant change of HOMOs on the TPT moiety after alkylation, while the LUMO remains almost unchanged. This is obvious due to the fact that the electron-rich pyridine ring with one nonbonding lone pair of electrons becomes electron-poor after alkylation. Therefore, while the HOMOs of TPT reside on the pyridine ring, the alkylation yielded a total shift from pyridinium ring to comparatively electron-rich triazine core. This theoretical HOMO-LUMO observation is also supported by the Hirschfeld plot (Fig. 6b).




Fig. 6. a) Contour plots of the calculated HOMO and LUMO of TPT and $[1]^{3+}$. Calculations were performed using b3lyp/6-311+g(d,p) level of theory. b) Hirschfeld plot of TPT and trialkylTPT \cdot 3PF $_6$. c) Energy profile diagram showing the energy difference between the planar and bowl conformations of $[1]^{3+}$ in $\mathbf{1}(\text{PF}_6)_3$ using the identical function and basis set b3lyp/6-311+g(d,p) (opt freq, solvent: acetonitrile).

Notably, the geometry-optimized structure of triethyltrispyridiniumtriazine $[1]^{3+}$ gave a perfectly planar conformation, which contrasts with the isolated ‘bowl’-shaped structure of $[1]^{3+}$ (Fig. 1, *vide supra*), obtained from the single-crystal XRD analysis. The reason behind the stabilization of bowl-structure is not very clear, therefore, we performed the single-point energy calculation using the identical functional and the basis set to realize the difference in energy between the two conformations. The calculation revealed that the ground state energy of the bowl structure is 0.05743124 HF unit (~ 36 kcal/mole, HF = Hartree-Fock) higher than the planar structure (Fig. 6c). This energy difference falls perfectly in line with the bending



vibration (elastic molecules) for such a moderately sized cationic species with four rigid and planar six-membered rings.^{44–46} However, we assume that the uneven hydrophobic interactions between the cationic $[1]^{3+}$ and fifteen counter anions (PF_6^-) play a significant influence for its stabilization in the bowl form (Fig. 1e,f).


Conclusion

We have described the design and synthesis of a stimuli-responsive molecular system that exhibits a unique chemically induced structural flexibility on a rigid ligand scaffold. Unlike traditional supramolecular systems where guest-induced transformation occurs through metal-ligand reassembly, this work showcases a rare example of conformational switching in a ‘free’ ligand 2,4,6-tri(4-pyridyl)-1,3,5-triazine (TPT) *via* *N*-alkylation. The resultant trialkylated tricationic ligand, $1(\text{PF}_6)_3$, adopts a bowl-shaped conformation, which flattens upon binding planar electron-rich guests, enabled by charge-transfer and π – π interactions. This bowl-to-disc conversion opens new possibilities for dynamic host-guest systems and redox-tunable molecular devices. This structural adaptability is particularly important in supramolecular systems, leading to emergent properties such as all-or-nothing templation effects and varying degrees of cooperativity.^{1,47–50} The binding of different guest molecules to the host can either enhance or diminish the effective interactions between them, allowing for precise control over molecular recognition.

Furthermore, the pronounced electron deficiency of $1(\text{PF}_6)_3$ enables its use as a selective organo-oxidant, offering an efficient metal-free platform for oxidative transformations. Its performance in encryption-decryption experiments demonstrates the receptor’s capability for recyclable, stimuli-responsive, molecular-level information security. The rapid and reversible generation of charge-transfer signals, combined with the intrinsic chemical reconfigurability of the system, positions $1(\text{PF}_6)_3$ as a highly versatile molecular device for secure data storage, visual encoding, and emerging stimuli-gated information technologies.

Author contributions

Conceptualization, supervision and original draft preparation: Rajorshi Das, Investigation and data curation: Logeshwari Seethapathy and Rohith M, Experimental support: Chinmoy K. Hazra, Crystallographic analysis: Jayanta Samanta and Rajorshi Das, Theoretical studies: Susnata Pramanik and Logeshwari Seethapathy, Writing – review and editing: all authors.



Conflicts of interest

There are no conflicts to declare.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgements

The authors gratefully acknowledge SRM Institute of Science and Technology (SRMIST) for the funding support under the Selective Excellence Research Initiative (SERI) (Grant no. SRMIST/R/AR(A/SERI2024/174/17–342). The authors also thank SRM-Central Instrumentation Facility (SCIF) and SRM Nanotechnology Research Centre (NRC) for providing essential instrumentation facilities. L. S. and R. M. are grateful to SRMIST for their doctoral fellowship. The authors also thank Dr. S. Anandhakumar and Mr. Abilash S. for the measurement of electrochemical data.

References

- 1 D. D. Boehr, R. Nussinov and P. E. Wright, *Nat. Chem. Biol.*, 2009, **5**, 789–796.
- 2 H. R. Bosshard, *Physiology*, 2001, **16**, 171–173.
- 3 M. Nocker and P. Cozzini, *Curr. Top. Med. Chem.*, 2011, **11**, 133–147.
- 4 K. A. Wreggett and J. W. Wells, *J. Biol. Chem.*, 1995, **270**, 22488–22499.
- 5 S. Erbas-Cakmak, S. Kolemen, A. C. Sedgwick, T. Gunnlaugsson, T. D. James, J. Yoon and E. U. Akkaya, *Chem. Soc. Rev.*, 2018, **47**, 2228–2248.
- 6 C. Emanuelson, A. Bardhan and A. Deiters, *ACS Synth. Biol.*, 2024, **13**, 538–545.
- 7 J. Krämer, R. Kang, L. M. Grimm, L. De Cola, P. Picchetti and F. Biedermann, *Chem. Rev.*, 2022, **122**, 3459–3636.
- 8 A. Patel and P. Peralta-Yahya, *Biochemistry*, 2023, **62**, 187–195.
- 9 K.-Y. Wang, J. Zhang, Y.-C. Hsu, H. Lin, Z. Han, J. Pang, Z. Yang, R.-R. Liang, W. Shi and H.-C. Zhou, *Chem. Rev.*, 2023, **123**, 5347–5420.
- 10 M. Scherer, D. L. Caulder, D. W. Johnson and K. N. Raymond, *Angew. Chem. Int. Ed.*, 1999, **38**, 1587–1592.
- 11 K. Kumazawa, Y. Yamanoi, M. Yoshizawa, T. Kusukawa and M. Fujita, *Angew. Chem. Int. Ed.*, 2004, **43**, 5936–5940.
- 12 T. Sawada, H. Hisada and M. Fujita, *J. Am. Chem. Soc.*, 2014, **136**, 4449–4451.
- 13 S. Wang, T. Sawada, K. Ohara, K. Yamaguchi and M. Fujita, *Angew. Chem. Int. Ed.*, 2016, **55**, 2063–2066.
- 14 K. Iizuka, H. Takezawa and M. Fujita, *J. Am. Chem. Soc.*, 2024, **146**, 32311–32316.
- 15 R. Zhu, J. Lübben, B. Dittrich and G. H. Clever, *Angew. Chem. Int. Ed.*, 2015, **54**, 2796–2800.
- 16 A. Walther, I. Regeni, J. J. Holstein and G. H. Clever, *J. Am. Chem. Soc.*, 2023, **145**, 25365–25371.



- 17S. Sudan, F. Fadaei-Tirani, R. Scopelliti, K. E. Ebbert, G. H. Clever and K. Severin, *Angew. Chem. Int. Ed.*, 2022, **61**, e202201823. View Article Online
DOI: 10.1039/D3MH02406D
- 18I. A. Riddell, M. M. J. Smulders, J. K. Clegg, Y. R. Hristova, B. Breiner, J. D. Thoburn and J. R. Nitschke, *Nat. Chem.*, 2012, **4**, 751–756.
- 19D. M. Wood, W. Meng, T. K. Ronson, A. R. Stefankiewicz, J. K. M. Sanders and J. R. Nitschke, *Angew. Chem. Int. Ed.*, 2015, **54**, 3988–3992.
- 20C. T. McTernan, J. A. Davies and J. R. Nitschke, *Chem. Rev.*, 2022, **122**, 10393–10437.
- 21F. J. Rizzuto and J. R. Nitschke, *Nat. Chem.*, 2017, **9**, 903–908.
- 22R. Banerjee, P. Bhandari, N. Hickey and P. S. Mukherjee, *J. Am. Chem. Soc.*, 2025, **147**, 23049–23059.
- 23M. Aggarwal, R. Banerjee, N. Hickey and P. S. Mukherjee, *Angew. Chem. Int. Ed.*, 2024, **63**, e202411513.
- 24H. Lee, P. Elumalai, N. Singh, H. Kim, S. U. Lee and K.-W. Chi, *J. Am. Chem. Soc.*, 2015, **137**, 4674–4677.
- 25L.-X. Cai, Y.-H. Hu, L.-P. Zhou, P.-M. Cheng, X.-Q. Guo, Y.-T. Chan and Q.-F. Sun, *Chem. Sci.*, 2025, **16**, 7956–7962.
- 26A. V. Virovets, E. Peresyphkina and M. Scheer, *Chem. Rev.*, 2021, **121**, 14485–14554.
- 27X.-Q. Guo, L.-P. Zhou, S.-J. Hu, L.-X. Cai, P.-M. Cheng and Q.-F. Sun, *J. Am. Chem. Soc.*, 2021, **143**, 6202–6210.
- 28M. Fujita, M. Tominaga, A. Hori and B. Therrien, *Acc. Chem. Res.*, 2005, **38**, 369–378.
- 29R. Chakrabarty, P. S. Mukherjee and P. J. Stang, *Chem. Rev.*, 2011, **111**, 6810–6918.
- 30N. Hafezi, J. M. Holcroft, K. J. Hartlieb, E. J. Dale, N. A. Vermeulen, C. L. Stern, A. A. Sarjeant and J. F. Stoddart, *Angew. Chem. Int. Ed.*, 2015, **54**, 456–461.
- 31L.-X. Cai, S.-C. Li, D.-N. Yan, L.-P. Zhou, F. Guo and Q.-F. Sun, *J. Am. Chem. Soc.*, 2018, **140**, 4869–4876.
- 32J. Samanta and R. Natarajan, *Org. Lett.*, 2016, **18**, 3394–3397.
- 33S. Ibáñez, P. Salvà, L. N. Dawe and E. Peris, *Angew. Chem. Int. Ed.*, 2024, **63**, e202318829.
- 34R. Taheri-Ledari, J. Rahimi and A. Maleki, *Mater. Res. Express*, 2020, **7**, 015067.
- 35M. R. M. Izawa, D. M. Applin, L. Norman and E. A. Cloutis, *Icarus*, 2014, **237**, 159–181.
- 36H. Zhang, Q. Li, Y. Yang, X. Ji and J. L. Sessler, *J. Am. Chem. Soc.*, 2021, **143**, 18635–18642.
- 37S. D. Dahlhauser, C. D. Wight, S. R. Moor, R. A. Scanga, P. Ngo, J. T. York, M. S. Vera, K. J. Blake, I. M. Riddington, J. F. Reuther and E. V. Anslyn, *ACS Cent. Sci.*, 2022, **8**, 1125–1133.
- 38Y. Shen, X. Le, Y. Wu and T. Chen, *Chem. Soc. Rev.*, 2024, **53**, 606–623.
- 39W. Zhong, J. Zhang, Y. Lin, S. Li, Y. Yang, W.-J. Wang, C. Si, F. E. Kühn, Z. Zhao, X.-M. Cai and B. Z. Tang, *Chem. Sci.*, 2024, **15**, 3920–3927.
- 40N. G. Connelly and W. E. Geiger, *Chem. Rev.*, 1996, **96**, 877–910.
- 41R. Das, M. Linseis, S. M. Schupp, L. Schmidt-Mende and R. F. Winter, *Chem. Eur. J.*, 2022, **28**, e202104403.
- 42R. Das, M. Linseis, S. Scheerer, K. Zoller, L. Senft, I. Ivanović-Burmazović and R. F. Winter, *Inorg. Chem.*, 2022, **61**, 12662–12677.
- 43B. Su, R. P. Nia, F. Li, M. Hojeij, M. Prudent, C. Corminboeuf, Z. Samec and H. H. Girault, *Angew. Chem. Int. Ed.*, 2008, **47**, 4675–4678.
- 44B. Zhu, B. Cheng, L. Zhang and J. Yu, *Carbon Energy*, 2019, **1**, 32–56.
- 45V. Mehmeti and M. Sadiku, *Computation*, 2022, **10**, 68.
- 46S. Ehrlich, J. Moellmann and S. Grimme, *Acc. Chem. Res.*, 2013, **46**, 916–926.
- 47U. B. Choi, H. Sanabria, T. Smirnova, M. E. Bowen and K. R. Weninger, *Biomolecules*, DOI:10.3390/biom9030114.
- 48E. Lopez-Fontal, L. Milanesi and S. Tomas, *Chem. Sci.*, 2016, **7**, 4468–4475.



- 49 S. Whitlam, C. Rogers, A. Pasqua, C. Paavola, J. Trent and P. L. Geissler, *Nano Lett.*, 2009, **9**, 292–297. Article Online
DOI: 10.1039/D3MH02406D
- 50 T. M. Nguyen and T. N. Truong, *npj Comput Mater*, 2025, **11**, 372.

