



Cite this: *Mater. Horiz.*, 2026, 13, 3127

DOI: 10.1039/d6mh90015a

rsc.li/materials-horizons

Correction: Moderately polarized carborane-MOF with inverse C₂ selectivity for one-step polymer-grade ethylene purification

Changhong Liu,^a Guangzu Xiong,^a Yuzhe Wang,^c Wenlei Yang,^a Yiwen Yuan,^a Hui Wang,^a Lingyao Wang^{*a} and Yuanbin Zhang^{*ab}

Correction for 'Moderately polarized carborane-MOF with inverse C₂ selectivity for one-step polymer-grade ethylene purification' by Changhong Liu *et al.*, *Mater. Horiz.*, 2026, **13**, 473–479, <https://doi.org/10.1039/D5MH01641J>.

The authors regret that an incorrect ratio of C₂H₂/C₂H₄/C₂H₆ was used to generate the data shown in Fig. 5b in the published article. The corrected version of Fig. 5 is shown in this notice. This correction does not affect any of the conclusions of the article.




Fig. 5 CB-Zn-DPG breakthrough curves for (a) a C₂H₄/C₂H₆ (50/50) mixture and (b) a C₂H₂/C₂H₄/C₂H₆ (1/90/9) mixture at 1 mL min⁻¹.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Key Laboratory of the Ministry of Education for Advanced Catalysis Materials, College of Chemistry and Materials Sciences, Zhejiang Normal University, Jinhua 321004, China. E-mail: lywang@zjnu.edu.cn, ybzhang@zjnu.edu.cn

^b Zhejiang Key Laboratory of Advanced Catalysis and Adsorption Materials, Zhejiang Normal University, Jinhua 321004, China

^c Department of Chemical and Environmental Engineering, University of Nottingham Ningbo China, Ningbo, 315100, China

