

CORRECTION

[View Article Online](#)
[View Journal](#)

Cite this: DOI: 10.1039/d6mh90059c

Correction: A silk fibroin-based hydrogel desensitizer achieving 660 μm dentin tubule occlusion for dentin hypersensitivity treatmentKuangdi Xin,^{ab} Wanshan Gao,^{abcd} Yulu Xie,^a Xiaohua Dong,^a Lingshan Ran,^a Tianyi Xia,^b Jing Xie,^b Jiaojiao Yang,^c Tao Hu,^{cd} Jianshu Li,^{bc} Xiaoyu Huang,^{*e} Rongmin Qiu^{*a} and Jun Luo^{*b}

DOI: 10.1039/d6mh90059c

rsc.li/materials-horizonsCorrection for 'A silk fibroin-based hydrogel desensitizer achieving 660 μm dentin tubule occlusion for dentin hypersensitivity treatment' by Kuangdi Xin *et al.*, *Mater. Horiz.*, 2025, **12**, 4177–4192, <https://doi.org/10.1039/D4MH01927J>.

The authors regret the inadvertent inclusion of an incorrect image as Fig. 5E2 in the published article. The corrected Fig. 5 is shown in this notice. The authors confirm that this does not affect any results, data interpretation or conclusions of the published article.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a College of Stomatology, Hospital of Stomatology Guangxi Medical University, Guangxi Key Laboratory of Oral and Maxillofacial Rehabilitation and Reconstruction, Guangxi Clinical Research Center for Craniofacial Deformity, Guangxi Key Laboratory of Oral and Maxillofacial Surgery Disease Treatment, Guangxi Health Commission Key Laboratory of Prevention and Treatment for Oral Infectious Diseases, Nanning 530021, P. R. China. E-mail: rongminqiu@hotmail.com

^b College of Polymer Science and Engineering, State Key Laboratory of Advanced Polymer Materials, Sichuan University, Chengdu, 610065, P. R. China. E-mail: luojuncd@scu.edu.cn

^c State Key Laboratory of Oral Diseases, West China Hospital of Stomatology, Sichuan University, Chengdu 610041, P. R. China

^d National Center for Stomatology, National Clinical Research Center for Oral Diseases, Frontier Innovation Center for Dental Medicine Plus, Sichuan University, Chengdu 610041, P. R. China

^e Department of Preventive Dentistry, School and Hospital of Stomatology, Fujian Medical University, Fuzhou, Fujian Province, P. R. China. E-mail: hxy@fjmu.edu.cn






Fig. 5 Long-term assessment of DT occlusion efficacy by STU. (A) Schematic illustration of the pH cycling assessment of dentin slices. The model involved 8 h of exposure to artificial saliva (AS, pH 7.0) followed by 16 h in a demineralization solution (DS, pH 4.8) each day, in the subgroup, the demineralization solution was replaced with 2% EDTA solution. (B) Air pressure changes in the airtightness test over time. The error bars represent the mean \pm SD for $n = 5$, *** $p < 0.001$, **** $p < 0.0001$. (C)–(E) SEM images of dentin slices after ST or STU hydrogel treatment, with or without the brushing challenge. (C1) and (C3) Surface morphology of ST/STU-treated dentin; (C2) and (C4) Surface morphology of ST/STU-treated dentin with brushing; (D1) and (D3) Cross-sectional morphology of ST/STU-treated dentin; (D2) and (D4) Cross-sectional morphology of ST/STU-treated dentin with brushing; (E1) and (E2) Surface and cross-sectional morphology of brushed dentin treated with ST. (E3) and (E4) Surface and cross-sectional morphology of brushed dentin treated with STU.

