

RSC Mechanochemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. Irfan, P. Caboni, A. Porcheddu and F. Basoccu, *RSC Mechanochem.*, 2026, DOI: 10.1039/D5MR00141B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Reimagining Sulfohydroxamic Acid Formation through Mechanochemical Solid-State Activation

Mohammad Irfan,^{†a} Pietro Caboni,^a Andrea Porcheddu,^{*a} and Francesco Basoccu^{†a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Sulfohydroxamic acids ($-\text{SO}_2-\text{NHOH}$), including classical Piloty's acid derivatives, are key NO/HNO donors and valuable building blocks, but their synthesis still relies on hazardous hydroxylamine solutions and solvent-intensive protocols. Here we report a simple, solvent-free mechanochemical route based on milling sulfonyl chlorides with hydroxylamine hydrochloride in the presence of an inorganic base. Magnesium oxide (MgO) emerges as an ideal promoter, combining sufficient basicity with low nucleophilicity to release hydroxylamine in situ while preserving both the sulfonyl chloride and the sulfohydroxamic product. Under optimised conditions, a broad range of aromatic sulfonyl chlorides bearing electron-donating and electron-withdrawing substituents are converted into the corresponding sulfohydroxamic acids. The method efficiently produces sulfohydroxamic acids with yields ranging from 18% to 81% under mild conditions and is adaptable to standard milling equipment, enabling scalable synthesis. The protocol can be readily extended to O- and N,O-substituted hydroxylamine derivatives by using appropriately functionalized hydroxylamine salts. This mechanochemical strategy drastically reduces solvent consumption and streamlines work-up, providing a safer, cleaner, and scalable approach to sulfohydroxamic acids and their NO-donor analogues.

Introduction

Sulfohydroxamic acids constitute a unique and enduring class of compounds that have achieved significant status at the confluence of synthetic, medicinal, and biological chemistry.¹⁻⁶ Their structural motif, defined by the N–O–S linkage, imparts a remarkable combination of stability and reactivity, facilitating their participation in diverse chemical transformations and controlled decomposition into bioactive nitrogen species.⁶⁻¹⁰

The prototypical member of this family, benzenesulfohydroxamic acid, commonly known as Piloty's acid, was first reported by Oskar Piloty in the late 19th century.¹¹ His groundbreaking research not only introduced a novel functional group into the chemist's repertoire but also provided a straightforward synthetic framework that remains the primary method for producing these derivatives over a century later. Piloty's initial synthesis, which involved reacting a sulfonyl chloride dissolved in methanol with an aqueous solution of hydroxylamine, was sophisticated and effective. This transformation elucidates the fundamental reactivity of sulfonyl electrophiles with nitrogen nucleophiles, culminating in a stable N–O–S linkage amenable to further manipulation under

mild conditions. The simplicity and broad applicability of this method have facilitated its swift adoption and adaptation, catalysing extensive research into the reactivity and coordination behavior of sulfohydroxamic acids and their derivatives (Figure 1).¹²⁻¹⁶ Over the subsequent decades, this class of compounds has become integral to both synthetic methodology and biological chemistry, owing to their ability to serve as donors of reactive nitrogen species, such as nitric oxide (NO) and nitroxyl (HNO).^{13, 17-19}

The biological significance of these two species is profound. Nitric oxide, a small, diffusible free radical, is integral to numerous physiological processes, including vasodilation, neurotransmission, and immune modulation.²⁰ Consequently, the controlled release of NO from chemical precursors, such as sulfohydroxamic acids, has attracted sustained interest from medicinal chemists seeking reliable and tunable NO donors for therapeutic and diagnostic applications. In contrast, the one-electron-reduced form of NO, HNO, demonstrates a distinct and complementary reactivity profile, particularly in redox and metalloprotein chemistry.²⁰⁻²²

Sulfohydroxamic acids capable of generating either NO or HNO under mild conditions have thus emerged as potent chemical tools for investigating biological redox processes and as versatile intermediates for the synthesis of nitrogen-containing compounds.¹⁸

*Corresponding authors

^aDipartimento di Scienze Chimiche e Geologiche, Università degli Studi di Cagliari, Cittadella Universitaria, S5554 bivio per Sestu, 09042-Monserrato, CA, Italy
E-mail: porcheddu@unica.it

†F.B. and M.I. contributed equally to this study.






Figure 1. Different uses of sulfohydroxamic acids in scientific fields.

Despite its enduring significance, the synthesis of sulfohydroxamic acids has undergone minimal evolution since Piloty's era.^{8, 13, 23, 24} The traditional solution-phase conversion of sulfonyl chlorides with hydroxylamine remains the predominant methodology; however, it suffers from several inherent limitations. The high reactivity and limited stability of both reagents and products frequently result in competing side reactions, leading to reduced yields and complex purification processes.^{25–27} Furthermore, hydroxylamine presents significant safety concerns owing to its volatility, toxicity, and propensity for explosive decomposition, particularly in concentrated or heated solutions.^{28–32} These hazards are amplified when reactions are scaled up, posing challenges to industrial applications.³² Even contemporary modifications of the original method, such as those utilizing mixed solvent systems (e.g., MeOH–H₂O–THF) and solid bases like magnesium oxide, have not completely overcome these challenges. They continue to rely on substantial volumes of organic solvents, generate considerable waste, and necessitate the careful handling of hazardous reagents, all of which contravene the principles of green and sustainable chemistry.^{33–35}

In response to these challenges, mechanochemistry has emerged as a highly promising alternative platform for the synthesis of sulfohydroxamic acids. Mechanochemistry is characterized by the utilization of mechanical energy, typically through grinding, milling, or shearing, to drive chemical reactions, signifying a genuine shift in synthetic philosophy.^{36–41} Unlike conventional solution-based approaches, mechanochemical methods operate under solvent-free or solvent-minimized conditions, significantly reducing chemical waste generation and enhancing the overall process safety.^{42–49} Beyond these ecological advantages, mechanochemical reactions often exhibit accelerated reaction rates, improved selectivity, and distinct mechanistic behaviours compared to their solution-phase counterparts. In numerous instances, novel products or reactivity patterns emerge that are otherwise inaccessible in traditional media, highlighting the fundamentally unique solid-state environment.^{8, 50–56}

Mechanochemistry has already demonstrated a transformative impact in diverse fields, including organic synthesis, coordination chemistry, and materials science.^{57–63} Its advantages, such as minimal solvent use, simplified work-up, and often superior atom economy, render it particularly attractive for the preparation of sensitive or unstable compounds. These features are especially valuable in the context of sulfohydroxamic acids due to the inherent instability of both hydroxylamine and sulfohydroxamic intermediates

under aqueous or alcoholic conditions suggests that a solvent-free or solid-assisted approach could significantly enhance both safety and efficiency. Furthermore, the mechanical activation of sulfonyl chlorides in the presence of solid hydroxylamine salts offers the prospect of controlled and selective conversion under mild conditions, eliminating the need for hazardous liquid reagents and cumbersome solvent systems.

Building on our previous work on the synthesis of hydroxamic acids, we sought to extend this methodology to the preparation of the corresponding sulfonic derivatives by identifying an alternative to carbonyl diimidazole (CDI).³⁶ The use of CDI proved unsuccessful, as the higher acidity of sulfonic acids, together with the intrinsic instability of the sulfonyl intermediate, resulted in no productive activation and led to the exclusive recovery of the starting material. Therefore, we opted for the direct milling of sulfonyl chlorides with hydroxylamine hydrochloride in the presence of a benign inorganic base, such as magnesium oxide, enables the clean and selective formation of sulfohydroxamic acids at room temperature. Importantly, this method not only enhances laboratory safety and practicality but also exemplifies how mechanochemical activation can fundamentally reshape established reactivities. By bridging Piloty's pioneering work on sulfohydroxamic acids with the principles of contemporary green chemistry, the present methodology redefines a century-old reaction through the lens of safety and environmental stewardship. The results underscore the capacity of solid-state reactivity to deliver practical, scalable, and inherently safer synthetic routes to valuable NO-donor compounds, thereby advancing both synthetic methodology and medicinal chemistry applications.

Results and Discussion

The initial approach for synthesizing sulfohydroxamic acids involved using equimolar quantities of *p*-tosyl chloride (**1j**) and hydroxylamine hydrochloride as the hydroxylamine source. Various bases were assessed under neat conditions; however, the reactivity was notably limited when only one equivalent of the base was employed (Table 1, entries 1–5). To overcome this limitation, a liquid component was incorporated into the reaction mixture. This adjustment resulted in a significant enhancement in the formation of target sulfohydroxamic acid **2j** (Table 1, entries 6–7), although an undefined impurity was also detected under these conditions. We postulated that this byproduct may arise from secondary transformations of nascent sulfohydroxamic acid, indicative of its inherent reactivity.⁶⁴

To enhance process efficiency of the process, the methodology was refined by increasing the equivalents of both hydroxylamine hydrochloride and the base. This modification led to a significant improvement in the conversion to **2j** (Table 1, entries 8 and 9). Notably, doubling the amount of base further increased the reactivity, achieving a conversion rate of 65% (Table 1, entry 10). However, subsequent efforts to optimize the outcome, such as using three equivalents of hydroxylamine and base along with extended reaction times of up to 3 h, did not yield additional improvements in conversion



(Table 1, entries 11–14). Collectively, these findings suggest that the most efficient conditions are attained with two equivalents of hydroxylamine hydrochloride and four equivalents of MgO, balancing reactivity and process simplicity. These optimized parameters were adopted as standard conditions for subsequent investigations.

With the optimized conditions established, we proceeded to evaluate the generality and robustness of the protocol by extending its application to a wide array of sulfonyl chlorides, thereby facilitating the synthesis of a diverse library of sulfohydroxamic acids (Scheme 1). The methodology demonstrated considerable versatility, accommodating a broad range of substituents with notable efficiencies. Substrates containing halogen substituents exhibited consistently good reactivity, allowing to potentially use these moieties as valuable handles for subsequent synthetic elaboration. Conversely, the incorporation of either electron-withdrawing groups or

electron-donating groups adversely affected the reaction outcome, indicating that, instead of specific electronic balance, other factors such as the rheology of the mixture might negatively affect the protocol. Notably, substrates containing an additional aromatic ring were particularly suitable for the transformation, as the extended conjugation appeared to facilitate the reaction pathway, resulting in the production of the corresponding sulfohydroxamic acids in commendable yields. In addition to synthesizing primary sulfohydroxamic acids, we further broadened the method's applicability to encompass the synthesis of structurally diverse derivatives featuring alkyl substituents on either the oxygen or nitrogen atoms. This was accomplished by substituting hydroxylamine hydrochloride with appropriately functionalized hydroxylamine derivatives, pre-installed with the desired substituents on the hydroxylamine scaffold (Scheme 1).




Table 1. Optimization of the synthesis of **2j**.

Entry	<i>p</i> -Tosyl chloride (eq.)	NH ₂ OH·HCl (eq.)	Base	Base (eq.)	LAG	Time (h)	2j %
1	1	1	NaHCO ₃	1	/	2	14%
2	1	1	Na ₂ CO ₃	1	/	2	27%
3	1	1	NaOH	1	/	2	11%
4	1	1	Na ₃ PO ₄	1	/	2	29%
5	1	1	MgO	1	/	2	25%
6	1	1	Na ₂ CO ₃	1	MeOH	2	40%
7	1	1	Na ₃ PO ₄	1	H ₂ O	2	27%
8	1	2	Na ₂ CO ₃	2	MeOH	2	51%
9	1	2	MgO	2	/	2	49%
10	1	2	MgO	4	/	2	65%
11	1	3	Mg(OH) ₂	3	/	2	66%
12	1	3	MgO	3	/	2	68%
13	1	3	Li ₂ CO ₃	3	ⁱ PrOH	3	63%
14	1	3	MgO	3	/	3	66%

The reactions were performed on a 1.0 mmol scale using a Retsch MM500 ball milling machine. The conversion rates were calculated using ¹H-NMR analysis. The amount of liquid additive had an η of 0.3 $\mu\text{L mg}^{-1}$, which represents the ratio of the liquid volume to the weight of the reaction mixture.


This strategic modification of the original protocol facilitated the efficient preparation of a diverse range of *O*- and *N,O*-substituted sulfohydroxamic acids, exemplified by *O*-

benzylhydroxylamine hydrochloride and *N, O*-dimethylhydroxylamine hydrochloride. Notably, the overall efficiency of the process was markedly enhanced, which was attributed to



both the increased stability of the resulting scaffolds and the absence of collateral pathways that affect the formation of the final product. The ability to access a diverse range of sulfohydroxamic acid derivatives not only illustrates the versatility of the methodology but also highlights its potential utility in the synthesis of tailored molecules relevant to both

synthetic and medicinal chemistry applications. For completeness, the synthesis of **2j** was also performed on a gram scale, resulting in the final 60% conversion of the starting material to the corresponding sulfohydroxamic acid.




Scheme 1. Scope of sulfohydroxamic acids without further functionalization of the nitrogen and oxygen atoms. The reactions were performed on a 1.0 mmol scale using a Retsch MM500 ball milling machine. The yields refer to the pure isolated compounds. ^aThe reaction was run on a 0.5 mmol scale.

To examine the reaction mechanism, it is essential to analyse the characteristics of this process under mechanochemical conditions. The transformation occurs through two sequential steps: an acid–base reaction between hydroxylamine hydrochloride and the chosen base, followed by the

nucleophilic attack of the liberated hydroxylamine on the sulfonyl chloride (Scheme 2).



ARTICLE



Scheme 2. General reaction mechanism concerning the release of gaseous hydroxylamine and its reaction with sulfonyl chloride present in the reaction environment.

The selection of the base and the potential inclusion of a liquid additive significantly impact the transformation outcome.⁶⁵ Sulfonyl chlorides and the resultant sulfohydroxamic acids are inherently sensitive to nucleophilic species and strongly basic environments. Consequently, inappropriate conditions can readily compromise the stability of both the starting material and product, leading to degradation pathways or competing side reactions. Therefore, the selected base (and, if applicable, the liquid additive) must meet a crucial dual criterion: it must possess sufficient basicity to effectively deprotonate hydroxylamine hydrochloride, thereby enabling the in situ release of free hydroxylamine within the milling jar to initiate the reaction. To exclude the formation of gaseous hydroxylamine or volatile by-products, the mass loss during milling of hydroxylamine hydrochloride and magnesium oxide was analysed (see ESI for details). Under the optimized conditions, no significant mass loss was detected, indicating that free hydroxylamine predominantly operates without developing any gaseous species. Although the reaction mixture remained macroscopically solid at the end of milling, a transient liquid phase arising from the low melting point of the free base (33 °C) cannot be completely ruled out. Concurrently, it must

not exhibit excessive reactivity to avoid undesirable interactions with sulfonyl chloride or alterations to the newly formed sulfohydroxamic acid.

Among all the bases examined, magnesium oxide (MgO) emerged as the most reliable promoter, uniquely combining adequate Brønsted basicity with a mild nucleophilic profile. Under neat, solvent-free milling conditions, MgO enables the controlled in situ liberation of hydroxylamine from its hydrochloride salt, while simultaneously maintaining a reaction environment that preserves as much as possible both the integrity of the sulfonyl chloride and the stability of the newly formed sulfohydroxamic acid (Figure 2, a further analysis of the stability of sulfonyl chlorides can be checked in the ESI file). Beyond its fundamental chemical role, the solid-state characteristics of MgO, including its high surface area, heterogeneous distribution of basic sites, and mechanical robustness, are ideally suited for mechanochemical activation.^{66, 67}

Notably, alternative bases either failed to initiate the reaction or induce undesirable side processes, underscoring the unique dual functionality of MgO as both a mild base and a structural moderator of the milling medium.




Figure 2. General steps regarding the mechanochemical process.

Lastly, the green metrics of the mechanochemical methodology were calculated, revealing an improvement in process mass intensity (PMI) of more than 20% (see the ESI file for additional details). Furthermore, the elimination of toxic volatile organic compounds (VOCs), such as THF, further enhances the overall safety of the process.

Conclusions

This study establishes mechanochemistry as a robust, user-friendly, and inherently safe method for synthesizing sulfohydroxamic acids. The straightforward, solvent-free milling of sulfonyl chlorides with hydroxylamine hydrochloride in the presence of a benign inorganic base, such as MgO, facilitates the



production of target compounds under mild conditions. This approach avoids hazardous, waste-intensive organic solvents and simplifies the workup.

In addition to operational simplicity, mechanochemical activation directly addresses the persistent challenges in this area of chemistry. This reduces the handling risks associated with volatile hydroxylamines and mitigates the instability of sensitive substrates and products, which are often problematic under solution conditions. Consequently, this method enhances practical access to sulfohydroxamic scaffolds, which are significant in both synthetic and medicinal chemistry applications.

Overall, these findings position mechanochemistry as a versatile and sustainable platform for the preparation of sulfohydroxamic acids, aligning with green chemistry principles by combining safety, efficiency, and environmental sustainability. We anticipate that this approach will be readily applicable to a diverse range of sulfonyl chloride frameworks and related *N*-functionalization reactions. Furthermore, scaling up using contemporary enabling technologies, such as RAM or extrusion, is expected to facilitate its adoption in research and development.

Author contributions

Conceptualisation: A.P.; Investigation: M.I., F.B.; Data Curation: M.I. Supervision: A.P., P.C.; Writing, original draft: M.I., A.P.; Writing, review, and editing: all authors. F.B. and M.I. have contributed equally to this work.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Data availability

The data supporting this article have been included as part of the ESI. All experimental data relevant to the manuscript's preparation and formulation of theories and/or hypotheses, including conclusions, etc. are provided in the ESI accompanying the manuscript.

Acknowledgements

The authors acknowledge financial support under the National Recovery and Resilience Plan (NRRP), Mission 4, Component 2, Investment 1.1, Call for Tender No. 1409 published on 14. 9. 2022 by the Italian Ministry of University and Research (MUR), funded by the European Union – NextGenerationEU-Project Title: The Second Life of Posidonia Oceanica: Sustainability-guided Isolation and Valorisation of its Main Constituents – CUP J53D23014680001- Grant Assignment Decree No. 1064 adopted on 18. 07. 2023 by the Italian Ministry of University and Research (MUR). The financial support of the European Union NextGenerationEU under the National Recovery and Resilience Plan (NRRP) of Ministero

dell'Università e della Ricerca (MUR) (Project code PE0000021, CUP no. F53C22000770007, Network 4-0 Energy Sustainable Transition, NEST) is acknowledged. Fondazione Banco di Sardegna (FDS) also supported the project financially, with grant number F73C23001600007.

Notes and references

1. A. Y. Jeng; M. Chou; D. T. Parker *Bioorganic & Medicinal Chemistry Letters* 1998, **8**, 897-902.
2. Z. Huang; C. Velázquez; K. Abdellatif; M. Chowdhury; S. Jain; J. Reisz; J. DuMond; S. B. King; E. Knaus *Organic & Biomolecular Chemistry* 2010, **8**, 4124-4130.
3. F. N. Shirota; E. G. DeMaster; M. J. C. Lee; H. T. Nagasawa *Nitric Oxide* 1999, **3**, 445-453.
4. Z. Huang; C. A. Velázquez; K. R. A. Abdellatif; M. A. Chowdhury; J. A. Reisz; J. F. DuMond; S. B. King; E. E. Knaus *Journal of Medicinal Chemistry* 2011, **54**, 1356-1364.
5. Y. Long; Z. Xia; A. M. Rice; S. B. King *Molecules* 2022, **27**, 5305.
6. M. N. Hughes; R. Cammack, in *Methods in Enzymology*, Academic Press, **1999**, vol. 301, pp. 279-287.
7. M. A. Alam *Curr Org Chem* 2019, **23**, 978-993.
8. A. Porcheddu; L. De Luca; G. Giacomelli *Synlett* 2009, **2009**, 2149-2153.
9. R. Smulik-Izydorczyk; J. Pięta; R. Michalski; M. Rola; K. Kramkowski; A. Artelska; J. Zielonka; A. B. Sikora *Redox Biochemistry and Chemistry* 2024, **8**, 100031.
10. A. Citarella; D. Moi; L. Pinzi; D. Bonanni; G. Rastelli *ACS Omega* 2021, **6**, 21843-21849.
11. O. Piloty *Berichte der deutschen chemischen Gesellschaft* 1896, **29**, 1559-1567.
12. P. C. Wilkins; H. K. Jacobs; M. D. Johnson; A. S. Gopalan *Inorganic chemistry* 2004, **43**, 7877-7881.
13. K. Sirsalmath; S. A. Suárez; D. E. Bikiel; F. Doctorovich *J. Inorg. Biochem.* 2013, **118**, 134-139.
14. K. Aizawa; H. Nakagawa; T. Suzuki; N. Miyata *Free Radical Biology and Medicine* 2012, **53**, S141.
15. J. Polaczek; H. Subedi; Ł. Orzeł; L. S. Lisboa; R. B. Cink; G. y. Stochel; N. E. Brasch; R. Van Eldik *Inorganic Chemistry* 2021, **60**, 2964-2975.
16. A. M. Confer; A. C. Vilbert; A. Dey; K. M. Lancaster; D. P. Goldberg *Journal of the American Chemical Society* 2019, **141**, 7046-7055.
17. S. Mondal; S. Malakar *Tetrahedron* 2020, **76**, 131662.
18. X.-R. Huang; Y.-M. Zhang; T.-B. Wan; P. Zhang; X.-X. Zhang; F.-M. Wang; D. Xu; M.-H. Shen; H.-D. Xu *Tetrahedron* 2019, **75**, 130336.
19. R. Zamora; A. Grzesiok; H. Weber; M. Feelisch *Biochemical Journal* 1995, **312**, 333-339.
20. D. Sanna; G. Rocchitta; M. Serra; M. Abbondio; P. A. Serra; R. Migheli; L. De Luca; E. Garribba; A. Porcheddu *Pharmaceuticals* 2017, **10**, 74.
21. H. Subedi; N. E. Brasch *Dalton Transactions* 2016, **45**, 352-360.
22. J. F. DuMond; S. B. King *Antioxidants & Redox Signaling* 2011, **14**, 1637-1648.
23. T. Nakano; N. Kousaka; A. Nakayama; Y. Kato; K. Takashima; G. Tanabe; M. Yoshimatsu *Organic Letters* 2024, **26**, 1753-1757.
24. F. J. Geffers; F. R. Kurth; P. G. Jones; D. B. Werz *Chemistry – A European Journal* 2021, **27**, 14846-14850.



25. M. Medeiros; E. H. Wanderlind; J. R. Mora; R. Moreira; A. J. Kirby; F. Nome *Organic & Biomolecular Chemistry* 2013, **11**, 6272-6284.
26. J. B. Domingos; E. Longhinotti; T. A. S. Brandão; C. A. Bunton; L. S. Santos; M. N. Eberlin; F. Nome *The Journal of Organic Chemistry* 2004, **69**, 6024-6033.
27. Y. Zeng; Y. Zhao; L. Yang; Y. Jiang *ChemistrySelect* 2023, **8**, e202303345.
28. C. T. A. Evelo; A. A. M. G. Spooren; R. A. G. Bisschops; L. G. M. Baars; J. M. Neis *Blood Cells, Molecules, and Diseases* 1998, **24**, 280-295.
29. S. C. Gad, in *Encyclopedia of Toxicology (Third Edition)*, ed. P. Wexler, Academic Press, Oxford, **2014**, pp. 982-983.
30. J. M. Desesso; G. C. Goeringer *Toxicology and Industrial Health* 1990, **6**, 109-121.
31. L. A. Long *Process Safety Progress* 2004, **23**, 114-120.
32. F. N. A. f. t. E. a. E. M. (ADEME), Hydroxylamine Explosion in a Chemical Plant - ARIA Accident Report No. 14952, https://www.aria.developpement-durable.gouv.fr/accident/14952_en/?lang=en.
33. I. T. Horvath; P. T. Anastas *Chemical reviews* 2007, **107**, 2169-2173.
34. M. Tobiszewski; M. Marć; A. Gałuszka; J. Namieśnik *Molecules* 2015, **20**, 10928-10946.
35. S. Trakhtenberg; J. C. Warner *Chemical reviews* 2007, **107**, 2174-2182.
36. R. Mocchi; L. D. Luca; F. Delogu; A. Porcheddu *Advanced Synthesis & Catalysis* 2016, **358**, 3135-3144.
37. F. Basoccu; L. De Luca; A. Porcheddu *European Journal of Organic Chemistry* 2024, **27**, e202400425.
38. N. Fantozzi; J.-N. Volle; A. Porcheddu; D. Virieux; F. García; E. Colacino *Chemical Society Reviews* 2023, **52**, 6680-6714.
39. S. L. James; T. Friščić *Chemical Society Reviews* 2013, **42**, 7494-7496.
40. J.-L. Do; T. Friščić *ACS central science* 2017, **3**, 13-19.
41. Joseph L. Howard; Q. Cao; D. L. Browne *Chemical Science* 2018, **9**, 3080-3094.
42. T. Friščić; C. Mottillo; H. M. Titi *Angewandte Chemie* 2020, **132**, 1030-1041.
43. J. Batteas; K. G. Blank; E. Colacino; F. Emmerling; T. Friščić; J. Mack; J. Moore; M. E. Rivas; W. Tysoe *RSC Mechanochemistry* 2025, **2**, 10-19.
44. K. Floyd; L. Gonnet; T. Friščić; J. Batteas *RSC Mechanochemistry* 2024, **1**, 289-295.
45. J. G. Hernández; C. Bolm *The Journal of organic chemistry* 2017, **82**, 4007-4019.
46. G. N. Hermann; P. Becker; C. Bolm *Angewandte Chemie* 2015, **127**, 7522-7525.
47. A. Gallego; N. Pétry; J. Pinaud; O. Giani; F. Lamaty *ChemistryEurope* 2025, e202500232.
48. M. Perez-Venegas; E. Juaristi *ACS Sustainable Chemistry & Engineering* 2020, **8**, 8881-8893.
49. S. L. James; C. J. Adams; C. Bolm; D. Braga; P. Collier; T. Friščić; F. Grepioni; K. D. Harris; G. Hyett; W. Jones *Chemical Society Reviews* 2012, **41**, 413-447.
50. T. Nakano; N. Kousaka; A. Nakayama; Y. Kato; K. Takashima; G. Tanabe; M. Yoshimatsu *Organic Letters* 2024, **26**, 1753-1757.
51. G. Kirby; G. Prestat; F. Berhal *The Journal of Organic Chemistry* 2023, **88**, 4720-4729.
52. G. Giacomelli; A. Porcheddu; M. Salaris *Organic Letters* 2003, **5**, 2715-2717.
53. I. S. Sow *European Journal of Chemistry* 2024, **15**, 345-354.
54. E. Juaristi; C. G. Avila-Ortiz *Synthesis* 2023, **55**, 2439-2459.
55. F. Basoccu, **2025**.
56. K. Kubota; H. Ito *Trends in chemistry* 2020, **2**, 1066-1081.
57. F. Basoccu; F. Cuccu; A. Porcheddu *ChemSusChem* 2024, **17**, e202301034.
58. F. Fischer; K.-J. Wenzel; K. Rademann; F. Emmerling *Physical Chemistry Chemical Physics* 2016, **18**, 23320-23325.
59. A. A. Michalchuk; F. Emmerling *Angewandte Chemie International Edition* 2022, **61**, e202117270.
60. P. P. Mazzeo; M. Prencipe; T. Feiler; F. Emmerling; A. Bacchi *Crystal Growth & Design* 2022, **22**, 4260-4267.
61. A. M. Constantin; F. Mele; M. Lanzi; G. Maestri; R. Maggi; N. Della Ca'; L. Capaldo *ChemSusChem* 2025, 2500211.
62. D. E. Crawford; S. L. James; T. McNally *ACS Sustainable Chemistry & Engineering* 2018, **6**, 193-201.
63. A. Krusenbaum; S. Grätz; G. T. Tigineh; L. Borchardt; J. G. Kim *Chemical Society Reviews* 2022, **51**, 2873-2905.
64. It was not possible to fully elucidate the exact chemical structures of the impurities detected in the 1H NMR spectra of para-methylbenzenesulfohydroxamic acid, even with the use of additional analytical techniques. This area remains underexplored in the literature, as impurities arising from sulfohydroxamic acid synthesis are rarely characterized. Consequently, this aspect of our study remains an ongoing area of investigation.
65. F. Basoccu; S. Piermarini; T. Angelini; M. Mari; E. Mariani; A. Falchi; A. Porcheddu *ChemSusChem* 2025, **18**, e202500457.
66. C. Sousa; J. Mejias; G. Pacchioni; F. Illas *Chemical physics letters* 1996, **249**, 123-129.
67. S. Ardizzone; C. Bianchi; B. Vercelli *Applied surface science* 1998, **126**, 169-175.



View Article Online
DOI: 10.1039/D5MR00141B

Data availability

The data supporting this article have been included as part of the ESI. All experimental data relevant to the manuscript's preparation and formulation of theories and/or hypotheses, including conclusions, etc. are provided in the ESI accompanying the manuscript.

Cagliari, 24/02/2026

Andrea Porcheddu
Prof. Andrea Porcheddu

RSC Mechanochemistry Accepted Manuscript

