



Cite this: DOI: 10.1039/d6mr00013d

Received 28th January 2026
Accepted 4th April 2026

DOI: 10.1039/d6mr00013d

rsc.li/RSCMechanochem

Mechanoredox radical annulation for the solvent-free synthesis of 2-substituted benzothiophenes

Ji Ho Song,^{ab} Eunsil Kim,^{ab} Do Hyun Kang,^{cde} Ji Young Hyun,^{ab} Hwan Jung Lim^{ID} ^{*ab}
and Seong Jun Park^{ID} ^{*ab}

A mechanoredox-enabled radical annulation under ball-milling conditions affords 2-substituted benzothiophenes **3** from 2-(methylthio)benzenediazonium salts **1** and alkynes **2**. Mechanical energy drives single-electron transfer without photoredox or electrochemical systems, providing a practical and environmentally benign approach to heterocycle synthesis.

Introduction

Radical intermediates can be generated under mild, functional-group-tolerant conditions, enabling diverse bond-forming reactions.¹ Recent advances, particularly in photoredox, electrochemical and mechanoredox strategies, have greatly expanded the synthetic toolbox for controlled radical chemistry.^{2,3}

Mechanochemistry has emerged as a sustainable approach to bond construction, enabling solvent-free access to unique and efficient reactivity.⁴⁻⁷ Building on these advances, we sought to develop a mechanoredox-based radical transformation that addresses limitations of conventional solution-phase methods.

In this context, the synthesis of 2-substituted benzothiophenes was chosen as a benchmark transformation, given the prominence of this heterocyclic scaffold in pharmaceuticals, functional materials and agrochemicals,⁸⁻¹⁹ and its established accessibility *via* radical annulation strategies (Fig. 1).^{20,21}



Unlike solution-based approaches that require organic solvent (typically DMSO), specialised photochemical or

electrochemical equipment, and in some cases stoichiometric additives, the mechanoredox protocol exploits mechanical energy alone to drive single-electron transfer, consistent with a mechanoredox SET process,^{3,22} enabling a solvent-free and promoter-dependent alternative with competitive efficiency, selectivity, and sustainability. We acknowledge that ball milling itself requires dedicated equipment and that Al₂O₃ is used in stoichiometric quantities; these aspects are weighed against the advantages when evaluating the practical utility of the method.

Results and discussion

Milling was carried out using a planetary micro mill (Pulverisette 7 Premium Line, Fritsch GmbH, Germany) equipped with a 20 mL ZrO₂ bowl²³ and ZrO₂ balls (10 mm in diameter).²⁴ When the planetary micro mill was equipped with 10 mm ZrO₂ balls, the maximum rotational speed was set to 850 rpm. Due to

<Previous work (solution-based approaches)>

(a) Koenig and coworkers (photoredox method)²⁰(b) Zeng and coworkers (electrochemical method)²¹

<This work (solution-free approach, mechanoredox method)>




Fig. 1 Synthesis of 2-substituted benzothiophenes **3** *via* radical annulation strategies.

^aData Convergence Drug Research Center, Therapeutics & Biotechnology Division, Korea Research Institute of Chemical Technology (KRICT), 141 Gajeong-ro, Yuseong-gu, Daejeon 34114, Republic of Korea. E-mail: sjunpark@kRICT.re.kr; Fax: +82 42 860 7160; Tel: +82 42 860 7175

^bDepartment of Medicinal Chemistry and Pharmacology, University of Science & Technology, Daejeon 34113, Republic of Korea

^cA Nano-Lithography & Manufacturing Research Center, Korea Institute of Machinery and Materials (KIMM), Daejeon 34103, Republic of Korea

^dAdvanced Bioconvergence, University of Science & Technology, Daejeon 34113, Republic of Korea

^eDepartment of Chemistry, Chung-Ang University, Seoul 06974, Republic of Korea



the maximum continuous milling time of 1 h, the process was conducted in two cycles of 1 h milling separated by 1 h cooling (Fig. 2).²⁵

The reaction conditions were optimised, as summarised in Table 1. The reaction was conducted using five ZrO₂ balls at the maximum speed of 850 rpm. An optimization study of the number of ZrO₂ balls was performed, and detailed results are provided in the SI.²⁷

As shown in Table 1, the absence of a promoter completely suppressed the reaction (entry 2), highlighting the essential role of the promoter. A reaction time of 2 h was identified as optimal (entry 4). To probe the influence of surface acidity, three commercially available Al₂O₃ samples of defined acidity (acidic, neutral, and basic; Brockmann Activity I, Sigma-Aldrich) were evaluated under otherwise identical ball-milling conditions (entries 4–7). The acidic Al₂O₃ afforded the highest yield (53%, entry 4), while both neutral and basic Al₂O₃ gave a reduced but still appreciable yield of 42% (entries 6 and 7), indicating that surface Lewis acidity enhances, but is not solely responsible for, the mechanoredox SET process. Furthermore, conducting the reaction in DMSO solution in the presence of Al₂O₃ (entry 8) afforded no detectable product, confirming that mechanical

activation—rather than chemical interaction with the solid additive alone—is the key driving force for the reaction. In contrast, replacement of Al₂O₃ with BaTiO₃ led to a reduced yield (entry 9), while TiO₂ was largely ineffective, affording only a very low yield (entry 10).

These results indicate that the presence of a promoter is essential for the reaction under mechanochemical conditions. A reaction time of 2 h was sufficient to achieve optimal conversion. In comparison, BaTiO₃ exhibited lower activity as a promoter, while TiO₂ was largely ineffective under the same conditions, suggesting that the activity as a promoter is strongly dependent on the nature of the promoter. The superior activity of Al₂O₃ relative to BaTiO₃ and TiO₂ argues against a purely piezoelectric mechanism³ and is instead consistent with a mechanoredox SET process in which mechanical activation enables single-electron reduction of the diazonium substrate. The Lewis acidic surface sites of Al₂O₃ are further proposed to activate the diazonium substrate toward SET.

To explore the scope of the optimized reaction, 2-(methylthio)benzenediazonium salt **1** was reacted with a variety of ethynyl compounds, including aryl-, heterocyclic-, and alkyl-substituted ethynes **2**. Detailed results are provided in the SI. Among the ethynyl substrates examined, ethynylbenzene, 3-ethynylpyridine, and 3-ethynylthiophene, which afforded relatively high yields, were selected and subsequently reacted with variously substituted derivatives of 2-(methylthio)benzenediazonium salt **1** to synthesise a series of 2-substituted benzothiophenes **3** (Fig. 3). Most of the compounds were isolated in moderate yields; notably, hetero-heterocyclic compounds (**3b** and **3c**) could also be readily obtained. These results compare favourably with those of related solution-phase methods: for the benchmark substrate **3a** (R² = Ph), the photoredox method of Koenig and co-workers²⁰ affords 70–85% in 14 h in DMSO, and the electrochemical method of Zeng and co-



Fig. 2 Planetary ball mill used in this study: Fritsch Pulverisette 7 premium line (left), ZrO₂ bowl before grinding (right).

Table 1 An initial reaction optimization study²⁶

Entry	2a (eq)	Promoters ^a	Reaction time (h)	3a Yield (%) ^b
1	15	Al ₂ O ₃ (acidic, 10 eq)	0.5	27
2	15	None	0.5	<1
3	10	Al ₂ O ₃ (acidic, 10 eq)	1	40
4	10	Al ₂ O ₃ (acidic, 10 eq)	2	53
5	10	Al ₂ O ₃ (acidic, 10 eq)	3	53
6	10	Al ₂ O ₃ (neutral, 10 eq)	2	42
7	10	Al ₂ O ₃ (basic, 10 eq)	2	42
8	10	Al ₂ O ₃ (10 eq), DMSO, no ball milling	2	<1 ^c
9	10	BaTiO ₃ (10 eq)	2	35
10	10	TiO ₂ (10 eq)	2	4

^a Commercially available Al₂O₃ (Brockmann Activity I, Sigma-Aldrich); acidic (pH ~4.5), neutral (pH ~7.0), basic (pH ~10.0). ^b After column chromatography. ^c Solution-phase control: reaction performed in DMSO without ball milling; confirms that mechanical activation is essential.






Fig. 3 Scope of the optimized reaction.




Fig. 4 Proposed reaction mechanism.

workers²¹ affords 60–75% in DMSO, compared with 53% in 2 h under solvent-free mechanochemical conditions in the present work. Notably, the hetero–heterocyclic products (3b and 3c series, $\text{R}^2 = 3\text{-pyridyl}$ and 3-thienyl) are, to the best of our knowledge, novel compounds not previously reported by either solution-phase method, further demonstrating the synthetic utility of the present approach. A comprehensive comparison for all directly comparable substrates is provided in SI Table S2.

To gain insight into the mechanism of the mechanochemical reaction, radical trapping experiments were conducted. Ball milling of 1a with TEMPO completely suppressed formation of 3a (3% yield) and afforded the TEMPO adduct 4 in 80% yield, unambiguously confirming the intermediacy of a free aryl radical.²⁸ The formation of the TEMPO adduct 4 was confirmed by ¹H NMR spectroscopy. Based on literature reports,^{3,20–22,28–30} together with our experimental results, a plausible reaction mechanism is proposed in Fig. 4; more detailed and rigorous mechanistic investigations are currently underway.

Conclusions

In summary, we have developed a mechanochemical-enabled radical annulation strategy for the synthesis of 2-substituted benzothiophenes 3 under solvent-free and promoter-dependent ball-milling conditions. This protocol exploits mechanical energy to drive single-electron transfer, providing a sustainable alternative to established photoredox and electrochemical

approaches. A range of aryl- and heterocyclic-substituted ethynes 2 were successfully employed, affording the desired benzothiophenes 3 in generally moderate yields, including hetero–heterocyclic derivatives (3b and 3c). Control experiments and promoter screening confirmed the crucial role of the promoter under mechanochemical conditions, while radical trapping studies support the involvement of radical intermediates. Overall, this study demonstrates the potential of mechanochemical chemistry as a practical and environmentally benign platform for radical heterocycle synthesis, expanding the synthetic toolbox for sustainable bond-forming reactions.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the supplementary information (SI). Supplementary information is available. See DOI: <https://doi.org/10.1039/d6mr00013d>.

Acknowledgements

This research was supported by the Korea Drug Development Fund funded by Ministry of Science and ICT, Ministry of Trade, Industry and Energy, and Ministry of Health and Welfare (RS-2025-02223190, Republic of Korea), by Korea Research Institute of Chemical Technology (KRICT, KK2531-20 and KK2631-10). D. H. K. acknowledges the financial support from National Research Council of Science & Technology (NST) grant by the in Korea government (MSIT) (Grant No. CAP22013-200).

Notes and references

- 1 Y. Zhang, Y. Zhang and C. Li, *Chem. Soc. Rev.*, 2025, **54**, 10427.
- 2 J. A. Leitch and D. L. Browne, *Chem.–Eur. J.*, 2021, **27**, 9721.
- 3 K. Kubota, Y. Pang, A. Miura and H. Ito, *Science*, 2019, **366**, 1500.
- 4 J. L. Howard, Q. Cao and D. L. Browne, *Chem. Sci.*, 2018, **9**, 3080.
- 5 T. Friščić, S. L. Childs, S. A. A. Rizvi and W. Jones, *CrystEngComm*, 2009, **11**, 418.
- 6 D. Tan and F. García, *Chem. Soc. Rev.*, 2019, **48**, 2274.
- 7 J. Andersen and J. Mack, *Green Chem.*, 2018, **20**, 1435.
- 8 Z. Gai, B. Yu, X. Wang, Z. Deng and P. Xu, *Microbiology*, 2008, **154**, 3804.
- 9 J. Konishi, T. Onaka, Y. Ishii and M. Suzuki, *FEMS Microbiol. Lett.*, 2000, **187**, 151.
- 10 C. Yun, J. You, J. Kim, J. Huh and E. J. Kim, *Photochem. Photobiol. C*, 2009, **10**, 111.
- 11 M. S. Malamas, J. Sredy, C. Moxham, A. Katz, W. Xu, R. McDevitt, F. O. Adebayo, D. R. Sawicki, L. Seestaller, D. Sullivan and J. R. Taylor, *J. Med. Chem.*, 2000, **43**, 1293.



- 12 J. W. Ellingboe, T. R. Alessi, T. M. Dolak, T. T. Nguyen, J. D. Tomer, F. Guzzo, J. F. Bagli and M. L. McCaleb, *J. Med. Chem.*, 1992, **35**, 1176.
- 13 L. F. Tietze, J. K. Lohmann and C. Stadler, *Synlett*, 2004, **6**, 1113.
- 14 L. F. Tietze, K. Thede, R. Schimpf and F. Sanniccolo, *Chem. Commun.*, 2000, 583.
- 15 L. F. Tietze and K. Thede, *Chem. Commun.*, 1999, 1811.
- 16 I. Fouad, Z. Mechbal, K. I. Chane-Ching, A. Adenier, F. Maurel, J.-J. Aaron, P. Vodicka, K. Cernovska, V. Kozmik and J. Svoboda, *J. Mater. Chem.*, 2004, **14**, 1711.
- 17 A. J. Seed, K. J. Toyne, J. W. Goodby and M. Hird, *J. Mater. Chem.*, 2000, **10**, 2069.
- 18 S. Pu, M. Li, C. Fan, G. Liu and L. Shen, *J. Mol. Struct.*, 2009, **919**, 100.
- 19 K. H. Jung, K. H. Kim, D. H. Lee, D. S. Jung, C. E. Park and D. H. Choi, *Org. Electron.*, 2010, **11**, 1584.
- 20 D. P. Hari, T. Hering and B. König, *Org. Lett.*, 2012, **14**, 5334.
- 21 L.-Y. Lan, Y.-Y. Jiang, R. D. Little and C.-C. Zeng, *J. Electrochem.*, 2024, **30**, 2313002.
- 22 D. Chen, S. Li, Z. Ren, C. Wang, R. Ding, J. Wang, A. Li, J. Chen and Z. Wang, *Nat. Commun.*, 2026, **17**, 1222.
- 23 A 20 mL milling bowl was used, as this was the smallest size commercially available from the manufacturer.
- 24 As comparable yields were obtained with 5 mm ZrO₂ balls, 10 mm ZrO₂ balls were used; larger balls could not be accommodated in the 20 mL milling bowl.
- 25 Safety note: aryl diazonium tetrafluoroborate salts should be handled with appropriate care. Although these materials are substantially more thermally stable than diazonium chloride or nitrate salts, they should not be heated or subjected to friction in the dry state in large quantities. All milling experiments described herein were performed on a small scale (≤ 2.1 mmol of diazonium salt), using a sealed ZrO₂ milling jar at ambient temperature with the standard cooling cycle (1 h milling, 1 h rest, 1 h milling). Appropriate personal protective equipment (safety glasses, laboratory coat, and gloves) was worn at all times. No exothermic events or decomposition were observed during any experiment.
- 26 Among the conditions not included in Table 1, the use of 5 equiv. of ethynylbenzene **2a** led to damage of the milling bowl, preventing further experimentation. In addition, increasing the loading of Al₂O₃ to 15 or 20 equiv. did not result in a significant change in yield.
- 27 SI Table S2 compares the yields obtained in this work with those reported by Koenig and co-workers (photoredox) and Zeng and co-workers (electrochemical) for directly comparable substrates.
- 28 I. V. Alabugin and M. Manoharan, *J. Am. Chem. Soc.*, 2005, **127**, 12583.
- 29 J. Yan, J. Xu, Y. Zhou, J. Chen and Q. Song, *Org. Chem. Front.*, 2018, **5**, 1483.
- 30 D. Zhang, J. Cai, J. Du, X. Wang, W. He, Z. Yang, C. Kiu, Z. Fang and K. Guo, *J. Org. Chem.*, 2021, **86**, 2593.

