



RESEARCH ARTICLE

 View Article Online
View Journal


Cite this: DOI: 10.1039/d6qi00434b

Silicon–carbon bond cleavage from a hydroboration sequence

Manjur O. Akram,  Ayesha Begum and Caleb D. Martin  *

The reactions of internal pyridyl/trialkylsilyl alkynes with the potent hydroboration reagent bis(1-methyl-*ortho*-carboranyl)borane ($\text{HB}^{\text{Me}}\text{oCb}_2$) results in unanticipated Si–C_{alkyl} cleavage following a regioselective 1,1-hydroboration under mild conditions. The rare Si–Me bond rupture of the trimethylsilyl group occurs when using an intramolecular frustrated Lewis pair system forming a silylium heterocycle with a pendant borate bearing the methyl group. Investigating a methyl/ethyl mixed silyl substituent, SiEtMe_2 , revealed that the reaction is completely selective in breaking the Si–Me bond over the Si–Et bond. Exposure of the silylium/borate zwitterion to methanol resulted in O–H bond cleavage and ring opening to a zwitterionic pyridinium/borate. These findings provide insight into the challenging selective functionalization of Si–C_{alkyl} bonds.

 Received 5th March 2026,
Accepted 27th March 2026

DOI: 10.1039/d6qi00434b

rsc.li/frontiers-inorganic

Introduction

The hydroboration of alkynes is a powerful synthetic method to access building blocks with versatile boryl functional groups.^{1–4} The reaction can be mediated by a metal catalyst or proceed uncatalyzed, with the latter being possible with very Lewis acidic boranes or activated alkynes. The reaction can lead to either 1,2- or 1,1-hydroboration products.⁵ Both regioisomers are desirable and highlight the versatility of alkyne hydroboration. The classical uncatalyzed 1,2-hydroboration pioneered by Brown occurs *via* a concerted 4-membered cyclic transition state **I** to furnish the vinyl borane **II** with the boryl group and hydride on adjacent carbon atoms (Fig. 1).^{6,7} Contrarily, the 1,1-hydroboration gives the alternative vinyl borane isomer with the boryl and hydride on the same carbon.^{8–18} The accepted reaction mechanism proceeds by isomerization to form a reactive zwitterionic vinyl borate intermediate (**III**) that undergoes a 1,2-hydride shift from the borate to the carbocation to furnish the vinyl borane (**IV**).^{11,17} Wrackmeyer, Piers, Stephan, Erker, and others have investigated intermediates of type **III**, also referred to as boravinylidenes, but they were unable to characterize them.^{19–21,22–29} Challenges occur due to the extremely reactive vinyl cation adjacent to the borate center.

In hydroboration reactivity, a consequence of the reactive boravinylidene is often poor regioselectivity giving mixtures of products. The highly Lewis acidic secondary borane, bis(1-methyl-*ortho*-carboranyl)borane ($\text{HB}^{\text{Me}}\text{oCb}_2$), is an effective

1,1-hydroboration reagent for phenyl ethynylsilanes ($\text{PhC}\equiv\text{CSiR}_3$).³⁰ Regioselective 1,1-hydroboration reactions are rare and in this instance is attributed to the high hydride affinity of the carborane substituted $-\text{B}^{\text{Me}}\text{oCb}_2$ boryl group that attenuates the hydride transfer as well as hyperconjugative stabilization of the vinyl cation from the silyl group.^{31–33} In fact, the hydroboration attempt of a bis(trimethylsilyl)alkyne with $\text{HB}^{\text{Me}}\text{oCb}_2$ enabled the crystallographic characterization of boravinylidene **V**, as the additional silyl group sufficiently stabilized the carbocationic center (Fig. 2a).³⁴

Given that silyl hyperconjugative interactions were effective for characterizing a boravinylidene, we envisioned that an internal alkyne with a trimethylsilyl and pyridyl group (**VI**) could provide mechanistic insight into 1,1-hydroboration reactions and lead to the isolation of an intramolecularly stabilized boravinylidene (**VII**, Fig. 2b). Instead, the reaction of the alkyne with $\text{HB}^{\text{Me}}\text{oCb}_2$ led to the unexpected cleavage of a Si–Me bond of the $-\text{SiMe}_3$ group (**VIII**). Generally, $-\text{SiMe}_3$ groups are considered inert, underscored by their use as protecting




Fig. 1 1,2- and 1,1-hydroboration reactions of alkynes.







Fig. 2 (a) Characterized boravinylidene intermediate **V** and (b) reactions targeting base-stabilized boravinylidenes in this study.

groups in synthesis. The selective activation and cleavage of unactivated Si–Me bonds offer a potential means for the functionalization of alkylsilanes, moieties prominent in medicinal chemistry, as well as in polymer and materials science. Herein, we investigate the reactivity of pyridyl/silyl alkynes with $\text{HB}^{\text{Me}}\text{oCb}_2$.

Results and discussion

The equimolar reaction of commercially available 2-((trimethylsilyl)ethynyl)pyridine (**A1**) and $\text{HB}^{\text{Me}}\text{oCb}_2$ at 23 °C in benzene gave a major product after 24 hours that was isolated in 92% yield (Scheme 1a). The characteristic tricoordinate ^{11}B { ^1H } NMR signal for $\text{HB}^{\text{Me}}\text{oCb}_2$ at 71.0 ppm is consumed and the identity of the product was determined by a single crystal X-ray diffraction study as the pyridine/borane adduct **1** (Fig. 3). Heating a benzene solution of **1** at 70 °C for 24 hours did not result in any conversion to a different compound.

It has been established that $\text{HB}^{\text{Me}}\text{oCb}_2$ is an effective Lewis acid and binds irreversibly with phosphines (*i.e.* PMe_3 and PPh_3), triethylphosphine oxide, 4-fluorobenzonitrile, ethylacetate, tri-*n*-butylamine, and carbene (IMes).^{30,35–37} Bulky phosphines like PCy_3 , PtBu_3 , and PMes_3 do not make adducts due to the steric demands. Regarding pyridines, 2,6-lutidine coordinates to boron; thus, to preclude adduct formation to access hydroboration reactivity, we believed that the pyridine moiety in **A1** would require a substituent bulkier than methyl. Accordingly, we prepared an analogue of **A1** that features a *tert*-butyl group on the pyridine carbon adjacent to nitrogen, 2-((*tert*-butyl)-6-((trimethylsilyl)ethynyl)pyridine (**A2**), by a Sonagashira coupling reaction. The equimolar reaction of **A2** with $\text{HB}^{\text{Me}}\text{oCb}_2$ at 23 °C in benzene gave a major product after 24 hours that was isolated in 88% yield (Scheme 1b). The identity of the product was established as the silylium/borate zwitterion **2** by a single crystal X-ray diffraction study (Fig. 3). Studies by Piers, Gandelman, Alcarazo, Gagné, and our group have demonstrated Si–H bond splitting under mild conditions using borane/phosphine or borane/amine FLP systems.^{38–54} Though the Si–C bond energies are similar to



Scheme 1 Reactions of (a) pyridine-, (b) *tert*-butyl-pyridine-, and (c) quinoline-tethered (trimethylsilyl)ethynyl species with $\text{HB}^{\text{Me}}\text{oCb}_2$.

Si–H bond energies ($\sim 90 \text{ kcal mol}^{-1}$), their activation is less common.⁵⁵ The compound features an intramolecular pyridine-silylium adduct in an SiNC_2 heterocycle consisting of the nitrogen, *ortho*-carbon of the pyridine, the *ipso*-alkyne carbon, and the silicon bearing only two methyl groups. The borate center bears the two carborane substituents as well as the migrated methyl group and is bound to the distal alkyne carbon. The hydride is introduced on the same carbon as the borate, consistent with a 1,1-hydroboration pathway.

The ^1H NMR spectrum features a singlet at 1.13 ppm integrating to 6 for the SiMe_2 protons and a singlet at 0.24 ppm with an integration of 3 for the migrated BMe group. The ^{29}Si { ^1H } NMR signal for **2** is shifted significantly downfield from **A2** (59.9 ppm, *cf.* -17.0 ppm) and is in the range of known pyridine-stabilized alkylsilylium species (45–65 ppm).^{56–59} In the solid-state structure of **2**, the carbon–carbon bond is consistent with an alkene [C2–C3 1.336(3) Å] that is corroborated in the FT-IR spectrum with a C=C stretching frequency of 1600 cm^{-1} . The geometries at C2 and C3 are distorted trigonal planar due to the bulk around the double bond and the strain from the four membered ring. Compound **2** is stable under a nitrogen atmosphere at -40 °C for over 12 months with no detectable decomposition. The synthesis of **2** also works smoothly on a larger scale (236 mg, 85% yield) by reacting 0.50 mmol of **A2** and $\text{HB}^{\text{Me}}\text{oCb}_2$ in C_6H_6 (5 mL) at 23 °C for 24 hours. The reactions of **A2** with HBpin or HBCat did not result in any reaction, even after heating at 70 °C for 24 hours, indicating the importance of the electrophilic $\text{HB}^{\text{Me}}\text{oCb}_2$ reagent.






Fig. 3 Solid state structures of **1–3** and **3'**. Ellipsoids depicted at the 50% probability level. Hydrogen atoms, except on the vinylic carbon or central boron atom, and solvates are omitted for clarification.

To determine if a quinoline-tethered ethynylsilane would alter the reaction outcome, the corresponding alkyne (**A3**) was reacted with $\text{HB}^{\text{Me}}\text{oCb}_2$ at 23 °C for 24 hours. Adduct **3'** was generated as confirmed by single crystal X-ray diffraction (Fig. 3). The B1–N1 bond length in adducts **1** and **3'** are similar [1.623(3) Å, *cf.* 1.6080(19) Å] and comparable to that of the only other N-ligated $\text{HB}^{\text{Me}}\text{oCb}_2$ species, a 2,6-lutidine adduct [1.631(4) Å].³⁵ Heating a benzene solution of **3'** at 70 °C for 24 hours led to conversion to the Si–Me split product akin to **2** with a 5-membered sila-cycle (**3**) isolated in 83% yield (Scheme 1c). The structure of **3** was confirmed by single crystal X-ray diffraction. The $^{29}\text{Si}\{^1\text{H}\}$ NMR resonance of **3** appears downfield from the alkyne but upfield from that of **1** (**3**: 29.5 ppm *cf.* **A3**: –17.4, and **2**: 59.9 ppm).

1,1-Hydroboration reactions often suffer from poor selectivity leading to mixtures of both *E*- and *Z*-isomers.⁶⁰ The proposed mechanism for **2** involves silane to boron transmetalation between the –SiMe₃ group and $\text{HB}^{\text{Me}}\text{oCb}_2$ to generate the

ethynylborate with a pyridine ligated silylium cation (**Int1**, Fig. 4). Hydride and silylium migration generates **Int2** that is guided by the tethered pyridine to achieve the regioselective *trans*-1,1-hydroboration. The pyridine and borane serve as an intramolecular FLP system in **Int2** to enable the Si–Me bond cleavage to form **2**. Monitoring the reaction by ^1H and ^{11}B NMR spectroscopy in C_6D_6 at 23 °C did not enable the detection of intermediates.

The controlled cleavage of Si–Me bonds is rare, often requiring harsh reaction conditions and struggles with selectivity. Thus, the selective activation of unactivated Si–Me bonds under mild reaction conditions is desirable. Transition-metals have had minimal success through oxidative addition or σ -bond metathesis pathways.^{61–70} In transition-metal free approaches, potent electrophilic reagents have been effective including some boron systems.^{71–82} Notably, Wang recently disclosed Si–C/B–H to Si–H/B–C bond metathesis of the carbene stabilized hydroborenium ion, $[\text{IME}_4\text{B}(\text{H})\text{oCb}][\text{B}$




Fig. 4 Plausible reaction mechanism for the formation of **2**.




(C₆F₅)₄], with alkyl and aryl silanes to furnish the hydrosilane and organoborenium species (HSiR₃ and [IMe₄B(R)_oCb][B(C₆F₅)₄]).⁸³ Stephan and Inoue independently reported methyl abstraction of activated N-SiMe₃ groups in silyl-phosphinimides R₃P=N-SiMe₃ (R = *t*Bu, *i*Pr, Ph) and the N-heterocyclic imine IPr=N-SiMe₃ by B(C₆F₅)₃ to generate [MeB(C₆F₅)₃⁻] anions [IPr = bis(2,6-diisopropylphenyl)imidazoline] with coordinated silylium cations.^{84,85} The hydroboration sequence reported herein is a new pathway in boron mediated cleavage of unactivated Si-C bonds.

To investigate the selectivity of the Si-C bond activation, a variant of **A2** featuring a -SiEtMe₂ group in place of the -SiMe₃ group (**A4**) was reacted with HB^{Me}_oCb₂ at 23 °C for 24 h (Scheme 2). The reaction showed exclusive selectivity for Si-Me bond cleavage with no evidence of Si-Et activation by NMR spectroscopy. The structure was identified as silylium/borate zwitterion **4** by a single crystal X-ray diffraction study, the analogue of **2** differing in the ethyl group on silicon (Fig. 5). In the ²⁹Si{¹H} NMR spectrum, the resonance for **4** at 64.3 ppm is comparable to that for **2** (*cf.* **2**: 59.9 ppm).

Examining the X-ray diffraction structures of the silylium/borate species **2-4** reveals that the metrical parameters of the four membered products, **2** and **4**, are essentially within the error of measurement (Table 1). The N-Si bond length in **3** is slightly shorter than those of **2** and **4** attributed to the reduced ring strain [**2**: 1.8896(17) Å, **3**: 1.858(2) Å, **4**: 1.862(4) Å, Table 1].⁵⁷ The bond angles in the silacycle are more compressed in **2** and **4** compared to those in **3**, consistent with the 4- and 5-membered rings, respectively.

To probe the role of the pyridine moiety in the Si-Me bond cleavage in the generation of **2-4**, an alkyne analogue of **A1** with thiophene in place of pyridine was examined as it is a weaker Lewis base. The reaction of trimethyl(thiophen-2-ylethynyl)silane (**A5**) with HB^{Me}_oCb₂ in C₆D₆ at room temperature was monitored by ¹H NMR spectroscopy that revealed the 1,1-hydroboration product (**5**) within 10 minutes. The diagnostic vinyl proton in **5** appears at 6.42 ppm.³⁰ The vinyl carbon attached to the boron center appears broad at 152.1 ppm due to coupling with a quadrupolar ¹¹B-atom. A 2D ¹H-¹³C HSQC correlation experiment indicates that the hydrogen and boron atoms are attached to the same carbon, confirming the 1,1-hydroboration. Stirring a solution of **5** in C₆D₆ for 8 hours did not lead to any conversion for Si-Me bond rupture, suggesting the crucial role of the pyridine moiety for the Si-C bond cleavage (Scheme 3a).



Scheme 2 Selective Si-Me bond activation of pyridine-tethered (ethyldimethylsilyl)ethynyl **A4**.





Fig. 5 Solid state structure of **4**. Ellipsoids depicted at the 50% probability level. Hydrogen atoms (except on the vinylic carbon) and solvates are omitted for clarity.

Table 1 Selected bond lengths (Å) and angles (°) of cyclized species **2-4**

	2	3	4
B1-C1	1.619(3)	1.618(4)	1.624(6)
B1-C2	1.616(3)	1.614(4)	1.623(5)
C2-C3	1.336(3)	1.345(3)	1.352(5)
C3-Si1	1.852(2)	1.871(3)	1.862(4)
N1-Si1	1.8896(17)	1.858(2)	1.898(3)
∠C1-B1-C2	111.61(16)	112.2(2)	110.2(3)
∠B1-C2-C3	129.51(18)	131.9(2)	128.4(3)
∠C2-C3-Si1	144.29(16)	130.86(19)	142.9(3)
∠C3-Si1-N1	73.42(8)	89.93(10)	73.48(16)



Scheme 3 Reaction of (a) **A5** with HB^{Me}_oCb₂ and (b) **2** with MeOH.

To examine the lability of the N-Si bond in **2**, it was reacted with acetonitrile and pyridine at 23 °C. No reaction was observed with either reagent, even after 24 h, indicating that intramolecular N-Si coordination is resilient. Attempts to abstract the -Me group from boron using MeOTf failed. At 23 °C, no reaction occurred with 10 equivalents of MeOTf and heating to 50 °C for 24 h only led to slow decomposition and a complex mixture as determined by NMR spectroscopy. Addition of MeOH to **2** resulted in O-H bond cleavage at the N/Si unit rather than B-Me bond rupture, leading to pyridinium borate zwitterionic species **6** as confirmed by NMR spectroscopy and X-ray diffraction studies (Scheme 3b and Fig. 6).






Fig. 6 Solid state structure of **6**. Ellipsoids depicted at the 50% probability level and hydrogen atoms (except on the nitrogen and vinylic carbon) are omitted for clarity. Selected bond lengths (Å) and angles (°): B1–C1 1.616(2), B1–C2 1.631(2), C2–C3 1.354(2), C3–Si1 1.903(15), Si1–C4 1.857(2), Si1–C5 1.848(19), Si1–O1 1.651(13), \angle C1–B1–C2 114.08(12), \angle B1–C2–C3 136.26(13), \angle C2–C3–Si1 132.07(11).

The resulting product was isolated in 88% yield. In the ^1H NMR spectrum in CDCl_3 , the pyridinium proton appears at 12.75 ppm. The SiMe_2 unit shows noticeable upfield shifts in the ^1H (2: 1.13 ppm, **6**: 0.57 ppm) and $^{29}\text{Si}\{^1\text{H}\}$ (2: 59.9 ppm, **6**: 14.1 ppm) NMR spectra. The ring strain by N/Si breakage is reflected in the decrease of the \angle C2–C3–Si1 angle in the solid-state structure [2: 144.29(16)°, **6**: 132.07(11)°]. Additionally, the pyridinium ring is not coplanar with the alkene as it is oriented 58.2° relative to the Si1–C3–C2–B1 plane.

Conclusions

The reactions of $\text{HB}^{\text{Me}}\text{oCb}_2$ with internal alkynes with alkyl silyl and pyridine substitution give diverse outcomes. The alkyne featuring 2-pyridyl and trimethylsilyl groups forms a Lewis acid/base adduct that is thermally stable. Installing a *tert*-butyl group on the carbon adjacent to nitrogen resulted in a 1,1-hydroboration sequence followed by –Me group migration from silicon to boron to generate a zwitterionic compound featuring a pyridine-silylium SiNC_2 ring with a pendant alkenylborate. Replacing an ethyl group in place of one of the methyl groups on the trimethylsilyl substituent revealed that the reaction is selective for Si–Me rupture over Si–Et. For an alkyne bearing quinoline and a trimethylsilyl group, adduct formation occurred that thermally isomerized to the zwitterionic product with a SiNC_3 heterocycle. The reaction outcomes are rationalized by the tremendous Lewis acidity of the $-\text{B}^{\text{Me}}\text{oCb}_2$ unit and the opposing nitrogen donor acting as a frustrated Lewis pair to induce the Si–C bond cleavage. This study focused on silyl alkyne systems, but it may be possible to extend the element- CH_3 cleavage reactivity to other main group elements such as germanium. In sum, these findings reveal insight into hydroboration processes and ability of

potent Lewis acidic bis(*ortho*-carboranyl)boryl groups to facilitate the cleavage of unactivated Si– C_{alkyl} bonds.

Author contributions

M. O. A. carried out the synthetic experiments. A. B. performed the single crystal X-ray diffraction analyses. M. O. A. and C. D. M. conceived the project with supervision by C. D. M. M. O. A. and C. D. M. wrote the manuscript with edits from A. B.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the supplementary information (SI). Supplementary information: experimental procedures, NMR spectra, and X-ray crystallographic data. See DOI: <https://doi.org/10.1039/d6qi00434b>.

CCDC 2523192–2523197 contain the supplementary crystallographic data for this paper.^{86a–f}

Acknowledgements

We are grateful to the National Science Foundation (Award No. 2349851) and Welch Foundation (Grant No. AA-2203-20240404) for their generous support of this work.

References

- N. Miyaura and A. Suzuki, Palladium-Catalyzed Cross-Coupling Reactions of Organoboron Compounds, *Chem. Rev.*, 1995, **95**, 2457–2483.
- E. R. Burkhardt and K. Matos, Boron Reagents in Process Chemistry: Excellent Tools for Selective Reductions, *Chem. Rev.*, 2006, **106**, 2617–2650.
- V. B. Saptal, R. Wang and S. Park, Recent advances in transition metal-free catalytic hydroelementation (E = B, Si, Ge, and Sn) of alkynes, *RSC Adv.*, 2020, **10**, 43539–43565.
- S. J. Geier, C. M. Vogels, J. A. Melanson and S. A. Westcott, The transition metal-catalysed hydroboration reaction, *Chem. Soc. Rev.*, 2022, **51**, 8877–8922.
- B. Wrackmeyer, 1,1-organoboration of alkenylsilicon, -germanium, -tin and -lead compounds, *Coord. Chem. Rev.*, 1995, **145**, 125–156.
- H. C. Brown and B. C. S. Rao, A New Technique for the Conversion of Olefins into Organoboranes and Related Alcohols, *J. Am. Chem. Soc.*, 1956, **78**, 5694–5695.
- H. C. Brown, *Boranes in Organic Chemistry*, Cornell University Press, 2019.



- 8 F. Meyer, M. U. Schmidt and P. Paetzold, 1,1-Hydroborierung von Alkinen mit 6-Aza-nido-decaboranen, *Chem. Ber.*, 1995, **128**, 947–951.
- 9 P. Paetzold and E. Leuschner, Hydroboration of Alkenes and Alkynes with Aza-nido-decaboranes, *Z. Anorg. Allg. Chem.*, 2002, **628**, 658–660.
- 10 J. Ugolotti, G. Dierker, G. Kehr, R. Fröhlich, S. Grimme and G. Erker, Five-Membered Metallacyclic Allenoids: Synthesis and Structure of Remarkably Stable Strongly Distorted Cyclic Allene Derivatives, *Angew. Chem., Int. Ed.*, 2008, **47**, 2622–2625.
- 11 A. Boussounière, X. Pan, S. J. Geib and D. P. Curran, Borenum-Catalyzed Hydroborations of Silyl-Substituted Alkenes and Alkynes with a Readily Available N-Heterocyclic Carbene–Borane, *Organometallics*, 2013, **32**, 7445–7450.
- 12 D.-T. Yang, S. K. Mellerup, X. Wang, J.-S. Lu and S. Wang, Reversible 1,1-Hydroboration: Boryl Insertion into a C-N Bond and Competitive Elimination of HBR₂ or R-H, *Angew. Chem., Int. Ed.*, 2015, **54**, 5498–5501.
- 13 B. R. Barnett, C. E. Moore, A. L. Rheingold and J. S. Figueroa, Frustrated Lewis pair behavior of monomeric (boryl)iminomethanes accessed from isocyanide 1,1-hydroboration, *Chem. Commun.*, 2015, **51**, 541–544.
- 14 T. H. Allen, S. J. Geib and D. P. Curran, Radical and Thermal Reactions of N-Heterocyclic Carbene Boranes with Diazo Compounds, *Organometallics*, 2016, **35**, 2975–2979.
- 15 C. Tang, Q. Liang, A. R. Jupp, T. C. Johnstone, R. C. Neu, D. Song, S. Grimme and D. W. Stephan, 1,1-Hydroboration and a Borane Adduct of Diphenyldiazomethane: A Potential Prelude to FLP-N₂ Chemistry, *Angew. Chem., Int. Ed.*, 2017, **56**, 16588–16592.
- 16 K. Škoch, C. G. Daniliuc, G. Kehr and G. Erker, Alkyne 1,1-Hydroboration to a Reactive Frustrated P/B-H Lewis Pair, *Angew. Chem., Int. Ed.*, 2021, **60**, 6757–6763.
- 17 Z.-L. An, C. Chen, A.-X. Yu, D.-X. Cao, Y.-Y. Jiang and B. Xiao, trans-Ge/B 1,1-Hydroboration of Alkynylgermanes with 9-BBN, *Org. Lett.*, 2025, **27**, 4079–4084.
- 18 X. Wang, X. Jing, G. Tang, D. Pan and H. Wang, Reaction of Diazo Compounds with Borenum Ions: C–H Bond Activation/Annulation vs 1,1-Carboboration, *Inorg. Chem.*, 2026, **65**, 761–766.
- 19 C. Chen, G. Kehr, R. Fröhlich and G. Erker, Carbon–Carbon Bond Activation by 1,1-Carboboration of Internal Alkynes, *J. Am. Chem. Soc.*, 2010, **132**, 13594–13595.
- 20 G. Kehr and G. Erker, 1,1-Carboboration, *Chem. Commun.*, 2012, **48**, 1839–1850.
- 21 G. Kehr and G. Erker, Advanced 1,1-carboboration reactions with pentafluorophenylboranes, *Chem. Sci.*, 2016, **7**, 56–65.
- 22 B. Wrackmeyer, K. H. von Locquenghien and S. Kundler, Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations, *J. Organomet. Chem.*, 1995, **503**, 289–295.
- 23 B. Wrackmeyer, G. Kehr and S. Willbold, Reactions of zwitterionic η^2 -(alkyn-1-yl-borate)alkenyltin compounds with Lewis bases, *J. Organomet. Chem.*, 1999, **590**, 93–103.
- 24 C. Fan, W. E. Piers, M. Parvez and R. McDonald, Divergent Reactivity of Perfluoropentaphenylborole with Alkynes, *Organometallics*, 2010, **29**, 5132–5139.
- 25 C. Jiang, O. Blacque and H. Berke, Activation of Terminal Alkynes by Frustrated Lewis Pairs, *Organometallics*, 2010, **29**, 125–133.
- 26 X. Zhao, L. Liang and D. W. Stephan, Facile synthesis of electrophilic vinyl boranes: reactions of alkynyl-borates and diazonium salts, *Chem. Commun.*, 2012, **48**, 10189–10191.
- 27 J. R. Lawson, V. Fasano, J. Cid, I. Vitorica-Yrezabal and M. J. Ingleson, The carboboration of Me₃Si-substituted alkynes and allenes with boranes and borocations, *Dalton Trans.*, 2016, **45**, 6060–6070.
- 28 B. Wrackmeyer, G. Kehr, A. Sebald and J. Kümmerlen, Organotin cations stabilized by π coordination – synthesis and NMR studies in solution and in the solid state, *Chem. Ber.*, 1992, **125**, 1597–1603.
- 29 B. Wrackmeyer, G. Kehr and R. Boese, π -Coordination of Diorganotin Dications, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 1370–1372.
- 30 M. O. Akram, J. R. Tidwell, J. L. Dutton and C. D. Martin, Bis(1-Methyl-ortho-Carboranyl)Borane, *Angew. Chem., Int. Ed.*, 2023, **62**, e202307040.
- 31 M. O. Akram, J. R. Tidwell, J. L. Dutton and C. D. Martin, Tris(ortho-carboranyl)borane: An Isolable, Halogen-Free, Lewis Superacid, *Angew. Chem., Int. Ed.*, 2022, **61**, e202212073.
- 32 M. O. Akram, C. D. Martin and J. L. Dutton, The Effect of Carborane Substituents on the Lewis Acidity of Boranes, *Inorg. Chem.*, 2023, **62**, 13495–13504.
- 33 R. A. Thornton, K. French, K. Shuford and C. D. Martin, Reactivity of gold and copper acetylide with a secondary borane, *Chem. Commun.*, 2026, **62**, 1193–1196.
- 34 M. O. Akram, K. A. French, K. L. Shuford and C. D. Martin, A Crystalline Boravinylidene from an Arrested Hydroboration Reaction, *J. Am. Chem. Soc.*, 2025, **147**, 33120–33126.
- 35 K. Vashisth and C. D. Martin, Frustrated Lewis Pair Reactivity of Bis(1-methyl-ortho-carboranyl)borane, *Inorg. Chem.*, 2025, **64**, 24192–24200.
- 36 S. Ranasinghe, Y. Li, M. E. Andrews, M. O. Akram, R. A. Thornton and C. D. Martin, Müller versus Gutmann–Beckett for assessing the Lewis acidity of boranes, *Chem. Commun.*, 2025, **61**, 10182–10185.
- 37 A. Begum, M. O. Akram and C. D. Martin, Dearomatic C2-borylation of indoles, *Dalton Trans.*, 2025, **54**, 5664–5667.
- 38 D. W. Stephan and G. Erker, Frustrated Lewis Pair Chemistry: Development and Perspectives, *Angew. Chem., Int. Ed.*, 2015, **54**, 6400–6441.
- 39 M. Oestreich, J. Hermeke and J. Mohr, A unified survey of Si–H and H–H bond activation catalysed by electron-deficient boranes, *Chem. Soc. Rev.*, 2015, **44**, 2202–2220.
- 40 M. C. Lipke, A. L. Liberman-Martin and T. D. Tilley, Electrophilic Activation of Silicon–Hydrogen Bonds in Catalytic Hydrosilations, *Angew. Chem., Int. Ed.*, 2017, **56**, 2260–2294.



- 41 T. Hackel and N. A. McGrath, Tris(pentafluorophenyl) borane-Catalyzed Reactions Using Silanes, *Molecules*, 2019, **24**, 432.
- 42 P. Ríos, A. Rodríguez and S. Conejero, Activation of Si–H and B–H bonds by Lewis acidic transition metals and p-block elements: same, but different, *Chem. Sci.*, 2022, **13**, 7392–7418.
- 43 D. J. Parks and W. E. Piers, Tris(pentafluorophenyl)boron-Catalyzed Hydrosilylation of Aromatic Aldehydes, Ketones, and Esters, *J. Am. Chem. Soc.*, 1996, **118**, 9440–9441.
- 44 M. Alcarazo, C. Gomez, S. Holle and R. Goddard, Exploring the Reactivity of Carbon(0)/Borane-Based Frustrated Lewis Pairs, *Angew. Chem., Int. Ed.*, 2010, **49**, 5788–5791.
- 45 A. Y. Houghton, J. Hurmalainen, A. Mansikkamäki, W. E. Piers and H. M. Tuononen, Direct observation of a borane–silane complex involved in frustrated Lewis-pair-mediated hydrosilylations, *Nat. Chem.*, 2014, **6**, 983–988.
- 46 K. Chulsky and R. Dobrovetsky, B(C₆F₅)₃-Catalyzed Selective Chlorination of Hydrosilanes, *Angew. Chem., Int. Ed.*, 2017, **56**, 4744–4748.
- 47 A. Gudź, P. R. Payne and M. R. Gagné, Phosphines as Silylium Ion Carriers for Controlled C–O Deoxygenation: Catalyst Speciation and Turnover Mechanisms, *Organometallics*, 2017, **36**, 4047–4053.
- 48 N. Zwitter, S. P. Walg, F. Belaj and N. C. Mösch-Zanetti, Heterolytic Si–H Bond Cleavage at a Molybdenum-Oxido-Based Lewis Pair, *Chem. – Eur. J.*, 2018, **24**, 7149–7160.
- 49 Y. Liu, B. Su, W. Dong, Z. H. Li and H. Wang, Structural Characterization of a Boron(III) η^2 - σ -Silane-Complex, *J. Am. Chem. Soc.*, 2019, **141**, 8358–8363.
- 50 I. Avigdori, A. Pogoreltsev, A. Kaushanski, N. Fridman and M. Gandelman, Frustrated Lewis Pairs Comprising Nitrogen Lewis Acids for Si–H Bond Activation, *Angew. Chem., Int. Ed.*, 2020, **59**, 23476–23479.
- 51 C. Zhang, J. Wang, W. Su, Z. Lin and Q. Ye, Synthesis, Characterization, and Density Functional Theory Studies of Three-Dimensional Inorganic Analogues of 9,10-Diboraanthracene—A New Class of Lewis Superacids, *J. Am. Chem. Soc.*, 2021, **143**, 8552–8558.
- 52 C. Zhang, X. Liu, J. Wang and Q. Ye, A Three-Dimensional Inorganic Analogue of 9,10-Diazido-9,10-Diboraanthracene: A Lewis Superacidic Azido Borane with Reactivity and Stability, *Angew. Chem., Int. Ed.*, 2022, **61**, e202205506.
- 53 K. Vashisth, S. Dutta, M. O. Akram and C. D. Martin, Examining the reactivity of tris(ortho-carboranyl)borane with Lewis bases and application in frustrated Lewis pair Si–H bond cleavage, *Dalton Trans.*, 2023, **52**, 9639–9645.
- 54 C. Dong, X. Tan, Z. H. Li and H. Wang, C–H Silylation of Unactivated Arenes Catalyzed by a Borenium/Pyridine Lewis Pair, *Angew. Chem., Int. Ed.*, 2025, **64**, e20852.
- 55 R. Walsh, Bond dissociation energy values in silicon-containing compounds and some of their implications, *Acc. Chem. Res.*, 1981, **14**, 246–252.
- 56 S. Bähr and M. Oestreich, A Neutral RuII Hydride Complex for the Regio- and Chemoselective Reduction of N-Silylpyridinium Ions, *Chem. – Eur. J.*, 2018, **24**, 5613–5622.
- 57 A. Fernandes, C. Laye, S. Pramanik, D. Palmeira, Ö.Ö. Pekel, S. Massip, M. Schmidtman, T. Müller, F. Robert and Y. Landais, Chiral Memory in Silyl-Pyridinium and Quinolinium Cations, *J. Am. Chem. Soc.*, 2020, **142**, 564–572.
- 58 H. F. T. Klare, L. Albers, L. Süsse, S. Keess, T. Müller and M. Oestreich, Silylium Ions: From Elusive Reactive Intermediates to Potent Catalysts, *Chem. Rev.*, 2021, **121**, 5889–5985.
- 59 N. Kumar, C. Laye, F. Robert and Y. Landais, Quinoline-Based Silylium Ions: Synthesis, Structure and Lewis Acidity, *Eur. J. Org. Chem.*, 2021, 3613–3621.
- 60 Q. Feng, H. Wu, X. Li, L. Song, L. W. Chung, Y.-D. Wu and J. Sun, Ru-Catalyzed Geminal Hydroboration of Silyl Alkynes via a New gem-Addition Mechanism, *J. Am. Chem. Soc.*, 2020, **142**, 13867–13877.
- 61 A. F. Heyduk, J. A. Labinger and J. E. Bercaw, Catalytic Alcoholysis of Tetramethylsilane via Pt-Mediated C–H Bond Activation, *J. Am. Chem. Soc.*, 2003, **125**, 6366–6367.
- 62 M. Tobisu, M. Onoe, Y. Kita and N. Chatani, Rhodium-Catalyzed Coupling of 2-Silylphenylboronic Acids with Alkynes Leading to Benzosiloles: Catalytic Cleavage of the Carbon–Silicon Bond in Trialkylsilyl Groups, *J. Am. Chem. Soc.*, 2009, **131**, 7506–7507.
- 63 Y. Liang, W. Geng, J. Wei and Z. Xi, Palladium-Catalyzed Intermolecular Coupling of 2-Silylaryl Bromides with Alkynes: Synthesis of Benzosiloles and Heteroarene-Fused Siloles by Catalytic Cleavage of the C(sp³)-Si Bond, *Angew. Chem., Int. Ed.*, 2012, **51**, 1934–1937.
- 64 M. Onoe, K. Baba, Y. Kim, Y. Kita, M. Tobisu and N. Chatani, Rhodium-Catalyzed Carbon–Silicon Bond Activation for Synthesis of Benzosilole Derivatives, *J. Am. Chem. Soc.*, 2012, **134**, 19477–19488.
- 65 Q.-W. Zhang, K. An and W. He, Rhodium-Catalyzed Tandem Cyclization/Si–C Activation Reaction for the Synthesis of Siloles, *Angew. Chem., Int. Ed.*, 2014, **53**, 5667–5671.
- 66 L. Li, Y. Zhang, L. Gao and Z. Song, Recent advances in C–Si bond activation via a direct transition metal insertion, *Tetrahedron Lett.*, 2015, **56**, 1466–1473.
- 67 Y. Liang, S. Zhang and Z. Xi, Palladium-Catalyzed Synthesis of Benzosilolo[2,3-*b*]indoles via Cleavage of a C(sp³)-Si Bond and Consequent Intramolecular C(sp²)-Si Coupling, *J. Am. Chem. Soc.*, 2011, **133**, 9204–9207.
- 68 M. Yu and X. Fu, Visible Light Promoted Hydroxylation of a Si–C(sp³) Bond Catalyzed by Rhodium Porphyrins in Water, *J. Am. Chem. Soc.*, 2011, **133**, 15926–15929.
- 69 J. Huang, F. Liu, X. Wu, J.-Q. Chen and J. Wu, Recent advances in the reactions of silacyclobutanes and their applications, *Org. Chem. Front.*, 2022, **9**, 2840–2855.
- 70 A. Sivaramakrishna, S. Pete, C. Mandar Mhaskar, H. Ramann, D. Venkata Ramanaiah, M. Arbaaz, M. Niyaz, S. Janardan and P. Suman, Role of hypercoordinated silicon(IV) complexes in activation of carbon–silicon



- bonds: An overview on utility in synthetic chemistry, *Coord. Chem. Rev.*, 2023, **485**, 215140.
- 71 L. H. Sommer, W. P. Barie and J. R. Gould, Kinetics and Mechanism of Methyl–Silicon Cleavage by Sulfuric Acid¹, *J. Am. Chem. Soc.*, 1953, **75**, 3765–3767.
- 72 H. Sakurai, K. Tominaga, T. Watanabe and M. Kumada, Aluminium chloride-catalyzed reactions of organosilicon compounds II. Facile syntheses of alkylchlorosilanes, -germanes, and -stannanes, *Tetrahedron Lett.*, 1966, **7**, 5493–5497.
- 73 D. H. O'Brien and C. M. Harbord, Cleavage of alkylsilanes by strong acids, *J. Organomet. Chem.*, 1970, **21**, 321–328.
- 74 D. D. Hopf and D. H. O'Brien, Cleavage of alkylsilanes by strong acids: IV. The reactions of alkyl dimethyl-fluorosilanes with fluorosulfonic acid, *J. Organomet. Chem.*, 1976, **111**, 161–169.
- 75 K. Barlos and H. Nöth, Eine einfache Methode zur Darstellung von Methylbisdibromid, *Chem. Ber.*, 1977, **110**, 3460–3461.
- 76 W. Haubold, A. Gemmler and U. Kraatz, Tetramethylsilan – ein Methylierungsreagenz für Bortrihalogenide/ Tetramethylsilane – a Methylating Agent for Borontrihalides, *Z. Naturforsch., B*, 1978, **33**, 140–141.
- 77 C. Wang, Q. Luo, H. Sun, X. Guo and Z. Xi, Lithio Siloles: Facile Synthesis and Applications, *J. Am. Chem. Soc.*, 2007, **129**, 3094–3095.
- 78 Q. Wu, Z.-W. Qu, L. Omann, E. Irran, H. F. T. Klare and M. Oestreich, Cleavage of Unactivated Si–C(sp³) Bonds with Reed's Carborane Acids: Formation of Known and Unknown Silylium Ions, *Angew. Chem., Int. Ed.*, 2018, **57**, 9176–9179.
- 79 Q. Wu, A. Roy, E. Irran, Z.-W. Qu, S. Grimme, H. F. T. Klare and M. Oestreich, Catalytic Difunctionalization of Unactivated Alkenes with Unreactive Hexamethyldisilane through Regeneration of Silylium Ions, *Angew. Chem., Int. Ed.*, 2019, **58**, 17307–17311.
- 80 H. F. T. Klare and M. Oestreich, The Power of the Proton: From Superacidic Media to Superelectrophile Catalysis, *J. Am. Chem. Soc.*, 2021, **143**, 15490–15507.
- 81 K. Matsuoka, N. Komami, M. Kojima, T. Mita, K. Suzuki, S. Maeda, T. Yoshino and S. Matsunaga, Chemoselective Cleavage of Si–C(sp³) Bonds in Unactivated Tetraalkylsilanes Using Iodine Tris(trifluoroacetate), *J. Am. Chem. Soc.*, 2021, **143**, 103–108.
- 82 T. He, H. F. T. Klare and M. Oestreich, Arenium-ion-catalysed halodealkylation of fully alkylated silanes, *Nature*, 2023, **623**, 538–543.
- 83 Z.-H. L. Xi Wang and H. Wang, Alkyl Exchange between Alkylsilanes and Hydroborenum Ion, *Chin. J. Org. Chem.*, 2023, **43**, 1852–1856.
- 84 S. Courtenay, C. M. Ong and D. W. Stephan, Phosphinimido Complexes of Silicon, Tin, and Germanium, *Organometallics*, 2003, **22**, 818–825.
- 85 T. Ochiai, T. Szilvási and S. Inoue, Facile Access to Stable Silylium Ions Stabilized by N-Heterocyclic Imines, *Molecules*, 2016, **21**, 1155.
- 86 (a) CCDC 2523192: Experimental Crystal Structure Determination, 2026, DOI: [10.5517/ccdc.csd.cc2qpl9b](https://doi.org/10.5517/ccdc.csd.cc2qpl9b); (b) CCDC 2523193: Experimental Crystal Structure Determination, 2026, DOI: [10.5517/ccdc.csd.cc2qplbc](https://doi.org/10.5517/ccdc.csd.cc2qplbc); (c) CCDC 2523194: Experimental Crystal Structure Determination, 2026, DOI: [10.5517/ccdc.csd.cc2qpplcd](https://doi.org/10.5517/ccdc.csd.cc2qpplcd); (d) CCDC 2523195: Experimental Crystal Structure Determination, 2026, DOI: [10.5517/ccdc.csd.cc2qppldf](https://doi.org/10.5517/ccdc.csd.cc2qppldf); (e) CCDC 2523196: Experimental Crystal Structure Determination, 2026, DOI: [10.5517/ccdc.csd.cc2qpplfg](https://doi.org/10.5517/ccdc.csd.cc2qpplfg); (f) CCDC 2523197: Experimental Crystal Structure Determination, 2026, DOI: [10.5517/ccdc.csd.cc2qplgh](https://doi.org/10.5517/ccdc.csd.cc2qplgh).

