



INORGANIC CHEMISTRY

FRONTIERS

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: V. Motornov and N. Limberg, *Inorg. Chem. Front.*, 2026, DOI: 10.1039/D6QI00530F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Well-defined bridged dicopper(III) chloride, azide, and amino complexes

Vladimir Motornov*^a and Niklas Limberg^aReceived 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

The biologically relevant bis-(μ -L)-dicopper(III) core with modified bridging ligands (L = Cl, N₃, NMe₂) mimicking the enzyme oxidation bis- μ -oxo-copper(III) intermediates, was prepared from the heterocubane [Cu(CF₃)₂(OH)]₄ and fully characterized, including X-ray crystallography. The electronic structure, effective oxidation state of copper, and natural charge distribution in these complexes were studied by DFT calculations. The oxidation state localized orbital (OSLO) method and XPS spectroscopy support a +3 oxidation state of copper in these complexes, albeit with a highly covalent Cu–C bonding, which is the most profound in the *chlorido*-complex. Potential of these complexes in synthetic chemistry was demonstrated on azidotrifluoromethylation of an alkyne by the azido complex [Cu(CF₃)₂(N₃)]₂ furnishing a vinyl azide, and alcohol *O*-difluoromethylation by the copper chloride complex.

Introduction

Stabilization of copper in the high oxidation state (+3) represents a fundamental challenge, which is essential for advancing the understanding of aerobic oxidation and metalloenzyme systems,¹ as well as cross-coupling reactions.² In particular, bridged dicopper(III) species are key intermediates in biochemistry.^{1,3} Here, dimeric bis- μ -oxo-Cu(III) complexes are the key scaffold in aerobic enzyme oxidation processes and oxygen activation, forming via rearrangement of the primary Cu(II)-peroxo adducts (Scheme 1, A).³ Mimicking these intermediates by gaining insights into similar dimeric high-valent copper species is a timely and challenging task for inorganic and organometallic chemists, given the elusiveness of the high-valent copper. In particular, well-defined dimeric copper(III) hydroxide compounds L_n-Cu(III)- μ ₂-OH-Cu(III)-L_n (where L_n = 2,6-pyridinedicarboxamide ligands) reported by Tolman are key models for putative intermediates in oxidation processes.⁴ Recently, direct formation of the bridged oxo-dicopper(III) core in aerobic oxidation was observed and studied by spectroscopic methods.⁵ However, due to low stability and propensity of amine- and azole-ligated bridged dicopper(III) complexes to reduction (including intramolecular charge transfer),^{1,5} stabilization of well-defined dicopper(III) species is still difficult to realize.

The trifluoromethyl group is known to stabilize high oxidation states due to stabilizing metal-to-ligand π -back donation and contraction of σ -orbitals on a high-valent metal.^{2c,6} These factors preclude the reductive elimination. We have envisioned the use of a tetrameric copper(III) trifluoromethyl hydroxide

recently discovered by us⁷ for accessing novel dimeric Cu(III) species bearing two trifluoromethyl groups. Very recently, we have used this starting material to prepare copper(III) carboxylates,^{7a} 1,3-diketonates,^{7a} bench-stable synthetically useful scorpionates,⁸ and dimeric bridged copper(III) hydroxides.^{7b} To the best of our knowledge, dicopper(III) cores with other bridging ligands apart from oxo- and hydroxide groups are very rare. Only a single example of well-defined bridged formally copper(III) chloride complex was reported in 2022, where this unique core was stabilized in a metallacycle (Scheme 1, B).⁹ Nitrogen-bridged dicopper(III) complexes could be stabilized by a complex macrocyclic pyrazole-imidazolium NHC ligand,¹⁰ whereas formation of elusive phenoxide-bridged mixed Cu(II)-Cu(III) complex was observed spectroscopically from the well-defined Cu(II) dimer.¹¹ A common theme around all these species is presence of bulky substituents (such as mesityl)⁹ or macrocyclic environment¹⁰ to shield the high-valent copper centre. Use of trifluoromethyl groups for stabilization is a conceptually different alternative strategy. Herein we report isolation of novel elusive μ ₂-chlorido-, μ ₂-azido- and μ ₂-amino-dicopper(III) complexes prepared from easily accessible tetrameric hydroxide [Cu(CF₃)₂(OH)]₄ (**1**) with their characterization including X-ray crystallography and oxidation state analysis (Scheme 1, C).

^a Dr V. Motornov, N. Limberg, Freie Universität Berlin; Fabockstraße 34-36, 14195 Berlin, Germany. e-mail: cuprate51@gmail.com, motornov@zedat.fu-berlin.de
Supplementary Information available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x






ARTICLE

View Article Online
DOI: 10.1039/D6QI00530F

E

Journal Name

A. Enzyme oxidation pathway: formation of μ_2 -oxo-Cu(III) coreB. Formation of dimeric μ_2 -Cl-Cu(III) complexC. This work - formation of bridged μ_2 -(Cl, N₃, NMe₂)-Cu(III) analogs

Scheme 1. Key binuclear dicopper(III) core in enzyme oxidation, its chlorine-bridged analogue stabilized in bulky environment, and this work

Results and discussion


Inspired by the reactivity of the easily available high-valent copper hydroxide $[Cu(CF_3)_2(OH)]_4$ (**1**) as a hydroxide base,⁷ we attempted substitution of the hydroxyl group with a strong acid (HCl) or a secondary amine (HNMe₂), which could result in dehydration and formation of corresponding formally copper(III) chloride “ $Cu(CF_3)_2Cl$ ” or amino species. However, for reaction products formed by this direct method decomposition occurred even below $-40^\circ C$. Moreover, use of other common acidic chloro-dehydrative agents such as $SOCl_2$ and $POCl_3$ also resulted in decomposition. Therefore, we envisioned the use of trimethylsilyl derivatives as transfer reagents. To our delight, reaction of the heterocubane **1** with 2 equivalents of TMSCl (**2a**) at $-40^\circ C$ afforded a new product **3** formed as yellow crystals, which after recrystallization could be identified as di- μ_2 -chlorido-diaqua-tetrakis(trifluoromethyl)dicopper(III), a dimeric formally copper(III) complex with bridging μ_2 -chlorido ligands and two water molecules coordinated in apical positions (Scheme 2). X-ray crystallography revealed that the geometry of the well-known binuclear μ_2 -oxo-dicopper(III) core is

significantly altered when two hydroxyl ligands are replaced with chlorine atoms as in the compound **3**. Two copper centers with two chloride ligands form a rectangle with two nearly equal $Cl-Cu-Cl$ ($87.32(3)^\circ$) and two $Cu-Cl-Cu$ ($91.37(3)^\circ$) angles. There are two short $Cu-Cl$ bonds ($2.2181(8) \text{ \AA}$) at the opposite sides of the $Cu-Cl-Cu$ rectangle, and two significantly longer ($2.7572(9) \text{ \AA}$) contacts in the apical positions to the copper centers. This is remarkably different from the known dimeric square-shaped copper(III) chloride core, with nearly equal (2.318 and 2.327 \AA) $Cu-Cl$ distances.⁹ Water molecules are lying nearly in the same plane as two trifluoromethyl groups and one of the chloride ligands, in cis-configuration to each other. The $Cu-O$ bond lengths are $1.946(2) \text{ \AA}$, which indicates stronger coordination in contrast to previously reported copper(III) complex with water as a weakly coordinated ligand ($Cu-O = 2.316(1) \text{ \AA}$).⁷

The success of substitution of hydroxyl group with chloride prompted us to explore substitution of a hydroxyl with a chemically similar pseudohalide. Importantly, coordination of azido groups to transition metals can take place via either one or both terminal nitrogen atoms, thus formation of different isomers could be envisioned. To our delight, reaction of $TMSN_3$ (**2b**) with heterocubane **1** at $-40^\circ C$ afforded a bridged neutral copper(III) azide $[Cu(CF_3)_2(\mu-N_3)_2Cu(CF_3)_2]$ compound **4** with μ_2 - κ_1 - coordination of azide groups. It can be crystallized from a DCM/pentane mixture and can exist both in dimeric and polymeric form. If the crystallization is done in the presence of acetone, units of **4** with and without solvent form a coordination polymer of the composition $[Cu(CF_3)_2(\mu-N_3)_2Cu(CF_3)_2 \cdot 0.5C_3H_6O]_n$. The monomer unit of **4** is rhombus-shaped with four nearly equal $Cu-N$ distances of 1.950 - 1.956 \AA , which are slightly elongated in the solvated unit in the presence of coordinated acetone molecules. Surprisingly, compound **4** was found to be more stable than the respective chloride complex **3** and thus can be safely handled even at ambient conditions.

With these neutral halide and pseudohalide complexes in hand, we attempted to substitute one of the hydroxyl ligands with an amino group, which is more challenging due to a stronger $Si-N$ bond and instability of the formally copper(III) amino species. Application of (*N,N*-dimethylamino)trimethylsilane as a reaction partner at $-78^\circ C$ afforded a very unstable binuclear complex **5** with bridging dimethylamino- and hydroxyl groups (Scheme 2). Attempts to substitute both hydroxyl groups in excess of silane led to decomposition, which can be attributed to very low stability of bis(amino)dicopper(III) complex.





Scheme 2. Synthesis of bridged Cu(III) dimers chlorido-Cu(III) **3**, azido-Cu(III) **4**, and μ_2 -(amino)- μ_2 -(hydroxy)-Cu(III) **5**. Hydrogen atoms are omitted for clarity. Ellipsoids are shown at 50% probability level.

The crystal structure of **5** with the μ_2 -(dimethylamino)- μ_2 -(hydroxy)dicopper(III) core with square-planar copper centers revealed that both dimethylamino and hydroxyl groups are strongly coordinated, as shown by 1.850(4) and 1.878(4) Å copper-oxygen distances, whereas the Cu–N bonds lengths are 1.949(5) and 1.960(5) Å. The two N–Cu–O angles are 80.2(2)° and 79.3(2)° respectively.

With the structure of the complexes established, we investigated the bonding situation in these compounds by DFT calculations, as well as their structure in solution by optimization of geometry from XRD on B3LYP-D3(BJ)/ZORA-def2-TZVP level with a CPCM(DCM) solvation model. Geometry of **3** in DCM solution features more symmetrical tetragonal unit with Cu–Cl distances of 2.330 and 2.353 Å compared to more significant difference in the crystal structure, and elongated Cu–O bonds for coordinated water molecules (Figure 1). Both calculated frontier orbitals for all three complexes are mixed with predominantly ligand-centered character. HOMO of the

complex **5** has the most significant contribution of NMe₂ ligand, which could be the most reactive part of the molecule. LUMOs of the both complexes are centered along the Cu–C bonds, thus with a strong σ^* (Cu–CF₃) character, which resembles known copper trifluoromethyl complexes.^{6–8} The natural bonding orbital (NBO) analysis at ω B97X-D3(BJ)/def2-QZVPPD//B3LYP-D3(BJ)/ZORA-def2-TZVP level of theory revealed the positive NPA charge of +0.99 on copper for the complex **3** and +1.06 and +1.10 for the complexes **4** and **5** respectively, slightly higher than copper(III) species with three trifluoromethyl groups^{6b,c} and slightly lower than in the initial hydroxide **1**.⁷ Similar result (NPA +1.026) was observed recently for a nitrogen-coordinated Cu(III)-pseudopeptide complex.¹² In contrast to two Cu–CF₃ bonds with high degree of covalency (WBI = 0.50 in all three complexes), the Cu–Cl bonds in **3** display relatively low degree of covalency (WBI = 0.19–0.22 for all contacts) comparable to both Cu–N (WBI = 0.23) and Cu–O (WBI = 0.18) coordination in the case of **5**.



ARTICLE




bis- μ_2 -chloro complex **3**bis- μ_2 -azido complex **4** μ_2 -amino- μ_2 -hydroxy complex **5**

Figure 1. Frontier orbitals of complexes **3**, **4** and **5** (left) and their natural charge distribution (right). Minimum structures (ω B97X-D3(BJ)/def2-QZVPPD//B3LYP-D3(BJ)/ZORA-def2-TZVP, CPCM(DCM)) with blue numbers corresponding to the NPA charges

With insights into the structure and natural charge distribution in these dimeric formally copper(III) cores, we analysed the oxidation state in these complexes. The oxidation state assignment in formally copper(III) complexes has been a subject of long debates, since the occurring inverted ligand field, predominantly ligand-centered LUMOs, and increased d-shell population point to assignment of high-valent copper in these complexes as copper(I).¹³ However, the concept of oxidation states is formal by IUPAC definition,¹⁴ and re-assessment of this

assignment by XAS,¹⁵ XPS^{2d,8,12,16} and the recently developed OSLO (Oxidation State Localized Orbitals)¹⁷ method provides more evidence for existence of Cu(III),^{6b,8} despite the fact that d^8 -configuration is almost never the case in these compounds according to DFT calculations of orbital population.^{13,12,8} X-ray photoelectron spectroscopy for the more stable chloride **3** and azide **4** complexes supports the +3 oxidation state (see SI for details). The binding energy of $\text{Cu}2p_{3/2}$ was found to be 936.2 eV for the chloride (**3**) and 935.8 eV for the azide (**4**), which



excellently matches the literature reports, where the binding energies of 936 ± 1 eV indicate the copper(III) oxidation state, ca. 2 eV higher than the respective Cu(II) species.^{2d,8,12,16} The binding energies of Cu2p_{1/2} were found at ca. 956 eV (956.2 eV for **3** and 955.4 eV for **4**). These XPS results also corroborate the characteristics of the inorganic cuprate(III) species.¹⁸ Due to the instability of the complex **3** under XPS irradiation, Cu(I) decomposition product with 932 eV binding energy was detected under inert conditions.

According to the OSLO calculations (Table 1), Cu(III) oxidation state is also supported for all three complexes. However, for the chloride **3** the assignment is the least clear, which is reflected by high FOLI value close to the most ambiguous case (FOLI = 2), as well as low Δ -FOLI values below the confidence value of 0.5 indicating clear OSLO assignment.¹⁷ This discrepancy can be explained by strong π -donor properties and polarizability of chloride, where π -donation significantly increases the electron density on copper atom, which additionally results in a lower NPA charge in the chloride complex (see above). Amino- and azido-complexes **4-5** are less electron-rich, and are more clearly described as Cu(III) compounds by OSLO method, in contrast to the predominantly ligand-centered LUMO suggesting ligand field inversion for all complexes.^{13a}

Table 1. Determination of effective oxidation state in dimers **3-5** using OSLO method. Single point energies were calculated on ω B97X-D3(BJ)/def2-TZVPPD//B3LYP-D3(BJ)/ZORA-def2-TZVP (level 1); or DLPNO-CCSD(T)/ZORA-def2-TZVP//B3LYP-D3(BJ)/ZORA-def2-TZVP (level 2) levels of theory.

| Compound | %Cu(3d) in LUMO ^a | OS(Cu) ^b OSLO | FOLI (level 1) | Δ -FOLI (level 1) | FOLI (level 2) | Δ -FOLI (level 2) |
|---|---------------------------------|-----------------------------|-------------------|-----------------------------|-------------------|-----------------------------|
| 3 | 32.8 | +3 | 1.95 | 0.30 | 1.86 | 0.46 |
| 4 | 38.0 | +3 | 1.92 | 0.41 | 1.69 | 0.74 |
| 5 | 39.5 | +3 | 1.82 | 0.54 | 1.54 | 1.07 |
| [Cu(CF ₃) ₄] ⁻ | 36.2 | +3 | 1.56 | 1.11 | 1.26 | 2.50 |


^a – Determined by Löwdin population analysis on ω B97X-D3(BJ)/def2-TZVPPD//B3LYP-D3(BJ)/ZORA-def2-TZVP level of theory. ^b – The determined oxidation state is the same for both levels used

Finally, we compared the correlation of the OSLO oxidation state assignments with the Cu(3d) contribution from the Löwdin population analysis in LUMO orbital,^{13a} which are very similar (32.8-39.5%) for all species considered here. While a slightly diminished Cu(3d) character was indeed observed for the chloride **3** with the most ambiguous situation (Δ -FOLI < 0.5), which is reflected by a fragment population of the last OSLO of 43% on copper and 57% on the CF₃ group (see SI for full details). No strong correlation was observed between %Cu(3d) and FOLI values in OSLO system. Consistently higher FOLI and lower Δ -FOLI for neutral complexes than for anionic species^{6c} such as [Cu(CF₃)₄]⁻ complex suggests higher bond covalency, whereas comparing Cu(3d) contributions directly does not reflect this tendency.


With these insights, reactivity and potential of complexes **3-5** in synthetic organic and organometallic chemistry was

demonstrated (Scheme 3). The azide complex **4**, most stable of all three, was capable of trifluoromethylazidation of alkyne **6** under photochemical conditions (Scheme 3, A) to furnish the vinyl azide **7** in 71% yield. Control experiments with TEMPO and detection of TEMPO-CF₃ adduct supported radical mechanism of this transformation (see SI for full details), which is also reflected in the $\sigma^*(\text{Cu}-\text{CF}_3)$ character of LUMO for the complex **4** (Figure 2 above). Moreover, a different mode of reactivity based on very recently reported difluorocarbene transfer from Cu-CF₃ complexes,¹⁹ was realized for the chloride complex **3**, to afford alcohol difluoromethylation product **8** (Scheme 3, B). While a direct difluorocarbene transfer is feasible in the presence of HCl,¹⁹ electrophilicity of copper(III) or copper(II) chloride intermediates can enable both HCl generation due to fast ligand exchange, and formation of the difluorocarbene intermediate.²⁰ Finally, formation of *N*-trifluoromethyl amine **9** was observed during decomposition of the pre-formed amino-complex **5** at room temperature. Formation of binuclear Cu^{III}-Cu^I and Cu^{II}-Cu^I complexes upon reductive elimination can be envisioned, which undergo quick degradation to form a mixture of [Cu^{III}(CF₃)₄]⁻ and Cu^I(CF₃) by-products, detected by ¹⁹F NMR (Scheme 3, C).


A. Azidotrifluoromethylation of alkyne



B. O-difluoromethylation of alcohol



C. Formation of *N*-(trifluoromethyl)amine



Scheme 3. Transformations and synthetic potential of dimeric formally copper(III) complexes **3-5** in alkyne difunctionalization (A), alcohol difluoromethylation (B), and amine *N*-trifluoromethylation (C).



Conclusions

In conclusion, novel well-defined dimeric formally Cu(III) complexes with bridging chloro, azido- and amino-ligands were prepared and fully characterized. The molecular structures in the solid state, the natural charge distribution, and the effective oxidation state of these complexes were studied by DFT calculations, including the OSLO method, which support an assignment of a +3 oxidation state of copper in these complexes. We believe the structure of these complexes will open new avenues in the use of high-valent copper dimers in organic synthesis and biochemistry and encourage further studies of this unique family of copper(III) complexes derived from the parent cubane $[\text{Cu}(\text{CF}_3)_2(\text{OH})_4]$.

Experimental

Preparation of di- μ_2 -chlorido-dicopper(III) complex 3

In a 10 ml round-bottomed Schlenk flask under argon atmosphere to the solid heterocubane **1** (21.8 mg, 0.025 mmol [0.1 mmol monomer]) dry dichloromethane (1 ml) was added and the mixture was cooled to -40°C . Trimethylsilyl chloride (25 μl , 0.2 mmol, 2 equiv.) was added with vigorous stirring. The mixture was stirred at -40°C for 1 hour; during this time the solution turned intensive yellow. It was concentrated to ca. 0.2 ml volume under vacuum, then layered with pentane (ca. 2 ml) containing 0.5% of acetone. Crystallization at -40°C for 2 days afforded yellow crystals of **di- μ_2 -chlorido-diaqua-tetrakis(trifluoromethyl)dicopper(III) 3** with a composition $[\text{Cu}(\text{CF}_3)_2\text{Cl} \cdot \text{H}_2\text{O}]_2$ (11 mg, 43%), which were dried by a stream of argon at -40°C to -20°C and collected.

Preparation of di- μ_2 -azido-dicopper(III) complex 4

In a 10 ml round-bottomed Schlenk flask under argon atmosphere, to the solution of heterocubane **1** (21.8 mg, 0.025 mmol [0.1 mmol monomer]) dry dichloromethane (0.5 ml) was added and the mixture was cooled to -40°C . TMSN_3 (26.3 μl , 0.2 mmol, 2 equiv.) was added with vigorous stirring. The mixture was additionally stirred for 1 hour at -40°C , then connected to high vacuum and evaporated to dryness while warming up to room temperature to completely remove silanol by-products and afford complex **4** $[\text{Cu}(\text{CF}_3)_2(\mu_2\text{-N}_3)]_2$ as a bright yellow solid (23.5 mg, 96%). Single crystals for X-ray crystallography were grown from 1:1 pentane/DCM with or without an addition of 0.5% of acetone at -20°C .

Preparation of μ_2 -(dimethylamino)- μ_2 -hydroxy-dicopper(III) complex 5

In a 10 ml Schlenk tube under argon atmosphere to the solid heterocubane **1** (21.8 mg, 0.025 mmol [0.1 mmol monomer]) dry dichloromethane (1 ml) was added and the mixture was cooled to -78°C (acetone/dry ice). (N,N-

Dimethylamino)trimethylsilane (8 μl , 0.05 mmol, 0.5 equiv.) was added with vigorous stirring via syringe. The resulting orange mixture was stirred at -78°C for 30 minutes, then layered with pentane (ca. 3 ml). Crystallization in the freezer at -80°C for 2 days afforded orange crystals of dimeric **μ_2 -dimethylamino- μ_2 -hydroxy-tetrakis(trifluoromethyl)dicopper(III) 5** (5.6 mg, 24%), which were immediately collected for X-ray and other measurements. The compound undergoes slow decomposition even at -78°C , and decomposes within minutes at ambient conditions to form dark green residue.

Author contributions

V. Motornov conceived an idea, led the project, acquired the funding, conducted the experiments and most calculations, and wrote the manuscript. N. Limberg measured X-ray crystal structures and partially conducted the calculations. Both authors contributed into the final version of this manuscript.

Conflicts of interest

There are no conflicts to declare.

Data availability

Data is available in supplementary material of this article. Crystallographic data for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre. CCDC 2533527 (for **3**), 2551392 (for **4**), 2539161 (for the acetone solvate of **4**), and 2533528 (for **5**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/structures.

Acknowledgements

Alexander von Humboldt Foundation is gratefully acknowledged for the financial support of this project. We thank Ahmet Toraman (FU Berlin) for the assistance with low temperature NMR measurements, Dr Philip Nickl (SupraFAB FU Berlin) for the XPS service and the HPC Service of FUB-IT for the computational resources.

Notes and references

- a) C. E. Elwell, N. L. Gagnon, B. D. Neisen, D. Dhar, A. D. Spaeth, G. M. Yee, W. B. Tolman, Copper-oxygen complexes revisited:



- structures, spectroscopy, and reactivity, *Chem. Rev* 2017, **117**, 2059–2107. b) J. Serrano-Plana, I. Garcia-Bosch, A. Company, M. Costas, Structural and reactivity models for copper oxygenases: cooperative effects and novel reactivities, *Acc. Chem. Res* 2015, **48**, 2397–2406. c) Bioinorganic and bioinspired chemistry of copper (Ed. X. Ottenwaelder, S. Herres-Pawlis), *Inorg. Chim. Acta*, 2018, **481**, 1–200.
- a) A. Casitas, X. Ribas, A reflection on recent advances in organometallic copper(III) chemistry, *Chem. Sci*, 2026, **17**, 694. b) A. Casitas, X. Ribas, The role of organometallic copper (III) complexes in homogeneous catalysis, *Chem. Sci*, 2013, **4**, 2301. c) Y. Luo, Y. Li, J. Wu, X. S. Xue, J. F. Hartwig, Q. Shen, Oxidative addition of an alkyl halide to form a stable Cu (III) product *Science*, 2023, **381**, 1072–1079. d) Q. Zhang, S. Tong, M.-X. Wang, Unraveling the Chemistry of High Valent Arylcopper Compounds and Their Roles in Copper-Catalyzed Arene C–H Bond Transformations Using Synthetic Macrocycles *Acc. Chem. Res* 2022, **55**, 2796–2810. e) A. J. Hickmann, M. S. Sanford, High-valent organometallic copper and palladium in catalysis, *Nature*, 2012, **484**, 177.
 - a) J. De Tovar, R. Leblay, Y. Wang, L. Woicik, A. Thibon-Pourret, M. Reglier, A. J. Simaan, N. Le Poul, C. Belle, Copper–oxygen adducts: new trends in characterization and properties towards C–H activation, *Chem. Sci*, 2024, **15**, 10308. b) B. Kim, K. D. Karlin, Ligand–Copper(I) Primary O₂-Adducts: Design, Characterization, and Biological Significance of Cupric-Superoxides, *Acc. Chem. Res*, 2023, **56**, 2197–2212. c) L. Siebe, C. Butenuth, A. Stammler, H. Bogge, S. Walleck, T. Glaser, Generation and Reactivity of μ -1,2-Peroxo Cu^{II}Cu^{II} and Bis- μ -oxo Cu^{III}Cu^{III} Species and Catalytic Hydroxylation of Benzene to Phenol with Hydrogen Peroxide, *Inorg. Chem*, 2024, **63**, 2627–2639.
 - M. Reza Halvagar, P. V. Solntsev, H. Lim, B. Hedman, K. O. Hodgson, E. I. Solomon, C. J. Cramer, W. B. Tolman, Hydroxo-Bridged Dicopper(II,III) and -(III,III) Complexes: Models for Putative Intermediates in Oxidation Catalysis, *J. Am. Chem. Soc.* 2014, **136**, 7269–7272.
 - a) T. A. G. Large, W. Keown, J. Brannon Gary, L. Chiang, T. D. P. Stack, Imidazolate-Stabilized Cu (III): Dioxygen to Oxides at Type 3 Copper Sites, *Angew. Chem. Int. Ed*, 2025, **64**, e202416967. b) W. Keown, T. A. G. Large, L. Chiang, E. C. Wasinger, T. D. P. Stack, Exclusive imidazole ligation to Cu III₂O₂ and Cu III Cu II₂O₂ cores, *Chem. Commun*, 2019, **55**, 7390–7393.
 - a) H. Liu, Q. Shen, Well-defined organometallic Copper (III) complexes: Preparation, characterization and reactivity, *Coord. Chem. Rev*, 2021, **442**, 213923. For examples of Cu-CF₃ complexes, see: b) A. M. Romine, N. Nebra, A. I. Konovalov, E. Martin, J. Benet-Buchholz, V. Grushin, Easy access to the copper (III) anion [Cu(CF₃)₄][–], *Angew. Chem. Int. Ed* 2015, **54**, 2745–2749. c) V. Motornov, M. Procházka, N. Alpuente, P. Salvador, P. Slavíček, B. Klepetářová, X. Ribas, P. Beier, Introducing Weakly Ligated Tris (trifluoromethyl) copper (III), *ChemistryEurope* 2024, **2**, e202400004. d) M. A. P. Ball, P. J. Myers, G. D. Ritch, J. K. Bower, C. E. Moore, N. K. Szymczak, S. Zhang, The Role of Electron Transfer in Copper-Mediated C(sp²)–H Trifluoromethylation, *Angew. Chem. Int. Ed*, 2024, **64**, e202420677. e) V. Motornov, L. Ackermann, Well-Defined Highly-Coordinated Copper (III) Iodide and Pincer Tris (trifluoromethyl) copper Complexes, *Chem. Eur. J*, 2024, **30**, e202401791. f) V. Motornov, B. Klepetářová, P. Beier, Cu (III) Trifluoromethyl Complexes with 1, 3-Diketonate Ligands and Their Versatile Reactivity in C–H Trifluoromethylation, *Adv. Synth. Catal.* 2023, **365**, 2858–2864. g) S.-L. Zhang, C. Xiao, H.-X. Wan, Diverse copper (iii) trifluoromethyl complexes with mono-, bi-and tridentate ligands and their versatile reactivity. *Dalton Trans* 2018, **47**, 4779–4784. h) D. Joven-Sancho, A. Echeverri, N. Saffon-Merceron, J. Contreras-García, N. Nebra, An Organocopper (III) Fluoride Triggering C–CF₃ Bond Formation, *Angew. Chem. Int. Ed* 2024, **136**, e202319412. i) D. Naumann, T. Roy, K.-F. Tehhe, W. Crump, Multifaceted Lewis Acidity of a High-Valent Copper(III) Hydroxide Complex, *ChemistryEurope*, 2026, accepted, DOI: 10.1002/ceur.70319.
 - V. Motornov, Copper(III) trifluoromethyl tris(pyrazolyl)borates, *Inorg. Chem. Front*, 2026, **13**, 3924–3930.
 - H. M. Bergman, D. D. Beattie, R. C. Handford, E. Rossomme, B. A. Suslick, M. Head-Gordon, T. R. Cundari, Y. Liu, D. Tilley, Copper(III) Metallacyclopentadienes via Zirconocene Transfer and Reductive Elimination to an Isolable Phenanthrocylobutadiene, *J. Am. Chem. Soc*, 2022, **144**, 9853–9858.
 - K. Hua, F. Xie, S. Ye, M.-T. Zhang, Three Distinct Oxidation States (II/II, II/III, and III/III) of Diorganocopper Complexes, *JACS Au* 2024, **4**, 4406–4414.
 - A. Thibon-Pourret, et al. Effect of Mono-electronic Oxidation of an Unsymmetrical Phenoxido Hydroxido Bridged Dicopper(II) Complex, *Inorg. Chem*, 2018, **57**, 12364.
 - R. Eerlapally, S. K. Vechalapu, S. Duraisamy, D. Lakshmi Hareendran, J. Debnath, D. Allimuthu and A. Draksharapu, A Cu(II) pseudopeptide complex: a surrogate to a formal Cu(III) species in water and an inhibitor of cancer cell proliferation, *Inorg. Chem. Front*, 2024, **11**, 5172.
 - a) I. M. DiMucci, J. T. Lukens, S. Chatterjee, K. M. Carsch, C. J. Titus, S. J. Lee, D. Nordlund, T. A. Betley, S. N. MacMillan, K. M. Lancaster, The myth of d⁸ copper (III), *J. Am. Chem. Soc*, 2019, **141**, 18508; b) R. C. Walroth, J. T. Lukens, S. N. MacMillan, K. D. Finkelstein, K. M. Lancaster, Spectroscopic evidence for a 3d¹⁰ ground state electronic configuration and ligand field inversion in [Cu(CF₃)₄]^{1–}, *J. Am. Chem. Soc*, 2016, **138**, 1922–1931; c) I. F. Leach, R. M. A. Havenith, J. E. M. N. Klein, Revisiting Formal Copper (III) Complexes: Bridging Perspectives with Quasi-d¹⁰ Configurations, *Eur. J. Inorg. Chem*, 2022, e202200247; d) W. Yan, A. T. Poore, L. Yin, S. Carter, Y.-S. Ho, C. Wang, S. C. Yachuw, Y.-H. Cheng, J. A. Krause, M.-J. Cheng, S. Zhang, S. Tian, W. Liu, Catalytically relevant organocopper (III) complexes formed through aryl-radical-enabled oxidative addition, *J. Am. Chem. Soc*, 2024, **146**, 15176; e) R. Hoffmann, T. Alvarez, C. Mealli, A. Falceto, T. J. Cahill III, T. Zeng, G. Manca, *Chem. Rev*, 2016, **116**, 8173; f) J. P. Snyder, Elusiveness of Cu^{III} Complexation; Preference for Trifluoromethyl Oxidation in the Formation of [Cu^I(CF₃)₄][–] Salts, *Angew. Chem. Int. Ed*, 1995, **34**, 80–81; g) M. Kaupp, H. G. von Schnering, *Angew. Chem. Int. Ed*, 1995, **34**, 986–987.
 - “Oxidation state” in *IUPAC Compendium of Chemical Terminology*, 5th ed. International Union of Pure and Applied Chemistry; 2025. Online version 5.0.0, 2025. <https://doi.org/10.1351/goldbook.O04365>.



E

Journal Name

- 15 B. L. Geoghegan, Y. Liu, S. Peredkov, S. Dechert, F. Meyer, S. DeBeer, G. E. Cutsail III, Combining Valence-to-Core X-ray Emission and Cu K-edge X-ray Absorption Spectroscopies to Experimentally Assess Oxidation State in Organometallic Cu(I)/(II)/(III) Complexes, *J. Am. Chem. Soc.*, 2022, **144**, 2520.
- 16 T. Takeyama, R. Masui, D. Shibata, K. Mitamura, F. Tani, T. Yamaguchi, T. Yajima, Y. Shimazaki, T. Ohta, N. Hamamoto, Revisiting the Oxidation State of [Cu(HIO₆)₂]⁵⁻: A Copper(III) Complex Between Classical Werner-Type and Inverted Ligand Fields, *Inorg. Chem.* 2026, **65**, 8210–8215.
- 17 M. Gimferrer, A. Aldossary, P. Salvador, M. Head-Gordon, Oxidation State Localized Orbitals: A Method for Assigning Oxidation States Using Optimally Fragment-Localized Orbitals and a Fragment Orbital Localization Index, *J. Chem. Theor. Comput.*, 2022, **18**, 309-322.
- 18 a) P. Steiner, V. Kinsinger, I. Sander, B. Siegwald, S. Hufner, The Cu Valence in the High T_c Superconductors and in Monovalent, Divalent and Trivalent Copper Oxides Determined from XPS Core Level Spectroscopy, *Z. Phys. B*, 1987, **67**, 497-502. b) D. G. Brown, U. Weser, X-ray Photoelectron Spectra of Copper(III) Complexes, *Inorg. Chem.*, 1980, **19**, 264-266.
- 19 S. Dai; C. Xie; S. Long; M. Luo; C. Chen; T. Li; W.-X. Lv; Y. R. Chi, Repurposing trifluoromethyl copper (III) complexes for difluoromethylation of saccharides and complex alcohols, *Nat. Commun.* 2026, *in press*, DOI: 10.1038/s41467-026-71706-3.
- 20 X. Zeng, Y. Li, Q.-Q. Min, X.-S. Xue, X. Zhang, Copper-catalysed difluorocarbene transfer enables modular synthesis, *Nat. Chem.* 2023, **15**, 1064-1073.



Data availability

All data (experimental procedures, computational and crystallographic details, analytical data) is available in supplementary material of this article. Crystallographic data for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre. CCDC 2533527 (for **3**), 2533528 (for **5**), and 2539161 (for **4**), contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/structures.

[View Article Online](#)

DOI: 10.1039/D6QI00530E

