


 Cite this: *RSC Adv.*, 2026, 16, 23217

A comprehensive review of the medicinal properties of *Terminalia arjuna* and its major constituent arjunolic acid: a natural product source for new drug leads

 Manohar Bhujel,^a Sai Giridhar Sarma Kandanur,^b Lakshminath Sripada^a and Nageswara Rao Golakoti^{*,a}

Plants have long been a primary source of natural products for medicine. *Terminalia arjuna*, a deciduous tree native to the Indian subcontinent, is renowned for its diverse biological properties, such as anti-cancer, anti-diabetic, and cardioprotective effects. Various parts of this tree have been used to treat numerous ailments such as cancer, diabetes, inflammation, and microbial infections, as mentioned in ancient medicinal texts such as *Ayurveda* and *Charaka Samhita*. Arjunolic acid, a pentacyclic triterpene, is a major constituent of *T. arjuna*, and exhibits a plethora of biological properties such as anti-cancer, anti-diabetic, anti-inflammatory, and antioxidant activities. Diverse research groups have utilized arjunolic acid as a template for synthesizing various semi-synthetic derivatives, which have been evaluated for their anti-cancer, anti-diabetic, and antioxidant properties. This review comprehensively compiles and discusses the literature related to *T. arjuna*, arjunolic acid and its derivatives, focusing on their medicinal properties and the mechanism of their biological activities.

 Received 24th March 2026
 Accepted 29th March 2026

DOI: 10.1039/d6ra02419j

rsc.li/rsc-advances

1. Introduction

Natural products, classified into alkaloids, carbohydrates, flavonoids, lipids, steroids, terpenes, *etc.*, have been a rich and consistent source of biologically important and chemically diverse molecules.^{1,2} In view of the recent surge in FDA approvals for natural-product-based drug leads,^{3,4} terpenes present a promising opportunity to identify new drug candidates, as they are the largest class among natural products. The major pentacyclic triterpene constituent of *T. arjuna*, arjunolic acid, has received widespread attention from various research groups^{5,6} for its diverse biological properties. The structural feasibility of arjunolic acid to undergo various semi-synthetic transformations has resulted in analogues with interesting *in vitro* and *in vivo* biological activity.^{7–9} Although efforts have been made to review the literature on *T. arjuna*^{10–16} and arjunolic acid,^{5,6,17} the present review aims to comprehensively compile and discuss the following areas: (a) *T. arjuna* and its medicinal properties, (b) the isolation of arjunolic acid from various sources and its biological activities, and (c) various semi-synthetic derivatives of arjunolic acid, along with their biological properties and action mechanisms, reported to date.

1.1 Morphology of *Terminalia arjuna* (*T. arjuna*)

Terminalia arjuna is a species in the genus *Terminalia* of the family Combretaceae, which includes approximately 20 genera and 600 species.¹⁸ The genus name *Terminalia* refers to the terminal arrangement of leaves on many species in this group. *T. arjuna* is found throughout the sub-Himalayan regions of Bihar, Delhi, Madhya Pradesh, Uttar Pradesh, and the Deccan Plateau in India, as well as in the forests of Sri Lanka, Bangladesh, Mauritius, and Myanmar.^{19,20} It is a deciduous tree with buttressed roots, a sturdy trunk, and horizontally spreading branches and can reach heights of 60–70 feet (Fig. 1). The bark has a single-layered epidermis with hair-like projections, along with periderm and secondary phloem in mature bark. Its bark is usually white-grey, turning pink depending on the season and age, and it flakes off in large, flat pieces.²¹

The leaves are conical, oblong or elliptic, typically 10–15 cm long and 4–7 cm wide, with a green or dark-green upper surface and a light-brown underside. The petioles have one or two prominent glands at the top.^{21,22} The flowers are pale yellow to milky white, bisexual, and grow in dense clusters at the branch tips or as axillary spikes. Each flower has 10 stamens and an ovary covered with reddish or yellow hairs, while the calyx is glabrous. The woody fruit is 2.5–3.5 cm long with five angled wings that curve upward. *T. arjuna* flowers from March to June, with fruit appearing between September and November.²¹ *T. arjuna* contains many phytochemicals that have been reported to exhibit biological properties (Table 1 and Fig. 2). Among

^aDepartment of Chemistry, Sri Sathya Sai Institute of Higher Learning, Prasanthi Nilayam, Puttaparthi, A.P., 515134, India. E-mail: gnageswararao@sssihl.edu.in
^bIndependent Researcher, Hyderabad-500094, Telangana, India




Fig. 1 Images of *T. arjuna*, flowers, fruits, leaves and bark.

them, arjunolic acid is the major constituent, and this compound has been the focus of diverse research groups as it was reported to exhibit a plethora of biological properties.⁵⁶

Apart from the major constituent arjunolic acid, many phytochemicals, as described in Fig. 2, have been reported to exhibit biological activities. This further affirms that *T. arjuna* has been a source of bioactive molecules, and it would be interesting to explore the potential of these other phytochemicals in various drug discovery programs.

1.2 Uses of *T. arjuna* in traditional medicine

T. arjuna has been valued since the Vedic era, referred to as *Arjun* in Ayurvedic texts. It appears in ancient medicinal literature, including the *Astang Hridayam*, *Charaka Samhita*, and *Sushruta Samhita*. The Ayurvedic scholar Vagbhattacharya was the first to mention the use of bark powder prepared from its stem for heart-related ailments, a use later supported by Chakradattam and Bhavasisram. In Ayurveda, *T. arjuna* is considered a cardi tonic and is used to treat conditions such as *kshata* (wounds), *kshaya* (emaciation), *visham* (poison), *raktavikara* (styptic), *medoroga* (diabetes), *prameha* (urinary disorders), and *vrana* (ulcers). It is also known to relieve painful urination in diabetic conditions. A decoction of *T. arjuna* bark in conjunction with white sandalwood is used to boost sperm count in men and to relieve menstrual pain and leukorrhoea in women. It is applied for skin ailments, fever, and general vitality enhancement.⁷³

Root paste is applied to the forehead to alleviate headaches. Leaf paste mixed with milk and sugar is given to treat spermatorrhoea, while water-soaked leaves are used for dyspepsia and burning sensations. In traditional veterinary medicine, leaves combined with *Syzygium cumini* and *Acacia catechu* are used to treat cattle diarrhoea.^{74,75} Healers in Tamil Nadu's Kanchipuram district use fruit paste topically for wounds and inhale boiled bark for headaches and toothache relief.⁷⁶ The Malamalar tribe in Parambikulam uses fresh leaf juice for earache, while the powdered bark is employed for heart issues.⁷⁷ In Madhya Pradesh's Surguja district, bark is used for treating fever and hypertension. In the Malkangiri district of Orissa, fresh bark is chewed as an antacid.⁷⁸ Rajasthan communities combine bark powder with ghee and milk to improve heart health.⁷⁹

Throughout India, pastes of the bark are applied to wounds for faster healing and to stop bleeding. The bark decoction is useful for treating asthma, dysentery, ulcers, and acne (when

mixed with honey).⁸⁰ Bark ash is used to treat snakebites and scorpion stings, while a bark paste is applied to the area of a fracture for quicker recovery and to reduce swelling from injuries.^{77,81} It is also employed for gleet, urinary issues, as an expectorant, aphrodisiac, tonic, and diuretic.⁸²

1.3 Medicinal properties of *T. arjuna*

1.3.1 Cardioprotective activity.

T. arjuna has been used for the treatment of heart disease since ancient times. A study on the effects of an aqueous extract of *T. arjuna* bark to understand the mechanisms involved in cardiovascular disorders was conducted, and the oleanane triterpenoids present in this extract were found to be primarily responsible for the plant's cardioprotective effects.⁸³ The bark decoction also significantly lowered body mass index and systolic blood pressure in angina patients without affecting normal kidney or liver function.⁸⁴ Studies showed that administering bark powder to angina patients significantly reduced the frequency of angina in both ischemic cardiomyopathy and post-myocardial infarction angina. This treatment improved the left ventricular ejection fraction. It reduced left ventricular mass, providing relief from coronary heart failure, and long-term use of the drug did not alter normal kidney or liver physiology.⁸⁵

The herbal formulation "Lipistat," which includes the bark of the arjuna tree, reduced the loss of myocardial high-energy phosphate stores and lactate accumulation in myocardial tissue in cases of isoproterenol-induced myocardial necrosis, also offering protection in ischemic heart disease. Additionally, *T. arjuna* bark extract improved symptoms and signs in New York Heart Association (NYHA) class patients, reducing left ventricular end-diastolic and end-systolic volume indices, while increasing left ventricular ejection fractions. Long-term treatment and evaluation of NYHA class patients showed improved symptoms, signs, and exercise tolerance.⁸⁶ It was observed that the ethanolic extract of the bark reduced total cholesterol, HDL cholesterol, triglyceride levels, LDL cholesterol, and LDL/HDL ratios, without impacting renal or hepatic function.⁸⁷

The hypolipidemic activity of *T. arjuna* was evaluated in experimentally induced atherosclerosis in rabbits fed on a cholesterol-rich diet, where it acted as a hypolipidemic agent, partially inhibiting atheroma formation and indicating an antiatherogenic role.⁸⁸ A study on "Hartone," an arjuna-based herbal formulation, found that 80% of patients receiving Hartone treatment experienced relief from stable angina



Table 1 Phytochemicals present in *T. arjuna* and their medicinal properties

Part	Phytochemicals	Medicinal properties	
Stem bark	Arjunolic acid (1) ^{23,24}	Cardioprotective, anti-cancer ^{7,25}	
	Arjunin (2) ^{23,24,26}	Anti-cancer ²⁷	
	Arjungenin (3) ^{23,24}	Cytotoxic ²⁸	
	Terminic acid (4) ²⁹	—	
	Terminolitin (5) ^{23,24}	—	
	Arjunic acid (6) ²⁶	Antioxidant ³⁰	
	2 α ,3 β -Dihydroxyurs-12,18-dien-28-oic acid-28-O- β -D-glucopyranosyl ester (7) ³¹	—	
	2 α ,3 β ,23-Trihydroxyurs-12,18-dien-28-oic acid-28-O- β -D-glucopyranosyl ester (8) ³¹	—	
	Quadranoside VIII (9) ³¹	—	
	Kajiichigoside F1 (10) ³¹	—	
	2 α ,3 β ,23-Trihydroxyurs-12,19-dien-28-oic acid-28-O- β -D-glucopyranosyl ester (11) ³¹	—	
	Arjunetin (12) ^{23,24}	Anti-viral, ³² antioxidant ³³	
	Arjunoside I (13), arjunoside II (14), arjunoside III(15), arjunoside IV (16) ^{34,35}	—	
	Arjunolone (17) ³⁶	—	
	Arjunolitin (18) ³⁷	—	
	Arjunaphthanolide (19) ^{38,39}	—	
	Arjunoglycoside I (20), arjunoglycoside II (21), arjunoglycoside III (22), arjunoglycoside IV (23), arjunoglycoside V (24) ^{40,41}	—	
	Olean-3 β ,22 β -diol-12-en-28-oic acid-28- β -D-glucopyranoside (25) ⁴²	—	
	Terminarjunoside I (26) and terminarjunoside II (27) ⁴³	—	
	Terminoside A (28) ⁴⁴	Inhibits NO production ⁴⁵	
	Terminoic acid (29) ⁴⁴	—	
	Arjunone (30) ³⁶	—	
	Luteolin (31) ³⁶	Antioxidant, antimicrobial, anticancer ⁴⁶	
	Baicalein (32) ³⁶	Anticancer, antimicrobial, anti-biofilm ⁴⁷	
	Ethyl gallate (33) ³⁶	—	
	Galic acid (34) ³⁶	Antioxidant ⁴⁸⁻⁵⁰	
	Kempferol (35) ³⁶	Anti-cancer, anti-inflammatory ^{51,52}	
	Oligomeric proanthocyanidin (36) ³⁶	—	
	Pelargonidin (37) ³⁶	—	
	Quercetin (38) ³⁶	Antimicrobial, anticancer ⁴⁷	
	(+) catechin (39), (+) gallo catechin (40) and (–) epigallocatechin (41) ⁵³	Antioxidant ⁵⁴	
	Galic acid (34), ellagic acid (42) ⁵³	Antioxidant, hepatoprotective. ⁴⁸⁻⁵⁰	
	3-O-Methyl-ellagic acid-4-O- β -D-xylopyranoside (43) ⁵³	—	
	3-O-Methyl ellagic acid-3-O- β -D-rhamnoside (44) ⁵³	—	
	3-O-Methyl ellagic acid-4'-O- α -L-rhamnopyranoside (45) ³¹	—	
	(–)-Epicatechin (46) ³¹	Antioxidant ⁵⁴	
	Pyrocatechol (47) ⁵⁵	Antioxidant ⁵⁴	
	Punicallin (48) ⁵⁶	Anti-cancer ⁵⁷	
	Castalagin (49) ⁵⁸	Antibacterial, antiviral ⁵⁹	
	Casuarinin (50) ⁵⁸	Anti-cancer, antioxidant ^{60,61}	
	Casuarinin (51) ⁵⁸	Anti-cancer, anti-inflammatory ^{60,61}	
	Puncicalagin (52) ⁵⁸	Antioxidant ⁵⁷	
	Terchebulin (53) ⁵⁸	Antibacterial ⁶²	
	Terflavin C (54) ⁵⁸	Antimicrobial ⁶³	
	β -Sitosterol (55) ²⁹	Antimicrobial ⁶⁴	
	Roots	Arjunoside I-IV (13–16) ²⁹	—
		Arjunolic acid (1) ²⁹	Cardioprotective, anti-cancer ^{7,25}
Oleanolic acid (56) ²⁹		Antimicrobial, anti-cancer ⁶⁵	
Terminic acid (4) ²⁹		—	
2 α ,19 α -Dihydroxy-3oxo-olean-12-en-28-oic acid-28-O- β -D-glucopyranoside (57) ⁶⁶		—	



Table 1 (Contd.)

Part	Phytochemicals	Medicinal properties
Fruits	Arjunic acid (6) ^{23,24}	Antioxidant ³⁰
	Arjunetoside (3-O-β-D-glucopyranosyl-2α,3β,19α-trihydroxyolean-12-en-28-oic acid-28-O-β-D-glucopyranoside) (58) ⁶⁷	—
	Arjunic acid (6) ⁶⁸	Antioxidant ³⁰
	Arjunone (30) ⁶⁸	—
	Arachidic acid (59) and stearic acid (60) ⁶⁸	Anti-inflammatory ⁶⁹
	Cerasidin (61) ⁶⁸	—
	Ellagic acid (42) ⁶⁸	Antioxidant. ⁴⁸⁻⁵⁰
	Friedelin (62) ⁶⁸	Anti-inflammatory, antibacterial ⁷⁰
	Gallic acid (34) ⁶⁸	Antioxidant ⁴⁸⁻⁵⁰
	Hentriacontane (63) ⁶⁸	—
Leaves	Methyl oleate (64) ⁶⁸	—
	Myristyl oleate (65) ⁶⁸	—
	β-Sitosterol (55) ⁶⁸	Anti-inflammatory ⁶⁴
	Gallic acid (34) ⁷¹	Antioxidant ⁴⁸⁻⁵⁰
	Luteolin (31) ⁷¹	Antimicrobial, anticancer. ⁴⁶
	Quercetin (38) ⁷¹	Antimicrobial, anticancer ⁴⁷
	Catechin (39) ⁷¹	Antioxidant ⁵⁴
	Epicatechin (46) ⁷¹	Antioxidant ⁵⁴
Apigenin (66) ⁷¹	Antibacterial, anticancer ⁷²	

pectoris compared to those on isosorbide mononitrate (ISMN), with a higher reduction in heart attack rates than ISMN. This formulation also improved blood pressure response to stress without disturbing renal or hepatic function.⁸⁹


In rabbits, oral administration of a suspension of bark powder of *T. arjuna* for 12 weeks increased myocardial antioxidants, superoxide dismutase, catalase, and glutathione levels, while inducing heat-shock protein-72. In cases of ischemic reperfusion injury, oxidative stress and heart tissue injury were prevented in bark powder-administered rabbit hearts. In these treated rabbits, left ventricular end-diastolic pressure normalized within 10 minutes of reperfusion, and myocardial lipid peroxidation levels were lower.⁹⁰ Bark powder treatment significantly reduced total cholesterol and LDL cholesterol levels and decreased lipid peroxide levels.⁹¹ The bark powder of *T. arjuna* reduced ischemic mitral regurgitation (IMR), improved the E/A ratio (indicating early and late phases of ventricular filling), and reduced angina frequency.⁹⁰ In acute myocardial infarction, a 500 mg oral dose of *T. arjuna* bark powder administered for three months decreased ischemic mitral regurgitation and significantly reduced angina frequency.⁹²

Another study based on the hypothesis that the aqueous stem bark extract of *T. arjuna* (TAE) could inhibit IL-18-induced atherosclerosis through the NF-κB/PPAR-γ-mediated pathway in ApoE^{-/-} mice was conducted. Twelve-week-old male ApoE^{-/-} mice, fed with a standard chow diet, were divided into four groups (*n* = 6 per group): Group I (control): Phosphate-buffered saline (PBS) for 2 months. Group II: Recombinant IL-18 (rIL-18) for 1 month, followed by PBS for 1 month. Group III: rIL-18 for 1 month, followed by TAE for 1 month. Group IV: rIL-18 for 1 month, followed by atorvastatin for 1 month. Treatment with IL-

18 significantly elevated pro-inflammatory IL-18 levels (1178.66 ± 8.08 pg mL⁻¹ vs. 170 ± 9.16 pg mL⁻¹ in controls, *p* < 0.001) and downregulated the cholesterol efflux gene PPAR-γ (0.6-fold vs. 1.00 in control mice). Administration of TAE significantly reduced IL-18 levels (281.66 ± 9.60 pg mL⁻¹ vs. 1178.66 ± 8.08 pg mL⁻¹) and upregulated PPAR-γ expression (1.5-fold vs. 0.6-fold in the IL-18-treated group). TAE also decreased atherogenic lipids and reduced the percentage of atherosclerotic lesion area, showing comparable efficacy to atorvastatin. The findings suggest that TAE mitigates IL-18-induced atherosclerosis via the NF-κB/PPAR-γ-mediated pathway, highlighting its potential as a therapeutic agent.⁹³

Another study investigated the therapeutic and prophylactic potential of *T. arjuna* bark extract in a rat model of isoproterenol (ISO)-induced chronic heart failure (CHF). Rats were administered isoproterenol (ISO) at a dose of 85 mg kg⁻¹ subcutaneously, twice at an interval of 24 hours, to induce CHF. Fifteen days post-injection, the rats exhibited reduced cardiac function, including declines in the maximal rate of rise and fall of left ventricular pressure (LV (dP/dt) max and LV (dP/dt) min), cardiac contractility index (LV (dP/dt) max/LVP), and cardiac output. Additionally, an increase in left ventricular end-diastolic pressure was observed. The CHF rats also showed significantly elevated serum levels of creatine kinase isoenzyme-MB (CK-MB) and malondialdehyde (MDA), along with reduced activities of antioxidant enzymes, such as superoxide dismutase (SOD), and reduced glutathione (GSH). Alterations in lipid profiles, elevated levels of the pro-inflammatory cytokine tumor necrosis factor-alpha (TNF-α), and histological heart damage were also evident. Treatment with *T. arjuna* bark extract (500 mg kg⁻¹, orally) both before and for 15 days after ISO administration significantly improved cardiac function and reduced



Fig. 2 Structures of various chemical constituents isolated from *T. arjuna*.




Fig. 2 (Contd.)






Fig. 2 (Contd.)







Fig. 2 (Contd.)






Fig. 2 (Contd.)

myocardial injury. The cardioprotective effects of *T. arjuna* were comparable to those of fluvastatin, a synthetic drug. The bark extract effectively attenuated cardiac dysfunction and myocardial damage by restoring endogenous antioxidant enzyme activity, reducing lipid peroxidation, and lowering cytokine levels.⁹⁴

An alcoholic bark extract at a dose of 0.75 mg kg⁻¹ body weight normalized *in vivo* myocardial injury in rats. In a double-blind, placebo-controlled study on chronic stable angina patients, *T. arjuna* bark extract improved treadmill and clinical exercise parameters compared to placebo, yielding results similar to isosorbide mononitrate therapy.⁹⁵ A 6 mg kg⁻¹ dose of a 70% alcoholic extract of the bark produced a hypotensive effect in anesthetized dogs, which was blocked by propranolol, suggesting that active constituents in the extract may have adrenergic beta-2 receptor potential, directly affecting heart muscle and supporting its cardiovascular use in Ayurveda.⁷³

Dilated cardiomyopathy with reduced left ventricular ejection fraction (LVEF), whether due to idiopathic or ischemic causes, showed significant improvement in left ventricular parameters and functional capacity with standard therapy supplemented by 50% ethanol extract of *T. arjuna* bark. In

diabetic rats, 50% ethanol bark extract significantly mitigated cardiac dysfunction and myocardial injury, reducing oxidative stress, endothelin-1 (ET-1), and inflammatory cytokine levels. However, the decreased body weight, heart rate, blood pressure, and elevated blood sugar levels in diabetic rats did not improve with 50% ethanol bark extract therapy. These findings suggest that *T. arjuna* bark extract may enhance altered myocardial function in diabetic rats, possibly by maintaining endogenous antioxidant enzyme activities and reducing ET-1 and cytokine levels.⁹⁶

1.3.2 Anti-cancer activity. Tannins, terpenoids and flavonoids from the leaves, bark, and stems of *T. arjuna* are reported as the primary contributors to its anti-cancer properties.²⁸ Acetone extract of bark containing tannins has been shown to inhibit mutagenicity in *Salmonella typhimurium*.⁹⁷ Methanolic and acetone extracts have been reported to inhibit the growth of transformed cells in normal human fibroblasts, osteosarcoma, and glioblastoma.⁹⁸ Additional studies suggest that ethyl acetate and acetone extracts, which contain both non-polar and polar compounds, display the highest mutagenic activity against the mutagen 4-nitroquinoline-N-oxide (4NQO) compared to chloroform and methanolic-HCl extracts, highlighting their anti-



cancer potential.⁹⁹ Comet assays and micronuclei tests indicate that acetone and methanolic extracts are more effective than others in reducing DNA damage caused by 4NQO.⁹⁹ Leaf extracts of *T. arjuna* also exhibit moderate cytotoxic activity against BT-20 human breast cancer cells.²⁷ Additionally, bark extracts were reported to protect DNA from ADR-induced damage, and aqueous stem bark extracts demonstrate antioxidant effects that reduce oxidative stress and inhibit anaerobic metabolism, contributing to anti-carcinogenic activity.¹⁰⁰ *In vitro* screening of petroleum ether bark extracts against two human cancer cell lines, HEP2 (liver) and HT29 (colon), using the Sulforhodamine B (SRB) assay showed growth inhibition rates of 78% and 79.33%, respectively.¹⁰¹

1.3.3 Anti-bacterial activity. Studies have shown that *T. arjuna* bark exhibits antibacterial properties. Methanolic leaf extracts of *T. arjuna* demonstrated strong antibacterial activity against multi-drug-resistant *Salmonella typhi*.¹⁰² Research on the leaves and bark also indicated their potential to treat ear infections caused by bacteria.¹⁰³ Aqueous and methanolic extracts of *T. arjuna* effectively inhibited the growth of various bacterial species, including *E. coli*, *Klebsiella* sp., *Pseudomonas* sp., and *Staphylococcus* sp., in a dose-dependent manner.¹⁰⁴ Methanolic extracts from the fruit of *T. arjuna* also showed significant antibacterial activity against *Staphylococcus aureus* and *Pseudomonas aeruginosa*.¹⁰⁵ The *n*-butanol fraction of the methanolic extract of bark and the *n*-butanol fraction of the methanol extract of leaves exhibited strong antimicrobial effects against *Bacillus subtilis*, *E. coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Salmonella typhi*.¹⁰⁶

Shreya *et al.* studied the bark extract of *T. arjuna* for its antimicrobial properties. Methanolic extracts of bark exhibited a stronger inhibitory effect against Gram-negative bacteria compared to Gram-positive bacteria.¹⁰⁷

The ethanolic bark extract inhibited *Vibrio cholera* growth at all concentrations, with zone diameters increasing proportionally to concentration. Regression coefficients for the relationship between concentration and zone diameter ranged from 0.75 to 0.984 for most isolates, suggesting a linear relationship.¹⁰⁸

Methanolic and water extracts of *T. arjuna* bark showed significant inhibition zones against 22 tested bacteria, including eight uropathogens, with minimum inhibitory concentration (MIC) values between 0.16 and 2.56 mg mL⁻¹. The chloroform extract showed no antibacterial activity. The aqueous extract of *T. arjuna* bark also demonstrated strong antifungal effects against eight *Candida* species, with MIC values ranging from 0.16 to 0.64 mg mL⁻¹. The high antimicrobial efficacy of polar extracts correlated with their elevated polyphenol content, in contrast to the non-polar chloroform fraction.¹⁰⁹

Aqueous extracts of *Terminalia arjuna* (fruit, bark, root, and leaves) demonstrated notable antibacterial properties. The bark extract effectively inhibited the growth of both Gram-positive and Gram-negative bacteria. In contrast, the inhibitory effect of the leaf extract varied depending on the microorganism. Aqueous extracts from the leaves and fruits showed greater

activity against Gram-negative bacteria compared to Gram-positive bacteria. Among the microorganisms tested, the highest inhibition was observed in *Escherichia coli*, followed by *Pseudomonas aeruginosa*, *Bacillus cereus*, *Bacillus subtilis*, and *Lactobacillus bulgaricus*. However, *Micrococcus glutamicus* and *Micrococcus luteus* exhibited lower sensitivity to the bark extract. Overall, the bark extract showed the strongest antibacterial activity across most strains, except for *M. glutamicus*. However, leaf and fruit extracts were not active toward any of the Gram-positive strains used in the study.¹¹⁰

Meena *et al.* evaluated the antimicrobial efficacy of sequentially fractionated solvent extracts from *T. arjuna* (Roxb) against key pathogens in aquaculture. The study examined solvent extracts from the bark, fruit, and leaves of *T. arjuna* against 17 bacterial isolates from various species and the fungus *Aphanomyces invadans*. The minimum inhibitory concentration (MIC) for the bacterial isolates ranged from 0 to 25 µg mL⁻¹, with an average of 11.39 ± 0.48 µg mL⁻¹. Among the extracts, the ethanolic bark extract demonstrated the largest inhibition zone (20.83 ± 0.10 mm), followed by the ethanolic fruit extract (19.50 ± 0.06 mm) against *Edwardsiella tarda*. The solvent extracts showed efficacy at concentrations of 0.5 mg mL⁻¹ and 1 mg mL⁻¹ for bacterial isolates and fungal strains, respectively. The chemical analysis identified prominent bioactive compounds, including ellagic acid (*m/z* 302.01), 18-glycyrrhetic acid (*m/z* 470.34), azelaic acid (*m/z* 188.10), and caffeic acid (*m/z* 180.04), with a fit of over 80% in mzCloud analysis. Additionally, naringenin (*m/z* 272.06) was detected in the acetone bark extract, while kanamycin (*m/z* 426.35) and tetracycline (*m/z* 444.415) were present in the ethanol bark extract. Orientin (*m/z* 448.10) and phloretin (*m/z* 274.08) were identified in the leaf extracts. Results show that the ethanolic bark extract exhibited the most significant antimicrobial activity, followed by the methanolic bark extract.¹¹¹

1.3.4 Antioxidant activity. The protective effects of ethanolic extracts of *T. arjuna* bark and its fractions (dichloromethane (TAD), ethyl acetate (TAE), butanol (TAB), and water (TAW)) were examined for their ability to counteract free radicals, protein oxidation, and DNA damage. The TAE fraction demonstrated the highest inhibition of DPPH, hydroxyl, ABTS, nitric oxide radicals, and metal chelation, with IC₅₀ values of 270 ± 2 µg mL⁻¹, 175 ± 11 µg mL⁻¹, 25 ± 1.2 µg mL⁻¹, 82 ± 4 µg mL⁻¹, and 405 ± 9 µg mL⁻¹, respectively. According to the study, *T. arjuna* bark extracts ameliorated various impairments linked to DNA damage and free radical formation.¹¹²

The stem bark alcoholic extract of *T. arjuna* (ALTA) was also tested for antioxidant and antimutagenic (anticlastogenic) activities. ALTA displayed potent antioxidant properties with EC₅₀ values of 2.491 ± 0.160 µM, 50.110 ± 0.150 µM, and 71.00 ± 0.250 µM in DPPH, lipid peroxidation, and superoxide radical scavenging assays, respectively, comparable to ascorbic acid (EC₅₀ of 2.471 ± 0.140, 40.500 ± 0.390, and 63.00 ± 0.360). In a micronucleus test, ALTA (100 and 200 mg kg⁻¹, orally) significantly reduced the percentage of micronuclei in both normochromatic erythrocytes (NCE) and polychromatic erythrocytes (PCE) and showed a significant reduction in the P/N



ratio. These findings suggest that ALTA has notable antioxidant and antimutagenic effects.¹¹³

The free-radical scavenging capacity and antioxidant activity of bark extracts of *T. arjuna* (aqueous : ethanol (20 : 80 v/v) and aqueous : methanol (20 : 80 v/v)) were also evaluated, with extraction yields ranging from 6.66–19.09 g/100 g (w/w) on a dry-weight basis. The extracts contained substantial amounts of total phenolic content (TPC) at 6.02–11.00 g/100 g (as gallic acid equivalent) and total flavonoid content (TFC) at 1.75–5.96 g/100 g (as catechin equivalent). DPPH radical-scavenging activity had IC₅₀ values of 2.71–7.68 μg mL⁻¹, inhibition of peroxidation ranged from 64.79 to 71.43%, and reducing power ranged from 0.001 to 1.584 mg mL⁻¹. These results indicate that *T. arjuna* bark extracts are a naturally rich source of antioxidants.¹¹⁴ An *in vitro* model using goat red blood cells subjected to Cu²⁺-ascorbate-induced oxidative stress demonstrated that aqueous *T. arjuna* bark extract reduced lipid peroxidation, increased the reduced glutathione levels, and decreased protein carbonyl content in treated RBCs. The aqueous extract also protected the activity of antioxidant enzymes catalase and superoxide dismutase (SOD) and exhibited scavenging effects on hydroxyl and superoxide anion radicals.¹¹⁵ The effects of ethanolic extract (at doses of 250 and 500 mg kg⁻¹ body weight) of *T. arjuna* stem bark were evaluated in alloxan-induced diabetic rats, specifically examining lipid peroxidation and enzymatic and non-enzymatic antioxidant activity in liver and kidney tissues. The extract significantly ($P < 0.05$) reduced lipid peroxidation (LPO), with the 500 mg kg⁻¹ dose showing a greater effect than the 250 mg kg⁻¹ dose. Additionally, the extract significantly ($P < 0.05$) increased levels of superoxide dismutase, catalase, glutathione peroxidase, glutathione-S-transferase, glutathione reductase, glucose-6-phosphate dehydrogenase, reduced glutathione, vitamins A, C, and E, as well as total sulphhydryl groups (TSH) and non-protein sulphhydryl groups (NPSH) in the liver and kidney of diabetic rats. These findings suggest that the extract exhibits antioxidant activity by mitigating oxidative stress, supporting the traditional use of this plant in managing diabetes.¹¹⁶

The aqueous extract of *T. arjuna* bark demonstrated protective effects on liver and kidney tissues against CCl₄-induced oxidative stress.¹¹⁷ Methanol and ethanol extracts of *T. arjuna* seeds showed high antioxidant activity by inhibiting DPPH (88% ± 1.52% and 87% ± 0.057% inhibition, respectively), which was significant, although slightly lower than that of standard ascorbic acid (93.73% ± 1.12%).¹¹⁸ Alcoholic extracts of the bark exhibited notable radical-scavenging activity against DPPH and nitric oxide radicals.¹¹⁹

In another study, oral administration of CCl₄ at a dose of 1 mL kg⁻¹ body weight for 2 days significantly reduced the activities of antioxidant enzymes—catalase (CAT), glutathione-S-transferase (GST), and superoxide dismutase (SOD)—and depleted reduced glutathione (GSH) levels in cardiac tissue. Lipid peroxidation and oxidized glutathione (GSSG) levels were increased under these conditions. However, oral administration of the active constituents of *T. arjuna* (50 mg kg⁻¹ body weight) for 7 days before CCl₄ exposure significantly restored antioxidant enzyme activities, increased GSH levels, and reduced lipid

peroxidation products. The FRAP assay showed that the active constituents also enhanced intracellular antioxidant activity in cardiac tissue. Histological studies further supported the cardioprotective effects of these active constituents, which were comparable to those of vitamin C, a known antioxidant.¹²⁰

Shreya *et al.* used thin-layer chromatography to separate the flavonoid components of *T. arjuna* and evaluated the phytochemical composition, antimicrobial activity, antioxidant activity, and total flavonoid content. Phytochemical screening was conducted using standard and commonly available tests. Antioxidant activity and the presence of flavonoid compounds were analyzed using thin-layer chromatography (TLC). Total antioxidant capacity was determined using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) colorimetric assay. The methanolic extract showed significant antioxidant activity, as evidenced by a decrease in DPPH radical absorption during the scavenging assay. Flavonoid components with antioxidant properties were present in the methanolic extract at a concentration of 199.00 mg quercetin equivalent per gram of dried extract, as determined by the colorimetric method.¹⁰⁷

Meena *et al.* investigated the antioxidant, free-radical scavenging, and DNA-nicking inhibition activities of 21 solvent extracts of *T. arjuna* under hydrogen peroxide (H₂O₂) conditions. The study revealed a strong positive correlation and interaction (explaining 84.54% variation in PCA 1) between 2,2-diphenyl-1-picrylhydrazyl (DPPH), ferric reducing antioxidant power (FRAP), 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulphonic acid, and nitrous oxide scavenging (%) with ethanol bark extract (EBE) and methanol fruit extract (MFE). The total phenolic content (TPC) ranged from 144.67 to 1794 μg mL⁻¹ (gallic acid equivalent, GAE), and total flavonoid content (TFC) ranged from 2.5 to 34 μM Fe(II)/g, with EBE exhibiting the highest values. Among the solvent extracts, EBE demonstrated the greatest antioxidant capacity, followed by ELE, MBE, and DWFE. The DNA nicking inhibition analysis showed significant positive correlations ($p < 0.01$) for relative front, relative quantity, band intensity (%), and lane intensity (%), with an R^2 value of 0.94. EBE was found to nick supercoiled, circular plasmid DNA effectively, showing comparable activity to standard antioxidants such as butylated hydroxyanisole, butylated hydroxytoluene, ascorbic acid, and gallic acid. Notably, gallic acid (m/z 170.0208) and ellagic acid (m/z 302.0063) were present in high concentrations in the extracts, with an effective concentration of 0.48 mg required for DNA nicking inhibition. This work also reports higher steroid concentrations in acetone, ethanol, and methanol bark fractions. It provides significant insights into the phytochemical composition, antioxidant potential, and DNA nicking inhibition properties of *T. arjuna* solvent extracts, which could aid in developing remedies for geriatric chronic diseases.¹²¹

1.3.5 Anti-inflammatory activity. The anti-inflammatory effect of *T. arjuna* was evaluated using the carrageenan-induced paw oedema method, revealing significant efficacy ($p < 0.001$) in acute inflammatory conditions at both 250 mg kg⁻¹ and 500 mg kg⁻¹ doses. The extract was also assessed for centrally acting analgesic effects through formalin and hot-plate tests and peripheral analgesic effects with the acetic acid-



induced writhing test in mice. At oral doses of 250 mg kg⁻¹ and 500 mg kg⁻¹, 50% ethanolic bark extract of *T. arjuna* demonstrated significant analgesic activity ($p < 0.01$; $p < 0.001$) across these models. In the hot-plate test, both dose levels showed a significant increase in latency period ($p < 0.001$) compared to the control group. Additionally, in the acetic acid-induced writhing and formalin tests, *T. arjuna* significantly reduced the number of writhes at both dose levels ($p < 0.05$; $p < 0.001$) relative to the control.¹²²

In vitro studies suggested that methanol extracts of *T. arjuna* leaves and fruits had a stronger anti-inflammatory effect than the methanol extract of the bark, potentially due to high levels of saponins, tannins, and flavonoids, which may contribute to HRBC membrane stabilization and anti-inflammatory activity.¹²³ Oral administration of *T. arjuna* fruit aqueous extract demonstrated an acute anti-inflammatory effect in carrageenan-induced paw oedema, along with analgesic effects in acetic acid-induced writhing and heat-induced pain tests, indicating both central and peripheral mechanisms by inhibiting the biosynthesis and/or release of pain mediators.¹²⁴

Studies found that bark powder of *T. arjuna* in formulations such as Arjuna Ksheera Paka (AKP) and hydroalcoholic (HA) extracts exhibited significant anti-inflammatory activity ($p < 0.05$) in reducing paw oedema in mice. HA was more effective in the early phase of inflammation, while AKP demonstrated greater efficacy in the late phase, potentially due to the presence of milk solids in AKP.¹²⁵

Pre-treatment with methanolic extracts of *T. arjuna* leaves (200 and 400 mg kg⁻¹) reduced paw oedema by 38.05% and 64.95%, respectively, and decreased granulomatous tissue weight to 115.33 mg and 73.67 mg. Haematological analysis of pre-treated animals showed decreased white blood cells (WBC) and platelet counts with increased Hb and RBC levels. Methanolic leaf extract pre-treatment also reduced nitrite and lipid peroxidation (LPO) levels while increasing antioxidant enzymes, such as superoxide dismutase (SOD), glutathione synthetase (GSH), and glutathione peroxidase (GPx). Histopathology revealed a decrease in inflammatory cell accumulation with methanolic leaf extract treatment.¹²⁶ Additionally, *T. arjuna* bark powder (400 mg kg⁻¹, orally) significantly minimized formalin-induced paw oedema at 24 hours.¹²⁷

1.3.6 Anti-diabetic activity. *T. arjuna* has been explored for its anti-diabetic roles since ancient times.¹²⁸ The aqueous stem bark extract of *T. arjuna* demonstrated anti-diabetic activity by increasing granulocyte count and reducing free haemoglobin and total cellular content in diabetic human blood and plasma samples.¹²⁹ In an *in vivo* study, the acetone extract of *T. arjuna* bark showed protective effects on blood glucose levels, oral glucose tolerance (OGTT), and insulin tolerance in streptozotocin (STZ)-treated rats. Rats fed with 500 mg kg⁻¹ body weight of the bark extract exhibited significantly ($P < 0.05$) improved parameters compared to diabetic control rats and those fed with 250 mg kg⁻¹, with no significant difference compared to glimepiride-treated rats. The 500 mg kg⁻¹ dosage showed greater anti-hyperglycemic and anti-diabetic effects ($P < 0.05$) than the 250 mg kg⁻¹ dosage.¹³⁰ Treatment with the acetone extract of the root of *T. arjuna* significantly reduced elevated

levels of total cholesterol (TC) and triglycerides (TG) compared to negative controls in diabetic animals. This study indicated that both extract and glibenclamide protected diabetic rats from substantial body weight loss when administered orally for 15 days.¹³¹

Further evaluation of *T. arjuna*'s anti-diabetic effects showed that its root extract has potent antidiabetic effects in alloxan-induced diabetic rats. A 50% ethanol extract of the stem bark, administered at 1.25 g kg⁻¹ body weight for 21 days to type 2 diabetic Long-Evans rats, significantly ($p < 0.05$) improved glucose tolerance by the end of this study compared to controls. Fasting serum glucose levels also decreased significantly ($p < 0.05$) after 21 days of treatment, though no changes were observed in liver glycogen content or serum insulin levels. Additionally, *T. arjuna* stem bark 50% ethanolic extract showed beneficial effects on lipid profiles, significantly reducing serum total cholesterol and triglyceride levels ($p < 0.01$ and $p < 0.001$, respectively). Glibenclamide (5 mg kg⁻¹) similarly produced a significant reduction ($p < 0.01$) in serum glucose in type 2 diabetic rats.¹³²

The hydroethanolic extract of *T. arjuna* bark exhibited α -amylase inhibitory activity, with an IC₅₀ of 145.90 $\mu\text{g mL}^{-1}$ compared to the IC₅₀ of acarbose 62.35 $\mu\text{g mL}^{-1}$, suggesting that α -amylase inhibition might contribute to its antidiabetic effect through the synergistic action of its phytochemicals.¹³³

T. arjuna also showed efficacy in managing diabetes with cardiovascular complications, helping to restore body weight in treated rats compared to controls. The hydroalcoholic extract of *T. arjuna* bark (HATAB) treatment significantly reduced hyperglycaemia compared to the diabetic ISP control (D-ISP) group, though Vildagliptin (VIL) showed a superior effect. Rats treated with both VIL and (HATAB) had a significantly lower heart-to-body weight ratio compared to the D-ISP group. Combined VIL and HATAB extract treatment reversed STZ/ISP-induced increases in CPK-MB and hs-CRP levels. Serum CPK-MB isoenzyme and hs-CRP levels were significantly lower in treated rats, indicating reduced cardiac damage. Moreover, VIL and treatment with HATAB extract significantly reduced serum DPP-4 levels in D-ISP rats, with a positive correlation between cardiac marker CPK-MB and serum DPP-4 confirmed by histopathology (VIL, $r = 0.899$; $p < 0.01$; bark, $r = 0.848$; $p < 0.05$).¹³⁴

T. arjuna bark aqueous extract also stimulated insulin release alone ($p < 0.001$) and in combination with other modulators, though this effect required extracellular Ca²⁺. The extract increased intracellular calcium without affecting depolarized cells and significantly enhanced glucose uptake ($P < 0.001$). At higher concentrations, it inhibited both starch digestion and protein glycation ($p < 0.001$).¹³⁵

1.3.7 Hepatoprotective activity. *T. arjuna* has been reported to exhibit hepatoprotective activity against paracetamol-induced hepatic injuries.¹³⁶⁻¹³⁹ Administration of aqueous extracts of *T. arjuna* bark before paracetamol/CCl₄ administration led to a significant reduction in sGOT, sGPT, sALP, and SB levels, almost comparable to silymarin in CCl₄-induced hepatocarcinoma. Histopathological examination of liver tissue from control and treated animals further confirmed the hepatoprotective effect.¹⁴⁰ Another study indicated that methanolic



extracts of *T. arjuna* stems and their phytochemicals also exhibit hepatoprotective activity. CCl_4 treatment caused a marked increase in serum glutamate pyruvate transaminase (GPT) and alkaline phosphatase (ALP) levels, as well as in thiobarbituric acid reactive substances (TBARS) levels, while reducing catalase (CAT), glutathione-S-transferase (GST) and superoxide dismutase (SOD) levels in liver and kidney tissue homogenates of treated mice. The aqueous extract of *T. arjuna* bark successfully prevented these adverse changes in experimental animals.¹⁴¹

1.3.8 Insect repellent activity. The repellent and anti-feedant effects of *Saraca asoca* and *T. arjuna* bark extracts were evaluated for controlling *Sitophilus oryzae*. The methanol extract of *T. arjuna* bark exhibited a stronger repellent effect against *S. oryzae* adults compared to the *Saraca asoca* bark extract. Additionally, the methanol extract of *T. arjuna* bark demonstrated greater antifeedant activity against *S. oryzae* adults than the *Saraca asoca* bark extract. These insecticidal, repellent, and antifeedant properties in *Saraca asoca* and *T. arjuna* bark extracts may be attributed to their bioactive compounds.¹⁴²

1.3.9 Wound-healing activity. The wound-healing properties of two herbal formulations (Himax ointment and lotion) containing *Indradaru* extract, specifically *T. arjuna* bark (Family: Combretaceae), were evaluated in two wound models in rats: (i) excision and (ii) incision. Both formulations showed significant activity in both wound models. Their effects were comparable to those of the standard drug nitrofurazone, particularly in terms of wound contraction, epithelialization period, tensile strength, and tissue regeneration at the wound site.¹⁴³

Meenakshi *et al.* prepared a 50% aqueous alcohol extract of *T. arjuna* bark and fractionated this extract to obtain a tannin-rich fraction. The effects of topical application of this tannin-rich fraction were also assessed on rat dermal wounds *in vivo*. The findings showed a statistically significant increase in tensile strength in incision wounds and in epithelialization percentage in excision wounds compared to the control ($p < 0.05$).¹⁴⁴

Additionally, wound-healing effects of a 50% ethanolic extract of *T. arjuna* bark and tannins isolated from the bark were investigated in incision and excision models following oral or topical application as a hydrogel. The results showed a statistically significant increase in tensile strength in incision wounds and a greater reduction in wound size in excision wounds compared to the control. Topical treatment with tannins was particularly effective in both models, with an increase in hydroxyproline content in the granulation tissue of excision wounds, indicating enhanced collagen turnover and accelerated wound healing.¹⁴⁵

1.3.10 Food and nutrition. Meena *et al.* assessed the nutritional value of *T. arjuna* through a 90-days experiment designed to optimize the use of *T. arjuna* bark powder (TABP) in the diet of *Labeo rohita*. Four iso-nitrogenous diets (305.74 g crude protein per kg diet) and iso-caloric diets (18.15 MJ gross energy per kg diet) were formulated, incorporating TABP at levels of 0.0, 5.0, 10.0, and 15.0 g kg^{-1} feed, designated as TABP0, TABP5, TABP10, and TABP15, respectively. A total of 540 juvenile fish (average weight 20.7 g, 45 fish per tank) were randomly distributed into 12 fiber-reinforced plastic (FRP)

tanks, each with a 500-Liter water capacity. Fish were fed to satiation twice daily, at 9:30 AM and 5:00 PM. The results indicated that diets containing 10 g TABP per kg feed produced the highest growth rates (weight gain percentage and specific growth rate), nutrient utilization (protein efficiency ratio, apparent net protein utilization, fat retention, and energy retention), and survival rate. In contrast, the feed conversion ratio (FCR) displayed an inverse relationship ($P < 0.05$) with growth rates. Growth rates, feed conversion, and nutrient utilization were comparable between the TABP0 and TABP15 groups ($P > 0.05$). The viscerosomatic index (VSI) and craniosomatic index (CSI) were significantly affected ($P < 0.05$), while the hepatosomatic index (HSI) showed no significant changes ($P > 0.05$). Whole-body composition was significantly influenced ($P < 0.05$), except for ash content ($P > 0.05$). Furthermore, the RNA content and the RNA : DNA ratio were positively correlated with growth rates and nutrient retention ($P < 0.05$). The study concluded that TABP, at an optimal inclusion level of 12.3 g kg^{-1} feed, can effectively produce healthy, high-quality fish for human consumption, termed “Green Fish.” This research also provides a foundation for developing TABP-based medicated feeds on a larger scale, which could be tested under field conditions alone or in combination with commercial feeds.¹⁴⁶

Meena *et al.* also assessed the impact of processing and pre-treatment on the biochemical composition, mineral profiling, and yield (%) of selected solvent extracts of *T. arjuna* (Arjuna). The analysis revealed that the maximum yield was obtained from the ethanolic bark extract in both individual ($23.6\% \pm 0.026\%$) and serial ($22.23\% \pm 0.017\%$) fractions. The yield of ethanolic and methanolic bark extracts showed no significant difference ($p \geq 0.05$), whereas non-polar solvent extracts exhibited significant differences ($p \leq 0.05$). Mineral profiling demonstrated substantial variation between dry powders and their solvent extracts. Fractionation increased the zinc (Zn) content in fruit extracts, with the methanolic extract recording the highest Zn level (45.29 mg L^{-1}). An inverse relationship between ash and moisture content was observed across all solvent extracts. The maximum ash content was recorded in the bark powder, with values of $28.95\% \pm 0.001\%$ (serial fraction) and $28.19\% \pm 0.008\%$ (individual fraction). However, ash content did not directly correlate with mineral profiling, suggesting higher acid-insoluble ash levels in bark compared to leaves and fruit. This may contribute to the bio-efficacy of the solvent extracts.

The findings indicated that the inherent medicinal properties of *T. arjuna* bark can be leveraged, and its solvent extracts could be utilized in drug formulation. The study highlights the potential to enhance mineral profiling, particularly zinc—a potent neurotransmitter—which could be used to develop suitable feeds for livestock and fisheries. Furthermore, these findings provide a pathway for addressing nervous disorders in human health through dietary supplementation or therapeutic formulations.¹⁴⁷

Pankaj *et al.* developed a process utilizing *T. arjuna* bark extract to produce herbal ghee from buffalo milk. Three types of extracts were evaluated: commercially available aqueous *arjuna* extract powder, laboratory-prepared aqueous extract, and



alcoholic extract. Among these, ghee produced with the alcoholic extract (at a 4% fat-to-extract ratio) was found to be superior in quality. To optimize the level of the alcoholic extract to maximize phytosterol retention in ghee, samples were prepared using extract concentrations of 5%, 6%, and 7% relative to the weight of fat. No significant difference in sensory acceptability scores was observed among these levels. However, phytosterol content was highest (0.38 mg g⁻¹ with cream as the fat source and 0.47 mg g⁻¹ with butter) at the 7% extract level. The optimized herbal ghee was prepared using 7% alcoholic *arjuna* extract with the creamery butter method. Its chemical composition included 99.92% fat, 0.08% moisture, 0.22% free fatty acid (as oleic acid), a butyro-refractometer reading of 41.5 at 40 °C, a Reichert-Meisssl value of 31.5, and a phytosterol content of 0.39 mg g⁻¹. This process demonstrates the potential of incorporating *arjuna* extract into ghee for enhanced phytosterol content and associated health benefits.¹⁴⁸

Suman *et al.* developed buffalo meat rolls by incorporating extracts from *T. arjuna* bark and *Aloe vera* gel at varying levels (2%, 4%, and 6%) to determine the optimal incorporation rate and evaluate their effects on the texture profile of the products. Meat rolls containing 4% *Aloe vera* gel scored slightly higher in juiciness and tenderness compared to those with 2% *T. arjuna* bark extract. However, no significant differences were observed between these two samples in overall sensory scores, which were comparable to the control. Based on sensory evaluations, 4% *Aloe vera* gel and 2% *T. arjuna* bark extract were identified as the optimal levels for incorporation and were selected for further study. The texture attributes of the developed products, such as hardness, springiness, and cohesiveness, were comparable with those of the control samples. The study concluded that buffalo meat rolls with desirable sensory and textural properties can be successfully developed by incorporating 4% *Aloe vera* gel and 2% *T. arjuna* bark extract, enhancing the product's overall quality.¹⁴⁹

Pravin *et al.* evaluated the hypolipidemic and antioxidant properties of a vanilla-chocolate dairy drink fortified with encapsulated *T. arjuna* bark powder (1.8%) in Wistar rats fed a high-cholesterol diet for 60 days. By the conclusion of the experimental period, a significant lowering in body weight gain was observed in rats administered with the encapsulated herb extract when compared to those on a high-cholesterol diet without supplementation. Additionally, microencapsulation of the herb resulted in a significant reduction in organ weights, particularly epididymal fat and liver. The supplemented group exhibited notable reductions in serum lipid levels, including total cholesterol, triglycerides, low-density lipoprotein (LDL), very-low-density lipoprotein (VLDL), and the atherogenic index. Encapsulation also enhanced antioxidant activity, evidenced by increased reduced glutathione levels and decreased thiobarbituric acid reactive substances (TBARS) in liver tissues and red blood cell lysates. The findings highlighted that the bioactive compounds in encapsulated *T. arjuna*—such as phytosterols, flavonoids, saponins, and tannins—remained stable during processing and were effectively released in the intestine. These compounds contributed to its hypolipidemic and

antioxidant effects, offering therapeutic potential for managing cardiovascular diseases.¹⁵⁰

Various groups have also formulated herbal tea with the extracts of *T. arjuna*, along with other herbal extracts. These products are commercially available and known to have several health benefits.^{151,152}

1.3.11 Immunomodulatory effects. Meena *et al.* explored the dietary immunomodulatory effects of *T. arjuna* bark powder (TABP) in *Labeo rohita*, a freshwater fish model. Fish were fed four iso-nitrogenous and iso-caloric diets containing graded levels of TABP (0, 5, 10, and 15 g kg⁻¹) for 90 days, followed by a 10-days challenge with the pathogenic bacteria *Aeromonas hydrophila* and *Edwardsiella tarda*. The study employed an Integrated Biomarker Response (IBR) approach to evaluate hematological, adaptive, and humoral immune parameters alongside liver histomorphology. Dietary inclusion of TABP at 10 g kg⁻¹ significantly improved ($p < 0.05$) hematological indices (hemoglobin levels, red blood cell count, and hematocrit), specific immune responses (lysosomal enzyme activity, phagocytosis, and respiratory burst), and non-specific immune parameters (serum lysozyme and alternative complement activity). Additionally, liver histomorphology showed structural improvements indicative of enhanced immune function. A broken-line regression analysis identified 11.5 g kg⁻¹ as the optimal dose, while 15 g kg⁻¹ resulted in a slight decline in some parameters, suggesting a dosage threshold for TABP's efficacy. These findings highlight TABP's potential as a natural immunomodulator in fish aquaculture.¹⁵³

In a related study by the same group, a nutritional fish feed incorporating TABP was administered to *Labeo rohita* at varying levels: 0% (0 g kg⁻¹, CT), 0.5% (5 g kg⁻¹, T1), 1% (10 g kg⁻¹, T2), and 1.5% (15 g kg⁻¹, T3). A comprehensive biomarker response strategy was used to assess the genomic impact of TABP inclusion. Following bacterial infection, the expression of immunogenic genes—STAT1 (signal transducer and activator of transcription 1), ISG15 (interferon-stimulated gene 15), and Mx (myxovirus resistance gene)—was elevated. Densitometry analysis revealed a dose-dependent increase in STAT1 and ISG15, with Mx reaching its highest levels at 10 g kg⁻¹ TABP.¹⁵⁴

1.3.12 Toxicology. Khurram *et al.* examined the toxicity profile of arjunolic acid (AA) in healthy female Sprague Dawley rats. In this study, AA was administered orally in single doses of 300 mg kg⁻¹ and 2000 mg kg⁻¹ body weight to groups 1 and 2, respectively, while group 3 served as the control. The animals were monitored for two weeks for any behavioural or physical changes. On the 15th day, blood samples were collected for hematological and biochemical analysis, and the animals were euthanized to harvest organs for histopathological examination. Histological alterations in tissues were assessed using four staining methods: hematoxylin and eosin, Masson trichrome, periodic acid-Schiff (PAS), and Oil Red O. The study observed no morbidity or mortality in the treated groups at either dose level. Daily food and water intake, body weight, relative organ weight, hematological, and biochemical parameters remained within normal ranges. Histopathological analysis revealed no significant structural changes in tissues. These findings highlight the



safety profile of AA, demonstrating that it is well tolerated even at high doses.¹⁵⁵

Meena *et al.* conducted an *in vitro* toxicity study to validate, standardize, and supplement data on the toxicity assessment of various solvent extracts of *T. arjuna* using *Artemia salina* as a model organism. Experimental conditions were optimized, including the use of 100 W yellow light, 5% salinity, pH 8.0–8.5, and a temperature of 30 °C, over 48 hours. Nauplii were collected using a modified dropper and incubated for 24 hours under the same conditions before being observed with a 50× magnifying glass. Functional screening of solvent extracts and their respective mother solvents revealed LC₅₀ values for hexane (118.50 ppm), ethyl acetate (101.75 ppm), chloroform (93.36 ppm), acetone (278.32 ppm), ethanol (528.78 ppm), and methanol (477.67 ppm). These values categorized the solvents as medium, medium, high, medium, low, and medium toxicity, respectively. Interestingly, the toxicity of the solvent extracts differed from their mother solvents, demonstrating the extracts' effectiveness. Universal solvents, such as DMSO and distilled water (DW), were classified as non-toxic according to Meyers' toxicity index and Clarkson's toxicity criterion. Among the solvent extracts of *T. arjuna*, all were toxic per Meyers' toxicity index but varied in toxicity classification under Clarkson's criterion, ranging from low to high toxicity. Principal Component Analysis (PCA) revealed strong correlations, with PCA1 and PCA2 explaining 69.46% and 19.74% of the variation, respectively. The study concluded that LC₅₀ values could be categorized as relative LC₅₀ (linked to the mother solvent) or absolute LC₅₀ (actual extract potential). Relationships between relative and absolute LC₅₀ values were linear, with their percentage fractions and inversely related to their counterparts.¹⁵⁶

1.3.13 Generation of nanoparticles using leaves and bark of *T. arjuna* and their biological properties. Several studies have explored the green synthesis and applications of nanoparticles using *T. arjuna* bark and leaf extracts.

Anuradha and colleagues reported the biosynthesis of copper (CuNPs) and zinc (ZnNPs) nanoparticles using aqueous *T. arjuna* bark extract as a reducing and stabilizing agent. The nanoparticles, characterized using UV-visible spectroscopy, FTIR, and SEM, were found to be spherical, with CuNPs ranging from 10 to 26 nm and ZnNPs from 15 to 25 nm. XRD analysis confirmed their structure. The antibacterial activity of the CuNPs was observed to be superior to that of ZnNPs against the tested organisms, although *Klebsiella pneumoniae* showed higher resistance. Both nanoparticles demonstrated anti-haemolytic activity, with CuNPs being more effective.¹⁵⁷

Amol and collaborators described the synthesis of gold nanoparticles (AuNPs) using *T. arjuna* leaf extract, with UV-visible spectroscopy showing a characteristic peak at 530 nm. TEM analysis revealed AuNPs with sizes of between 15 and 30 nm. The synthesized nanoparticles exhibited antibacterial activity against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Salmonella typhimurium*.¹⁵⁸

Rakhi *et al.* demonstrated the synthesis of AuNPs using bark extract under mild conditions at room temperature. Characterization through HRTEM, AFM, and XRD confirmed their structural properties. These nanoparticles were successfully

used to catalyze the reduction of 4-nitrophenol to 4-aminophenol.¹⁵⁹

Gopinath and his team synthesized spherical AuNPs (20–50 nm) using leaf extract and reported their characterization by UV-visible spectroscopy, FTIR, and TEM. These nanoparticles promoted mitotic cell division and pollen germination without cytotoxic effects.¹⁶⁰

Natarajan *et al.* utilized ethanolic bark extract to synthesize AuNPs (20–50 nm), which were found to possess neuro-protective properties. Characterization techniques, including UV-visible spectroscopy and TEM, confirmed their stability and size. The nanoparticles demonstrated antioxidant, anticholinesterase, and anti-amyloidogenic activities, highlighting their potential for treating neurodegenerative disorders such as Alzheimer's disease.¹⁶¹

Akther and colleagues developed silver nanoparticles (AgNPs) through microwave-assisted green synthesis using bark extract. These nanoparticles showed significant antibacterial and anti-biofilm activity against multidrug-resistant *Escherichia coli*, including strains harbouring the CTXM-15 gene. Docking studies revealed strong interactions of phytochemicals with CTXM-15, inhibiting its expression.¹⁶²

Koparde *et al.* synthesized AgNPs (40–50 nm) using bark extract and confirmed their formation through colorimetric and SEM analyses. These nanoparticles exhibited strong antibacterial activity, especially against *Pseudomonas aeruginosa*.¹⁶³

Yallappa *et al.* employed a microwave-assisted method to produce AgNPs (10–15 nm) using bark extract. The synthesized nanoparticles demonstrated excellent antioxidant and antibacterial properties against bacteria and yeast.¹⁶⁴

Sakheel and colleagues reported a simple “one-pot” synthesis of AgNPs using leaf extract, without additional stabilizers. FTIR analysis confirmed bio-capping by plant biomolecules. The nanoparticles exhibited antibacterial efficacy, making them suitable for biomedical applications such as wound treatment.¹⁶⁵

Chabbi and collaborators synthesized palladium nanoparticles (PdNPs) using *T. arjuna* bark extract. Characterized by UV-visible spectroscopy and TEM, these PdNPs demonstrated catalytic efficiency in C–C coupling reactions and dye degradation.¹⁶⁶

These findings collectively highlight the ability of *T. arjuna* extracts to facilitate the environmentally friendly synthesis of nanoparticles with significant applications in medicine, catalysis, and environmental remediation.

1.3.14 *T. arjuna* extracts in drug delivery. Gaikwad *et al.* investigated the potential of *T. arjuna* bark for transdermal drug delivery applications. Transdermal patches loaded with *T. arjuna* bark extract were developed using the solvent casting technique, incorporating varying amounts of chitosan and Eudragit RL 100, according to a 3² factorial design. These patches were evaluated for physicochemical properties, drug content, *in vitro* and *ex vivo* diffusion, skin irritation, and stability. Infrared spectroscopy confirmed that no chemical interactions or compositional changes occurred during patch preparation. The optimized batch (S3) exhibited drug release rates of 74.56% *in vitro* and 69.12% *ex vivo* over 12 hours. Skin irritation studies demonstrated that neither



the extract nor the excipients caused adverse effects. Stability testing revealed that the transdermal matrix formulations remained stable during storage. The study concluded that these formulations, containing different polymeric components, are suitable for transdermal delivery in treating chronic conditions such as cardiovascular disorders.¹⁶⁷

Sobia *et al.* developed and characterized a pH-triggered *in situ* gelling system using moxifloxacin HCl (MOX-HCl). The gelling system utilized gum extracted from *T. arjuna* bark resin,

blended with sodium alginate as the gelling agent. Sterilized formulations were prepared and assessed for their physico-chemical properties, microbiological activity, and eye irritation potential. The drug-loaded *in situ* gel appeared as a clear solution, which transformed into a gel in the presence of tear fluid. The optimized formulation was stable, therapeutically effective, non-irritating, and exhibited sustained drug release for up to 12 hours. In rabbit studies, the gel caused no irritation symptoms such as redness, inflammation, or excessive tearing compared




Fig. 3 Arjunolic acid, a pentacyclic triterpene acid, is a major constituent of *Terminalia arjuna*.




Fig. 4 Medicinal properties exhibited by arjunolic acid.



Table 2 Sources and methods of isolation of arjunolic acid with the corresponding yields

S. No	Source	Part	Method	Qty. of raw material used	Isolated quantity (% yield)	Ref.
1	<i>T. arjuna</i>	Heartwood	Column chromatography	—	—	169
2	<i>T. arjuna</i>	Stem bark	Column chromatography	1.5 kg	400 mg (0.04%)	170
3	<i>T. arjuna</i>	Stem bark	Column chromatography	2 kg	210 mg (0.01%)	171
4	<i>T. arjuna</i> ^a	Heartwood	Flash column chromatography	1 kg	15 g	172
5	<i>T. arjuna</i>	Bark	HPTLC	—	—	173
6	<i>T. arjuna</i>	Bark	HPLC-PDA	—	—	174
7	<i>T. arjuna</i>	Stem bark	Column chromatography	1 kg	250 mg (0.025%)	175
8	<i>T. arjuna</i>	Heartwood	Flash column chromatography	1 kg	11.5 g (1.15%)	7 and 176
9	<i>Akebiae fructus</i>	—	UHPLC-QTOF-MS	—	—	177
10	<i>Tilia kiusiana</i>	Leaves	Column chromatography	187 g	3.5 mg (0.002%)	178
11	<i>T. catappa</i>	Stem bark	MPLC	—	—	179
12	<i>Combretum griffithii</i>	Stem bark	Column chromatography	4.1 kg	127 mg (0.003%)	180
13	<i>Lagerstroemia indica</i>	Stem	Column chromatography	5 kg	2 mg (0.00004%)	181
14	<i>Shuang-Huang-Lian</i>	—	UHPLC-QTOF	—	—	182
15	<i>Wuzi-Yanzong-Wan</i>	—	UPLC-ESI-LTQ-orbitrap-MS	—	—	183
16	<i>Combretum racemosum</i>	Air dried pericarp	Column chromatography	9 kg	184 mg (0.002%)	184
17	<i>Premna odorata</i>	Leaves	LC-HRESIMS	—	—	185
18	<i>Folium eribotryae</i>	—	—	—	—	186
19	<i>Bixa orellana</i> L	Root	Extraction/Separation	—	—	187
20	<i>Quercus serrata</i> Thunb ^a	Seeds/acorn	Column chromatography	30 kg	8.6 mg (0.0003%)	188
21	<i>Quince (Cydonia vulgaris)</i>	Peels	Column chromatography	3.1 kg	7.9 mg (0.002%)	189
22	<i>Phlomis umbrosa</i>	Root	Column chromatography	15 kg	9.4 mg (0.006%)	190
23	<i>Juglans sinensis</i>	Leaves and twigs	Column chromatography	20 kg	500 mg (0.002%)	191
24	<i>Cleistocalyx operculatus</i> ^a	Leaves	Column chromatography	3.3 kg	60 mg (0.002%)	192
25	<i>Stauntonia hexaphylla</i>	Fruit	Column chromatography	—	—	193
26	<i>T. albida</i>	Root	Column chromatography	470.9 g	7.4 mg (0.002%)	194
27	<i>Poraqueiba sericea</i>	Stems	Column chromatography	856 mg	2 mg (0.23%)	195
28	<i>Combretum nelsonii</i> ^a	Leaves	Column chromatography	12.38 g	81 mg (0.65%)	196
29	<i>Myrica adenophora</i>	Root	MPLC	7.2 kg	110 mg (0.002%)	197
30	<i>Myrcia rubella</i> Cambess	Leaves	Extraction/HPLC	320 g	0.3 mg (0.0009%)	198
31	<i>T. bentzoe</i> L	Leaves	Flash column chromatography	1.5 kg	20 mg (0.0013%)	199
32	<i>Clidemia hirta</i>	Root	HPLC	—	—	200
33	<i>Psidium guajava</i> L ^a	Leaves	Column chromatography	—	—	201
34	<i>Mosla dianthera</i>	Whole herb	Column chromatography	4.4 kg	80 mg (0.0018%)	202
35	<i>Eugenia jambolana</i> Lam	Berry/fruit pulp	Prep HPLC/column chromatography	500 g	8.2 mg (0.0016%)	203
36	<i>Glochidion assamicum</i>	Aerial parts	Column chromatography	14 kg	3 mg (0.00002%)	204
37	<i>Melandrium firmum</i>	Aerial parts	Column chromatography	10 kg	13.5 mg (0.00013%)	205
38	<i>Duabanga grandiflora</i> L	Stem	Column chromatography	5 kg	162 mg (0.003%)	206
39	<i>Gentiana scarba</i>	Root and rhizomes	Column chromatography	2.5 kg	45.9 mg (0.0018%)	207
40	<i>Cyclocarya paliurus</i>	Leaves	HPLC-QTOF-MS	—	—	208
41	<i>Lagerstroemia. guilinensis</i>	Branches	Column chromatography	—	—	209
42	<i>Melaleuca alternifolia</i>	Bark and stem	Column chromatography	700 g	3.9 mg (0.0005%)	210
43	<i>Combretum leprosum</i>	Root	Column chromatography	—	—	211
44	<i>Vaccinium emarginatum</i>	Whole plant	Column chromatography	17.4 kg	36.4 mg (0.0002%)	212
45	<i>Juglans mandshurica</i> ^a	Green walnut husk	Column chromatography	10 kg	5.2 mg (0.0005%)	213
46	<i>Akebia trifoliata</i>	Aerial parts	Column chromatography	—	—	214
47	<i>Siphoneugena densiflora</i>	Stem	Column chromatography	60.1 g	12 mg (0.019%)	215
48	<i>Eucalyptus globulus</i> ^a	Stem bark	GC-MS	—	—	216
49	<i>Osmanthus fragrans</i> ^a	Fruits	UHPLC-TQUAD-MS	—	—	217
50	<i>Miconia trailii</i>	Twig/leaves	Column chromatography	370 g	5 mg (0.0013%)	218
51	<i>Psidium guajava</i> ^a	Leaves	Column chromatography	20 kg	3 mg (0.00001%)	219
52	<i>Combretum nelsonii</i> ^a	Leaves	Column chromatography	500 g	81 mg (0.016%)	220
53	<i>Rubus sanctus and Rubus ibericus</i>	Leaves	UHPLC-ESI/HRMS	—	—	221
54	<i>Sapium haematospermum</i>	Aerial parts	HPLC	850 g	3 mg (0.0003%)	222
55	<i>Carissa opaca</i>	Whole plant	Column chromatography	10 kg	17 mg (0.00017%)	223
56	<i>Cola rostara</i> K. Schum	Root	Column chromatography	2.5 kg	10 mg (0.004%)	224



Table 2 (Contd.)

S. No	Source	Part	Method	Qty. of raw material used	Isolated quantity (% yield)	Ref.
57	<i>Heteropyxis canescens</i>	Stems and root	Column chromatography	1.5 kg	17 mg (0.001%)	225
58	<i>Durio zibethinus</i>	Fruit shell	Column chromatography	20 kg	5 mg (0.00002%)	226
59	Hawthorn berries	Berries	Column chromatography	8 kg	4.7 mg (0.0006%)	227
60	<i>Potentilla chinensis</i>	Whole plant	CC/MPLC	100 g	5.2 mg (0.05%)	228
61	<i>Terminalia mollis</i> ^a	Stem bark	Column chromatography	—	—	229
62	<i>Myrtus communis</i> ^a	Leaves	HPLC/HRMS	—	—	230
63	<i>Solanum melongena</i> L	Root	Column chromatography	12 kg	6.5 mg (0.005%)	231
64	<i>Eucalyptus globulus</i> Wood ^a	Wood	GC-MS	—	—	232
65	<i>Cochlospermum tinctorium</i>	Rhizomes	Column chromatography	75 g	75 mg (0.1%)	233
66	<i>Symplocos lancifolia</i>	Leaves	HPLC/Column chromatography/prep HPLC	600 g	17 mg (0.002%)	234
67	<i>Syzygium samarangense</i>	Aerial parts	Column chromatography	—	—	235
68	<i>Myrica arborea</i>	Twigs	Column chromatography	1 kg	70 mg (0.007%)	236
69	<i>Cecropia schreberiana</i>	Leaves	Column chromatography	590 g	2 mg (0.003%)	237
70	<i>Premna fulva</i>	Stalks	Column chromatography	—	—	238
71	<i>Rhododendron hainanense</i>	Aerial parts	Column chromatography	1 kg	2 mg (0.0002)	239
72	<i>Campsis grandiflora</i> K. Schum	Flower	Column chromatography	1 kg	33 mg (0.003)	240
73	<i>Isodon yuennanensis</i>	Rhizomes	Column chromatography	—	—	241
74	<i>Momordica cochinchinensis</i>	Seeds	Column chromatography	50 kg	86 mg (0.001%)	242
75	<i>Rhodomyrtus tomentosa</i>	Roots	Column chromatography	—	—	243
76	<i>Averrhoa carambola</i>	Fruit	Column chromatography	6.8 kg	13.3 mg (0.001%)	244
77	<i>Rhododendron colletianum</i>	Aerial parts	Column chromatography	20 kg	15 mg (0.00007%)	245
78	<i>Datisca cannabina</i> linn	Whole plant	Column chromatography	15 kg	8.4 mg (0.00005%)	246
79	<i>Cornus capita</i>	Leaves	Column chromatography	—	—	247

^a Mixture of arjunolic acid and asiatic acid.

to controls. The study concluded that MOX-HCl-loaded *in situ* gel could serve as an alternative to conventional eye drops, offering extended precorneal retention, enhanced corneal permeability, and improved ocular bioavailability.¹⁶⁸

In the following section, the various isolation methods of arjunolic acid, its biological activities, and its mechanism of action are presented and thoroughly discussed.

2. Arjunolic acid

Arjunolic acid (AA) (Fig. 3) is a pentacyclic triterpene acid, a major constituent of *T. arjuna* with the IUPAC name 2 α ,3 β ,23 α -trihydroxy-olean-12-en-28-oic acid. It is reported to exhibit a plethora of biological activities (Fig. 4). It was first isolated by King *et al.* from the heartwood of *T. arjuna*.^{5,6,169}

Although arjunolic acid is readily available in *T. arjuna*, it has also been isolated from various other plant sources (Table 2). Along with arjunolic acid, other known reported minor constituents (Fig. 2 and Table 1) of *T. arjuna* include arjunic acid, arjuntenin, baicalein, and quercetin (Fig. 2). Due to the diverse biological profile exhibited by arjunolic acid, many groups have focused their research efforts on this compound.^{6,17}

2.1 Isolation of arjunolic acid from *T. arjuna*

The limited number of literature reports on the isolation of arjunolic acid from *T. arjuna* encouraged us to explore novel isolation processes that could furnish arjunolic acid from *T. arjuna* in higher yields.¹⁷⁶ In general, the isolation process

adopted by various groups involved the extraction of the plant material with various organic solvents (ethyl acetate, acetone, *etc.*), followed by either chromatographic purification or a recrystallisation process to furnish the pure product.^{25,169,170,248,249} However, it was seen that the aforementioned isolation processes were low-yielding (milligram quantities). In another study, Bag *et al.*²⁵⁰ reported the extraction and separation of arjunolic acid and asiatic acid from the heartwood of *T. arjuna* by employing cumbersome chemical methods involving several synthetic transformations followed by crystallisation. However, Manohar *et al.*¹⁷⁶ were able to efficiently separate arjunolic acid and asiatic acid by a simple flash chromatography method, affording the former in high purity and in large quantities (gram quantities, Table 2, Entry 8). Thus, an efficient process for the isolation of arjunolic acid from the crude plant material was developed in-house by our co-worker Manohar *et al.*, affording arjunolic acid in high yields when compared with the isolated yields reported to date.¹⁷⁶

3. Biological activities of arjunolic acid

3.1 Cardioprotective activity

Arjunolic acid (AA) has been extensively studied for its cardioprotective effects across various *in vitro* and *in vivo* models. Research by Ghosh *et al.* suggests that doxorubicin-induced cardiac damage results primarily from two mechanisms: first, doxorubicin exposure generates H₂O₂ in myocytes, which



regulates the p53 and p38-JNK signaling pathways in cardiomyocytes; second, activation of p38 and JNK promotes Bax translocation to the mitochondria, disrupting the balance between proapoptotic and antiapoptotic factors. These imbalances trigger mitochondrial permeabilization, leading to cytochrome c release, caspase activation, and cell death. Arjunolic acid supplementation (25 mg kg⁻¹ body weight every two days for three treatments) offers protection by detoxifying reactive oxygen species and inhibiting p38 and JNK MAPKs phosphorylation, preventing Bax translocation to the mitochondria and mitochondrial permeabilization.²⁵¹

Sumitra *et al.* further demonstrated the cardioprotective effects of arjunolic acid, showing that it reduces platelet aggregation, coagulation, and myocardial necrosis. In isoproterenol-induced myocardial necrosis, arjunolic acid restored elevated serum enzyme levels (creatinine kinase and lactate dehydrogenase) and normalized electrocardiographic parameters, preserving heart architecture pre- and post-isoproterenol exposure.¹⁷¹

Trisha *et al.* explored the mechanisms of arjunolic acid and identified its intracellular target in ameliorating hemodynamic load-induced cardiac fibrosis. Results show that AA binds and stabilizes the ligand-binding domain of peroxisome proliferator-activated receptor gamma (PPAR γ), increasing its expression during cardiac hypertrophy. In hypertrophic samples, PPAR γ knockdown reveals that AA's cardioprotective effects are primarily due to PPAR γ agonism. This upregulation of PPAR γ inhibits TGF- β signaling by suppressing TAK1 phosphorylation. AA-bound PPAR γ interacts with TAK1, masking its kinase domain and decreasing TAK1 phosphorylation, which reduces p38 MAPK and NF- κ Bp65 activation, ultimately reducing excess collagen synthesis in cardiac hypertrophy.²⁵²

Prasenjit *et al.* studied the cardioprotective property of arjunolic acid in arsenic-induced oxidative cardiac damage. NaAsO₂ exposure (10 mg kg⁻¹ body weight) increased arsenic levels in cardiac tissue and reduced antioxidant enzyme activities (superoxide dismutase, catalase, glutathione-S-transferase, glutathione reductase, and glutathione peroxidase). Treatment with arjunolic acid (20 mg kg⁻¹ body weight for four days before NaAsO₂ exposure) protected cardiac tissues by maintaining antioxidant enzyme activities and normalizing oxidative stress markers. Additionally, arjunolic acid prevented arsenic-induced hyperlipidemia and preserved cardiac architecture, as shown in histological studies.²⁵³

Hassan *et al.* investigated AA's effects on TLR4 downstream signaling in response to lipopolysaccharide (LPS) treatment. LPS exposure upregulated TLR4, MyD88, MAPK, JNK, and NF- κ B markers (2–6 times higher than the control), whereas AA treatment downregulated MyD88, NF- κ B, p38, and JNK in H9C2 myotubes. AA also decreased TLR4 expression in both H9C2 and C2C12 myotubes, suggesting potential cardioprotective effects through modulation of TLR4 expression.²⁵⁴

Gayyar *et al.* highlighted AA's protective effects on sodium nitrite-induced cardiac toxicity. Sodium nitrite increased pro-inflammatory cytokines (TNF- α , IL-1 β), C-reactive protein, and caspase activity while reducing anti-inflammatory cytokines (IL-4, IL-10) and cytochrome C oxidase levels. AA treatment reduced

pro-inflammatory cytokines and apoptosis markers while increasing anti-inflammatory cytokines and antioxidant enzymes. AA also improved sodium nitrite-induced histopathological changes in cardiomyocytes, suggesting protection against both intrinsic and extrinsic cell death pathways.²⁵⁵

Hany *et al.* reported that AA attenuated LPS-induced cardiotoxicity. Pretreatment with AA reduced serum cardiac troponin I, lactate dehydrogenase, and creatine kinase levels in Swiss albino mice. AA pretreatment also increased antioxidant levels, reduced lipid peroxidation, decreased C-reactive protein, and normalized pro- and anti-inflammatory cytokine balance. These findings collectively indicate that AA offers cardioprotection through antioxidant activity, inflammation inhibition, and prevention of cardiac cell death.²⁵⁶

3.2 Anti-cancer activity

Arjunolic acid has demonstrated anticancer properties in various *in vitro* and *in vivo* studies. Ramesh *et al.* studied anticancer properties of arjunolic acid. Ehrlich ascites carcinoma (EAC) and Dalton's lymphoma (DAL) cell lines were incubated with different concentrations of arjunolic acid, and the results showed that a 100 μ g concentration induced approximately 66% cell death in DAL cells and 70% in EAC cells, primarily through membrane disruption.²⁵⁷

Arjunolic acid (AA) was also seen to exhibit significant anticancer activities against MCF-7 and HeLa cell lines, with minimal toxicity observed in normal lymphocytes. Studies have highlighted the self-assembly properties of AA, which contribute to its anticancer effects. Fluorescence-based analyses, such as ROS generation and EtBr/AO and DAPI staining, showed that AA at 50 μ g mL⁻¹ disrupts redox balance in cancer cells by generating reactive oxygen species. This ROS-induced apoptosis is mediated by regulatory proteins TNF- α and NF- κ B, with elevated levels of pro-inflammatory cytokines indicating the involvement of a ROS-driven apoptotic pathway.²⁵⁸

In another study reported by Elsherbiny *et al.*,⁸⁹ AA showed antitumor effects by enhancing cytotoxicity and apoptosis, partly by blocking TGF- β R1 and modulating inflammatory cytokine levels. In this study, Ehrlich Ascites Carcinoma (EAC) was induced in fifty female Swiss albino mice, and AA was administered at doses of 100 mg kg⁻¹ and 250 mg kg⁻¹. AA significantly reduced tumor volume and cell count, decreased cell viability, and increased cytotoxicity in EAC cells. Additionally, AA lowered the levels of TNF- α , IL-1 β , TGF- β , TGF- β type I receptor, and latency-associated peptide, while increasing the IL-10 levels in both *in vitro* and *in vivo* conditions.²⁵⁹

3.3 Anti-diabetic activity

Arjunolic acid (AA) has been widely studied for its antidiabetic effects across several models. Prasenjit *et al.* investigated AA's prophylactic effects against streptozotocin (STZ)-induced diabetes in the pancreatic tissue of Swiss albino rats. In this study, it was observed that STZ administration (65 mg kg⁻¹ body weight *via* tail vein) increased reactive oxygen and nitrogen species (ROS and RNS) in pancreatic tissue. This oxidative stress reduced intracellular antioxidant defense, heightened lipid



peroxidation, protein carbonylation, serum glucose, and TNF- α levels. The signaling pathway analysis showed that STZ activated phospho-ERK1/2, phospho-p38, and NF- κ B, and disrupted mitochondrial transmembrane potential, causing cytochrome c release and caspase-3 activation without significantly affecting total ERK1/2 and p38 levels. AA treatment (20 mg kg⁻¹ body weight, orally), both before and after STZ administration, effectively mitigated these adverse effects by reducing excessive ROS and RNS formation and down-regulating phospho-ERK1/2, phospho-p38, NF- κ B, and mitochondria-dependent apoptosis pathways.²⁶⁰

AA's protective effects on hyperglycaemia-induced liver dysfunction were evaluated in experimental rats with type 1 hyperglycaemia induced by STZ. AA, administered orally at 20 mg kg⁻¹ before and after diabetes induction, was compared to an insulin-treated positive control. Hyperglycaemia led to body weight loss, reduced serum insulin, increased HbA1C, and elevated advanced glycation end products (AGEs). Markers of liver dysfunction included raised serum alanine transaminase (ALT), alkaline phosphatase (ALP), ROS, RNS, lipid peroxidation, and oxidative DNA damage (8-OHdG/2-dG ratio) along with decreased GSH content and antioxidant defenses. Hyperglycaemia also triggered iNOS, I κ B α /NF- κ B, and MAPK pathways and mitochondrial signals, which promoted apoptotic cell death. Increased PARP expression, NAD and ATP depletion, and DNA fragmentation in liver tissue confirmed apoptosis. Immunofluorescence with anti-caspase-3 and anti-Apaf-1, DAPI/PI staining, DNA laddering, and FACS analysis corroborated apoptotic death. Histology supported these findings, and AA treatment reduced diabetic liver complications, apoptosis, ROS, RNS, HbA1C, AGEs, and PARP-mediated DNA damage, highlighting AA's potential in diabetic care.²⁶¹

Gonçalves-Neto *et al.* examined whether AA could offset dexamethasone (DEX)-induced glucose imbalance in adult Wistar rats. Groups included vehicle-treated (Ctrl), DEX-treated (1 mg kg⁻¹ body weight intraperitoneally for 5 days) (DEX), AA-treated (30 mg kg⁻¹ by oral gavage twice daily) (AA s), AA plus DEX (AA DEX), and AA following DEX (DEX AA). AA treatment significantly improved glucose intolerance when administered before and during DEX (AA DEX) but did not ameliorate it when given after (DEX AA). AA failed to reduce the DEX-induced rise in hepatic glycogen and triacylglycerol, and prolonged AA with DEX increased hepatic steatosis. Liver function and oxidative stress markers were not notably altered among groups, suggesting that AA could partly prevent DEX-induced glucose intolerance but was less effective after DEX initiation.²⁶²

The antidiabetic potential of AA isolated from *T. arjuna* was tested in type 2 diabetic Sprague Dawley rats, with T2DM induced *via* a single STZ-nicotinamide injection. After 10 days, fasting and random blood glucose, body weight, food and water intake, serum C-peptide, insulin, and glycated haemoglobin (HbA1c) levels confirmed T2DM development. Over four weeks, AA (25 and 50 mg kg⁻¹ day⁻¹) significantly normalized fasting and random glucose, restored body weight, controlled excessive eating and drinking, and improved glucose tolerance. AA also reduced HbA1c, total cholesterol (TC), triglycerides (TG), and LDL levels, while raising HDL slightly. Serum and pancreatic

insulin and C-peptide levels increased with AA, while GDF-15 levels were stable, and AA lowered serum and pancreatic pro-inflammatory cytokines, including TNF- α , IL-1 β , and IL-6, in a dose-dependent manner.¹⁷⁰

In another study, AA isolated from *T. arjuna* selectively inhibited glucagon secretion in palmitate-induced α TC1-6 cells and isolated islets, improving hyperglycaemia and pancreatic islet morphology in STZ-induced type 2 diabetic mice. AA's selective inhibition of α -cell glucagon secretion involved ephrin-A1 and EphA4 interaction, which activated the PI3K and Akt pathways, suggesting AA as a promising therapeutic agent for T2DM.²⁶³

In an STZ-induced diabetic model, AA dosed at 10 and 30 mg kg⁻¹ orally over ten weeks was examined for its effect on retinal damage from diabetic retinopathy (DR). H₂O₂-induced ARPE-19 cells were used to assess anti-apoptotic and antioxidant effects. The results showed that AA reduced STZ-induced weight loss, increased retinal thickness and nuclei, and upregulated HO-1 protein levels *in vitro* and *in vivo*. AA protected retinal cells from apoptosis *via* the AMPK-mTOR-regulated autophagy pathway, with dorsomorphin dihydrochloride blocking AA's anti-apoptotic effects, indicating AA's potential as a treatment for diabetic retinal oxidative stress and inflammation.²⁶⁴

AA's effects on insulin signaling and TLR-4-Wnt crosstalk in the pancreas of type 2 diabetic rats were investigated by Khurram *et al.* using histology, immunohistochemistry, and RT-PCR. AA treatment reversed T2DM-induced pancreatic apoptosis and necrosis, reduced TLR-4, MyD88, NF- κ B, p-JNK, and Wnt/ β -catenin levels by inhibiting TLR-4/MyD88 and Wnt signaling, while IRS-1, PI3K, and pAkt were upregulated through NF- κ B and β -catenin crosstalk alteration. These findings indicate AA's potential as a therapeutic agent for T2DM-associated meta-inflammation.²⁶⁵

3.4 Antioxidant activity

Arjunolic acid is well-documented for its antioxidant activity. It plays an important role in maintaining levels of key antioxidant enzymes, such as catalase (CAT), glutathione-S-transferase (GST), glutathione reductase (GR), glutathione peroxidase (GPx), and superoxide dismutase (SOD), along with cellular metabolites such as reduced glutathione (GSH) and oxidized glutathione (GSSG), thereby reducing oxidative stress.

Ramesh *et al.* evaluated the antioxidant properties of arjunolic acid (AA) using four different *in vitro* methods: beta-carotene bleaching assay, DPPH radical scavenging assay, reducing power assay, and hydroxyl radical scavenging assay. The findings were compared with those of standard antioxidants, including ascorbic acid, butylated hydroxytoluene (BHT), and catechin. All four assays confirmed the antioxidant potential of AA, though it showed slightly lower efficacy than the standards.²⁶⁶

Maity *et al.* assessed the protective effect of arjunolic acid against arsenic-induced oxidative stress and reproductive dysfunction in female Wistar rats. Sodium arsenite (10 mg kg⁻¹) was administered along with arjunolic acid (10 mg kg⁻¹) for two estrous cycles. Electrozymographic analysis indicated that arjunolic acid co-treatment reduced arsenic-induced ROS production in uterine tissue by enhancing endogenous antioxidant enzyme



activities. Arjunolic acid also helped protect against DNA damage, necrosis, and ovarian and uterine tissue damage in arsenic-treated rats by promoting ovarian steroidogenesis. These protective mechanisms may be related to an enhanced antioxidant defense system, partly facilitated by the elimination of arsenic *via* the *S*-adenosyl methionine pathway, where levels of vitamin B12, folic acid, and homocysteine play crucial roles.²⁶⁷

Sinha *et al.* investigated the preventive effects of arjunolic acid on arsenic-induced oxidative damage in the murine brain. Sodium arsenite was used as the arsenic source. The *in vitro* free-radical scavenging activity and *in vivo* antioxidant power of arjunolic acid were assessed using the 2,2-diphenyl-1-picryl hydrazyl (DPPH) radical scavenging and ferric reducing/antioxidant assays, respectively. Oral administration of sodium arsenite (10 mg kg⁻¹) for two days significantly reduced the activity of antioxidant enzymes, including CAT, GST, GR, SOD, and GPx, as well as cellular metabolites, such as reduced glutathione and total thiols, while increasing oxidized glutathione levels. Lipid peroxidation products and protein carbonyl content were also elevated. Treatment with arjunolic acid (20 mg kg⁻¹) for four days before arsenic exposure nearly normalized these levels. Histological findings supported these biochemical results, showing that arjunolic acid mitigates arsenic-induced oxidative damage in the brain. The effects of vitamin C were included as a positive control. Overall, the results suggest that arjunolic acid can protect against arsenic-induced oxidative damage in the brain, likely due to its antioxidant properties.²⁶⁸

Arsenic-induced tissue damage is a major concern, particularly because oxidative stress and impaired antioxidant defense are primary causes of arsenic toxicity, which can lead to reproductive failure. Prasenjit *et al.* examined the protective role of arjunolic acid, isolated from *T. arjuna* bark, against arsenic-induced testicular damage in mice. Arsenic administration (10 mg kg⁻¹) over two days significantly reduced antioxidant enzyme activity and cellular metabolite levels, while increasing testicular arsenic content, lipid peroxidation, protein carbonylation, and glutathione disulfide (GSSG) levels. Arsenic exposure also caused degeneration in the seminiferous tubules, leading to necrosis and spermatocyte detachment. Pretreatment with arjunolic acid (20 mg kg⁻¹) for four days helped prevent oxidative stress and structural damage in the testes. Arjunolic acid demonstrated both *in vitro* free radical scavenging activity and *in vivo* antioxidant efficacy. Overall, the study suggests that arjunolic acid's antioxidant properties help prevent arsenic-induced testicular toxicity.²⁶⁹

In another study, rats pretreated with arjunolic acid showed significant reductions in neurological deficits and infarct size. Arjunolic acid mitigated ischemia/reperfusion (I/R)-induced neuronal damage by regulating levels of malondialdehyde (MDA), reduced glutathione (GSH), nitric oxide (NO), protein carbonyl content, and mitochondria-generated reactive oxygen species (ROS). It also maintained the activities of Na⁺-K⁺ ATPase, SOD, CAT, GPx, and GR enzymes, effectively preventing I/R-induced cerebral oxidative damage due to its antioxidant potential.²⁷⁰

Sumitra *et al.* examined the modulation of neutrophil function and ROS generation by arjunolic acid *in vitro*. Neutrophils were

isolated from normal and myocardial infarction (MI) mice to evaluate AA's efficacy in reducing oxidative stress. Stimulation of neutrophils with phorbol-12-myristate-13-acetate (PMA) resulted in an oxidative burst of superoxide anions (O₂⁻) and increased release of lysosomal enzymes. Treatment with arjunolic acid significantly inhibited phosphorylation of p47phox and ERK in both stimulated control and MI neutrophils, reduced oxidative phosphorylation activity, and lowered intracellular oxidative stress. This study highlights potential targets for arjunolic acid in ROS regulation through the MAPK pathway in neutrophils.²⁷¹

Prasenjit *et al.* investigated the benefits of arjunolic acid in modulating cytokine levels and oxidative stress signaling pathways in the spleen under hyperglycaemic conditions. Diabetes was induced by streptozotocin (STZ) injection (70 mg kg⁻¹), which increased IL-2 and IFN-γ levels and decreased TNF-α in diabetic animals. Hyperglycaemia was associated with elevated production of intracellular reactive intermediates, lipid peroxidation, and protein carbonylation, leading to reduced antioxidant defense. Increased expression of phosphorylated p65, its inhibitor phospho IκBα, and phosphorylated MAPKs was also observed in diabetic spleen tissue. Studies on isolated splenocytes showed mitochondrial membrane potential disruption, cytochrome c release, and caspase 3 activation, leading to apoptotic cell death. Histological examination indicated a depletion of white pulp, consistent with reduced immune response.²⁷²

3.5 Hepatoprotective activity

The hepatoprotective properties of arjunolic acid have been documented by multiple studies.⁶ Prasenjit *et al.* assessed arjunolic acid for its protective effects against arsenic-induced oxidative stress in mouse liver tissue. Administering sodium arsenite (10 mg kg⁻¹ body weight for two days) notably decreased the activity of antioxidant enzymes—superoxide dismutase, catalase, glutathione-*S*-transferase, glutathione reductase, and glutathione peroxidase—and depleted reduced glutathione and total thiols. It also elevated the levels of serum markers (alanine transaminase and alkaline phosphatase), DNA fragmentation, protein carbonylation, lipid peroxidation products, and oxidized glutathione. Arjunolic acid, known for its *in vitro* radical-scavenging and *in vivo* antioxidant properties, demonstrated protective effects when administered at 20 mg kg⁻¹ for four days before arsenic exposure. This pretreatment normalized antioxidant enzyme activities and minimized tissue damage. Histology confirmed reduced liver necrosis with arjunolic acid compared to arsenic alone. Vitamin C served as a positive control, reinforcing arjunolic acid's role in reducing arsenic-induced liver oxidative stress.²⁴⁹

Additionally, this study investigated arjunolic acid's potential to protect against acetaminophen (APAP)-induced acute liver toxicity. Exposure to a hepatotoxic APAP dose (700 mg kg⁻¹) increased oxidative stress biomarkers, reactive oxygen species (ROS), and cell death. Pretreatment with arjunolic acid (80 mg kg⁻¹) provided significant liver protection, preventing glutathione depletion and reducing APAP metabolite formation, without affecting hepatic glutathione levels. The findings suggest that arjunolic acid inhibits APAP bioactivation by



cytochrome P450 and reduces APAP-induced JNK and Bcl-2 phosphorylation, thus preventing mitochondrial damage.¹⁷⁵

In another model, arjunolic acid was tested for its protective effects against arsenic trioxide (ATO)-induced oxidative stress in mouse liver and kidney tissues. Administering ATO (30 mg kg⁻¹ day⁻¹ for eight weeks) elevated serum markers, ROS, and disrupted the prooxidant-antioxidant balance, triggering apoptotic pathways. Post-treatment with arjunolic acid (20 mg kg⁻¹ for four days) mitigated oxidative stress and apoptosis markers, including mitochondrial membrane potential loss, caspase activation, and cytochrome c release.²⁷³

In a study on cisplatin-induced liver toxicity, arjunolic acid demonstrated hepatoprotective effects by reducing oxidative stress, inflammation, and apoptosis. Rats pretreated with arjunolic acid (20 mg kg⁻¹ for 10 days) showed improved liver function, histology, and suppressed malondialdehyde and nitric oxide levels. Arjunolic acid significantly enhanced

antioxidant enzyme activities, reduced tumor necrosis factor α , and decreased caspase-3 expression.²⁷⁴

Arjunolic acid also reduced lipid accumulation in free fatty acid (FFA)-challenged hepatocytes without toxicity. *In vivo*, arjunolic acid reduced adiposity, hepatic inflammation, and lipid disorders in the non-alcoholic fatty liver disease (NAFLD) model, activating Sirt1/AMPK pathways, enhancing autophagy, and restoring gut barrier integrity.²⁷⁵

Finally, in NAFLD models, arjunolic acid showed promising results by reducing triglyceride accumulation, ALT, and AST levels. *In vivo*, it lowered liver enzyme markers and modified PPAR and FXR expression, indicating reduced liver steatosis and inflammation. Collectively, these studies suggest arjunolic acid as a promising agent for liver protection across various hepatic damage models due to its antioxidant, anti-inflammatory, and anti-apoptotic effects.¹⁷²




Fig. 5 Arjunolic acid analogues synthesized by Bag *et al.*^{285,286,291}



3.6 Anti-inflammatory activity

Arjunolic acid has been widely studied for its anti-inflammatory properties. The anti-arthritis effects of arjunolic acid were evaluated in a rat model of arthritis induced by complete Freund's adjuvant (CFA). Arthritis was induced in male Sprague Dawley rats through an intradermal injection of 0.1 mL CFA into the right footpad. Following arthritis induction, arjunolic acid was administered orally at doses of 40 and 80 mg kg⁻¹ once daily for 25 consecutive days. Indomethacin, a reference drug, was given at 3 mg kg⁻¹ twice weekly for the same duration. Standard procedures were used to assess paw swelling, serum haematology, antioxidant enzymes, inflammatory mediators, and histopathology. The results showed that arjunolic acid significantly ($p < 0.01$) reduced paw swelling and weight loss in arthritic rats. It also lowered malondialdehyde (MDA) levels, spleen index, and thymus index, while enhancing the activities of superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx). Furthermore, arjunolic acid downregulated TNF- α , IL-1 β , and IL-6 serum levels ($p < 0.01$), improving haematological and histopathological changes in CFA-induced arthritis due to its anti-inflammatory effects.²⁷⁶

Atopic dermatitis (AD), a chronic, pruritic, eczematous skin disorder associated with allergic inflammation, involves oxidative stress and immune imbalance. The anti-pruritic effects of oral arjunolic acid (AA) were studied in a mouse model of AD induced by 2,4-dinitrochlorobenzene (DNCB). Mice were treated with DNCB on their ears and shaved dorsal skin to induce AD-like symptoms, followed by oral administration of AA at 250 mg kg⁻¹ for three weeks. The results indicated that AA significantly improved AD symptoms, reducing skin lesions, dermatitis scores, ear thickness, and scratching behaviour. AA also decreased reactive oxygen species and modulated inflammatory markers, lowering serum

TNF- α , IL-6, IgE, and caspase-3 while maintaining IL-4 and IL-10 levels, thus mitigating DNCB-induced AD symptoms.²⁷⁷

Zining *et al.* investigated arjunolic acid's effect on a Crohn's disease (CD) model. CD was simulated in IL-10 knockout mice, with arjunolic acid's impact assessed through weight monitoring, histopathology, and intestinal barrier function. Arjunolic acid improved colitis symptoms, reducing weight loss, inflammation scores, and enhancing intestinal permeability. AA also inhibited apoptosis in intestinal epithelial cells by reducing Bax and C-caspase-3 levels. Mechanistic analysis revealed that AA suppressed TLR4 signaling and altered gut microbiome composition, increasing short-chain fatty acid-producing bacteria. Fecal transplantation from AA-treated mice alleviated CD-like colitis in recipient mice, highlighting AA's potential to inhibit key inflammatory pathways and promote gut microbiota balance in CD therapy.²⁷⁸

Edozie *et al.* evaluated the protective effects of arjunolic acid on fluoxetine (FXT)-induced testicular dysfunction in male rats. Thirty-six rats were assigned to six groups and treated with combinations of saline, arjunolic acid (1.0 or 2.0 mg kg⁻¹), fluoxetine (10 mg kg⁻¹), or fluoxetine plus arjunolic acid. The results showed that FXT treatment disrupted testicular steroidogenic enzymes (3 β -HSD and 17 β -HSD), ATPase enzymes, and testicular structure. FXT also increased oxidative inflammation markers (MDA, MPO, TNF- α , IL-1 β , caspase-3, and p53) and promoted apoptosis. Arjunolic acid treatment counteracted these effects, restoring steroidogenic enzyme activity, improving testicular structure, and increasing antioxidant levels (SOD, CAT, GSH) while reducing oxidative stress and inflammation.²⁷⁹

Ying *et al.* investigated the effects of arjunolic acid on depression and its potential mechanisms. *In vivo* and *in vitro* models were established by administering intraperitoneal injections of lipopolysaccharide (LPS) to mice and stimulating BV2 microglia with LPS. Behavioral tests, H&E staining, and





Fig. 6 Aromatic and aliphatic acetals/ketals of arjunolic acid synthesized by Bag *et al.*²⁹²



ELISA were conducted to assess arjunolic acid's effect on depression, while RT-qPCR, immunofluorescence, molecular docking, and Western blot analyses were performed to explore the molecular mechanisms. Arjunolic acid significantly improved depressive behaviors in LPS-treated mice, elevated BDNF and 5-HT levels in the hippocampus, decreased iNOS + IBA1 + cells, and increased Arg1 + IBA1 + positive cells in the brain. Furthermore, arjunolic acid facilitated the polarization of

BV2 microglia from the M1 to M2 phenotype. Notably, DARTS, CETSA, and molecular docking techniques identified SIRT1 as a target of arjunolic acid. Inhibition of SIRT1 with EX-527 blocked arjunolic acid's effects on reducing LPS-induced depressive behavior and promoted M2 microglia polarization. Additionally, arjunolic acid activated AMPK and reduced Notch1 expression; however, inhibiting AMPK weakened its effect on Notch1 downregulation. This study revealed that




Fig. 7 Arjunolic acid analogues synthesized by various groups exhibiting anti-oxidant, anti-cancer, anti-viral, and α -glucosidase inhibitory activities.^{9,233,288,289}






Fig. 8 Arjunolic acid derivatives synthesized by Manohar *et al.*, exhibiting anti-cancer activities.⁷

arjunolic acid mitigates neuroinflammation by modulating the SIRT1/AMPK/Notch1 signaling pathway, suggesting its potential as an antidepressant agent.²⁸⁰

3.7 Anti-asthmatic activity

Arjunolic acid has been reported to exhibit anti-asthmatic properties. In a study with rats, treatment with arjunolic acid

at doses of 50 and 100 mg kg⁻¹ body weight provided significant protection against mast cell disruption, showing 42% and 33% degranulation, respectively, compared to the standard degranulation level induced by disodium chromoglycate. Additionally, arjunolic acid at these doses offered 49% and 64% protection against histamine challenges and 40% and 51% protection against acetylcholine challenges. These anti-asthmatic effects of



Table 3 Structural modifications leading to various biological activities

Therapeutic action	Rings of arjunolic acid	Modifications
Anti-cancer	A	Selective conversion of hydroxyls (R-OH) attached at C-2, C-3 and C-23 to acetals and acetates
	A	Conversion of the A ring to lactol
	A	Simultaneous conversion of the A ring to a five-membered ring containing an α,β -unsaturated carbonyl moiety, and conversion of the acid (R-COOH) at C-28 to alkyl esters or alkyl amides
Anti-diabetic	D/E	Selective conversion of acid (R-COOH) at C-28 to amides
	D/E	Selective conversion of acid (R-COOH) at C-28 to aryl propargyl esters and isoxazole carbinol esters
Anti-viral	A	Simultaneous conversion of hydroxyls (R-OH) attached at C-2, C-3 and C-23 to acetates
Anti-oxidant	D/E	Selective conversion of acid (R-COOH) at C-28 to methyl ester
	A	Selective conversion of hydroxyls (R-OH) attached at C-2, C-3 and C-23 to acetates

arjunolic acid may be attributed to its ability to stabilize mast cells and inhibit the release of histamine and acetylcholine in response to antigens.²⁸¹

3.8 Anti-microbial activity

Arjunolic acid has demonstrated moderate antimicrobial activity. Isolated from the leaves of *Syzygium guineense*, arjunolic acid exhibited antibacterial effects against *Bacillus subtilis* (0.5 μg per spot), *Escherichia coli* (3 μg per spot), and *Shigella sonnei* (30 μg per spot).²⁸² It also showed inhibitory activity against *Cryptococcus neoformans*, with an IC_{50} of 20 $\mu\text{g mL}^{-1}$, and demonstrated significant antibacterial activity against *Brucella melitensis*, with a zone of inhibition of 9 mm and a minimum inhibitory concentration (MIC) of 30 $\mu\text{g mL}^{-1}$.^{283,284} Furthermore, a mixture of arjunolic acid and asiatic acid exhibited moderate antifungal effects against *Candida albicans* and *Candida parapsilosis*.²²⁹

3.9 Insecticidal activity

Bhakuni *et al.* identified the insecticidal activity of arjunolic acid. Arjunolic acid, isolated from the stem of *Cornus capitata* (Wall. ex Roxb.) showed significant inhibitory effects on the fourth instar larvae of *Spilarctia obliqua*. A dose-dependent effect was observed, with optimal concentrations for reducing larval feeding and growth found to be 617 ppm and 666 ppm, respectively.²⁴⁷

3.10 Physicochemical properties

Bag *et al.* studied the potential of arjunolic acid as a robust and functional molecular framework suitable for creating nanoscale supramolecular architectures and nanomaterials. This nano-sized triterpenoid demonstrated effective gelation of various organic solvents (methanol, ethanol, propanol, butanol, dichloromethane, chloroform, cyclohexane, benzene, cyclohexanone, diethylene glycol, and triethylene glycol) at low concentrations. In these solvents, low molecular weight gelator molecules self-assembled to form nanofibers, creating a fibrous network that restricted solvent movement, resulting in gel

formation.²⁸⁵ Arjunolic acid also exhibited hierarchical self-assembly in aqueous solutions, forming vesicular structures on nano-to micrometer scales that contribute to gel formation and have proven effective for encapsulating and releasing anticancer drugs such as doxorubicin at physiological pH.

Bag *et al.* also highlighted the self-assembly properties of sodium and potassium salts of arjunolic acid in water and aliphatic alcohols, showing that, at lower concentrations, the molecules form vesicular structures, while, at higher concentrations, they transition into large fibrillar assemblies to create stable gels. These vesicular assemblies are further utilized for the controlled release of various set fluorophores, including doxorubicin, at physiological pH.^{286,287}

4. Analogues of arjunolic acid and their biological activities

Diverse research groups^{7,9,233,288,289} have subjected arjunolic acid to semi-synthetic modifications by carrying out either selective or simultaneous transformations to various rings and functional groups present in arjunolic acid.

Bag *et al.* first reported the synthesis of an 18-crown-6 methyl ester derivative of arjunolic acid. Arjunolic acid was converted into its methyl ester (67). The primary hydroxy group of compounds (67) was protected using trityl chloride to yield compound (68). This compound was further reacted with sodium hydride and pentaethyleneglycol ditosylate in dry THF to yield 23-[(triphenylmethyl)oxy]-28-(methoxycarbonyl)-arjuna-18-crown-6 (69). This compound was further reacted with dichloro acetic acid to obtain compound (70), which was further treated with benzoyl chloride to yield the final product (71, Fig. 5). In the same study this group also reported the synthesis of benzylidene derivatives (72–73, Fig. 5) prepared by reacting arjunolic acid methyl ester with benzaldehyde in the presence of perchloric acid as a catalyst. The hydroxy group at the 2-position of the compound was esterified with 3,5-dinitrobenzoyl chloride, and these derivatives were examined for organogelator properties in various organic solvents.^{285,290}



Subsequently Bag's team synthesized ten aliphatic and aromatic acetals and ketals of arjunolic acid (Fig. 6) and studied their self-assembly properties in diverse organic solvents. Except for the *p*-nitrobenzylidene derivative, these ketals, at low concentrations, formed gel-like dispersions in several organic solvents. Using optical, electron, and atomic-force microscopies, the morphologies were found to consist of fibrillar networks and vesicles capable of entrapping 5(6)-carboxy-fluorescein. X-ray diffractograms showed that the fibrillar structures were crystalline. Additionally, a charge-transfer complex was formed from a 1:1 mixture of ketal derivatives with electron-donating and electron-accepting groups, and the fibrillar network of the 9-anthrylidene derivative dimerized under irradiation. This work demonstrates that minor

structural variations in the ketals significantly influence their aggregation pathways.²⁹²

Reddy *et al.* reported the synthesis and HPLC validation of acetylated arjunolic acid (**84**, Fig. 7), which showed higher antioxidant activity than arjunolic acid, BHT, and vitamin E, as measured by the thiobarbituric acid method (TBA).²⁸⁹

Diallo and co-workers reported the synthesis of the triacetate of arjunolic acid and its methyl ester. Compound **1** was treated with acetic anhydride and pyridine at room temperature for 12 hours to yield compound **84**. Methylation of **1** and **84** was achieved by treating them with diazomethane in diethyl ether at room temperature to yield compounds **85** and **86** (Fig. 7). These compounds were tested for Epstein-Barr virus (EBV) inhibition and further assessed in a two-stage carcinogenesis assay in





Fig. 9 Mechanism of arjunolic acid and its derivatives for various biological activities.



mouse skin. The results indicated that mice treated with the triacetate and triacetate methyl ester of arjunolic acid showed delayed papilloma development compared to the controls.^{293,294}

Bolatito *et al.* synthesized new phenyl acetylene and isoxazole analogues of arjunolic acid. The acid functional group was converted into the propargyl ester using propargyl bromide. This propargyl ester of arjunolic acid was reacted with substituted aryl iodide in the presence of a palladium catalyst to obtain compounds **87–89**. Simultaneously, this propargyl ester was reacted with various bromoaloximes to yield compounds **90–92**. These analogues were evaluated for their tyrosinase and alpha-glucosidase inhibitory activity. All the tested analogues showed stronger inhibitory effects than the standard drug acarbose. One compound demonstrated tyrosinase inhibition with an IC_{50} of $14.3 \pm 7.6 \mu\text{M}$, about three times more potent than kojic acid (IC_{50} of $41.5 \pm 1.0 \mu\text{M}$). Additionally, compound **89** exhibited significant alpha-glucosidase inhibition activity with an IC_{50} close to that of the standard acarbose (10.4 ± 0.06).⁹

Bruno *et al.* synthesized novel arjunolic acid derivatives containing a five-membered A-ring with an enal moiety, combined with additional modifications of COOH (at C-28) as esters and aliphatic amides. Compounds **93–103** were reacted with NaIO_4 at room temperature to yield compounds **104–110**. These derivatives were further treated with acetic acid and piperidine and refluxed in dry benzene before being subjected to reflux with anhydrous magnesium sulphate to yield compounds **111–117**. These derivatives were tested for cytotoxicity on cancer and non-tumor human cell lines, aiming to identify promising compounds. Compound **115** showed the highest selectivity between malignant cells and non-malignant fibroblasts. This compound also induced cell cycle arrest in the G_0/G_1 phase and significantly inhibited the wound closure rate of PANC-1 cancer cells. At a concentration of $0.24 \mu\text{M}$, compound **115** synergistically enhanced gemcitabine cytotoxicity and showed no toxicity at lower doses in preliminary pharmacological studies.⁸

Manohar *et al.* synthesized various acetals (**118–136**, Fig. 8) and amides (**137–159**, Fig. 8) of arjunolic acid and evaluated their anticancer properties. All the synthesized compounds (**118–159**) were tested against 60 cancer cell lines, representing nine different cancer types, at the National Cancer Institute (NCI), USA. These acetals and amides of arjunolic acid exhibited significant activity, particularly against colon, leukemia, breast, and melanoma cancer types. From the library of acetals (**118–136**), compounds **119**, **121**, **126**, and **135** demonstrated activity against all nine cancer types, including breast cancers, colon, CNS, leukemia, non-small cell lung cancer, melanoma, ovarian, renal, and prostate cancer. Among these, compound **126** exhibited the most potent IC_{50} value of $1.56 \mu\text{M}$ against the CT-26 cancer cell line. Further studies on compound **126** revealed that it induces ROS generation, causing cell necrosis by arresting cells in the G_2/M phase of the cell cycle.

Tri-hydroxy amides (**151–155**, Fig. 8) showed greater activity than tri-acetate amides in the series of aniline-substituted amides, Fig. 8. Conversely, in the case of amides derived from heterocyclic amines, tri-acetate amides (**147–150**) were more active compared to tri-hydroxy amides (**156–159**). Substituted

benzylamine amides were inactive against all cancer cell lines. Among the amides, compound **147** exhibited the best IC_{50} values, with $1.88 \mu\text{M}$ against the CT-26 cell line and $0.4 \mu\text{M}$ against the HCT-116 colon cancer cell line. Further investigations revealed that compound **147** induces apoptosis through ROS generation and arrests cells in the G_0/G_1 phase of the cell cycle. *In vivo* studies also showed that compound **147** inhibited cancer growth in the case of the mouse model.⁷

Analysis of the structure–activity relationship (SAR) and modifications responsible for the therapeutic profile of arjunolic acid derivatives is tabulated in Table 3. These studies could help identify future hits, optimize lead molecules with drug-like properties and support future drug discovery efforts. We have also made efforts to capture the mechanisms of action of biological activities exhibited by arjunolic acid and its derivatives in Fig. 9.

5. Conclusion

In summary, natural products continue to be a constant source of interesting biologically active molecules. *T. arjuna*, a traditional medicinal plant native to the Indian subcontinent, was shown to exhibit several medicinal properties and was thus used in traditional medicine. Arjunolic acid, the major triterpenoid constituent of *T. arjuna*, has been reported to exhibit a plethora of biological activities. The structural amenability of arjunolic acid for diverse modifications has attracted the attention of various research groups, affording semi-synthetic derivatives with interesting therapeutic value. To date, many investigations have focused their research efforts on exploring the multi-therapeutic potential of arjunolic acid and delineating its corresponding mechanism of action. However, the limited reports on arjunolic acid analogues and their corresponding biological profile reveal that its therapeutic potential as a source of new drug leads has not yet been thoroughly explored. The continuous rise in FDA approvals for natural-product-based drugs further augments the need to look at arjunolic acid as a potential new source of natural-product-based drug leads. Hence, natural-product-based drug-discovery programs focused on arjunolic acid would build on the existing armamentarium of desirable molecules with drug-like properties and with a potential for clinical trials.

Author contributions

Manohar Bhujel: methodology, investigation, writing – original draft preparation. Sai Giridhar Sarma Kandanur: conceptualization, methodology, investigation, writing – original draft Preparation. Lakshminath Sripada: conceptualization, supervision, writing – reviewing and editing. Nageswara Rao Golakoti: conceptualization, supervision, writing – reviewing and editing.

Conflicts of interest

Authors M. B., L. S. and N. R. G. have a patent “A Process For Separating Arjunolic Acid And Asiatic Acid From Heartwood of Terminalia Arjuna” pending to the Sri Sathya Sai Institute of



Higher Learning, Prasanthi Nilayam-515134, India. The authors declare no additional conflicts of interest.

Data availability

No primary research results, software or codes have been included, and no new data were generated or analysed as part of this review.

Acknowledgements

The authors would like to express their deep sense of gratitude to the Founder Chancellor, Bhagawan Sri Sathya Sai Baba, for His constant guidance and inspiration throughout the work.

References

- S. B. Bharate and C. W. Lindsley, Natural Products Driven Medicinal Chemistry, *J. Med. Chem.*, 2024, **67**, 20723–20730.
- S. G. S. Kandam, N. Tamang, N. R. Golakoti and S. Nanduri, Andrographolide: A natural product template for the generation of structurally and biologically diverse diterpenes, *Eur. J. Med. Chem.*, 2019, **176**, 513–533.
- D. J. Newman and G. M. Cragg, Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019, *J. Nat. Prod.*, 2020, **83**, 770–803.
- D. Domingo-Fernández, Y. Gadiya, A. J. Preto, C. A. Kretzler, S. Mubeen, A. Allen, D. Healey and V. Colluru, Natural Products Have Increased Rates of Clinical Trial Success throughout the Drug Development Process, *J. Nat. Prod.*, 2024, **87**, 1844–1851.
- T. Hemalatha, S. Pulavendran, C. Balachandran, B. M. Manohar and R. Puvanakrishnan, Arjunolic acid: A novel phytomedicine with multifunctional therapeutic applications, *Indian J. Exp. Biol.*, 2010, **48**, 238–247.
- J. Ghosh and P. C. Sil, Arjunolic acid: A new multifunctional therapeutic promise of alternative medicine, *Biochimie*, 2013, **95**, 1098–1109.
- M. Bhujel, L. Sripada, K. Buvanavaragurathan, P. Perumal, D. Jain, N. Pandey, A. Bajaj and N. R. Golakoti, Synthesis and Anti-cancer activity of Acetals of Arjunolic acid, *New J. Chem.*, 2024, **48**, 16957–16967.
- B. M. F. Gonçalves, V. I. S. Mendes, S. M. Silvestre and J. A. R. Salvador, Design, synthesis, and biological evaluation of new arjunolic acid derivatives as anticancer agents, *RSC Med. Chem.*, 2022, **14**, 313–331.
- B. E. Olanipekun, M. G. Ponnappalli, H. K. Patel, K. Munipalle and K. Shaik, Design, synthesis of new phenyl acetylene and isoxazole analogues of arjunolic acid as potent tyrosinase and alpha glucosidase inhibitors, *Nat. Prod. Res.*, 2023, **37**, 1092–1097.
- S. Gupta, J. P. Bishnoi, N. Kumar and H. Kumar, Terminalia arjuna (Roxb .) Wight & Arn: Competent source of bioactive components in functional food and drugs, *Pharma Innov. J.*, 2018, **7**, 223–231.
- S. Thakur, H. Kaurav and G. Chaudhary, Terminalia arjuna: A Potential Ayurvedic Cardio Tonic, *Int. J. Res. Appl. Sci. Biotechnol.*, 2021, **8**, 227–236.
- N. Mishra and A. Kumar, Pharmacological Approach of Terminalia arjuna: A Review, *Plant Cell Biotechnol. Mol. Biol.*, 2021, **22**, 1–5.
- N. Soni and V. K. Singh, Efficacy and Advancement of Terminalia arjuna in Indian Herbal Drug Research: A Review, *Trends Appl. Sci. Res.*, 2019, **14**, 233–242.
- A. Amalraj and S. Gopi, Medicinal properties of Terminalia arjuna (Roxb.) Wight & Arn.: A review, *J. Tradit. Complement. Med.*, 2017, **7**, 65–78.
- S. K. Maulik and K. K. Talwar, Therapeutic Potential of Terminalia Arjuna in Cardiovascular Disorders, *Am. J. Cardiovasc. Drugs*, 2012, **12**, 157–163.
- P. Ramesh and A. Palaniappan, Terminalia arjuna , a Cardioprotective Herbal Medicine – Relevancy in the Modern Era of Pharmaceuticals and Green Nanomedicine — A Review, *Pharmaceuticals*, 2023, **16**, 2–24.
- A. Nanda, S. H. Ansari and S. Khatkar, Arjunolic Acid: A Promising Antioxidant Moiety with Diverse Biological Applications, *Curr. Org. Chem.*, 2017, **21**, 287–293.
- G. R. De Moraes Lima, I. R. P. De Sales, M. R. D. C. Filho, N. Z. T. De Jesus, H. De Sousa Falcão, J. M. Barbosa-Filho, A. G. S. Cabral, A. L. Souto, J. F. Tavares and L. M. Batista, Bioactivities of the genus Combretum (Combretaceae): A review, *Molecules*, 2012, **17**, 9142–9206.
- P. K. Warriar, V. P. K. Nambiar and C. Ramankutty, *Indian Medicinal Plants: A Compendium of 500 Species*, Orient Blackswan, 1st edn, 1993, vol. 5.
- S. Sharma, D. Sharma and N. Agarwal, Diminishing effect of arjuna tree (Terminalia arjuna) bark on the lipid and oxidative stress status of high fat high cholesterol fed rats and development of certain dietary recipes containing the tree bark for human consumption, *Res. Pharm.*, 2012, **2**, 22–30.
- C. Orwa, A. Mutua, R. Kindt, R. Jamnadass and S. Anthony, in *World Agroforestry Database*, 2009, vol. 4, pp. 1–5.
- M. R. Wani and S. S. Singh, Variation in leaf morphological traits of Terminalia arjuna Roxb. in natural population of lower parts of Achanakmar Amarkantak Biosphere Reserve (AABR) of Central India, *Int. J. Adv. Res.*, 2016, **4**, 484–491.
- D. V. Singh, R. K. Verma, S. C. Singh and M. M. Gupta, RP-LC determination of oleane derivatives in Terminalia arjuna, *J. Pharm. Biomed. Anal.*, 2002, **28**, 447–452.
- D. V. Singh, R. K. Verma, M. M. Gupta and S. Kumar, Quantitative determination of oleane derivatives in Terminalia arjuna by high performance thin layer chromatography, *Phytochem. Anal.*, 2002, **13**, 207–210.
- M. Sumitra, P. Manikandan, D. A. Kumar, N. Arutselvan, K. Balakrishna, B. M. Manohar and R. Puvankrishnan, Experimental myocardial necrosis in rats : Role of arjunolic acid on platelet aggregation , coagulation and antioxidant status, *Mol. Cell. Biochem.*, 2001, **224**, 135–142.
- L. R. Row, P. S. Murty, G. S. Rao, C. P. Sastry and K. V. J. Rao, Chemical examination of Terminalia species. XII. Isolation



- & structure determination of arjunic acid, a new trihydroxytriterpene carboxylic acid from Terminalia arjuna bark, *Indian J. Chem.*, 1970, **8**, 772–775.
- 27 F. E. Kandil and M. I. Nasar, A Tannin Anti-Cancer Promoter From Terminalia arjuna, *Phytochemistry*, 1998, **47**, 1567–1568.
- 28 M. Saxena, U. Faridi, R. Mishra, M. M. Gupta, M. P. Darokar, S. K. Srivastava, D. Singh, S. Luqman and S. P. S. Khanuja, Cytotoxic agents from Terminalia arjuna, *Planta Med.*, 2007, **73**, 1486–1490.
- 29 A. S. R. Anjaneyulu and A. V. R. Prasad, Structure of terminic acid, a dihydroxytriterpene carboxylic acid from Terminalia arjuna, *Phytochemistry*, 1983, **22**, 993–998.
- 30 F. Y. Sun, X. P. Chen, J. H. Wang, H. L. Qin, S. R. Yang and G. H. Du, Arjunic acid, a strong free radical scavenger from Terminalia arjuna, *Am. J. Chin. Med.*, 2008, **36**, 197–207.
- 31 W. Wang, Z. Ali, Y. Shen, X. C. Li and I. A. Khan, Ursane triterpenoids from the bark of Terminalia arjuna, *Fitoterapia*, 2010, **81**, 480–484.
- 32 G. S. Arumugam, A. Sen, S. S. Dash, K. Mitra, M. Doble, G. Rajaraman and S. N. Gummadi, Arjunetin as a promising drug candidate against SARS-CoV-2: molecular dynamics simulation studies, *J. Biomol. Struct. Dyn.*, 2022, **40**, 12358–12379.
- 33 M. S. Kim, D. Y. Lee, S. H. Sung and W. K. Jeon, Anti-cholinesterase activities of hydrolysable tannins and polyhydroxytriterpenoid derivatives from Terminalia chebula Retz. fruit, *Rec. Nat. Prod.*, 2018, **12**, 284–289.
- 34 M. Ishimatsu, T. Tanaka, G. Nonaka and I. N. J. Lin, The Structure of Arjungenin. A New Sapogenin from Terminalia arjuna, *Chem. Pharm. Bull.*, 1976, **24**, 178–180.
- 35 T. Honda, T. Murae, T. Tsuyuki, T. Takahashi and M. Sawai, Arjungenin, Arjunglucoside I, and Arjunglucoside II. A new Triterpene and New Triterpene Glucosides from Terminalia arjuna, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3213–3218.
- 36 P. N. Sharma, S. AB, K. RS and P. SP, Arjunolone—a new flavone from stem bark of Terminalia arjuna, *Indian J. Chem. B*, 1982, **21**, 263.
- 37 V. K. Tripathi, V. B. Pandey, K. N. Udupa and G. Rücker, Arjunolitin, a triterpene glycoside from Terminalia arjuna, *Phytochemistry*, 1992, **31**, 349.
- 38 A. Ali, G. Kaur, H. Hamid, T. Abdullah, M. Ali, M. Niwa and M. S. Alam, Terminoside A, a new triterpene glycoside from the bark of Terminalia arjuna inhibits nitric oxide production in murine macrophages, *J. Asian Nat. Prod. Res.*, 2003, **5**, 137–142.
- 39 A. Ali, G. Kaur, K. Hayat, M. Ali and M. Ather, A novel naphthanol glycoside from Terminalia arjuna with antioxidant and nitric oxide inhibitory activities, *Pharmazie*, 2003, **58**, 932–934.
- 40 W. Wang, Z. Ali, X. C. Li, Y. Shen and I. A. Khan, Triterpenoids from two terminalia species, *Planta Med.*, 2010, **76**, 1751–1754.
- 41 R. Hossain, R. Sultana, M. Din Islam, S. Zaman and M. I. Choudhary, Isolation of two new triterpene glycoside from the fruits of Terminalia arjuna and their *in vitro* and *in silico* studies, *Nat. Prod. Res.*, 2022, **37**, 1–9.
- 42 T. Patnaik, R. K. Dey and P. Gouda, Isolation of triterpenoid glycoside from bark of Terminalia arjuna using chromatographic technique and investigation of pharmacological behavior upon muscle tissues, *E-J. Chem.*, 2007, **4**, 474–479.
- 43 M. S. Alam, G. Kaur, A. Ali, H. Hamid, M. Ali and M. Athar, Two new bioactive oleanane triterpene glycosides from Terminalia arjuna, *Nat. Prod. Res.*, 2008, **22**, 1279–1288.
- 44 A. MU, M. K. JK and N. T. Ulla, Terminic acid, A new trihydroxytriterpene carboxylic-acid from bark of Terminalia-arjuna, *Indian J. Chem. Sect. B-Organic Chem. Incl. Med. Chem.*, 1983, **22**, 738–740.
- 45 I. E. Cock, The medicinal properties and phytochemistry of plants of the genus Terminalia (Combretaceae), *Inflammopharmacology*, 2015, **23**, 203–229.
- 46 M. F. Manzoor, N. Ahmad, Z. Ahmed, R. Siddique, X. A. Zeng, A. Rahaman, R. Muhammad Aadil and A. Wahab, Novel extraction techniques and pharmaceutical activities of luteolin and its derivatives, *J. Food Biochem.*, 2019, **43**, 1–19.
- 47 G. Chaudhari and R. T. Mahajan, Bioprospecting of Terminalia arjuna stem bark and its flavonoids for antimicrobial and anti-biofilm potential, *J. Chem. Pharm. Res.*, 2015, **7**, 834–841.
- 48 W. R. García-Niño and C. Zazueta, Ellagic acid: Pharmacological activities and molecular mechanisms involved in liver protection, *Pharmacol. Res.*, 2015, **97**, 84–103.
- 49 N. A. AL Zahrani, R. M. El-Shishtawy and A. M. Asiri, Recent developments of gallic acid derivatives and their hybrids in medicinal chemistry: A review, *Eur. J. Med. Chem.*, 2020, **204**, 112609.
- 50 D. A. Vattam and K. Shetty, Biological Functionality Of Ellagic Acid: A Review, *J. Food Biochem.*, 2005, **29**, 234–266.
- 51 S. P. Bangar, V. Chaudhary, N. Sharma, V. Bansal, F. Ozogul and J. M. Lorenzo, Kaempferol: A flavonoid with wider biological activities and its applications, *Crit. Rev. Food Sci. Nutr.*, 2023, **63**, 9580–9604.
- 52 J. K. Kim and S. U. Park, Recent Studies on Kaempferol and its Biological and Pharmacological Activities, *EXCLI J.*, 2020, **19**, 627–634.
- 53 A. Saha, V. M. Pawar and S. Jayaraman, Characterisation of polyphenols in Terminalia arjuna bark extract, *Indian J. Pharm. Sci.*, 2012, **74**, 339–347.
- 54 T. W. Sirk, E. F. Brown, M. Friedman and A. K. Sum, Molecular binding of catechins to biomembranes: relationship to biological activity, *J. Agric. Food Chem.*, 2009, **57**, 6720–6728.
- 55 S. Takahashi, H. Tanaka, Y. Hano, K. Ito, T. Nomura and K. Shigenobu, Hypotensive effect in rats of hydrophilic extract from Terminalia arjuna containing tannin-related compounds, *Phyther. Res.*, 1997, **11**, 424–427.
- 56 T. C. Lin, S. I. Chien, H. F. Chen and F. L. Hsu, Tannins and related compounds from Combretaceae plants, *Chinese Pharm. J.*, 2000, **52**, 1–26.
- 57 A. P. Kulkarni, H. S. Mahal, S. Kapoor and S. M. Aradhya, *In vitro* studies on the binding, antioxidant, and cytotoxic



- action of punicalagin, *J. Agric. Food Chem.*, 2007, **55**, 1491–1500.
- 58 P. L. Kuo, Y. L. Hsu, T. C. Lin, L. T. Lin, J. K. Chang and C. C. Lin, Casuarinin from the bark of *Terminalia arjuna* induces apoptosis and cell cycle arrest in human breast adenocarcinoma MCF-7 cells, *Planta Med.*, 2005, **71**, 237–243.
- 59 A. R. Arau̇jo, A. C. Arau̇jo, R. L. Reis and R. A. Pires, Vescalagin and Castalagin Present Bactericidal Activity toward Methicillin-Resistant Bacteria, *ACS Biomater. Sci. Eng.*, 2021, **7**, 1022–1030.
- 60 T. M. Souza-Moreira, J. A. Severi, K. Lee, K. Preechath, E. Santos, N. A. R. Gow, C. A. Munro, W. Vilegas and R. C. L. R. Pietro, Anti-Candida targets and cytotoxicity of casuarinin isolated from *Plinia cauliflora* leaves in a bioactivity-guided study, *Molecules*, 2013, **18**, 8095–8108.
- 61 D. J. Kwon, Y. S. Bae, S. M. Ju, A. R. Goh, S. Y. Choi and J. Park, Casuarinin suppresses TNF- α -induced ICAM-1 expression via blockade of NF- κ B activation in HaCaT cells, *Biochem. Biophys. Res. Commun.*, 2011, **409**, 780–785.
- 62 A. M. Muddathir, K. Yamauchi and T. Mitsunaga, Anti-acne activity of tannin-related compounds isolated from *Terminalia laxiflora*, *J. Wood Sci.*, 2013, **59**, 426–431.
- 63 Z. Chang, Q. Zhang, W. Liang, K. Zhou, P. Jian, G. She and L. Zhang, A Comprehensive Review of the Structure Elucidation of Tannins from *Terminalia* Linn, *Evid. Based Complement. Alternat. Med.*, 2019, **2019**, 1–26.
- 64 A. Sen, P. Dhavan, K. K. Shukla, S. Singh and G. Tejavathi, Analysis of IR, NMR and Antimicrobial Activity of β -Sitosterol Isolated from *Momordica charantia*, *Sci Secur. J Biotech*, 2012, **1**, 9–13.
- 65 J. M. Castellano, S. Ramos-Romero and J. S. Perona, Oleanolic Acid: Extraction, Characterization and Biological Activity, *Nutrients*, 2022, **14**, 1–29.
- 66 B. K. Chouksey and S. K. Srivasava, Antifungal agent from *Terminalia arjuna*, *Indian J. Chem. B.*, 2001, **40**, 354–356.
- 67 R. K. Upadhyay, M. B. Pandey, R. N. Jha, V. P. Singh and V. B. Pandey, Triterpene glycoside from *terminalia arjuna*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 207–212.
- 68 R. M. Rastogi, Compendium of Indian medicinal plants, *Cent. Drug Res. Inst.*, 1990, **1**, 388–389.
- 69 R. M. Lopes, T. Da Silveira Agostini-Costa, M. A. Gimenes and D. Silveira, Chemical composition and biological activities of *Arachis* species, *J. Agric. Food Chem.*, 2011, **59**, 4321–4330.
- 70 M. H. Radi, R. A. El-Shiekh, A. M. El-Halawany and E. Abdel-Sattar, Friedelin and 3β -Friedelinol: Pharmacological Activities, *Rev. Bras. Farmacogn.*, 2023, **33**, 886–900.
- 71 P. P. S. Ambika and S. M. S. Chauhan, Activity-guided isolation of antioxidants from the leaves of *Terminalia arjuna*, *Nat. Prod. Res.*, 2014, **28**(10), 760–763.
- 72 R. Liu, B. Zhao, D. E. Wang, T. Yao, L. Pang, Q. Tu, S. M. Ahmed, J. J. Liu and J. Wang, Nitrogen-containing apigenin analogs: Preparation and biological activity, *Molecules*, 2012, **17**, 14748–14764.
- 73 P. Udaysing and D. K. Gaikwad, Comprehensive Review of Arjuna: A Potent Drug in the Management of Cardiac Disorder, *Int. J. Drug Formul. Res.*, 2014, **2**, 166–184.
- 74 R. L. S. Sikarwar, B. Pathak and A. Jaiswal, Some unique ethnomedicinal perceptions of tribal communities of Chitrakoot, Madhya Pradesh, *Indian J. Tradit. Knowl.*, 2008, **7**, 613–617.
- 75 M. Anisuzzaman, A. H. M. M. Rahman, M. Harun-Or-Rashid, A. T. M. Naderuzzaman and A. K. M. R. Islam, An ethnobotanical study of Madhupur, Tangail, *J. Appl. Sci. Res.*, 2007, **3**, 519–530.
- 76 C. Muthu, M. Ayyanar, N. Raja and S. Ignacimuthu, Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India, *J. Ethnobiol. Ethnomed.*, 2006, **2**, 1–10.
- 77 K. Yesodharan and K. A. Sujana, Ethnomedicinal knowledge among Malamalasar tribe of Parambikulam wildlife sanctuary, Kerala, *Indian J. Tradit. Knowl.*, 2007, **6**, 481–485.
- 78 A. B. Prusti and K. K. Behera, Ethnobotanical Exploration of Malkangiri District of Orissa, India, *Ethnobot. Leaflet*, 2007, **11**, 122–140.
- 79 L. Sharma and A. Kumar, Traditional medicinal practices of Rajasthan, *Indian J. Tradit. Knowl.*, 2007, **06**, 531–533.
- 80 K. Bone, *Clinical Applications of Ayurvedic and Chinese Herbs. Monographs for the Western Herbal Practitioner*, Phyther. Press, Australia, 1996, pp. 137–141.
- 81 S. Nesamony, *Oushadha Sasyangal – 2*, State Institute of Languages, Tiruvananthapuram, 2001.
- 82 L. D. Kapoor, *CRC Handbook of Ayurvedic Medicinal Plants*, CRC press, 1st edn, 2018.
- 83 N. Singh, K. K. Kapur, S. P. Singh, K. Shanker, J. N. Sinha and R. P. Kohli, Mechanism of cardiovascular action of *Terminalia arjuna*, *Planta Med.*, 1982, **45**, 102–104.
- 84 S. Dwivedi and M. P. Agarwal, Antianginal and cardioprotective effects of *Terminalia arjuna*, an indigenous drug, in coronary artery disease, *J. Assoc. Physicians India*, 1994, **42**(4), 287–289.
- 85 S. Dwivedi and R. Jauhari, Beneficial effects of *Terminalia arjuna* in coronary artery disease, *Indian Heart J*, 1997, **49**(5), 507–510.
- 86 A. Bharani, A. Ganguly and K. D. Bhargava, Salutary effect of *Terminalia Arjuna* in patients with severe refractory heart failure, *Int. J. Cardiol.*, 1995, **49**, 191–199.
- 87 A. Ram, P. Lauria, R. Gupta, P. Kumar and V. N. Sharma, Hypocholesterolaemic effects of *Terminalia arjuna* tree bark, *J. Ethnopharmacol.*, 1997, **55**, 165–169.
- 88 H. P. Shaila, S. L. Udupa and A. L. Udupa, Hypolipidemic activity of three indigenous drugs in experimentally induced atherosclerosis, *Int. J. Cardiol.*, 1998, **67**, 119–124.
- 89 P. U. Kumar, P. Adhikari, P. Pereira and P. Bhat, Safety and efficacy of Hartone in stable angina pectoris—an open comparative trial, *J. Assoc. Physicians India*, 1999, **47**, 685–689.
- 90 K. Gauthaman, M. Maulik, R. Kumari, S. C. Manchanda, A. K. Dinda and S. K. Maulik, Effect of chronic treatment with bark of *Terminalia arjuna*: a study on the isolated



- ischemic-reperfused rat heart, *J. Ethnopharmacol.*, 2001, **75**, 197–201.
- 91 R. Gupta, S. Singhal, A. Goyle and V. N. Sharma, Antioxidant and hypocholesterolaemic effects of Terminalia arjuna tree-bark powder: a randomised placebo-controlled trial, *J. Assoc. Physicians India*, 2001, **49**, 231–235.
- 92 B. Chopra, Antianginal and cardioprotective effects of Terminalia arjuna, *J. Assoc. Physicians India*, 1994, **42**, 756.
- 93 O. M. Bhat, P. U. Kumar, K. R. Rao, A. Ahmad and V. Dhawan, Terminalia arjuna prevents Interleukin-18-induced atherosclerosis via modulation of NF- κ B/PPAR- γ -mediated pathway in Apo E $^{-/-}$ mice, *Inflammopharmacology*, 2018, **26**, 583–598.
- 94 A. Parveen, R. Babbar, S. Agarwal, A. Kotwani and M. Fahim, Mechanistic clues in the cardioprotective effect of Terminalia arjuna bark extract in isoproterenol-induced chronic heart failure in rats, *Cardiovasc. Toxicol.*, 2011, **11**, 48–57.
- 95 N. K. Mary, B. H. Babu and J. Padikkala, Antiatherogenic effect of Caps HT2, a herbal Ayurvedic medicine formulation, *Phytomedicine*, 2003, **10**, 474–482.
- 96 F. Khaliq, A. Parveen, S. Singh, R. Gondal, M. E. Hussain and M. Fahim, Improvement in myocardial function by Terminalia arjuna in streptozotocin-induced diabetic rats: Possible mechanisms, *J. Cardiovasc. Pharmacol. Ther.*, 2013, **18**, 481–489.
- 97 S. J. Kaur, I. S. Grover and S. Kumar, Modulatory effects of a tannin fraction isolated from Terminalia arjuna on the genotoxicity of mutagens in Salmonella typhimurium, *Food Chem. Toxicol.*, 2000, **38**, 1113–1119.
- 98 A. Nagpal, L. S. Meena, S. Kaur, I. S. Grover, R. Wadhwa and S. C. Kaul, Growth suppression of human transformed cells by treatment with bark extracts from a medicinal plant, Terminalia arjuna, *Vitr. Cell. Dev. Biol. – Anim.*, 2000, **36**, 544–547.
- 99 G. Scassellati-Sforzolini, L. M. Villarini, L. M. Moretti, L. M. Marcarelli, R. Pasquini, C. Fatigoni, L. S. Kaur, S. Kumar and I. S. Grover, Antigenotoxic properties of Terminalia arjuna bark extracts, *J. Environ. Pathol. Toxicol. Oncol.*, 1999, **18**, 119–125.
- 100 T. K. Reddy, P. Seshadri, K. K. R. Reddy, G. C. Jagetia and C. D. Reddy, Effect of Terminalia arjuna Extract on Adriamycin-induced DNA Damage, *Phyther. Res.*, 2008, **22**, 1188–1194.
- 101 S. Singh, S. K. Verma and S. K. Singh, Analysis of anti-cancer potential of Terminalia arjuna, *Int. J. Adv. Sci. Res. Manag.*, 2017, **2**, 82–87.
- 102 P. Rani and N. Khullar, Antimicrobial evaluation of some medicinal plants for their anti-enteric potential against multi-drug resistant Salmonella typhi, *Phyther. Res.*, 2004, **18**, 670–673.
- 103 K. R. Aneja, C. Sharma and R. Joshi, Antimicrobial activity of Terminalia arjuna Wight & Arn: An ethnomedicinal plant against pathogens causing ear infection, *Braz. J. Otorhinolaryngol.*, 2012, **78**, 68–74.
- 104 J. P. Jethinlalkhosh and A. Antony, Antibacterial and cytotoxic activity of aqueous and methanolic extracts of Terminalia arjuna, *Int. J. Res. Pharm. Sci.*, 2013, **4**, 36–39.
- 105 T. Javed, S. Riaz, M. Uzair, G. Mustafa, A. Mohyuddin and B. A. Ch., Biological activity of Terminalia arjuna on Human Pathogenic Microorganisms, *Pakistan J. Pharm. Res.*, 2016, **2**, 23.
- 106 V. Kumar, N. Sharma, A. Sourirajan, P. K. Khosla and K. Dev, Comparative evaluation of antimicrobial and antioxidant potential of ethanolic extract and its fractions of bark and leaves of Terminalia arjuna from north-western Himalayas, India, *J. Tradit. Complement. Med.*, 2018, **8**, 100–106.
- 107 S. Mandal, A. Patra, A. Samanta, S. Roy, A. Mandal, T. Das Mahapatra, S. Pradhan, K. Das and D. K. Nandi, Analysis of phytochemical profile of Terminalia arjuna bark extract with antioxidative and antimicrobial properties, *Asian Pac. J. Trop. Biomed.*, 2013, **3**, 960–966.
- 108 F. Murad, K. A. Alam, R. M. Mazumdar, S. Islam, M. N. Nipa, A. Iqbal and H. R. Bhuiyan, Anti-bacterial Activity of the Extract of Terminalia arjuna Against Multi Antibiotic Resistant *Vibrio cholerae*, *J. Sci. Res.*, 2010, **3**, 129.
- 109 S. Debnath, D. Dey and S. Hazra, Antibacterial and antifungal activity of Terminalia arjuna Wight & Arn. bark against multi-drug resistant clinical isolates, *J. Coast. Life Med*, 2014, **4**(4), 312–318.
- 110 S. Ramya, T. Kalaivani, C. Rajasekaran, P. Jephanderamohan, N. Alaguchamy, M. Kalayansundaram and R. Jayakumararaj, Antimicrobial Activity of Aqueous Extracts of Bark, Root, Leaves and Fruits of Terminalia arjuna Wight & Arn, *Ethnobot. Leaflet.*, 2008, **12**, 1192–1197.
- 111 D. K. Meena, N. P. Sahu, P. P. Srivastava, M. Jadhav, R. Prasad, R. C. Mallick, A. K. Sahoo, B. K. Behera, D. Mohanty and B. K. Das, Effective valorisation of facile extract matrix of Terminalia arjuna (Roxb) against elite microbes of aquaculture industry—a credence to bioactive principles: Can it be a sustainability paradigm in designing broad spectrum antimicrobials?, *Ind. Crops Prod.*, 2021, **171**, 113905.
- 112 G. Phani Kumar, K. Navya, E. M. Ramya, M. Venkataramana, T. Anand and K. R. Anilakumar, DNA damage protecting and free radical scavenging properties of Terminalia arjuna bark in PC-12 cells and plasmid DNA, *Free Radic. Antioxid.*, 2013, **3**, 35–39.
- 113 G. L. shastry Viswanatha, S. K. Vaidya, C. Ramesh, N. Krishnadas and S. Rangappa, Antioxidant and antimutagenic activities of bark extract of Terminalia arjuna, *Asian Pac. J. Trop. Med.*, 2010, **3**, 965–970.
- 114 S. A. Shahid Chatha, Bioactive Components and Antioxidant Properties of Terminalia arjuna L.Extracts, *J. Food Process. Technol.*, 2014, **5**(2), 1–5.
- 115 A. K. Ghosh, E. Mitra, M. Dutta, D. Mukherjee, A. Basu, S. B. Firdaus, D. Bandyopadhyay and A. Chattopadhyay, Protective effect of aqueous bark extract of Terminalia arjuna on Cu²⁺-ascorbate induced oxidative stress in



- vitro: Involvement of antioxidant mechanism(S), *Asian J. Pharm. Clin. Res.*, 2013, **6**, 196–200.
- 116 B. Raghavan and S. K. Kumari, Effect of Terminalia arjuna stem bark on antioxidant status in liver and kidney of alloxan diabetic rats, *Indian J. Physiol. Pharmacol.*, 2006, **50**, 133–142.
- 117 P. Manna and P. C. Sil, Arjunolic acid: beneficial role in type 1 diabetes and its associated organ pathophysiology, *Free Radic. Res.*, 2012, **46**, 815–830.
- 118 N. Aslam, M. Ahmad, K. Barkat and M. Abbas, Antioxidant and Anti-urease Activity of Various Solvent Extracts of “Terminalia arjuna” Seeds, *J. Pharm. Res. Int.*, 2018, **20**, 1–5.
- 119 N. Anbalagan, M. Mallika, S. Kuruvilla, M. V. V. Prasad, A. Patra and K. Balakrishna, Anti-oxidant and hepatoprotective activities of alcoholic extract of Terminalia arjuna, *Nat. Prod. Sci.*, 2007, **13**(2), 105–109.
- 120 P. Manna, M. Sinha and P. C. Sil, Phytomedicinal activity of Terminalia arjuna against carbon tetrachloride induced cardiac oxidative stress, *Pathophysiology*, 2007, **14**, 71–78.
- 121 D. K. Meena, A. K. Sahoo, P. P. Srivastava, N. P. Sahu, M. Jadhav, M. Gandhi, H. S. Swain, S. Borah and B. K. Das, On valorization of solvent extracts of Terminalia arjuna (arjuna) upon DNA scission and free radical scavenging improves coupling responses and cognitive functions under *in vitro* conditions, *Sci. Rep.*, 2021, **11**, 1–20.
- 122 M. A. Morshed, M. A. Uddin, T. Hasan, T. Ahmed, F. Uddin, M. Zakaria, A. Haque and A. K. Parvez, Evaluation of Analgesic and Anti-inflammatory Effects of Terminalia arjuna Ethanol Extract, *Int. J. Pharm. Sci. Res.*, 2011, **2**, 2577–2585.
- 123 F. Abo-Elghiet, A. Abd-elstara, A. Metwaly and A.-E. Mohammad, Antioxidant, anti-inflammatory, and antimicrobial evaluation of Terminalia arjuna leaves, fruits, and bark, *Azhar Int. J. Pharm. Med. Sci.*, 2022, **2**, 148–158.
- 124 L. Chularojanamontri, N. Chiruntanat, K. Jaijoi, N. Soonthornchareonnon and S. Sireeratawong, Anti-inflammatory, Analgesic and Antipyretic Activity of Terminalia arjuna Roxb. Extract in Animal Models, *J. Med. Assoc. Thail.*, 2017, **100**, S92–S97.
- 125 N. Dube, C. Nimgulkar and D. K. Bharatraj, Validation of therapeutic anti-inflammatory potential of Arjuna Ksheera Paka – A traditional Ayurvedic formulation of Terminalia arjuna, *J. Tradit. Complement. Med.*, 2017, **7**, 414–420.
- 126 R. M. Sarma, B. Subramanian, P. Tamilmaran and G. Ramakrishnan, Anti-inflammatory and Anti-granuloma effect of the extract of the leaf of Terminalia arjuna (Roxb.) Wight & Arn, *J. Pharm. Sci. Res.*, 2019, **11**, 3579–3586.
- 127 S. Halder, N. Bharal, P. K. Mediratta, I. Kaur and K. K. Sharma, Anti-inflammatory, immunomodulatory and antinociceptive activity of Terminalia arjuna roxb bark powder in mice and rats, *Indian J. Exp. Biol.*, 2009, **47**, 577–583.
- 128 S. Shweta Shaligram, P. Tabassum Arif, K. Surekha Babasaheb and P. Scholar, Review on Arjuna (Terminalia Arjuna Roxb.) With Special Reference To Prameha (Diabetes), *World J. Pharm. Res.*, 2019, **8**, 1477–1489.
- 129 A. Gupta and S. R. Chaphalkar, Haemolytic Activities and Antidiabetic Effect of Terminalia arjuna and Emblica officinalis, *Eur. J. Pharm. Med. Res.*, 2016, **3**, 334–338.
- 130 C. Kumar, R. Kumar and S. Nehar, Phytochemical properties, total antioxidant status of acetone and methanol extract of Terminalia arjuna Roxb. bark and its hypoglycemic effect on Type-II diabetic albino rats, *J. Pharmacogn. Phytochem.*, 2013, **2**, 199–208.
- 131 S. Singh, N. Shukla, M. Bhowmick and J. Rathi, Evaluation of Anti-Diabetic Activity of Terminalia Arjuna Root Extract in Alloxan Induced Diabetic Rats, *J. Biomed. Pharm. Res.*, 2019, **8**(1), 20–29.
- 132 M. Alam Morshed, A. Haque, B. Rokeya and L. Ali, Anti-Hyperglycemic and Lipid Lowering Effect of Terminalia arjuna Bark Extract on Streptozotocin induced Type 2 Diabetic Model Rats, *Int. J. Pharm. Pharm. Sci.*, 2011, **3**, 449–453.
- 133 S. A. Shengule, S. Mishra and S. Bodhale, Inhibitory effect of a standardized hydroethanolic extract of Terminalia arjuna bark on alpha-amylase enzyme, *Asian J. Pharm. Clin. Res.*, 2018, **11**, 366–369.
- 134 M. K. Borde, I. R. Mohanty, U. Maheswari, R. K. Suman and Y. Deshmukh, Natural dipeptidyl peptidase-4 inhibitor Terminalia arjuna mitigates myocardial infarction co-existing with diabetes in experimental rats, *J. Diabetes, Metab. Disord. Control*, 2018, **5**, 48–56.
- 135 H. A. J. Thomson, O. O. Ojo, P. R. Flatt and Y. H. A. Abdel-Wahab, Aqueous Bark Extracts of Terminalia arjuna Stimulates Insulin Release, Enhances Insulin Action and Inhibits Starch Digestion and Protein Glycation *in vitro*, *Austin J. Endocrinol. Diabetes*, 2014, **1**(1).
- 136 S. P. Sangamithira, J. Revathy, S. Sheik Abdullah and P. Sampath Kumar, The hepatoprotective effect of ethanolic bark extract of Terminalia arjuna on paracetamol induced liver damage, *Biosci. Biotechnol. Res. Asia*, 2011, **8**, 777–781.
- 137 M. Biswas, K. Biswas, T. K. Karan, S. Bhattacharya, A. K. Ghosh and P. K. Haldar, Evaluation of Analgesic and Anti-inflammatory Activities of Terminalia arjuna Leaf, *J. Phytol.*, 2011, **3**, 33–38.
- 138 S. Sivalokanathan, M. Ilayaraja and M. P. Balasubramanian, Anticancer potency of Terminalia arjuna bark on N-nitrosodiethylamine- induced hepatocellular carcinoma in rats, *Nat. Prod. Sci.*, 2004, **10**, 190–195.
- 139 G. M. Chaudhari and R. T. Mahajan, In Vitro Hepatoprotective Activity of Terminalia arjuna Stem Bark and Its Flavonoids Against CCl₄ Induced Hepatotoxicity in Goat Liver Slice Culture, *Asian J. Plant Sci. & Res.*, 2016, **6**, 10–17.
- 140 A. P. S. Vishwakarma, A. Vishwe, P. K. Sahu and A. K. Chaurasiya, Screening of Hepatoprotective Potential of Ethanolic and Aqueous Extract of Terminalia arjuna



- bark against paracetamol/CCl₄ Induced Liver Damage in Wistar Albino Rats, *Int. J. Pharm. Arch.*, 2013, **10**, 243–250.
- 141 P. Manna, M. Sinha and P. C. Sil, Aqueous extract of Terminalia arjuna prevents carbon tetrachloride induced hepatic and renal disorders, *BMC Complement. Altern. Med.*, 2006, **6**, 1–10.
- 142 S. Gomathi and K. M. S. Rathinam, Evaluation of repellency and antifeedant activities of Saraca asoca and Terminalia arjuna Bark extracts against Sitophilus oryzae, *Int. J. Zool. Stud.*, 2017, **2**, 282–285.
- 143 P. K. Mukherjee, K. Mukherjee, M. R. Kumar, M. Pal and B. P. Saha, Evaluation of wound healing activity of herbal formulation, *Asian J. Chem.*, 2007, **19**, 1273–1276.
- 144 M. Chaudhari and S. Mengi, Evaluation of phytoconstituents of Terminalia arjuna for wound healing activity in rats, *Phyther. Res.*, 2006, **20**, 799–805.
- 145 M. M. Rane and S. A. Mengi, Comparative effect of oral administration and topical application of alcoholic extract of Terminalia arjuna bark on incision and excision wounds in rats, *Fitoterapia*, 2003, **74**, 553–558.
- 146 D. K. Meena, A. K. Sahoo, M. Jayant, N. P. Sahu, P. P. Srivastava, H. S. Swain, B. K. Behera, K. Satvik and B. K. Das, Bioconversion of Terminalia arjuna bark powder into a herbal feed for Labeo rohita: Can it be a sustainability paradigm for Green Fish production?, *Anim. Feed Sci. Technol.*, 2022, **284**, 115132.
- 147 D. K. Meena, A. K. Sahoo, H. Chowdhury, H. S. Swain, N. P. Sahu, B. K. Behera, P. P. Srivastava and B. K. Das, Effects of extraction methods and solvent systems on extract yield, proximate composition and mineral profiling of Terminalia arjuna (Arjuna) dry powders and solvent extracts, *J. Innov. Pharm. Biol. Sci.*, 2020, **7**, 22–31.
- 148 P. Parmar and K. Khamrui, Development of process for the production of arjuna herbal ghee from buffalo milk, *Indian J. Anim. Sci.*, 2017, **87**, 203–207.
- 149 S. Bishnoi and S. S. Ahlawat, Development of Buffalo Meat Rolls Incorporated with Aloe Vera Gel and Arjun Tree Bark Extract, *Haryana Vetnary*, 2015, **54**, 174–177.
- 150 P. D. Sawale, R. Pothuraju, S. Abdul Hussain, A. Kumar, S. Kapila and G. R. Patil, Hypolipidaemic and anti-oxidative potential of encapsulated herb (Terminalia arjuna) added vanilla chocolate milk in high cholesterol fed rats, *J. Sci. Food Agric.*, 2016, **96**, 1380–1385.
- 151 K. Batta and H. Rajput, Production, Consumption and Benefits of Different Herbal Tea: A Review, *J. Food Ind. Microbiol.*, 2021, **7**, 128.
- 152 S. Verma, S. R. Murti, D. L. Singh, R. K. Tiwari and N. Quadri, Formulation of Herbal Green Tea Using Terminalia Arjuna Bark, Glycyrrhiza Glabra Root, Gymnema Sylvester Leaf And Green Cardamom Seeds: Its Nutritional & Phytochemical Analysis, *Br. J. Pharm. Med. Res.*, 2020, **4**, 2202–2211.
- 153 D. K. Meena, B. K. Das, A. K. Sahoo, N. P. Sahu, P. P. Srivastava and S. Borah, Terminalia arjuna Bark Powder as a Potential Immunomodulator in Labeo rohita: Enhanced Hematological, Adaptive, and Humoral Responses against Bacterial Pathogens and Concordant Liver Histomorphology, *Pathogens*, 2024, **13**(4), 295.
- 154 D. K. Meena, S. P. Panda, A. K. Sahoo, P. P. Srivastava, N. P. Sahu, M. Kumari, S. Samantaray, S. Borah and B. K. Das, Immunogenic Effects of Dietary Terminalia arjuna Bark Powder in Labeo rohita, a Fish Model: Elucidated by an Integrated Biomarker Response Approach, *Animals*, 2023, **13**(1), 39.
- 155 K. Aamir, H. U. Khan, C. F. Hossain, M. Rejina Afrin, I. Shaik, N. Salleh, N. Giribabu and A. Arya, Oral toxicity of arjunolic acid on hematological, biochemical and histopathological investigations in female Sprague Dawley rats, *PeerJ*, 2019, **7**, e8045.
- 156 D. K. Meena, A. K. Sahoo, H. S. Swain, S. Borah, P. P. Srivastava, N. P. Sahu and B. K. Das, Prospects and perspectives of virtual in-vitro toxicity studies on herbal extracts of Terminalia arjuna with enhanced stratagem in Artemia salina model: A panacea to explicit the credence of solvent system in brine shrimp lethality bioassay, *Emirates J. Food Agric.*, 2020, **32**, 25–37.
- 157 V. Anuradha, P. Shankar, P. Bhuvana, M. Syedali and N. Yogananth, Terminalia Arjuna Bark Assisted Biosynthesis, Characterization and Bioactivity of Metal oxide Nanoparticles, *J. Chem. Pharm. Res.*, 2017, **9**, 34–46.
- 158 A. A. Dudhane, S. R. Waghmode, L. B. Dama and P. Vaibhav, Synthesis and Characterization of Gold Nanoparticles using Plant Extract of Terminalia arjuna with Antibacterial Activity, *Int. J. Nanosci. Nanotechnol.*, 2019, **15**, 75–82.
- 159 M. Rakhi and B. B. Gopal, Terminalia Arjuna Bark Extract Mediated Size Controlled Synthesis of Polyshaped Gold Nanoparticles and Its Application in Catalysis, *Int. J. Res. Chem. Environ.*
- 160 K. Gopinath, K. S. Venkatesh, R. Ilangovan, K. Sankaranarayanan and A. Arumugam, Green synthesis of gold nanoparticles from leaf extract of Terminalia arjuna, for the enhanced mitotic cell division and pollen germination activity, *Ind. Crop. Prod.*, 2013, **50**, 737–742.
- 161 N. Suganthi, V. S. Ramkumar and A. Pugazhendhi, Biogenic synthesis of gold nanoparticles from Terminalia arjuna bark extract: assessment of safety aspects and neuroprotective potential via antioxidant, anticholinesterase, and antiamyloidogenic effects, *Environ. Sci. Pollut. Res.*, 2018, **25**, 10418–10433.
- 162 T. Akther, S. Priya, S. K. Sah, M. S. Khan and S. Hemalatha, Ta-AgNps are Potential Antimicrobial Resistance Breakers, *J. Nanostructure*, 2019, **9**, 376–383.
- 163 S. B. Koparde and D. K. Gaikwad, Antibacterial Potential of Green Silver Nanoparticles Synthesized from Medicinal Plant Terminalia arjuna, *Int. J. Res. Biosci. Agric. Technol.*, 2017, 258–260.
- 164 S. Yallappa and J. Manjanna, Biological Evaluation of Silver Nanoparticles Obtained from T. arjuna Bark Extract as Both Reducing and Capping Agent, *J. Clust. Sci.*, 2014, 1449–1462.
- 165 S. Ahmed and S. Ikram, Silver Nanoparticles: One Pot Green Synthesis Using Terminalia arjuna Extract for



- Biological Application, *J. Nanomed. Nanotechnol.*, 2015, **6**(4), 1–6.
- 166 C. Garai, S. Nurul, H. Abir, C. Barai, S. Ghorai, S. Kumar, P. Braja and G. Bag, Green synthesis of Terminalia arjuna - conjugated palladium nanoparticles (TA - PdNPs) and its catalytic applications, *J. Nanostructure Chem.*, 2018, **8**, 465–472.
- 167 D. Gaikwad and N. Jadhav, Terminalia arjuna Transdermal Matrix Formulation Containing Different Polymer Components, *Asian J. Pharm. Clin. Res.*, 2019, **12**, 1–6.
- 168 S. Noreen, S. A. Ghumman, F. Batool, B. Ijaz, M. Basharat, S. Noureen, T. Kausar and S. Iqbal, Terminalia arjuna gum/alginate *in situ* gel system with prolonged retention time for ophthalmic drug delivery, *Int. J. Biol. Macromol.*, 2020, **152**(4), 1056–1067.
- 169 F. E. King, J. R. Housley and T. J. King, The Chemistry of Extractives from Hardwoods, *J. Chem. Soc.*, 1954, **16**, 1392–1399.
- 170 K. Aamir, H. Ullah, C. Faiz, M. Rejina, P. Regina, I. Waheed, G. Sethi and A. Arya, Arjunolic acid downregulates elevated blood sugar and pro-inflammatory cytokines in streptozotocin (STZ) -nicotinamide induced type 2 diabetic rats, *Life Sci.*, 2022, **289**, 120232.
- 171 M. Sumitra, P. Manikandan, D. A. Kumar, N. Arutselvan, K. Balakrishna, B. M. Manohar and R. Puvanakrishnan, Experimental myocardial necrosis in rats: Role of arjunolic acid on platelet aggregation, coagulation and antioxidant status, *Mol. Cell. Biochem.*, 2001, **224**, 135–142.
- 172 E. Toppo, S. Sylvester Darvin, S. Esakkimuthu, K. Buvanavaragurunathan, T. P. Ajeesh Krishna, S. Antony Caesar, A. Stalin, K. Balakrishna, P. Pandikumar, S. Ignacimuthu and N. A. Al-Dhabi, Curative effect of arjunolic acid from Terminalia arjuna in non-alcoholic fatty liver disease models, *Biomed. Pharmacother.*, 2018, **107**, 979–988.
- 173 S. Khatkar, A. Nanda and S. H. Ansari, Microwave assisted extraction, optimization using central composite design, quantitative estimation of arjunolic acid and arjunolic acid using HPTLC and evaluation of radical scavenging potential of stem bark of Terminalia arjuna, *Nat. Prod. Sci.*, 2017, **23**, 75–83.
- 174 S. C. Verma, C. L. Jain, M. M. Padhi and R. B. Devalla, Microwave extraction and rapid isolation of arjunolic acid from Terminalia arjuna (Roxb. ex DC.) stem bark and quantification of arjunolic acid and arjunolic acid using HPLC-PDA technique, *J. Sep. Sci.*, 2012, **35**, 1627–1633.
- 175 J. Ghosh, J. Das, P. Manna and P. C. Sil, Arjunolic acid, a triterpenoid saponin, prevents acetaminophen (APAP)-induced liver and hepatocyte injury *via* the inhibition of APAP bioactivation and JNK-mediated mitochondrial protection, *Free Radic. Biol. Med.*, 2010, **48**, 535–553.
- 176 M. Bhujel, G. Nageswara Rao and L. Sripada, *Development of an efficient chromatographic method for the extraction and separation of arjunolic acid and asiatic acid from the heartwood of Terminalia arjuna*, 2024.
- 177 B. Ma, S. Yang, J. Li, H. Ouyang, M. He, Y. Feng and T. Tan, A four-step filtering strategy based on ultra-high-performance liquid chromatography coupled to quadrupole-time-of-flight tandem mass spectrometry for comprehensive profiling the major chemical constituents of Akebiae fructus, *Rapid Commun. Mass Spectrom.*, 2019, **33**, 1464–1474.
- 178 M. Shimada, M. Ozawa, K. Iwamoto, Y. Fukuyama, A. Kishida and A. Ohsaki, A lanostane triterpenoid and three cholestane sterols from Tilia kiusiana, *Chem. Pharm. Bull.*, 2014, **62**, 937–941.
- 179 D. Pertuit, A. C. Mitaine-Offer, T. Miyamoto, C. Tanaka, S. Delemasure, P. Dutartre and M. A. Lacaille-Dubois, A new aromatic compound from the stem bark of Terminalia catappa, *Nat. Prod. Commun.*, 2015, **10**, 1005–1007.
- 180 L. Wang, M. Wu, J. Wang, J. Jiang, Y. Chen and T. Li, A new chalcone glycoside from Combretum griffithii, *Chem. Nat. Compd.*, 2014, **50**, 258–260.
- 181 K. W. Woo, J. M. Cha, S. U. Choi and K. R. Lee, A new triterpene glycoside from the stems of Lagerstroemia indica, *Arch. Pharm. Res.*, 2016, **39**, 631–635.
- 182 F. X. Zhang, M. Li, Z. H. Yao, C. Li, L. R. Qiao, X. Y. Shen, K. Yu, Y. Dai and X. S. Yao, A target and nontarget strategy for identification or characterization of the chemical ingredients in Chinese herb preparation Shuang-Huang-Lian oral liquid by ultra-performance liquid chromatography–quadrupole time-of-flight mass spectrometry, *Biomed. Chromatogr.*, 2018, **32**(3), e4110.
- 183 D. Zou, J. Wang, B. Zhang, S. Xie, Q. Wang, K. Xu, R. Lin and C. W. K. Lam, Analysis of chemical constituents in Wuzi-Yanzong-Wan by UPLC-ESI-LTQ-Orbitrap-MS, *Molecules*, 2015, **20**, 21373–21404.
- 184 D. P. A. Gossan, A. Alabdul Magid, P. A. Yao-Kouassi, J. Josse, S. C. Gangloff, H. Morjani and L. Voutquenne-Nazabadioko, Antibacterial and cytotoxic triterpenoids from the roots of Combretum racemosum, *Fitoterapia*, 2016, **110**, 89–95.
- 185 J. Wang, H. Ren, Q. L. Xu, Z. Y. Zhou, P. Wu, X. Y. Wei, Y. Cao, X. X. Chen and J. W. Tan, Antibacterial oleanane-type triterpenoids from pericarps of Akebia trifoliata, *Food Chem.*, 2015, **168**, 623–629.
- 186 J. Zhang, Y. Li, S. S. Chen, L. Zhang, J. Wang, Y. Yang, S. Zhang, Y. Pan, Y. Wang and L. Yang, Systems pharmacology dissection of the anti-inflammatory mechanism for the medicinal herb Folium Eriobotryae, *Int. J. Mol. Sci.*, 2015, **16**, 2913–2941.
- 187 B. Zhai, J. Clark, T. Ling, M. Connelly, F. Medina-Bolivar and F. Rivas, Antimalarial evaluation of the chemical constituents of hairy root culture of Bixa orellana L, *Molecules*, 2014, **19**, 756–766.
- 188 Y. Mai, Z. Wang, Y. Wang, J. Xu and X. He, Anti-neuroinflammatory triterpenoids from the seeds of Quercus serrata Thunb, *Fitoterapia*, 2020, **142**, 104523.
- 189 D. Alesiani, A. Canini, B. D'Ambrosca, M. DellaGreca, A. Fiorentino, C. Mastellone, P. Monaco and S. Pacifico, Antioxidant and antiproliferative activities of phytochemicals from Quince (Cydonia vulgaris) peels, *Food Chem.*, 2010, **118**, 199–207.



- 190 D. H. Nguyen, D. D. Le, B. T. Zhao, E. S. Ma, B. S. Min and M. H. Woo, Antioxidant compounds isolated from the roots of *Phlomis umbrosa* Turcz, *Nat. Prod. Sci.*, 2018, **24**, 119–124.
- 191 H. Yang, E. J. Jeong, J. Kim, S. H. Sung and Y. C. Kim, Antiproliferative triterpenes from the leaves and twigs of *Juglans sinensis* on HSC-T6 cells, *J. Nat. Prod.*, 2011, **74**, 751–756.
- 192 C. Wang, P. Wu, S. Tian, J. Xue, L. Xu, H. Li and X. Wei, Bioactive Pentacyclic Triterpenoids from the Leaves of *Cleistocalyx operculatus*, *J. Nat. Prod.*, 2016, **79**, 2912–2923.
- 193 L. Ba Vinh, H. J. Jang, N. Viet Phong, G. Dan, K. Won Cho, Y. Ho Kim and S. Young Yang, Bioactive triterpene glycosides from the fruit of *Stauntonia hexaphylla* and insights into the molecular mechanism of its inflammatory effects, *Bioorganic Med. Chem. Lett.*, 2019, **29**, 2085–2089.
- 194 M. A. Baldé, E. Tuentler, A. Matheussen, M. S. Traoré, P. Cos, L. Maes, A. Camara, M. S. T. Diallo, E. S. Baldé, A. M. Balde, L. Pieters and K. Foubert, Bioassay-guided isolation of antiplasmodial and antimicrobial constituents from the roots of *Terminalia albida*, *J. Ethnopharmacol.*, 2021, **267**, 113624.
- 195 I. Zebiri, M. Haddad, L. Duca, M. Sauvain, L. Paloque, B. Cabanillas, E. Rengifo, J. B. Behr and L. Voutquenne-Nazabadioko, Biological activities of triterpenoids from *Poraqueiba sericea* stems, *Nat. Prod. Res.*, 2017, **31**, 1333–1338.
- 196 P. Masoko, L. K. Mdee, L. J. Mampuru and J. N. Eloff, Biological activity of two related triterpenes isolated from *Combretum nelsonii* (Combretaceae) leaves, *Nat. Prod. Res.*, 2008, **22**, 1074–1084.
- 197 Y. C. Ting, H. H. Ko, H. C. Wang, C. F. Peng, H. S. Chang, P. C. Hsieh and I. S. Chen, Biological evaluation of secondary metabolites from the roots of *Myrica adenophora*, *Phytochemistry*, 2014, **103**, 89–98.
- 198 R. de Cássia Lemos Lima, L. Kato, K. T. Kongstad and D. Staerk, Brazilian insulin plant as a bifunctional food: Dual high-resolution PTP1B and α -glucosidase inhibition profiling combined with HPLC-HRMS-SPE-NMR for identification of antidiabetic compounds in *Myrcia rubella* Cambess, *J. Funct. Foods*, 2018, **45**, 444–451.
- 199 H. E. Elsayed, M. R. Akl, H. Y. Ebrahim, A. A. Sallam, E. G. Haggag, A. M. Kamal and K. A. El Sayed, Discovery, optimization, and pharmacophore modeling of oleanolic acid and analogues as breast cancer cell migration and invasion inhibitors through targeting Brk/Paxillin/Rac1 axis, *Chem. Biol. Drug Des.*, 2015, **85**, 231–243.
- 200 S. El Abdellaoui, E. Destandau, I. Kroliekiewicz-Renimel, P. Cancellieri, A. Toribio, V. Jeronimo-Monteiro, L. Landemarre, P. André and C. Elfakir, Centrifugal partition chromatography for antibacterial bio-guided fractionation of *Clidemia hirta* roots, *Sep. Purif. Technol.*, 2014, **123**, 221–228.
- 201 Y. Li, J. Xu, D. Li, H. Ma, Y. Mu, D. Zheng, X. Huang and L. Li, Chemical Characterization and Hepatoprotective Effects of a Standardized Triterpenoid-Enriched Guava Leaf Extract, *J. Agric. Food Chem.*, 2021, **69**, 3626–3637.
- 202 Y.-H. Kuo and S.-L. Lin, Chemical Components of the Whole Herb of *Mosla dianthera*, *Chem. Pharm. Bull.*, 1999, **47**, 1152–1153.
- 203 Y. Li, J. Xu, C. Yuan, H. Ma, T. Liu, F. Liu, N. P. Seeram, Y. Mu, X. Huang and L. Li, Chemical composition and anti-hyperglycaemic effects of triterpenoid enriched *Eugenia jambolana* Lam. berry extract, *J. Funct. Foods*, 2017, **28**, 1–10.
- 204 J. M. Tian, X. Y. Fu, Q. Zhang, H. P. He, J. M. Gao and X. J. Hao, Chemical constituents from *Glochidion assamicum*, *Biochem. Syst. Ecol.*, 2013, **48**, 288–292.
- 205 C. H. Zhang, J. Luo, T. Li, Y. Cui, M. Jin, D. L. Yao, M. S. Zheng, Z. H. Lin, J. M. Cui and G. Li, Chemical constituents from the aerial parts of *Melandrium firmum*, *Arch. Pharm. Res.*, 2015, **38**, 1746–1751.
- 206 W. Auamcharoen, A. Chandrapatya, A. Kijjoa, A. M. S. Silva and W. Herz, Chemical constituents of *Duabanga grandiflora* (Lythraceae), *Biochem. Syst. Ecol.*, 2009, **37**, 535–537.
- 207 W. Li, W. Zhou, S. H. Shim and Y. H. Kim, Chemical constituents of the rhizomes and roots of *Gentiana scabra* (Gentianaceae), *Biochem. Syst. Ecol.*, 2015, **61**, 169–174.
- 208 Y. Cao, S. Fang, Z. Yin, X. Fu, X. Shang, W. Yang and H. Yang, Chemical fingerprint and multicomponent quantitative analysis for the quality evaluation of *Cyclocarya paliurus* leaves by HPLC-Q-TOF-MS, *Molecules*, 2017, **22**, 1–16.
- 209 H. Dou, R. Zhang, X. Lou, J. Jia, C. Zhou and Y. Zhao, Constituents of three species of *Lagerstroemia*, *Biochem. Syst. Ecol.*, 2005, **33**, 639–642.
- 210 T. R. Vieira, L. C. A. Barbosa, C. R. A. Maltha, V. F. Paula and E. A. Nascimento, Constituintes químicos de *Melaleuca alternifolia* (Myrtaceae), *Quim. Nova*, 2004, **27**, 536–539.
- 211 F. F. A. Fernandes, M. A. Tomaz, C. Z. El-Kik, M. Monteiro-Machado, M. A. Strauch, B. L. Cons, M. S. Tavares-Henriques, A. C. O. Cintra, V. A. Facundo and P. A. Melo, Counteraction of Bothrops snake venoms by *Combretum leprosum* root extract and arjunolic acid, *J. Ethnopharmacol.*, 2014, **155**, 552–562.
- 212 P. C. Tu, Y. C. Liang, G. J. Huang, M. K. Lin, M. C. Kao, T. L. Lu, P. J. Sung and Y. H. Kuo, Cytotoxic and Anti-inflammatory Terpenoids from the Whole Plant of *Vaccinium emarginatum*, *Planta Med.*, 2020, **86**, 1313–1322.
- 213 Y. Zhou, B. Yang, Z. Liu, Y. Jiang, Y. Liu, L. Fu, X. Wang and H. Kuang, Cytotoxicity of triterpenes from green walnut husks of *Juglans mandshurica* Maxim in HepG-2 cancer cells, *Molecules*, 2015, **20**, 19252–19262.
- 214 R. S. Pawar, P. Balachandran, D. S. Pasco and I. A. Khan, Cytotoxicity studies of triterpenoids from *Akebia trifoliata* and *Clematis ligusticifolia*, *Acta Hort.*, 2006, **720**, 171–178.
- 215 M. B. C. Gallo, A. S. F. Marques, P. C. Vieira, M. F. D. G. F. da Silva, J. B. Fernandes, M. Silva, R. V. Guido, G. Oliva, O. H. Thiemann, S. Albuquerque and A. H. Fairlamb, Enzymatic Inhibitory Activity and Trypanocidal Effects of Extracts and Compounds from *Siphoneugena densiflora* O. Berg and *Vitex polygama* Cham, *Zeitschrift für Naturforsch. - Sect. C J. Biosci.*, 2008, **63**, 371–382.



- 216 J. Gominho, R. A. Costa, A. Lourenço, T. Quilhó and H. Pereira, Eucalyptus globulus Stumps Bark: Chemical and Anatomical Characterization Under a Valorisation Perspective, *Waste Biomass Valoriz.*, 2021, **12**, 1253–1265.
- 217 F. Hu, X. Liao, Y. Guo, S. Yamaki, X. Li, N. Hamada, Y. Hashi and Z. Chen, Fast determination of isomeric triterpenic acids in *Osmanthus fragrans* (Thunb.) Lour. fruits by UHPLC coupled with triple quadrupole mass spectrometry, *Food Chem.*, 2020, **322**, 126781.
- 218 Z. Zhang, H. N. ElSohly, X. C. Li, S. I. Khan, S. E. Broedel, R. E. Raulli, R. L. Cihlar and L. A. Walker, Flavanone glycosides from *Miconia trailii*, *J. Nat. Prod.*, 2003, **66**, 39–41.
- 219 M. Shao, Y. Wang, X. J. Huang, C. L. Fan, Q. W. Zhang, X. Q. Zhang and W. C. Ye, Four new triterpenoids from the leaves of *Psidium guajava*, *J. Asian Nat. Prod. Res.*, 2012, **14**, 348–354.
- 220 P. Masoko, J. Picard, R. L. Howard, L. J. Mampuru and J. N. Eloff, In vivo antifungal effect of *Combretum* and *Terminalia* species extracts on cutaneous wound healing in immunosuppressed rats, *Pharm. Biol.*, 2010, **48**, 621–632.
- 221 G. Zengin, C. Ferrante, I. Senkardes, R. Gevrenova, D. Zheleva-Dimitrova, L. Menghini, G. Orlando, L. Recinella, A. Chiavaroli, S. Leone, L. Brunetti, C. M. N. Picot-Allain, K. R. Rengasamy and M. F. Mahomoodally, Multidirectional biological investigation and phytochemical profile of *Rubus sanctus* and *Rubus ibericus*, *Food Chem. Toxicol.*, 2019, **127**, 237–250.
- 222 G. M. Woldemichael, M. T. Gutierrez-Lugo, S. G. Franzblau, Y. Wang, E. Suarez and B. N. Timmermann, Mycobacterium tuberculosis Growth Inhibition by Constituents of *Sapium haematospermum*, *J. Nat. Prod.*, 2004, **67**, 598–603.
- 223 S. Parveen, M. Saleem, N. Riaz, M. Ashraf, Q. Ul-Ain, M. F. Nisar and A. Jabbar, New norterpenoids and a sphingolipid from *Carissa opaca*, *J. Asian Nat. Prod. Res.*, 2016, **18**, 222–231.
- 224 W. D. Tékapı Tsopgni, A. G. Blaise Azebaze, J. E. Mbosso Teinkela, B. L. Ndjakou, F. F. Bosityom, B. N. Tchaleu and J. C. Vardamides, New unsaturated fatty acid and other chemical constituents from the roots of *Cola rostrata* K. Schum. (Malvaceae), *Biochem. Syst. Ecol.*, 2019, **86**, DOI: [10.1016/j.bse.2019.103913](https://doi.org/10.1016/j.bse.2019.103913).
- 225 A. M. A. Mohammed, P. H. Coombes, N. R. Crouch and D. A. Mulholland, Non-volatile isolates of two *Heteropyxis* species: A first chemotaxonomic assessment of subfamily *Psiloxylloideae* (Myrtaceae), *Biochem. Syst. Ecol.*, 2009, **37**, 241–243.
- 226 J. Feng, X. Yi, W. Huang, Y. Wang and X. He, Novel triterpenoids and glycosides from durian exert pronounced anti-inflammatory activities, *Food Chem.*, 2018, **241**, 215–221.
- 227 A. Qiao, Y. Wang, L. Xiang, Z. Zhang and X. He, Novel triterpenoids isolated from hawthorn berries functioned as antioxidant and antiproliferative activities, *J. Funct. Foods*, 2015, **13**, 308–313.
- 228 J. S. Han, J. G. Kim, T. P. L. Le, Y. B. Cho, M. K. Lee and B. Y. Hwang, Pentacyclic triterpenes with nitric oxide inhibitory activity from *Potentilla chinensis*, *Bioorg. Chem.*, 2021, **108**, 104659.
- 229 M. Liu, D. R. Katerere, A. I. Gray and V. Seidel, Phytochemical and antifungal studies on *Terminalia mollis* and *Terminalia brachystemma*, *Fitoterapia*, 2009, **80**, 369–373.
- 230 C. Liang, D. Staerk and K. T. Kongstad, Potential of *Myrtus communis* Linn. as a bifunctional food: Dual high-resolution PTP1B and α -glucosidase inhibition profiling combined with HPLC-HRMS and NMR for identification of antidiabetic triterpenoids and phloroglucinol derivatives, *J. Funct. Foods*, 2020, **64**, DOI: [10.1016/j.jff.2019.103623](https://doi.org/10.1016/j.jff.2019.103623).
- 231 B. Y. Yang, X. Yin, Y. Liu, Y. Sun, W. Guan, Y. Y. Zhou and H. X. Kuang, Terpenes and lignans from the roots of *Solanum melongena* L., *Nat. Prod. Res.*, 2020, **34**, 359–368.
- 232 A. Lourenço, A. V. Marques and J. Gominho, The identification of new triterpenoids in *Eucalyptus globulus* wood, *Molecules*, 2021, **26**, 1–13.
- 233 B. Diallo, R. Vanhaelen-Fastré, M. Vanhaelen, T. Konoshima, M. Takasaki and H. Tokuda, In vivo inhibitory effects of arjunolic acid derivatives on two-stage carcinogenesis in mouse skin, *Phyther. Res.*, 1995, **9**(6), 444–447.
- 234 I. L. Acebey-Castellon, L. Voutquenne-Nazabadioko, H. Doan Thi Mai, N. Roseau, N. Bouthagane, D. Muhammad, E. Le Magrex Debar, S. C. Gangloff, M. Litaudon, T. Sevenet, N. Van Hung and C. Lavaud, Triterpenoid saponins from *Symplocos lancifolia*, *J. Nat. Prod.*, 2011, **74**, 163–168.
- 235 R. Srivastava, A. K. Shaw and D. K. Kulshreshtha, Triterpenoids and chalcone from *Syzygium samarangense*, *Phytochemistry*, 1995, **38**, 687–689.
- 236 M. Tene, P. Tane and J. D. Connolly, Triterpenoids and diarylheptanoids from *Myrica arborea*, *Biochem. Syst. Ecol.*, 2008, **36**, 872–874.
- 237 J. Li, C. M. Coleman, H. Wu, C. L. Burandt, D. Ferreira and J. K. Zjawiony, Triterpenoids and flavonoids from *Cecropia schreberiana* Miq., *Biochem. Syst. Ecol.*, 2013, **48**, 96–99.
- 238 Y. S. Wang, H. Zhang, K. W. Wang, T. T. Zhang and X. F. Fan, Triterpenoids and Steroids from *Premna fulva*, *Chem. Nat. Compd.*, 2020, **56**, 373–375.
- 239 J. Zhao, Triterpenoids from *Rhododendron hainanense*, *Chem. Nat. Compd.*, 2013, **49**, 467–469.
- 240 D. H. Kim, K. M. Han, I. S. Chung, D. K. Kim, S. H. Kim, B. M. Kwon, T. S. Jeong, M. H. Park, E. M. Ahn and N. I. Baek, Triterpenoids from the flower of *Campsis grandiflora* K. Schum. as human Acyl-CoA: Cholesterol acyltransferase inhibitors, *Arch. Pharm. Res.*, 2005, **28**, 550–556.
- 241 Z. Y. Huang, B. Huang, C. J. Xiao, X. Dong and B. Jiang, Two new labdane diterpenoids from the rhizomes of *Isodon yunnanensis*, *Nat. Prod. Res.*, 2015, **29**, 628–632.
- 242 R. Fan, R. R. Cheng, H. T. Zhu, D. Wang, C. R. Yang, M. Xu and Y. J. Zhang, Two new oleanane-type triterpenoids from



- methanolized saponins of *Momordica cochinchinensis*, *Nat. Prod. Commun.*, 2016, **11**, 725–728.
- 243 Y. B. Zhang, W. Li, Z. M. Zhang, N. H. Chen, X. Q. Zhang, J. W. Jiang, G. C. Wang and Y. L. Li, Two new triterpenoids from the roots of *rhodomirtus tomentosa*, *Chem. Lett.*, 2016, **45**, 368–370.
- 244 D. Yang, H. Xie, B. Yang and X. Wei, Two tetrahydroisoquinoline alkaloids from the fruit of *Averrhoa carambola*, *Phytochem. Lett.*, 2014, **7**, 217–220.
- 245 F. Ullah, H. Hussain, J. Hussain, I. A. Bukhari, M. Tareq, H. Khan, M. I. Choudhary, A. H. Gilani and V. U. Ahmad, Tyrosinase Inhibitory Pentacyclic Triterpenes and Analgesic and Spasmolytic Activities of Methanol Extracts of *Rhododendron collettianum*, *Phyther. Res.*, 2008, **21**, 1076–1081.
- 246 M. Ahmad, N. Muhammad, M. Ahmad, M. A. Lodhi, N. Jehan, Z. Khan, R. Ranjit, F. Shaheen and M. I. Choudhary, Urease inhibitor from *Datisca cannabina* Linn, *J. Enzyme Inhib. Med. Chem.*, 2008, **23**, 386–390.
- 247 R. S. Bhakuni, Y. N. Shukla, A. K. Tripathi, V. Prajapati and S. Kumar, Insect growth inhibitor activity of arjunolic acid isolated from *Cornus capitata*, *Phyther. Res.*, 2002, **16**, 2000–2002.
- 248 E. Toppo, S. Sylvester Darwin, S. Esakkimuthu, K. Buvanavaragurunathan, T. P. Ajeesh Krishna, S. Antony Caesar, A. Stalin, K. Balakrishna, P. Pandikumar, S. Ignacimuthu and N. A. Al-Dhabi, Curative effect of arjunolic acid from *Terminalia arjuna* in non-alcoholic fatty liver disease models, *Biomed. Pharmacother.*, 2018, **107**, 979–988.
- 249 P. Manna, M. Sinha and P. C. Sil, Protection of arsenic-induced hepatic disorder by arjunolic acid, *Basic Clin. Pharmacol. Toxicol.*, 2007, **101**, 333–338.
- 250 B. G. Bag, P. P. Dey, S. K. Dinda, W. S. Sheldrick and I. M. Oppel, A simple route for renewable nano-sized arjunolic and asiatic acids and self-assembly of arjuna-bromolactone, *Beilstein J. Org. Chem.*, 2008, **4**, 2–6.
- 251 J. Ghosh, J. Das, P. Manna and P. C. Sil, The protective role of arjunolic acid against doxorubicin induced intracellular ROS dependent JNK-p38 and p53-mediated cardiac apoptosis, *Biomaterials*, 2011, **32**, 4857–4866.
- 252 T. Bansal, E. Chatterjee, J. Singh, A. Ray, B. Kundu, V. Thankamani, S. Sengupta and S. Sarkar, Arjunolic acid, a peroxisome proliferator-activated receptor agonist, regresses cardiac fibrosis by inhibiting non-canonical TGF- signaling, *J. Biol. Chem.*, 2017, **292**, 16440–16462.
- 253 P. Manna, M. Sinha and P. C. Sil, Arsenic-induced oxidative myocardial injury: Protective role of arjunolic acid, *Arch. Toxicol.*, 2008, **82**, 137–149.
- 254 M. M. Hasan, P. Madhavan, N. A. Ahmad Noruddin, W. K. Lau, Q. U. Ahmed, A. Arya and Z. A. Zakaria, Cardioprotective effects of arjunolic acid in LPS-stimulated H9C2 and C2C12 myotubes via the My88-dependent TLR4 signaling pathway, *Pharm. Biol.*, 2023, **61**, 1135–1151.
- 255 M. M. H. Al-Gayyar, A. Al Youssef, I. O. Sherif, M. E. E. Shams and A. Abbas, Protective effects of arjunolic acid against cardiac toxicity induced by oral sodium nitrite: Effects on cytokine balance and apoptosis, *Life Sci.*, 2014, **111**, 18–26.
- 256 H. Elsayy, M. Almallki, O. Elmenshawy and A. Abdel-Moneim, In vivo evaluation of the protective effects of arjunolic acid against lipopolysaccharide-induced septic myocardial injury, *PeerJ*, 2022, **10**, 1–16.
- 257 A. S. Ramesh, J. G. Christopher, R. Radhika, C. R. Setty and V. Thankamani, Isolation, characterisation and cytotoxicity study of arjunolic acid from *Terminalia arjuna*, *Nat. Prod. Res.*, 2012, **26**, 1549–1552.
- 258 S. Manna, A. Dey, R. Majumdar, B. G. Bag, C. Ghosh and S. Roy, Self assembled arjunolic acid acts as a smart weapon against cancer through TNF- α mediated ROS generation, *Heliyon*, 2020, **6**, 1–9.
- 259 N. M. Elsherbiny and M. M. H. Al-gayyar, Anti-tumor activity of arjunolic acid against Ehrlich Ascites Carcinoma cells *in vivo* and *in vitro* through blocking TGF- b type 1 receptor, *Biomed. Pharmacother.*, 2016, **82**, 28–34.
- 260 P. Manna, M. Sinha and P. C. Sil, Protective role of arjunolic acid in response to streptozotocin-induced type-I diabetes via the mitochondrial dependent and independent pathways, *Toxicology*, 2009, **257**, 53–63.
- 261 P. Manna, J. Das, J. Ghosh and P. C. Sil, Contribution of type 1 diabetes to rat liver dysfunction and cellular damage via activation of NOS, PARP, I κ B α /NF- κ B, MAPKs, and mitochondria-dependent pathways: Prophylactic role of arjunolic acid, *Free Radic. Biol. Med.*, 2010, **48**, 1465–1484.
- 262 L. M. Goncalves-Neto, F. B. Ferreira, L. Souza, C. dos Santos, A. C. Boschero, V. A. Facundo, A. R. S. Santos, E. A. Nunes and A. Rafacho, Disruption of glucose tolerance caused by glucocorticoid excess in rats is partially prevented, but not attenuated, by arjunolic acid, *Indian J. Experimental Biol.*, 2014, **52**, 972–982.
- 263 C. Fang, Y. Teng, Y. Wang, Y. Zhao and X. Zheng, Arjunolic acid from *Cyclocarya paliurus* selectively inhibits glucagon secretion from α cells and ameliorates diabetes via ephrin-A1 and EphA4 interaction, *J. Funct. Foods*, 2022, **99**, 105323.
- 264 X. Zhang, Y. Ji, L. Zhu, Z. Wang and C. Fang, Arjunolic acid from *Cyclocarya paliurus* ameliorates diabetic retinopathy through AMPK/mTOR/HO-1 regulated autophagy pathway, *J. Ethnopharmacol.*, 2022, **284**, 1–15.
- 265 K. Aamir, G. Sethi, M. Rejina, C. Faiz, P. Regina, S. D. Sarker and A. Arya, Arjunolic acid modulate pancreatic dysfunction by ameliorating pattern recognition receptor and canonical Wnt pathway activation in type 2 diabetic rats, *Life Sci.*, 2023, **327**, 121856.
- 266 S. R. A, G. Christopher and V. Thankamani, In-vitro study on antioxidant property of Arjunolic acid, *J. Pharmacogn. Herb. Formul.*, 2010, **1**, 1–11.
- 267 M. Maity, H. Perveen, M. Dash, S. Jana, S. Khatun, A. Dey and A. K. Mandal, Arjunolic Acid Improves the Serum Level of Vitamin B 12 and Folate in the Process of the



- Attenuation of Arsenic Induced Uterine Oxidative Stress, *Biol. Trace Elem. Res.*, 2018, **182**, 78–90.
- 268 M. Sinha, P. Manna and P. C. Sil, Protective Effect of Arjunolic Acid against Arsenic-Induced Oxidative Stress in Mouse Brain, *J. Biochem Mol. Toxicol.*, 2008, **22**, 15–26.
- 269 P. Manna, M. Sinha and P. C. Sil, Protection of arsenic-induced testicular oxidative stress by arjunolic acid, *Redox Rep.*, 2008, **13**, 67–77.
- 270 L. Yaidikar and S. Thakur, Arjunolic acid, a pentacyclic triterpenoidal saponin of Terminalia arjuna bark protects neurons from oxidative stress associated damage in focal cerebral ischemia and reperfusion, *Pharmacol. Reports*, 2015, **67**, 890–895.
- 271 S. Miriyala, M. Chandra, B. Maxey, A. Day, D. K. St and M. Panchatcharam, Arjunolic acid ameliorates reactive oxygen species *via* inhibition of p47 phox -serine phosphorylation and mitochondrial dysfunction, *Int. J. Biochem. Cell Biol.*, 2015, **68**, 70–77.
- 272 P. Manna, J. Ghosh, J. Das and P. C. Sil, Streptozotocin induced activation of oxidative stress responsive splenic cell signaling pathways : Protective role of arjunolic acid, *Toxicol. Appl. Pharmacol.*, 2010, **244**, 114–129.
- 273 S. Pal, A. Sarkar, P. B. Pal and P. C. Sil, Protective effect of arjunolic acid against atorvastatin induced hepatic and renal pathophysiology *via* MAPK, mitochondria and ER dependent pathways, *Biochimie*, 2015, **112**, 20–34.
- 274 I. O. Sherif, Hepatoprotective effect of arjunolic acid against cisplatin-induced hepatotoxicity: Targeting oxidative stress, inflammation, and apoptosis, *J. Biochem. Mol. Toxicol.*, 2021, **35**, 1–8.
- 275 X. Zheng, X. G. Zhang, Y. Liu, L. P. Zhu, X. S. Liang, H. Jiang, G. F. Shi, Y. Y. Zhao, Z. W. Zhao, Y. Teng, K. Pan, J. Zhang and Z. Q. Yin, Arjunolic acid from Cyclocarya paliurus ameliorates nonalcoholic fatty liver disease in mice *via* activating Sirt1/AMPK, triggering autophagy and improving gut barrier function, *J. Funct. Foods*, 2021, **86**, 104686.
- 276 Y. Fan, C. Zhang, G. Zheng, S. Wu, Y. Wang and J. Bian, Ameliorative and anti-arthritis potential of arjunolic acid against complete Freund's adjuvant-induced arthritis in rats, *Trop. J. Pharm. Res.*, 2020, **19**, 1933–1939.
- 277 A. Alyoussef, Arjunolic acid protects against DNCB-induced atopic dermatitis-like symptoms in mice by restoring a normal cytokine balance, *Eur. Cytokine Netw.*, 2015, **26**, 38–45.
- 278 Z. Zhang, L. Zuo, X. Song, L. Wang, Y. Zhang, Y. Cheng, J. Huang, T. Zhao, Z. Yang, H. Zhang, J. Li, X. Zhang, Z. Geng, Y. Wang, S. Ge and J. Hu, Arjunolic acid protects the intestinal epithelial barrier, ameliorating Crohn's disease-like colitis by restoring gut microbiota composition and inactivating TLR4 signalling, *Phytomedicine*, 2024, **123**, 155223.
- 279 E. O. Lynda, N. E. Kingsley, O. M. Obukohwo, B. A. Benneth, E. Victor, O. I. Simon, E. Agbonifo-Chijiokwu and O. B. Oghenetega, Arjunolic acid reverses fluoxetine-induced alterations in testicular steroidogenic enzymes and membrane bound ionic pump imbalance through suppression of oxido-inflammatory stress and apoptosis, *J. Bras. Reprod. Assist.*, 2024, **28**, 66–77.
- 280 Y. Yang, Y. Lai, X. Tong, Z. Li, Y. Cheng and L. Tian, Arjunolic acid ameliorates lipopolysaccharide-induced depressive behavior by inhibiting neuroinflammation *via* microglial SIRT1/AMPK/Notch1 signaling pathway, *J. Ethnopharmacol.*, 2024, **330**, 118225.
- 281 M. V. V. Prasad, N. Anbalagan, A. Patra, G. Veluchamy and K. Balakrishna, Antiallergic and anti-asthmatic activities of the alcoholic extract of Terminalia arjuna and arjunolic acid, *Nat. Prod. Sci.*, 2004, **10**, 240–243.
- 282 J. D. Djoukeng, E. Abou-Mansour, R. Tabacchi, A. L. Tapondjou, H. Bouda and D. Lontsi, Antibacterial triterpenes from Syzygium guineense (Myrtaceae), *J. Ethnopharmacol.*, 2005, **101**, 283–286.
- 283 M. Yasasve, P. M. Kumaran, D. Manoj, M. Kanmani and A. S. Ramesh, Evaluation of arjunolic acid against Brucella melitenis and *in vitro* cytotoxic study of lung adenocarcinomic cell line (A549), *Indian J. Exp. Biol.*, 2022, **60**, 510–513.
- 284 Q. Zhang, H. N. ElSohly, X. C. Li and L. A. Walker, A new triterpene from Leandra chaetodon, *Planta Med.*, 2003, **69**, 582–584.
- 285 B. G. Bag, G. C. Maity and S. R. Pramanik, A terpenoid-based gelator: The first arjunolic acid-derived organogelator for alcohols and mixed solvents, *Supramol. Chem.*, 2005, **17**, 383–385.
- 286 B. G. Bag and R. Majumdar, Vesicular self-assembly of a natural triterpenoid arjunolic acid in aqueous medium: study of entrapment properties and *in situ* generation of gel-gold nanoparticle hybrid material, *RSC Adv.*, 2015, **95**, 53327–53334.
- 287 R. Majumdar and B. G. Bag, Evolution of Vesicular Self-Assemblies of the Salts of a Natural Triterpenoid Arjunolic Acid into Superstructured Ambidextrous Gels and Study of Their Entrapment Properties, *ChemistrySelect*, 2018, **3**, 951–957.
- 288 B. M. F. Gonçalves, J. A. R. Salvador, S. Marín and M. Cascante, Synthesis and biological evaluation of novel asiatic acid derivatives with anticancer activity, *RSC Adv.*, 2016, **6**, 3967–3985.
- 289 K. N. Devaraj Reddy, M. P. Sadashiva, M. Mahesh, B. K. Bettadaiah and N. P. Geetha, HPLC method for determination of acetylated arjunolic acid-a derivative of arjunolic acid from Terminalia arjuna and their antioxidant activity, *Int. J. Res. Phytochem. Pharmacol.*, 2012, **2**, 188–193.
- 290 B. G. Bag, S. R. Pramanik and G. C. Maity, Arjunolic acid in molecular recognition: First synthesis and cation binding studies of a novel arjuna-18-crown-6, *Supramol. Chem.*, 2005, **17**, 297–302.
- 291 B. G. Bag and S. K. Dinda, Arjunolic acid: A renewable template in supramolecular chemistry and nanoscience, *Pure Appl. Chem.*, 2007, **79**, 2031–2038.
- 292 B. G. Bag, R. Majumdar, S. K. Dinda, P. P. Dey, G. C. Maity, V. A. Mallia and R. G. Weiss, Self-assembly of ketals of



- arjunolic acid into vesicles and fibers yielding gel-like dispersions, *Langmuir*, 2013, **29**, 1766–1778.
- 293 B. Diallo, M. Vanhaelen, R. Vanhaelen-Fastre, T. Konoshima and M. Kozuka, Studies on inhibitors of skin tumour promotion. Inhibitory effects of triterpenes from *Cochlospermum tznctorzum* on Epstein-Barr virus activation., *J. Nat. Prod.*, 1989, **52**, 879–881.
- 294 B. Diallo, R. Vanhaelen-Fastré, M. Vanhaelen, T. Konoshima, M. Takasaki and H. Tokuda, In vivo inhibitory effects of arjunolic acid derivatives on two-stage carcinogenesis in mouse skin, *Phyther. Res.*, 1995, **9**, 444–447.

