



Cite this: DOI: 10.1039/d5sc05588a

 All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 25th July 2025

Accepted 2nd May 2026

DOI: 10.1039/d5sc05588a

rsc.li/chemical-science

Peptide modification *via* a mild carbonylative Suzuki–Miyaura reaction gives late-stage access to diaryl-ketones

James P. Kleppen,^{†ab} Neil W. J. Scott,^{†ab} Tom J. Phillips,^{ab}
Ksenia S. Stankevich,^{ab} James R. Donald,^{ab} Lydia J. Barber,^{abc} Katie M. Lewis,^{ab}
Ian J. S. Fairlamb^{†a} and Christopher D. Spicer^{†*ab}

Benzophenones and other diaryl ketones are common handles for photo-crosslinking and photo-affinity labelling. Here, we introduce the use of an operationally simple carbonylative Suzuki–Miyaura cross-coupling for the late-stage introduction of these handles into unprotected peptides. By using molybdenum hexacarbonyl, Mo(CO)₆, as a carbon monoxide surrogate, these reactions take place under mild aqueous conditions, under an open (air) atmosphere, requiring no specialist equipment or techniques.

Diaryl-ketones, typified most commonly by benzophenones, are an important molecular handle for biomedical research. Upon irradiation with UV light, these ketones are activated to form a highly reactive ‘C–O’ triplet diradical species, which can react with nearby biomolecules *via* a sequential abstraction–recombination process.^{1,2} Diaryl-ketones are therefore important reagents for photo-crosslinking and photo-affinity labelling, techniques which covalently trap interacting biomolecules, and allow the investigation of important processes in health and disease.³

In this context, benzophenone-containing peptides have proved invaluable in the study of diverse biological processes, ranging from protein-membrane interactions⁴ to post-translational modification.⁵ In almost all cases, the benzophenone motif is integrated into the peptide either by amide coupling at the ϵ -amine of lysine residues, or through the incorporation of the unnatural amino acid *p*-benzoyl-L-phenylalanine (Bpa) during solid phase peptide synthesis (SPPS).⁶ However, the photo-labelling efficiency of Bpa is often poor, due to its low absorptivity and quantum yield.⁷ Diversification of the diaryl-ketone motif can greatly increase this photo-labelling efficiency, while also offering potential routes to red-shift absorbance away from the damaging UV region and generate photo-orthogonal crosslinkers for multiplexed labelling.⁸ Indeed, the effects of aryl substituents on diaryl ketone photo-physical properties have been extensively studied,⁹ with Joiner *et al.* having demonstrated up to 50-fold enhancements in protein crosslinking using halogenated analogues of Bpa.⁷

This need for diversification requires the parallel synthesis of large numbers of diaryl-ketone containing peptides, with associated challenges in the synthesis and protection of diaryl-ketone based amino acids for SPPS. In contrast, a chemical strategy for late-stage installation of diaryl-ketones into unprotected peptides would be a more powerful and versatile tool for the biological and biomedical communities. However, to be viable, such a strategy should be efficient at ambient temperatures, near neutral pH, and in aqueous media. It would also need to be able to tolerate the myriad of reactive functional groups typically found in peptides. Finally, the reaction would benefit from being operational in an open atmosphere to maximise translatability to non-specialist labs and to facilitate small-scale reactions.

The development of such a strategy therefore represents a significant challenge, that has yet to be addressed. Indeed, these requirements rule out conventional synthetic routes towards diaryl-ketones, such as Friedel–Crafts acylations or carbonyl addition/oxidation reactions.^{9,10} While carbonylative Suzuki–Miyaura cross couplings (cSMCCs) provide a promising alternative, these reactions commonly require the use of highly toxic and flammable CO.¹¹ This can be limiting to the use of these reactions in general synthesis, a factor that is exacerbated for bioconjugation. There is therefore great interest in the use of CO-surrogates and CO delivery vehicles, including transition metal carbonyls, for cSMCC.¹² Indeed, in 2019 Sun *et al.* reported the use of Mo(CO)₆ and palladium(II) trifluoroacetate, ‘Pd(TFA)₂’, as a precatalyst for the cSMCC of aryl iodides in acetonitrile : water mixtures at 50 °C.¹³ Building on our previous experiences of translating Pd-mediated cross-couplings to peptide and protein substrates, we were drawn to the possibilities of this approach in bioconjugation.^{14–18}

^aDepartment of Chemistry, University of York, Heslington, YO10 5DD, UK. E-mail: chris.spicer@york.ac.uk

^bYork Biomedical Research Institute, University of York, Heslington, YO10 5DD, UK

^cDepartment of Biology, University of York, Heslington, YO10 5DD, UK

† These authors contributed equally and joint first authors.






Fig. 1 Overview of the carbonylative Suzuki–Miyaura cross-coupling (cSMCC) developed in this paper for the late-stage diversification of unprotected peptides.

In this paper, we report the development of operationally simple carbonylative Suzuki–Miyaura cross-couplings (cSMCCs) for peptide modification (Fig. 1). By using $\text{Mo}(\text{CO})_6$ as a carbon monoxide surrogate, these couplings take place at $< 37^\circ\text{C}$, under aqueous, ‘open-to-air’ conditions, and are therefore applicable to fully deprotected substrates. This has allowed us to demonstrate the selective late-stage formation of structurally and chemically diverse diaryl-ketones, for future applications in photo-labelling studies.


Results and discussion

We commenced our studies by investigating cSMCCs on the unnatural aryl iodide amino acid *para*-iodophenylalanine (*p*IPhe). *p*IPhe can be site-selectively incorporated into proteins *via* codon suppression, where we have previously shown it can act as a handle for Suzuki–Miyaura modification and the mimicry of natural post-translational modifications.^{14–16,19} Couplings were initially performed on Boc-*p*IPhe-OMe **1**, in the presence of 0.5 equiv. $\text{Mo}(\text{CO})_6$, 5 mol% ‘Pd(TFA)₂’, and 1.5 equiv. 4-methoxyphenyl boronic acid. Under these conditions, the carbonylated product **2** was preferentially delivered over the non-carbonylative side-product **3**. Notably, couplings proceeded readily at room temperature and in pH 8.0 phosphate buffer (Scheme 1a). Importantly, when no effort was made to degas the reagents or to ensure an inert atmosphere, little difference in conversion with respect to the starting material was observed. Analogous results were found for the modification of Boc-3-

iodotyrosine-OMe **4**, another protein-compatible iodinated amino acid²⁰ that also plays a key role in human biology as an intermediate during thyroid hormone synthesis.²¹ Again, cSMCC product **5** dominated (51% yield), though the non-carbonylative side-product **6** also formed in significant amounts (26% yield, Scheme 1b).


With these promising results in hand, we applied the cSMCC reaction conditions to a model iodinated peptide, $\text{H}_2\text{N-}p\text{IPhe-AEAE-CONH}_2$, **7**, using phenylboronic acid as a coupling partner. Pleasingly, conversion to the cSMCC product **8a** was observed by LC-MS, with the desired structure confirmed by ¹H NMR (see SI, Fig. S2) and comparison to an equivalent peptide synthesised *via* SPPS using Bpa directly. Further evidence for the presence of the diaryl ketone was provided by reduction to the corresponding alcohol upon treatment with sodium borohydride (See SI Section 5.12). We therefore set out to screen and optimise the reaction conditions for peptide modification (Fig. 2).

Under the conditions of the reaction, there are three possible pathways that can operate following initial oxidative addition of the aryl halide (see SI Fig. S1 for a depiction of the catalytic cycle): (i) when CO insertion is faster than transmetalation of the boronic acid, the desired cSMCC product **8a** can be formed; (ii) if transmetalation is faster than CO insertion, then the corresponding non-carbonylated Suzuki–Miyaura product **9a** will be formed; and (iii) in the presence of a nucleophile, in this case water, an acylpalladium–hydroxo complex may form which can ultimately lead to the corresponding carboxylic acid product, **10a**.^{22,23} Therefore, the relative conversions to these two



Scheme 1 Model cSMCCs on protected *p*IPhe and 3-ITyr substrates. Both reactions were undertaken ‘open-to-air’ at room temperature, with no attempt to exclude oxygen. Reaction conditions: $\text{Mo}(\text{CO})_6$ (0.5 equiv.), Pd(TFA)₂ (0.05 equiv.), boronic acid (1.5 equiv.), acetonitrile : PBS (pH 8, 5 : 1), rt, 20 h.





Entry	Pd(L) ₂	Equiv	CO source	Equiv	MeCN%	8a (%)	9a (%)	10a (%)
1	Pd(OPiv) ₂	7	Mo(CO) ₆	24	40	29	32	6
2	Pd(OAc) ₂	7	Mo(CO) ₆	24	40	65	35	0
3	Pd(TFA) ₂	7	Mo(CO) ₆	24	40	64	36	1
4	Pd(TFA) ₂	7	Mn ₂ (CO) ₁₀	24	40	28	57	0
5	Pd(TFA) ₂	7	Fe ₂ (CO) ₉	24	40	39	56	0
6	Pd(TFA) ₂	7	Cr(CO) ₆	24	40	0	57	0
7	Pd(TFA) ₂	0.4	Mo(CO) ₆	0.5	40	64	1	21
8	Pd(TFA) ₂	0.4	Mo(CO) ₆	2	40	65	1	25
9	Pd(TFA) ₂	0.4	Mo(CO) ₆	6	40	67	1	27
10	Pd(TFA) ₂	0.1	Mo(CO) ₆	2	40	53	1	17
11	Pd(TFA) ₂	0.8	Mo(CO) ₆	2	40	68	2	28
12	Pd(TFA) ₂	0.4	Mo(CO) ₆	2	0	10	89	0
13	Pd(TFA) ₂	0.4	Mo(CO) ₆	2	10	54	21	0

Fig. 2 Optimisation of cSMCC on the model peptide pI₁Phe-AAEE, 7.

by-products was monitored *via* LC-MS alongside the desired diaryl-ketone.

First, we screened alternative palladium precatalysts. 'Pd(TFA)₂', 'Pd(OAc)₂', and water soluble 2-amino-4,6-dihydropyrimidine (ADHP)-Pd complex^{14,24} showed comparable activity. In contrast, palladium(II) pivalate, 'Pd(OPiv)₂', gave significantly lower conversion (Entries 1–3, see SI Section 5.1 for further details). Of the metal carbonyl complexes screened, Mo(CO)₆ gave the highest conversion to **8a**, with both Mn₂(CO)₁₀ and Fe₂(CO)₉ giving lower cSMCC : SMCC ratios, and Cr(CO)₆ giving solely the SMCC product (Entries 4–6, SI Section 5.2). In the absence of a metal carbonyl complex, **9a** was formed exclusively.

Following this initial validation, we aimed to optimise the reagent loadings. Formation of the cSMCC product **8a** was found to be consistent across a range of 0.5–24 equiv. of Mo(CO)₆ with a significant drop-off at lower loadings (Entries 7–9, SI Section 5.3). Similarly, Pd loadings could be lowered to as little as 10 mol% (Entry 10, SI Section 5.4). However, at such small loadings of catalyst, the practical challenges of measuring such small quantities led to high variability in reaction efficiency, and so a loading of 40 mol% was used for further reactions. Although these loadings appear high, at the small scales required for peptide modification they equate to ~2.5 mg of precatalyst for preparative-scale peptide modification, as discussed later. The effect of lowering the arylboronic acid equivalents was more impactful. At low loadings, 4-carboxy peptide **10a** became the dominant product, as would be expected when less boronic acid was available for transmetalation of the carbonylated-Pd complex (See SI Fig. S1). Up to 8 equiv. of boronic acid, a steady increase in the amount of **8a**

formed, above which further increases had minimal effect on conversion (SI Section 5.5).

We next sought to reduce the amount of acetonitrile required for effective cSMCC. Though acetonitrile is tolerated at low concentrations by many biomolecules,²⁵ at higher concentrations it can disrupt folding within peptides and proteins.²⁶ We observed little difference in cSMCC efficiency or selectivity over a range of 20–50% acetonitrile in PBS. At 10% acetonitrile, a small decrease in both conversion and selectivity for the cSMCC product was observed. However, even under these stringent conditions, **8a** was still the major product (54% relative conversion, Entry 13). In the absence of acetonitrile, the non-carbonylated SMCC side-product **9a** was the dominant species (Entry 12). The reaction was also tolerant of a number of buffers, with pH 8 potassium phosphate buffer being found to be optimal (SI Section 5.8).

We next performed a screen to identify an optimal temperature for the reaction. Conversion was seen at as little as 5 °C, though a marked increase in the formation of **8a** was observed above 25 °C (SI Section 5.9). Notably, above 40 °C conversion to the non-carbonylated SMCC side-product **9a** appeared to be enhanced, and so a temperature of 37 °C was chosen to move forwards. At this temperature, a time-screen indicated that the reaction neared completion after ~4 h, and so reactions were typically left to progress overnight (SI Section 5.10). Under these optimised conditions, peptides containing *para*-bromophenylalanine residues were unreactive, with no evidence of any consumption of the starting peptide.

Cumulatively, these results indicate that under the conditions of the reaction, at weakly basic pH and with aryl iodide coupling partners, CO insertion occurs at a faster rate than



transmetallation. However, transmetallation remains faster than elimination from an acylpalladium–hydroxo complex which may form during the catalytic cycle from the corresponding halide complex, though this pathway was still found to be competitive under some conditions.²³

The hypothesis that CO insertion does not rely on simple CO release from Mo(CO)₆, is supported by a number of observations: (i) selectivity for the cSMCC vs. non-carbonylative product is consistent between a Mo : Pd ratio of ~1 : 1–15 : 1, indicating that increasing CO concentration does not drive selectivity within this range – this observation is caveated by the low solubility of CO in water and the possibility that the catalyst may sequester CO, and so is not in isolation sufficient to support the hypothesis; (ii) when Mo(CO)₆ was pre-incubated in acetonitrile : PBS (pH 8.0) for 30 min, prior to coupling, no change in conversion to **8a** was observed; and (iii) when reactions were performed in an open flask no differences in selectivity were found. Together, these results indicate that direct acetonitrile-induced CO release from the Mo(CO)₆ complex does not play a role in the coupling, while similar selectivities observed when reactions were run in the dark also rule out a photo-activated process of CO release.²⁷ This is in line with the findings of Sun *et al.*, who proposed that acetonitrile plays a key role in activating the Mo(CO)₆ for cSMCC, though the exact mechanism by which this takes place is not clear – in their work, replacing Mo(CO)₆ with a CO atmosphere failed to deliver any product, suggesting induced CO release is not a sole contributing factor, though they did observe some degree of cSMCC at elevated pressures. Our observation that the non-carbonylated product **9a** dominated in the absence of acetonitrile further supports this idea. We have recently reported the formation of Pd–Fe nanocatalysts as a key species during cSMCCs in organic solvents, and it is possible that a related Pd–Mo species may play an important role in the couplings reported here.²⁸

Having optimised conditions for cSMCC, we subsequently applied these to a diverse panel of coupling partners (Fig. 3 See SI Section 5.7 for further details). A screen of organoboronic acids revealed that both electron-rich and electron-deficient substrates were tolerated, with relative conversions ranging from 66–86% for *para*- and *meta*-substituted methoxy (**8b,c**) and fluoro (**8d**) phenyl boronic acids. Couplings with halogenated boronic acids (**8e**), heteroaromatics (**8f**), unprotected phenols (**8h**) and anilines (**8i**) were all tolerated, though 4-pyridinylboronic acid resulted in no conversion (**8g**). Potassium trifluoroborates were also viable coupling partners, giving comparable conversions to the parent boronic acid. Potassium vinyltrifluoroborate provided the related vinyl ketone **8j** with 35% relative conversion, albeit with a high degree of competition from the corresponding vinylbenzene SMCC product (37% conversion). These results demonstrate the potential of our methodology to generate a wide-range of functionalised diarylketones *via* late-stage modification of a common peptide precursor.

With these results in hand, we applied our optimised conditions to the preparative scale modification of peptides. Couplings were performed on a ~20 μmol scale and after completion were centrifuged to remove insoluble molybdenum




Fig. 3 Conversion of pPhe-AAEE, **7**, to carbonylative products, **8**, with varying boronic acids; Reaction conditions: Mo(CO)₆ (2 equiv.), Pd(TFA)₂ (0.4 equiv.), boronic acid (8 equiv.), acetonitrile : PBS (pH 8, 4 : 6), 37 °C, 20 h.

and palladium byproducts. The supernatants were then treated with a commercial dimercaptotriazine-scavenger resin to remove remaining palladium, before peptide purification *via* reverse-phase chromatography. On this scale, phenyl-diaryl ketone **8a** was isolated in an 81% yield.

4-Methoxy and 4-fluoro phenylboronic acids were also coupled successfully on this scale, to generate **8b** (34%) and **8d** (32%) respectively. The reduction in isolated yield relative to **8a** was a result of loss during purification, rather than reduced conversion, as indicated by LC-MS analysis of the crude reaction mixtures.

Couplings were subsequently successfully performed on a number of peptide substrates, on both preparative and analytical scale, demonstrating the tolerance of the reaction to the wide-range of functional groups present on unprotected peptides. Peptides based on the HIV entry inhibitor Peptide T (**11**), an algal RuBisCO-binding domain (**12**), the yeast pheromone alpha-factor (**13**), and the N-terminal fragment of human adrenocorticotrophic hormone (**14**) all successfully underwent cSMCC with 4-fluorophenyl boronic acid (Fig. 4). In the case of **14**, low conversions were observed under our standard conditions, which we attributed to the histidine residue in the sequence. This could be easily addressed, either through the addition of extra palladium precatalyst, or through the addition of sodium ascorbate as a reductant.²⁹

4-Fluorobenzophenones have been previously shown to exhibit greatly enhanced photo-affinity labelling efficiency relative to unsubstituted Bpa.⁷ Modification of these sequences demonstrated that coupling is effective at both internal positions and at peptide termini. In cases where yields were reduced, LC-MS of the crude reaction mixtures indicated high






Fig. 4 Isolated yields of cSMCC-modified peptides. Reactions performed on a 20 mmol scale with 40 mol% 'Pd(TFA)₂', 2 equiv. Mo(CO)₆, and 8 equiv. boronic acid in 6 : 4 potassium phosphate buffer (250 mM, pH 8):acetonitrile. Reactions were incubated at 37 °C for 20 h. Relative conversions for small scale reactions (1.5 μmol) are provided in superscript parentheses where relevant. ^aRun with 100 mol% 'Pd(TFA)₂'.







Fig. 5 Synthesis of peptides modified *via* cSMCCs at 3-iodotyrosine residues. Reactions performed on a 20 mmol scale with 40 mol% Pd(TFA)₂, 2 equiv. Mo(CO)₆, and 8 equiv. boronic acid in 6 : 4 potassium phosphate buffer (250 mM, pH 8):acetonitrile. Reactions were incubated at 37 °C for 20 h. Relative conversions for small scale reactions (1.5 μmol) from LC-MS analysis are provided in superscript parentheses where relevant.



Scheme 2 Carbonylative-Sonogashira cross-coupling applied to Peptide T on a 20 mmol scale. Reaction conditions: Mo(CO)₆ (2 equiv.), Pd(TFA)₂ (0.4 equiv.), phenylacetylene (8 equiv.), acetonitrile:PBS (pH 8, 4 : 6), 37 °C, 20 h.

conversion to the desired cSMCC products, and losses occurred primarily during purification. For example, for Peptide T 78% relative conversion to **11** was observed *via* LC-MS, with an isolated yield of 36%. An exception to the functional group tolerance was the presence of unprotected thiols, with no conversion being observed for a free cysteine-containing decapeptide derivative of laminin (H₂N-CDPGYIGSR-*p*IPhe-CONH₂, **15**, see SI Section 5.11).

Considering alternative halogenated amino acids, 3-iodotyrosine containing peptides based on the sortase-recognition sequence LPTYG and a model peptide H₂N-3ITyr-AAEE-CONH₂ also underwent successful coupling, to generate the corresponding 2-hydroxybenzophenones **16** and **17** (Fig. 5). The 3-iodotyrosine could be installed either directly during SPPS, or *via* *ortho*-iodination of tyrosine residues within cleaved peptides with Barluenga's reagent (IPy₂BF₄) prior to cross-coupling.³⁰ In this scenario, careful purification of the iodinated peptide was essential to remove the pyridine side-product of iodination, which may otherwise inhibit cSMCC. Finally, we also demonstrated that carbonylative copper-free Sonogashira couplings (carbonylative Heck-Cassar-Sonogashira type-reactions) were also viable under our optimised conditions, with alkynyl-aryl ketone **18** being isolated in a 70% yield following the coupling of phenylacetylene to Peptide T (Scheme 2).

Conclusions

In this study, we have demonstrated that carbonylative Suzuki-Miyaura cross-couplings offer a viable method for the late-stage diversification of peptide substrates, and access to peptide-based diaryl ketones for photo-affinity labelling. Key to this work is the use of Mo(CO)₆ as a carbon monoxide surrogate, which negates the need for CO atmospheres which would otherwise make cSMCCs challenging on the small scales required for peptide modification, and without the specialist equipment needed to work under such atmospheres safely. As such, our operationally simple methodology, which can be performed in an open atmosphere, under mild aqueous conditions, is translatable to diverse laboratory settings.

Optimisation of the cSMCC reaction conditions and reagent loadings allowed us to demonstrate the tolerance of the reaction to a broad range of boronic acid coupling partners, as well as applicability to two common unnatural iodinated amino acids, *p*IPhe and 3-ITyr. Both of these amino acids can be integrated into proteins, in the case of 3-ITyr by both codon expansion technologies and chemical modification. Though the reliance on an acetonitrile co-solvent is currently limiting for the translation of this chemistry onto such proteins, we were able to show that cSMCC was possible at as high as 90% water, offering possibilities for further optimisation to open up this pathway. We are currently working to develop the chemistry further with this goal in mind, as well as exploring the use of alternative carbonylative routes to access diverse 'on-peptide' chemical space.



Author contributions

James Kleppen and Neil Scott contributed equally to this manuscript. JPK, NWJS, TJP, and KML synthesised and modified the peptides described in this study. NWJS and KSS synthesised and modified amino acid models and substrates for SPPS. JRD undertook preliminary studies that led directly to this work. LJB supervised and supported student researchers involved in the project. IJSF and CDS developed the project. KSS and CDS wrote the manuscript. CDS supervised and managed the study. All authors contributed to the editing of the manuscript.

Conflicts of interest

The authors declare no competing financial interests.

Data availability

All datasets associated with this manuscript will be made freely accessible through the University of York's PURE research depository.

Supplementary information: all experimental details, including peptide synthesis, reaction protocols, and full screening results. See DOI: <https://doi.org/10.1039/d5sc05588a>.

Acknowledgements

We thank Prof. Steven Cobb for useful discussions about incorporation of 3-ITyr during SPPS. Dr Martin Fascione is thanked for supplying brominated peptides. NWJS, IJSF, and CDS acknowledge support from the University of York Priming Funds (50109438). KSS and CDS are grateful to the Leverhulme Trust for funding (RPG-2022-174). JKP, LJB and CDS acknowledge -PhD studentship support from the BBSRC White Rose Mechanistic Biology DTP. LJB's PhD was supported with generous industrial CASE support from Qkine. CDS acknowledges generous support through a Wellcome Trust Career Development Award (225257/Z/22/Z). We thank Dr Ed Bergstrom and The York Centre of Excellence in Mass Spectrometry for support with peptide mass spectrometry. This centre was created thanks to a major capital investment through Science City York, supported by Yorkshire Forward with funds from the Northern Way Initiative, and subsequent support from the EPSRC (EP/K039660/1; EP/M028127/1).

References

- 1 L. Raynal, N. C. Rose, J. R. Donald and C. Spicer, *Chem.–Eur. J.*, 2021, **27**, 69–88.
- 2 E. Deseke, Y. Nakatani and G. Ourisson, *Eur. J. Org. Chem.*, 1998, 243–251.
- 3 E. Smith and I. Collins, *Future Med. Chem.*, 2015, **7**, 159–183.
- 4 L. Bechtella, C. Kirschbaum, M. Cosset, G. Clodic, L. Matheron, G. Bolbach, S. Sagan, A. Walrant and E. Sachon, *Anal. Chem.*, 2019, **91**, 9102–9110.
- 5 K. Kyro, S. P. Manandhar, D. Mullen, W. K. Schmidt and M. D. Distefano, *Bioorg. Med. Chem.*, 2010, **18**, 5675–5684.
- 6 J. C. Kauer, S. Erickson-Viitanen, H. R. Wolfe and W. F. DeGrado, *J. Biol. Chem.*, 1986, **261**, 10695–10700.
- 7 C. M. Joiner, M. E. Breen and A. K. Mapp, *Protein Sci.*, 2019, **28**, 1163–1170.
- 8 M. M. Hassan and O. O. Olaoye, *Molecules*, 2020, **25**, 2285.
- 9 G. Dormán, H. Nakamura, A. Pulsipher and G. D. Prestwich, *Chem. Rev.*, 2016, **116**, 15284–15398.
- 10 J. Ohata, *Org. Biomol. Chem.*, 2024, **22**, 3544–3558.
- 11 I. Karakaya, K. Rizwan and S. Munir, *ChemistrySelect*, 2023, **8**, e202204005.
- 12 L. Antonella Aronica, *Inorg. Chim. Acta*, 2023, **558**, 121763.
- 13 N. Sun, Q. Sun, W. Zhao, L. Jin, B. Hu, Z. Shen and X. Hu, *Adv. Synth. Catal.*, 2019, **361**, 2117–2123.
- 14 C. D. Spicer and B. G. Davis, *Chem. Commun.*, 2011, **47**, 1698–1700.
- 15 C. D. Spicer, T. Triemer and B. G. Davis, *J. Am. Chem. Soc.*, 2012, **134**, 800–803.
- 16 M. K. Bilyard, H. J. Bailey, L. Raich, M. A. Gafitescu, T. Machida, J. Iglésias-Fernández, S. S. Lee, C. D. Spicer, C. Rovira, W. W. Wyatt and B. G. Davis, *Nature*, 2018, **563**, 235–240.
- 17 Y. Lin, M. Penna, C. D. Spicer, S. G. Higgins, A. Gelmi, N. Kim, S.-T. Wang, J. P. Wojciechowski, E. T. Pashuck, I. Yarovsky and M. M. Stevens, *ACS Nano*, 2021, **15**, 4034–4044.
- 18 A. J. Reay, L. A. Hammarback, J. T. W. Bray, T. Sheridan, D. Turnbull, A. C. Whitwood and I. J. S. Fairlamb, *ACS Catal.*, 2017, **7**, 5174–5179.
- 19 C. D. Spicer and B. G. Davis, *Chem. Commun.*, 2013, **49**, 2747–2749.
- 20 K. Sakamoto, K. Murayama, K. Oki, F. Iraha, M. Katomurayama, M. Takahashi, K. Ohtake, T. Kobayashi, S. Kuramitsu and M. Shirouzu, *Structure*, 2009, **17**, 335–344.
- 21 S. Gnidehou, B. Caillou, M. Talbot, R. Ohayon, J. Kaniewski, M.-S. Noel-Hudson, S. Morand, D. Agnangji, A. Sezan, F. Courtin, A. Virion and C. Dupuy, *FASEB J.*, 2004, **18**, 1574–1576.
- 22 T. Ishiyama, H. Kizaki, N. Miyaura and A. Suzuki, *Tetrahedron Lett.*, 1993, **34**, 7595–7598.
- 23 J. F. Hartwig, in *Organotransition Metal Chemistry: From Bonding to Catalysis*, University Science Books, 2010.
- 24 J. M. Chalker, C. S. C. Wood and B. G. Davis, *J. Am. Chem. Soc.*, 2009, **131**, 16346–16347.
- 25 S. A. Byrne, M. J. Bedding, L. Corcilius, D. J. Ford, Y. Zhong, C. Franck, M. Larance, J. P. Mackay and R. J. Payne, *Chem. Sci.*, 2021, **12**, 14159.
- 26 K. Griebenow and A. M. Klivanov, *J. Am. Chem. Soc.*, 1996, **118**, 11695–11700.
- 27 J. J. Turner, M. W. George, M. Poliakoff and R. N. Perutz, *Chem. Soc. Rev.*, 2022, **51**, 5300–5329.
- 28 Z. Zhu, Z. Wang, Y. Jian, H. Sun, G. Zhang, J. M. Lynam, C. R. McElroy, T. J. Burden, R. L. Inight, I. J. S. Fairlamb, W. Zhang and Z. Gao, *Green Chem.*, 2021, **23**, 920–926.
- 29 N. Li, C. P. Ramil, R. K. V. Lim and Q. Lin, *ACS Chem. Biol.*, 2015, **10**, 379–384.
- 30 G. Espuña, D. Andreu, J. Barluenga, X. Pérez, A. Planas, G. Arsequell and G. Valencia, *Biochemistry*, 2006, **45**, 5957–5963.

