

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Y. Zhong, J. Zeng, M. Li, Y. Liang, S. Guan, K. Zhao, J. Mu, P. Tang, H. Wang and F. Chen, *Chem. Sci.*, 2026, DOI: 10.1039/D5SC08431H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Stereodivergent Access to α - and β -Azanucleosides via Catalyst-Free, Achiral Modulator-Controlled Iodocyclization: A Concise Synthesis of ForodesineYangyang Zhong^{a†}, Jie Zeng^{c†}, Mingwei Li^a, Yuli Liang^c, Shuman Guan^a, Kehan Zhao^a, Jiayi Mu^a, Pei Tang^a, Huijing Wang^{*a}, Fener Chen^{*a,b,d,e}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Stereoselective glycosidic bond formation remains a major challenge in nucleoside synthesis. Azanucleosides, a prominent class of nucleoside analogs wherein the sugar oxygen is replaced by nitrogen, exhibit unique biological activities but hard to achieve anomeric selectivity in synthesis. We disclose a catalyst-free iodocyclization strategy that uses simple achiral molecules—NaI or 2-mercaptobenzimidazole—to stereodivergently access both α - and β -azanucleosides in high yields (up to 98%) with excellent stereocontrol (β : α up to β only, α : β up to 19:1). The utility of this method is demonstrated by a concise synthesis of forodesine in 8 steps with 20% overall yield and >20:1 β : α selectivity—the shortest route and highest stereoselectivity reported to date. DFT studies reveal that hydrogen bonding/Na–O coordination and π – π stacking interactions govern the stereochemical outcomes. This work provides an efficient, scalable platform for accessing diverse azanucleoside therapeutics.

Introduction

Nucleoside analogues (NAs) are an important source of antiviral, antitumor and antibacterial drugs.^{1–5} Among them, azanucleosides constitute a prominent group of structurally modified nucleosides, featured by a nitrogen-containing ring. Interestingly, β -azanucleosides usually exhibit unique physical, chemical, and biological properties.^{6–8} For example, forodesine (BCX-1777, immucillin H) and galidesivir (BCX-4430, immucillin A), two well-known β -azanucleosides, are potent inhibitors of human purine nucleoside phosphorylase, protozoan nucleoside hydrolases, and purine phosphoribosyl transferases (Figure 1).^{9–12} Forodesine has been approved in Japan for the treatment of relapsed or refractory peripheral T-cell lymphoma. Galidesivir demonstrates broad-spectrum antiviral activity by disrupting viral RNA-dependent RNA polymerase.^{13–20} Additionally, several other bioactive molecules structurally similar to forodesine also exhibit significant antiviral activity.^{21–23} On the other hand, α -nucleosides usually have remarkable biological activities




Figure 1. Structures of representative azanucleoside drugs and bioactive molecules.

high enzyme stabilities, as well as inhibitory activities against tumors, bacteria, and plasmodia.²⁴ However, investigations of α -azanucleosides are merely at the initial stage, due to limited synthetic methods. Given the importance of both β - and α -azanucleosides, it is essential to develop a stereocontrolled synthetic strategy applicable to both configurations.

C-glycosylation is the most widely used method for synthesizing C-nucleosides, including electrophilic addition,²⁵ nucleophilic addition,^{26–29} Friedel Craft reaction,^{30,31} metal cross coupling,^{32–35} and radical-mediated reaction.^{36–40} Current strategies for synthesizing β -azanucleosides, as exemplified by the synthesis of forodesine, primarily involve constructing the key glycosidic bond through coupling reactions (Figure 2A).^{41–47}

^a Sichuan Research Center for Drug Precision Industrial Technology, West China School of Pharmacy, Sichuan University, Chengdu, 610041, China.

^b Institute of Flow Chemistry and Engineering, College of Chemistry and Materials, Jiangxi Normal University, Nanchang, 330022, Jiangxi, China.

^c Pharmaceutical Research Institute, Wuhan Institute of Technology, 430205 Wuhan, China.

^d Engineering Center of Catalysis and synthesis for Chiral Molecules, Department of chemistry, Fudan University, Shanghai, 200433, China Shanghai Engineering Center of Industrial.

^e Asymmetric Catalysis for Chiral Drugs, Shanghai 200433, China, Shanghai, 200433, China.

[†] These authors contributed equally to this work.

Supplementary Information available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



To achieve milder reaction conditions, improved stereoselectivity and higher yields, various sugar donors have been developed, such as imines, nitrones, and azalactams. Among all reported synthetic routes to forodesine, the highest β : α selectivity is 8:1, with a maximum yield of 38%.⁴⁷ The synthesis of galidesivir follows a similar strategy, focusing mainly on introducing the amine group at the C4 position of the purine ring.⁴⁵ Nevertheless, existing intermolecular cross-coupling methods for β -azanucleosides face two major challenges: 1) the synthesis of azasugar donors is often complex, typically requiring at least five steps from furanose with low efficiency; 2) control over glycosidic bond stereoselectivity remains unsatisfactory and is highly dependent on auxiliary groups at the C2' position of the glycosyl donor.

Catalytic asymmetric halocyclization of alkenes has proven to be a powerful strategy for accessing stereodefined heterocycles while installing halogen handles for further functionalization.^{48–51} In our previous work, we established a chiral phosphoric acid-

catalyzed intramolecular iodocyclization system for the synthesis of furanose nucleosides, in which achiral additives (NaI or S=PPh₃) were employed to modulate anomeric stereoselectivity.⁵² However, this system suffered from two critical limitations: it proved ineffective for synthesizing azanucleosides where the nucleophile is an NHR group, failing to deliver either α - or β -configured products with satisfied stereocontrol (see Supplementary Table 1); moreover, its reliance on a chiral phosphoric acid catalyst severely limited practicality and scalability. Indeed, industrial adoption of chiral phosphoric acid catalysts is often hampered by the high cost associated with the six-step synthesis from BINOL. The Global Environmental Factor (E_G factor) highlights environmental drawbacks in synthetic systems by quantifying waste generation across the full lifecycle, including catalyst synthesis and reaction processes.^{53,54} Reducing the use of expensive chiral catalysts represents a straightforward approach to minimize the E_G factor.




Figure 2. (A). Current strategies to synthesize forodesine. (B). This work: catalyst-free, achiral modulator-controlled iodocyclization for stereoselective synthesis of α - and β -azanucleosides.




To address these challenges, we have developed a fundamentally distinct, catalyst-free iodocyclization strategy. Stereodivergent synthesis of azanucleosides is achieved in the absence of any chiral catalyst, using only simple achiral modulators: NaI for α -selectivity and 2-mercaptobenzimidazole for β -selectivity (Figure 2B). This approach not only avoids the cost and environmental burden associated with chiral catalysts but also successfully addresses the long-standing challenge of stereoselective azanucleoside formation. The resulting C2-iodinated products serve as versatile intermediates for further functionalization, enabling efficient access to both α - and β -azanucleosides, including a concise synthesis of forodesine. We believe this method provides a robust and scalable platform for diversifying azanucleoside, which remain underexplored in medicinal chemistry.

Results and discussion

Our synthesis commenced with the preparation of halocyclization substrate **5a** (Scheme 1). Asymmetric α -alkylation of glycine Schiff base catalyzed by chiral phase-

transfer catalysts (**PTC**) is a well-established method for accessing unnatural amino acids.⁵⁵⁻⁶¹ Quinuclidinium salts, derived from cinchonidine and cinchonine, are widely used to induce stereoselectivity.⁶²⁻⁶⁵ To enhance the enantioselectivity, we chose glycine Schiff base **1** bearing a benzyl ester group instead of the conventional *tert*-butyl ester. The performance of cinchonidine-derived quinuclidinium salts (**PTC-1 to 4**) in the enantioselective allylation of **1** was evaluated (entries 1-4). The dimeric cinchonidine derivative linked by a benzophenone group (**PTC-4**) proved highly effective, affording product **2** with excellent enantioselectivity (entry 4, 92% yield, 95% *ee*). Lowering the reaction temperature improved both reactivity and enantioselectivity, giving **2** in 95% yield and 99% *ee* (entry 5). This condition performed well even on a 30 mmol scale (entry 6, 93% isolated yield, 99% *ee*). Acid deprotection afforded the free amine intermediate, which was subsequently protected with an Ns (nitrobenzenesulfonyl) group to yield compound **3** in 85% yield. A Heck reaction between aryl iodine **4** and alkene **3** in the presence of Pd[P(*t*-Bu)₃]₂ and Cs₂CO₃ furnished the C-C coupling product **5a**, which served as the substrate for subsequent halocyclization.




Scheme 1. The synthesis of compound **5a**. ^aReaction conditions: **1** (0.1 mmol), allyl bromide (0.12 mmol), **PTC** (0.0005 mmol), 50% aq. KOH (0.25 mL) in PhMe:CHCl₃ (*v:v* = 7:3, 0.75 mL), 0 °C, 8 h. ^bIsolated yield, *ee* values were determined by chiral HPLC.



In our previous work, NaI and S=PPh₃ were identified as key additives for controlling the stereoselectivity of catalytic halocyclization by directing the reaction along specific pathways.⁵² Therefore, we first investigated the effect of S=PPh₃ on stereochemical control (Table 1). Pleasingly, when S=PPh₃ was used as an achiral modulator with NIS as the halogen source, β -nucleoside **6a** was obtained as the major product (entry 1; β : α = 5:1). In contrast, when NaI was used as the modulator, the configuration inverted, affording α -nucleoside **7a** with a β : α ratio of 1:10 (entry 3). The absolute configuration of **7a** was confirmed by single-crystal X-ray crystallography (CCDC No. 2378539). The configuration of **6a** was then assigned by comparing its NMR NOE (Nuclear Overhauser Effect) data with that of **7a**. In the absence of any modulator, no stereocontrol was observed (entry 2). These results suggested that the thiocarbonyl group might be crucial for β -selectivity. We then screened various thiocarbonyl-containing compounds as β -selective modulators (see

Supplementary Table 2). Among them, 2-mercaptobenzoheterocycles exhibited excellent reactivities and stereoselectivities (entries 4-6). Notably, 2-mercaptobenzimidazole **10** afforded product **6a** in high yield (up to 95%) with excellent β -selectivity (β : α = 25:1; entry 6). To verify the necessity of the thiocarbonyl group, control experiments with compounds **11**, **12**, and **13** were conducted (entries 7-9). These compounds showed almost no stereocontrol. We further optimized the reaction conditions to improve α -selectivity (see Supplementary Table 3). By fine-tuning the solvent, high α -selectivity was achieved (β : α = 1:19, entry 10). In the absence of NaI, the diastereomeric ratio was only β : α = 1:3 (entry 11). Loading studies revealed that both NaI and **10** could promote stereoselective iodocyclization catalytically. High α -selectivity was maintained (β : α = 1:19), while β -selectivity slightly decreased (β : α = 1:14) under reduced loading of **10** (see Supplementary Table 4).

Table 1. Optimization of reaction conditions.^a



Entry	Variation from labeled conditions	Yield (%) ^b	β : α (6a : 7a) ^c
1	none	92	5:1
2	no S=PPh ₃	95	1:1
3	NaI instead of S=PPh ₃	85	1:10
4	8 instead of S=PPh ₃	93	11:1
5	9 instead of S=PPh ₃	95	10:1
6	10 instead of S=PPh ₃	95	25:1
7	11 instead of S=PPh ₃	80	2:1
8	12 instead of S=PPh ₃	82	5:1
9	13 instead of S=PPh ₃	79	2:1
10	NaI instead of S=PPh ₃ , THF instead of MeCN	93	1:19
11	no S=PPh ₃ , THF instead of MeCN	90	1:3

^aReaction conditions: **5a** (0.05 mmol), NIS (0.10 mmol), S=PPh₃ (0.05 mmol) in MeCN (2.5 mL), 0 °C for 1 h. ^bisolated yield. ^c β : α values were determined by HPLC.



We systematically evaluated the substrate scope under both β - and α -selective conditions, using a series of halocyclization substrates **5a–u** bearing varied R^1 groups and nucleobase structures (Figure 3). Starting from model substrate **5a**, the effect of different ester groups ($R^1 = \text{CO}_2\text{Me}$, $\text{CO}_2t\text{-Bu}$) was examined. Both β -azanucleosides **6b–c** and α -azanucleosides **7b–c** were obtained in high yields (>90%) with excellent stereoselectivity (β : α up to 24:1, and α : β up to 17:1). To closely mimic the azanucleoside structure, a hydroxymethyl group protected with various groups (TBS, TBDPS, MOM, Bn) was introduced at the C4' position. Substrates **5d–g** performed well, affording β -azanucleosides **6d–g** in high yields (>80%) with good stereocontrol (β : α up to 18:1), and α -azanucleosides **7d–g** in high yields (>87%) with moderate stereocontrol (α : β up to 10:1). In the screening of nucleobase structures, we systematically examined the influence of substituents in 5H-pyrrolo[3,2-*d*]pyrimidine derivatives. Both mono-substituted derivatives at the C4-position (such as OEt, OtBu, Cl, H, NHBn, NH*i*-Pr, Ph, and 4-OMe-Ph) and disubstituted derivatives at the C2 and C4 positions generally exhibited good to excellent stereoselectivity under the optimized conditions. β -Azanucleosides **6h–r** were formed with high selectivity (β : α up to β only) and high yields (>86%), except for **6k** and **6o** (β : α = 3:1). Similarly, α -azanucleosides **7h–r** were mostly obtained with high selectivity (α : β up to 10:1), except for **7l** (α : β = 2.5:1). Evaluation of N5-protecting groups (such as MOM and Bn) showed that they could direct the formation of the corresponding β -azanucleosides **6s–t** and α -azanucleosides **7s–t** with excellent stereoselectivities. Finally, preliminary evaluation of the 7H-pyrrolo[2,3-*d*]pyrimidine scaffold confirmed the good compatibility of the reaction system.

To demonstrate scalability, β -nucleoside analogue **6a** and α -nucleoside analogue **7a** were synthesized on a gram scale from **5a** (Figure 4). Both yields and stereoselectivities were maintained: β -nucleoside **6a** was obtained in 93% yield with β : α > 20:1, and α -nucleoside **7a** in 90% yield with α : β > 15:1.

Using this β -selective iodocyclization as the key step, we developed an efficient asymmetric synthesis of forodesine and a formal synthesis of galidesivir. As shown in Scheme 2, forodesine was synthesized in 20% overall yield over eight steps from commercially available glycine Schiff base **1**. Starting from the β -selective iodocyclization product **6a**, one-pot reductive debenzoylation and halide elimination smoothly afforded compound **14** in 75% yield. Stereospecific *syn*-dihydroxylation of **14** with OsO_4 , followed by deprotection, gave compound **15**.

Treatment of **15** with HCl in methanol then furnished forodesine. Following literature procedures,⁴⁵ functional group modification of the base moiety in forodesine, followed by deprotection, provided the bioactive compound galidesivir. To the best of our knowledge, this work provides the shortest route to forodesine (8 steps vs. 10 in prior reports) and the highest β : α selectivity (>20:1 vs. 8:1).







Figure 4. Gram-scale synthesis of β -azanucleoside **6a** and α -azanucleoside **7a** via catalyst-free, achiral modulator-controlled iodocyclizations.

To demonstrate the versatility of our method and its potential for constructing compound libraries in medicinal chemistry, we performed diverse derivatizations on both β - and α -azanucleosides (Scheme 3). Specifically, β -nucleoside **6a** underwent reductive elimination with NaBH_4/DBU , affording alkene **14** in 75% yield. Removal of the Ns group under 1-mercaptooctane/*t*-BuOK conditions gave **17** in 87% yield. Subsequent reflux of **17** in concentrated HCl/MeOH furnished **18** in 72% yield. Alternatively, hydrogenation of **17** followed by deprotection afforded the C2' and C3'-unsubstituted azanucleoside **20** in 60% yield over two steps. For the α -azanucleoside series, treatment of **7a** with DIBAL-H and subsequent Ns deprotection provided **21** in 72% yield. Radical-mediated deiodination of **21** yielded **22**, while DBU-promoted elimination afforded alkene **23** in 74% yield. Treatment of **23** with concentrated HCl gave **24**. Furthermore, **23** served as a key intermediate for the stereoselective synthesis of target compound **25**. This was achieved via Boc protection of the C4' hydroxymethyl group, dihydroxylation of the C2'–C3' alkene (occurring exclusively from the β -face to give the corresponding diol), and final Boc deprotection.



Figure 3. Substrate scope of α - and β -azanuclioside synthesis.



To gain a thorough understanding of the reaction mechanism, especially on the effect of two achiral molecules, NaI and 2-mercaptobenzimidazole **10**, density functional theory (DFT) studies were performed at PBE0 Level,⁶⁶ serving alkene **5a** as a model substrate (Figure 5). The α -selective iodocyclization starts from **Int1** (-4.6 kcal mol⁻¹). We used the interaction region indicator (IRI)^{67,68} to analyze the interactions between atoms of **Int1** (IRI pic. of **Int1**). Interestingly, NaI in **Int1** is identified as a centered role, cooperating with Ns and ester carbonyl oxygen through Na-O interactions. And Ns can stabilize NIS through π ... π stacking, allowing NIS to attack substrates from top face. These interactions provide a favorable spatial environment for

α -selectivity. The electrophilic addition of I⁺ to **5a** and meanwhile H⁺ being transferred to the N atom result in **Int2** with a reaction barrier of 22.5 kcal mol⁻¹. The following nucleophilic cyclization occurs to generate **PS**, which is exergonic by 25.1 kcal mol⁻¹. In the β -selective iodocyclization pathway, the IRI result of **Int3** reveals that due to the hydrogen bonding interaction, the thiol **10** consistently occupies the region above the C=C bond throughout the reaction. Due to steric effect, the I⁺ from NIS attacks the alkene from the bottom face only, resulting in β -selectivity. Finally nucleophilic cyclization is exergonic by 16.9 kcal mol⁻¹, and the reaction barrier is 32.5 kcal mol⁻¹ (**Int3** to **PR**).




Figure 5. Calculated free energy profile for the formation of nucleosides with α - and β -configurations from substrate **5a** and other reactants.

Conclusion

In summary, we developed a catalyst-free iodocyclization strategy using two simple achiral molecules, NaI and 2-mercaptobenzimidazole **10**, for the stereoselective synthesis of α - and β -azanucleosides in high yields and stereoselectivities. DFT studies reveal that NaI directs α -selectivity through hydrogen bonding and Na-O coordination, while 2-mercaptobenzimidazole **10** controls β -selectivity via π - π stacking interactions. The resulting C2'-iodinated azanucleoside products serve as key intermediates for further functionalization into diverse azanucleoside analogues. The utility of this method is demonstrated by a concise synthesis of forodesine (β : α > 20:1, 8 steps, 20% overall yield). To our knowledge, few existing methods achieve such stereocontrol using solely achiral molecules in the absence of chiral catalysts, particularly in nucleoside synthesis. This work not only expands fundamental chemical understanding but also provides access to underexplored azanucleosides for therapeutic development.

Author contributions

F.C and H.W conceived the idea, guided the project. H.W wrote the manuscript with feedback from other authors. Y. Z. made the initial observations and analyzed the results. Y. Z., M. L., K. Z., J. M., S. G and P. T explored substrate scope and performed derivatizations. J. Z. and Y. L performed the density functional theory calculations on the reaction mechanism.

Conflicts of interest

A patent application (grant no. 202411109085.7, China) dealing with the synthesis of forodesine has been applied, and Huijing Wang and Fener Chen may benefit from royalty payments.

Data availability

Additional data supporting the findings described in this paper are available in the Supplementary Information and available



from the corresponding author upon reasonable request. CCDC 2378539 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, by emailing data_request@ccdc.cam.ac.uk, or by contacting the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, U.K.; fax: +441223 336033.

Acknowledgements

We are grateful for the financial support from the National Natural Science Foundation of China (Grant No. 22201192, U24A20482, 22301228).

Notes and references

- 1 M. Guinan, C. Benckendorff, M. Smith, G. J. Miller, Recent advances in the chemical synthesis and evaluation of anticancer nucleoside analogues. *Molecules* 2020, **25**, 2050.
- 2 G. Li, T. Yue, P. Zhang, W. Gu, L. J. Gao, L. Tan, Drug discovery of nucleos(t)ide antiviral agents: dedicated to Prof. Dr. Erik De Clercq on occasion of his 80th birthday. *Molecules* 2021, **26**, 923.
- 3 M. I. Elzagheid, Nucleosides and nucleoside analogues as emerging antiviral drugs. *Mini-Rev. Org. Chem.* 2021, **18**, 672–679.
- 4 S. Man, Y. Lu, L. Yin, X. Cheng, L. Ma, Potential and promising anticancer drugs from adenosine and its analogs. *Drug Discov. Today* 2021, **26**, 1490–1500.
- 5 Q. Wang, K. Chen, Y. Wang, Z. Rao, X. Zhang, Biotechnological synthesis of nucleoside analogs: recent progress and perspectives. *Green Synth. Catal.* 2025, doi:10.1016/j.gresc.2025.05.002.
- 6 P. Merino, T. Tejero, I. Delso, Current developments in the synthesis and biological activity of aza-C-nucleosides: immucillins and related compounds. *Curr. Med. Chem.* 2008, **15**, 954–967.
- 7 G. B. Evans, P. C. Tyler, V. L. Schramm, Immucillins in infectious diseases. *ACS Infect. Dis.* 2018, **4**, 107–117.
- 8 G. B. Evans, V. L. Schramm, P. C. Tyler, The immucillins: design, synthesis and application of transition-state analogues. *Curr. Med. Chem.* 2015, **22**, 3897–3909.
- 9 R. W. Miles, P. C. Tyler, R. H. Furneaux, C. K. Bagdassarian, V. L. Schramm, One-third-the-sites transition-state inhibitors for purine nucleoside phosphorylase. *Biochemistry* 1998, **37**, 8615–8621.
- 10 G. A. Kicska, L. Long, H. Hörig, C. Fairchild, P. C. Tyler, R. H. Furneaux, V. L. Schramm, H. L. Kaufman, Immucillin H, a powerful transition-state analog inhibitor of purine nucleoside phosphorylase, selectively inhibits human T lymphocytes. *PNAS* 2001, **98**, 4593–4598.
- 11 S. Bantia, P. J. Miller, C. D. Parker, S. L. Ananth, L. L. Horn, J. M. Kilpatrick, P. E. Morris, T. L. Hutchison, J. A. Montgomery, J. S. Sandhu, Purine nucleoside phosphorylase inhibitor BCX-1777 (Immucillin-H) a novel potent and orally active immunosuppressive agent. *Int. Immunopharmacol.* 2001, **1**, 1199–1210.
- 12 A. Korycka, J. Z. Błoński, T. Robak, Forodesine (BCX-1777, Immucillin H) – a new purine nucleoside analogue: mechanism of action and potential clinical application. *Mini Rev. Med. Chem.* 2007, **7**, 976–983.
- 13 T. K. Warren, J. Wells, R. G. Panchal, K. S. Stuthman, N. L. Garza, S. A. V. Tongeren, L. Dong, C. J. Retterer, B. P. Eaton, G. Pegoraro, S. Honnold, S. Bantia, P. Kotian, X. Chen, B. R. Taubenheim, L. S. Welch, D. M. Minning, Y. S. Babu, W. P. Sheridan, S. Bavari, Protection against filovirus diseases by a novel broad-spectrum nucleoside analogue BCX4430. *Nature* 2014, **508**, 402–405.
- 14 J. G. Julander, S. Bantia, B. R. Taubenheim, D. M. Minning, P. Kotian, J. D. Morrey, D. F. Smee, W. P. Sheridan, Y. S. Babu, BCX4430, a novel nucleoside analog, effectively treats yellow fever in a Hamster model. *Antimicrob. Agents Chemother.* 2014, **53**, 6607–6614.
- 15 E. De Clercq, C-nucleosides to be revisited. *J. Med. Chem.* 2016, **59**, 2301–2311.
- 16 R. Taylor, P. Kotian, T. Warren, R. Panchal, S. Bavari, J. Julander, S. Dobo, A. Rose, Y. El-Kattan, B. Taubenheim, Y. Babu, W. P. Sheridan, BCX4430 – a broad-spectrum antiviral adenosine nucleoside analog under development for the treatment of Ebola virus disease. *J. Infect. Public Health* 2016, **9**, 220–226.
- 17 J. G. Julander, V. Siddharthan, J. Evans, R. Taylor, K. Tolbert, C. Apuli, J. Stewart, P. Collins, M. Gebre, S. Neilson, A. Van Wettere, Y. M. Lee, W. P. Sheridan, J. D. Morrey, Y. S. Babu, Efficacy of the broad-spectrum antiviral compound BCX4430 against Zika virus in cell culture and in a mouse model. *Antiviral Res.* 2017, **137**, 14–22.
- 18 J. B. Westover, A. Mathis, R. Taylor, L. Wandersee, K. W. Bailey, E. J. Seifing, B. T. Hickerson, K. H. Jung, W. P. Sheridan, B. B. Gowen, *Antiviral Res.* 2018, **156**, 38–45.
- 19 J. G. Julander, J. F. Demarest, R. Taylor, B. B. Gowen, D. M. Walling, A. Mathis, Y. S. Babu, An update on the progress of galidesivir (BCX4430), a broad-spectrum antiviral. *Antiviral Res.* 2021, **195**, 105180.
- 20 Y. Xie, Chi, L. Y.; S. Q. Liu, W. Y. Zhu, BCX4430 inhibits the replication of rabies virus by suppressing mTOR-dependent autophagy invitro. *Virology* 2023, **585**, 21–31.
- 21 K. J. Sparrow, R. Shrestha, J. M. Wood, K. Clinch, B. L. Hurst, H. Wang, B. B. Gowen, J. G. Julander, E. B. Tarbet, A. M. McSweeney, V. K. Ward, G. B. Evans, L. D. Harris, An isomer of galidesivir that potently inhibits influenza viruses and members of the bunyavirales order. *ACS Med. Chem. Lett.* 2023, **14**, 506–513.
- 22 Y. S. Babu, P. L. Kotian, S. Bantia, M. Wu, V. S. Kumar, Preparation of antiviral aza-sugar-containing nucleosides. 2014, WO 2014078778.
- 23 Y. S. Babu, P. L. Kotian, S. Bantia, M. Wu, V. S. Kumar, Antiviral azasugar-containing nucleosides. 2015, US20150291596.
- 24 G. Ni, Y. Du, F. Tang, J. Liu, H. Zhao, Q. Chen, Review of α -nucleosides: from discovery, synthesis to properties and potential applications. *RSC Adv.* 2019, **9**, 14302–14320.
- 25 J. Bouton, S. Van Calenbergh, J. Hullaert, Sydnone ribosides as a platform for the synthesis of pyrazole C-nucleosides: a unified synthesis of Formycin B and Pyrazofurin. *Org. Lett.* 2020, **22**, 9287–9291.
- 26 Q. Li, E. Lescrinier, E. Groaz, L. Persoons, D. Daelemans, P. Herdewijn, S. De Jonghe, Synthesis and biological evaluation of pyrrolo[2,1-f][1,2,4]triazine C-nucleosides with a ribose, 2'-deoxyribose, and 2',3'-dideoxyribose sugar moiety. *ChemMedChem* 2018, **13**, 97–104.
- 27 T. Liu, Z. Zhu, H. Ren, Y. Chen, G. Chen, M. Cheng, D. Zhao, J. Shen, W. Zhu, B. Xiong, Y. L. Chen, Efficient syntheses of α - and β -C-nucleosides and the origin of anomeric selectivity. *Org. Chem. Front.* 2018, **5**, 1992–1999.
- 28 P. Nie, E. Groaz, D. Daelemans, P. Herdewijn, Xylo-C-Nucleosides with a pyrrolo[2,1-f][1,2,4]triazin-4-amine heterocyclic base: synthesis and antiproliferative properties. *Bioorg. Med. Chem. Lett.* 2019, **29**, 1450–1453.
- 29 P. Nie, E. Groaz, P. Herdewijn, Synthesis of a threosyl-C-nucleoside phosphonate. *Eur. J. Org. Chem.* 2019, **2019**, 6666–6672.
- 30 H. Marzag, M. Zerhouni, H. Tachallait, L. Demange, G. Robert, K. Bougrin, P. Auberger, R. Benhida, Modular synthesis of new



- C-aryl-nucleosides and their anti-CML activity. *Bioorg. Med. Chem. Lett.* 2018, **28**, 1931–1936.
- 31 H. Tachallait, M. Safir Filho, H. Marzag, K. Bougrin, L. Demange, A. R. Martin, R. Benhida, A Straightforward and versatile FeCl₃ catalyzed Friedel-Crafts C-glycosylation process. Application to the synthesis of new functionalized C-nucleosides. *New J. Chem.* 2019, **43**, 5551–5558.
 - 32 H. Gong, M. R. Gagné, Diastereoselective Ni-catalyzed Negishi cross-coupling approach to saturated, fully oxygenated C-alkyl and C-aryl glycosides. *J. Am. Chem. Soc.* 2008, **130**, 12177–12183.
 - 33 Q. Wang, S. An, Z. Deng, W. Zhu, Z. Huang, G. He, G. Chen, Palladium-catalysed C–H glycosylation for synthesis of C-aryl glycosides. *Nat. Catal.* 2019, **2**, 793–800.
 - 34 Y. Li, Z. Wang, L. Li, X. Tian, F. Shao, C. Li, Chemoselective and diastereoselective synthesis of C-Aryl nucleoside analogues by Nickel-catalyzed cross-coupling of furanosyl acetates with aryl iodides. *Angew. Chem. Int. Ed.* 2022, **61**, e202110391.
 - 35 R. Xie, J. Xu, H. Shi, C. Xiao, N. Wang, N. Huang, H. Yao, Stereocoupled Synthesis of aryl C-nucleosides under ambient conditions. *Org. Lett.* 2024, **26**, 5162–5166.
 - 36 S. O. Badir, A. Dumoulin, J. K. Matsui, G. A. Molander, Synthesis of reversed C-acyl glycosides through Ni/photoredox dual catalysis. *Angew. Chem. Int. Ed.* 2018, **57**, 6610–6613.
 - 37 A. Dumoulin, J. K. Matsui, Á. Gutiérrez-Bonet, G. A. Molander, Synthesis of non-classical arylated C-saccharides through Nickel/photoredox dual catalysis. *Angew. Chem. Int. Ed.* 2018, **57**, 6614–6618.
 - 38 Y. Ma, S. Liu, Y. Xi, H. Li, K. Yang, Z. Cheng, W. Wang, Y. Zhang, Highly stereoselective synthesis of aryl/heteroaryl-C-nucleosides via the merger of photoredox and Nickel catalysis. *Chem. Commun.* 2019, **55**, 14657–14660.
 - 39 Q. Wang, J. Duan, P. Tang, G. Chen, G. He, Synthesis of non-classical heteroaryl C-glycosides via Minisci-type alkylation of N-heteroarenes with 4-glycosyl-dihydropyridines. *Sci. China Chem.* 2020, **63**, 1613–1618.
 - 40 Y. Wei, B. Ben-zvi, T. Diao, Diastereoselective synthesis of aryl C-glycosides from glycosyl esters via C–O bond homolysis. *Angew. Chem. Int. Ed.* 2021, **6**, 9433–9438.
 - 41 G. B. Evans, R. H. Furneaux, G. J. Gainsford, V. L. Schramm, P. C. Tyler, Synthesis of transition state analogue inhibitors for purine nucleoside phosphorylase and N-riboside hydrolases. *Tetrahedron Lett.* 2000, **56**, 3053–3062.
 - 42 G. B. Evans, R. H. Furneaux, T. L. Hutchison, H. S. Kezar, P. E. Morris, V. L. Schramm, P. C. Tyler, Addition of lithiated 9-deazapurine derivatives to a carbohydrate cyclic imine: convergent synthesis of the aza- C-nucleoside immucillins. *J. Org. Chem.* 2001, **66**, 5723–5730.
 - 43 V. P. Kamath, J. Xue, J. J. Juarez-Brambila, P. E. Morris, Alternative route towards the convergent synthesis of a human purine nucleoside phosphorylase inhibitor – forodesine HCl. *Tetrahedron Lett.* 2009, **50**, 5198–5200.
 - 44 V. P. Kamath, J. Xue, J. J. Juarez-Brambila, Synthesis of analogs of forodesine HCl, a human purine nucleoside phosphorylase inhibitor – Part II. *Bioorg. Med. Chem. Lett.* 2009, **19**, 2627–2629.
 - 45 M. Zhang, F. Xue, J. Ou, Y. Huang, F. Lu, B. Zhou, Z. Zheng, X. Y. Liu, W. Zhong, Y. Qin, Practical synthesis of immucillins BCX-1777 and BCX-4430. *Org. Chem. Front.* 2020, **7**, 3675–3680.
 - 46 J. Ou, M. Zhang, P. Li, F. Xue, Y. Qin, Synthesis of purine nucleoside phosphorylase inhibitor Forodesine. *West China J. Pharm. Sci.* 2021, **36**, 495–500.
 - 47 K. A. Krishnakumar, R. S. Lankalapalli, Synthesis of immucillins BCX-1777 and BCX-4430 from a common precursor. *Eur. J. Org. Chem.* 2022, e202200428.
 - 48 S. E. Denmark, W. E. Kuester, M. T. Burk, Catalytic, asymmetric halofunctionalization of alkenes—a critical perspective. *Angew. Chem. Int. Ed.* 2012, **51**, 10938–10953.
 - 49 R. Kristianslund, J. E. Tungen, T. V. Hansen, Catalytic enantioselective iodolactonization reactions. *Org. Biomol. Chem.* 2019, **17**, 3079–3092.
 - 50 S. Liu, B. Zhang, W. Xiao, Y. Li, J. Deng, Recent advances in catalytic asymmetric syntheses of functionalized heterocycles via halogenation/chalcogenation of carbon-carbon unsaturated bonds. *Adv. Synth. Catal.* 2022, **364**, 3974–4005.
 - 51 J. Yan, Z. Zhou, Q. He, G. Chen, H. Wei, W. Xie, The applications of catalytic asymmetric halocyclization in natural product synthesis. *Org. Chem. Front.* 2022, **9**, 499–516.
 - 52 Q. Wang, J. Mu, J. Zeng, L. Wan, Y. Zhong, Q. Li, Y. Li, H. Wang, F. Chen, Additive-controlled asymmetric iodocyclization enables enantioselective access to both α - and β -nucleosides. *Nat. Commun.* 2023, **14**, 138.
 - 53 R. A. Sheldon, The E factor 25 years on: the rise of green chemistry and sustainability. *Green Chem.* 2017, **19**, 18–43.
 - 54 A. Antenucci, S. Dughera, Usefulness of the global E factor as a tool to compare different catalytic strategies: four case studies. *Catalysts* 2023, **13**, 102.
 - 55 K. Maruoka, T. Ooi, Enantioselective amino acid synthesis by chiral phase-transfer catalysis. *Chem. Rev.* 2003, **103**, 3013–3028.
 - 56 T. Ooi, K. Maruoka, Recent advances in asymmetric phase-transfer catalysis. *Angew. Chem. Int. Ed.* 2007, **46**, 4222–4266.
 - 57 S. Jew, H. Park, Cinchona-based phase-transfer catalysts for asymmetric synthesis. *Chem. Commun.* 2009, **46**, 7090–7103.
 - 58 S. Shirakawa, K. Maruoka, Recent developments in asymmetric phase-transfer reactions. *Angew. Chem. Int. Ed.* 2013, **52**, 4312–4348.
 - 59 H. J. Lee, K. Maruoka, Recent asymmetric phase-transfer catalysis with chiral binaphthyl-modified and related phase-transfer catalysts over the last 10 years. *Chem. Rec.* 2023, **23**, e202200286.
 - 60 H. J. Lee, K. Maruoka, Asymmetric phase-transfer catalysis. *Nat. Rev. Chem.* 2024, **8**, 851–869.
 - 61 P. Pecchini, M. Fochi, F. Bartoccini, G. Piersanti, L. Bernardi, Enantioselective organocatalytic strategies to access noncanonical α -amino acids. *Chem. Sci.* 2024, **15**, 5832–5868.
 - 62 E. J. Corey, F. Xu, M. C. Noe, Rational approach to catalytic enantioselective enolate alkylation using a structurally rigidified and defined chiral quaternary ammonium salt under phase transfer conditions. *J. Am. Chem. Soc.* 1997, **119**, 12414–12415.
 - 63 S. Woo, Y. G. Kim, B. Lim, J. Oh, Y. Lee, H. Gwon, K. Nahm, An unusual electronic effect of an aromatic-F in phase-transfer catalysts derived from cinchona-alkaloid. *Org. Lett.* 2002, **4**, 4245–4248.
 - 64 S. Woo, Y. G. Kim, B. Lim, J. Oh, Y. Lee, H. Gwon, K. Nahm, Dimeric cinchona ammonium salts with benzophenone linkers: enantioselective phase transfer catalysts for the synthesis of α -amino acids. *RSC Adv.* 2018, **8**, 2157–2160.
 - 65 J. Oh, J. Park, K. Nahm, Counter-rotatable dual cinchona quinuclidinium salts and their phase transfer catalysis in enantioselective alkylation of glycine imines. *Chem. Commun.* 2021, **57**, 6816–6819.
 - 66 J. P. Perdew, K. Burke, M. Ernzerhof, Generalized Gradient Approximation Made Simple. *Phys. Rev. Lett.* 1997, **78**, 1396.
 - 67 T. Lu, Q. Chen, Interaction region indicator: a simple real space function clearly revealing both chemical bonds and weak interactions. *Chem. Methods* 2021, **1**, 231–239.
 - 68 T. Lu, F. Chen, Multiwfn: a multifunctional wavefunction analyzer. *J. Comput. Chem.* 2012, **33**, 580–5.



Data available statement

Supporting Information is available and includes experimental details.

