

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: N. Spreckelmeyer, J. Kim, L. Jessika, E. S. Horst, J. Zhang and A. Studer, *Chem. Sci.*, 2026, DOI: 10.1039/D5SC08793G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Reductive Rearrangement of Substituted Quinolines to 2,3-Disubstituted Indoles Enabled by Water Activation

Nico Spreckelmeyer, Jieun Kim, Jessika Lammert, Elena Sophia Horst, Jingjing Zhang and Armido Studer*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Herein, we report a selective reductive rearrangement of substituted quinolines into indoles — a privileged structural motif widely found in natural products and bioactive molecules. This quinoline skeletal editing is accomplished through water activation mediated by photocatalytically generated phosphine radical cations. The developed protocol provides a robust and broadly applicable approach for synthesizing diverse indole derivatives from readily available quinoline substrates.

Introduction

N-Heterocycles play a major role in drug discovery.^{1–6} Quinolines — and even more so indoles — demonstrate their significance through their prevalence in various developed drugs and naturally occurring bioactive molecules. Prominent examples include indole alkaloids naturally derived from L-tryptophan — such as reserpine, yohimbine, and members of the ibogaine family — which have found broad use in medicine (Figure 1A).⁷

Late-stage functionalization of heterocyclic compounds — such as quinolines — plays a crucial role in rapidly expanding the chemical space of potential drug candidates (Figure 1B, top). A wide range of reactions has been developed to expand the structural diversity within (hetero)aromatic scaffolds. Along these lines, C–H activation has emerged as a powerful strategy for the peripheral diversification of N-heterocycles, allowing access to an expanded spectrum of molecular structures.^{8–11} In this context, the meta-functionalisation of pyridines and quinolines has recently gained great attention.^{12–14} Furthermore, escaping the “flatland” of planar aromatic structures toward non-aromatic three-dimensional frameworks might be even more valuable. This can be achieved through various approaches, including hydrogenation and other dearomatization strategies.^{15–19} Hydrogenations can be carried out, for example, via transition-metal-catalyzed reductions with H₂ and other approaches; however, challenges often arise in controlling both stereoselectivity and chemoselectivity.^{20–30} Arene hydrogenation can also proceed via radical intermediates, employing established methods such as the

Birch reduction or using Sml₂ as a reductant, among others.^{31–43}

In recent years, photomediated dearomatization via triplet excitation, energy transfer (EnT) or single-electron transfer (SET) processes has attracted growing attention.^{44,45} Notably, beyond hydrogenation, photochemical approaches can also be employed to construct bi- and tricyclic structures through cycloadditions and coupling reactions.^{46–53}

Incorporating skeletal editing steps into (hetero)arene hydrogenation cascades can increase the complexity of the resulting structures and further expand the accessible chemical space, as demonstrated herein for the reductive transformation of quinolines to indoles (Figure 1B, bottom).^{54–56} Highlighting the different properties of these molecular scaffolds — quinolines are electron-poor aromatic systems with basic properties while indoles are rather electron rich. Quinolines can be accessed from indoles by reaction with carbenes through cyclopropanation followed by ring expansion, known as the *Ciamician–Dennstedt* reaction.^{57–60} Carbon-atom deletion in quinolines to access indoles can be accomplished by pre-functionalizing the nitrogen, for example via N-oxides or N-acylimides.^{61–64} In this context, *Kaneko's* groundbreaking photochemical rearrangement of quinoline N-oxides to N-acylindoles was subsequently advanced by *Levin*,⁶⁵ which upon further exploration led to a general and selective direct carbon deletion at either the 2- or 3-position, yielding the corresponding indole products.⁶⁶ This represents divergent skeletal editing, allowing access to multiple products from a single substrate, an area of increasing attention.^{67,68}

Rather unexplored remains the strategy to selectively induce a reductive rearrangement of non-pre-functionalized quinolines to give indoles via controlled structural reorganization of the heteroarene scaffold. This was shown by our group through photocatalytic water activation and also in joint elegant efforts by *Pratt* and *Chiba* through polysulfide anion catalysis especially

Organisch-Chemisches Institut, Universität Münster, Corrensstraße 40, 48149 Münster, Germany. E-mail: studer@uni-muenster.de


* Footnotes relating to the title and/or authors should appear here.

Supplementary Information available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x




ARTICLE


A) Indole Alkaloids



B) Late Stage Functionalisation of Quinolines & their Conversion to Indoles



C) Studer 2023 – Water Activation



D) Quinolines as Substrates




Figure 1: A) Examples of indole alkaloids derived from L-tryptophan. B) Late-Stage functionalisation of quinolines and their conversion to indole scaffolds by skeletal editing. C) Phosphine mediated water activation for hydrogenation. D) Employing quinolines as substrates in the system leads to a mixture of products & further optimization towards selective reductive rearrangement.



ARTICLE

covering C4 substituted quinolines.^{69,52} In our work, we found that photocatalytically generated phosphine radical cations immediately react with water (Figure 1C).⁶⁹ Deprotonation of such adducts then provides highly reactive phosphoranyl radicals with very low O–H bond dissociation energy (<15 kcal/mol) enabling the reduction of alkenes as well as naphthalenes through intermolecular hydrogen atom transfer in combination with an arylthiol cocatalyst. Of note, deprotonation of such reactive HO-phosphoranyl radicals leads to phosphine oxide radical anions that can be employed as potent ground state single electron reductants (measured –3.1 V vs SCE).⁷⁰

Encouraged by our initial studies on the reductive rearrangement of quinolines to indoles we decided to

investigate this highly valuable transformation in greater detail, as only moderate yields, limited selectivity, and a requirement for a 2-aryl substituent was reported in our first disclosure (Figure 1D).⁶⁹ The main challenges arise from the selective formation of the indole core (**2**) and accordingly the suppression of “simple” hydrogenation to give tetrahydroquinoline **3**.³⁰ Moreover, the formation of another undesired byproduct **4**, which arises from rearrangement followed by addition to a second quinoline, must also be suppressed.⁷¹ Herein, we report reaction conditions that enable the selective formation of indoles from quinolines via reductive rearrangement using our water-activation system.

Table 1: Optimization of the reaction conditions

Entry	Deviation from above	Yield of 2a	Conversion
1	TripSH (20 mol%) & no pTsOH	Traces + 3a 14%	28%
2[a]	150 μ L water	68% (55%) + 3a 15%	Full Conversion
3[a]	with 150 μ L water & TripSH (20 mol%)	57% + 3a 30%	Full Conversion
4[a]	with P2 instead of P1 & MeCN/THF (3/1)	96% (81%)	Full Conversion
5	–	>99% (86%)	Full Conversion
6	with 0.5 eq. pTsOH	46%	53%
7	with 2.0 eq. P	90%	Full Conversion
8	with 50 μ L H ₂ O	88%	Full Conversion
9	with 150 μ L H ₂ O	92%	Full Conversion
10	16 h reaction time	>99% (92%)	Full Conversion
11	no light / no PC / no P	– / – / –	No Conversion
12	no H ₂ O / no pTsOH	36% / Traces 3a	43% / 21%

General information: **1a** was used for optimizing & the pTsOH was dried at 100 °C under high vacuum. The yields and conversion were determined by ¹H-NMR with CH₂Br₂ as internal standard & the isolated yields are given in parentheses. [a] In these cases the HCl salt of quinoline **1a** was used and no pTsOH was submitted to the reaction mixture.

PC: [Ir(dFCF₃ppy)₂(dtbbpy)]PF₆

P1: tris(4-methoxyphenyl)phosphine **P2:** XPhos

TripSH: 2,4,6-Triisopropylbenzenethiol

pTsOH: p-toluenesulfonic acid

Results and Discussion

For reaction optimization, we selected the model substrate **1a**. Extensive experimentation revealed that a quantitative yield of

2a (86% isolated) can be achieved for the reductive skeletal editing with phosphine **P1** (P(p-MeOC₆H₄)₃, 2.5 equiv.), the Ir-based photocatalyst **PC** ([Ir(dFCF₃ppy)(dtbbpy)]PF₆, 2.5 mol%),



in the presence of *p*-toluenesulfonic acid (pTsOH, 1 equiv.) in acetonitrile/water (40:1) upon blue LED irradiation for 24 hours at 20 °C (Table 1, entry 5). The tetrahydroquinoline **3a** and the dimerization product **4a** were not formed under these conditions. We later found that the isolated yield of **2a** (92%) can be slightly further improved by shortening the reaction time to 16 hours, likely as a result of a slow decomposition of **2a** under the applied condition (entry 10). Comparing these results with the previously reported “acid-free” conditions (entry 1), a far higher conversion (99% versus 28%) could be achieved and the yield of **2a** could be significantly improved (92% isolated versus traces).⁶⁹ Surprisingly, reaction with the HCl-salt as the substrate (**1a**-HCl) in the absence of pTsOH provided the indole **2a** (68%) besides the tetrahydroquinoline **3a** (15%), indicating that the counter anion of the quinoline salt somehow influences chemodivergency (entry 2). Repeating the experiment with the HCl-salt in the presence of an aryl thiol cocatalyst (TripSH) further increased the yield of the tetrahydroquinoline byproduct **3a** (30%, entry 3). This observation further indicates that the use of a HAT cocatalyst biases product selectivity in favor of the hydrogenation product. However, using the HCl-salt as the substrate in combination with phosphine **P2** (XPhos), a high yield of the targeted rearrangement product **2a** was achieved (96%, entry 4). For the screening of other phosphines, we refer to the Supporting Information (SI). Since the preformation of the HCl-salt requires a precipitation step, the protocol using simple addition of pTsOH to the reaction mixture was considered more practical. Moreover, **P2** is costlier than **P1**. Quinoline protonation is required, as with 0.5 equiv. of pTsOH, only a 53% conversion was achieved, revealing that the unprotonated quinoline does likely only show very little reactivity under these conditions (entry 6). Lowering the amount of phosphine also had a detrimental effect on the reaction outcome (entry 7) and the amount of water also slightly affected reactivity (entries 8 & 9). Control reactions gave negative results showing the necessity of all reagents (entries 11 & 12).

To elucidate the generality of our reaction we submitted a wide range of quinolines to the optimized conditions (Figure 2A). The 3-alkyl-substituent was varied keeping the 2-phenyl substituent and we found that linear 3-alkyl groups led to slightly lower yields as compared to the branched isopropyl-substituted congener (see **2a**-**2c**). Of note, the disubstituted 3-ethyl-2-phenylquinoline (**1c**) and the monosubstituted 2-phenylquinoline (**1d**) provided the same yield, showing that monosubstituted quinolines are eligible substrates. We then investigated the electronic-effect that is exerted by the 2-aryl substituent on the reaction outcome. Installing a trifluoromethyl group in para-position resulted in a slightly reduced yield (**2e**, 76%). However, with the monosubstituted quinoline **1f** carrying the *p*-CF₃C₆H₄-substituent, the yield significantly decreased to 27% (**2f**). In comparison, the presence of an electron-donating para-methoxy group on the 2-aryl substituent had only a minor effect on the reaction efficiency, see **2g** (88%) and **2h** (62%). Similar yields were achieved with the para-methyl and para-fluoro congeners (**2i** and **2j**)

demonstrating that except for the trifluoromethyl compound, all products in this series were obtained in yields greater than 50%. A meta-trimethylsilyl (**2k**, 51%) and a meta-hydroxy (**2l**, 65%) substituent were well tolerated in the 2-aryl group, however, the 2-meta-methoxyphenyl-quinoline (**1m**) delivered the targeted indole **2m** (28%) in low yield only. A slightly reduced yield was also noted for the ortho-methoxy-substituted 2-aryl quinoline **1n**, likely for steric reasons (**2n**, 40%).

We extended our studies to examine the substituent effects on the non-heterocyclic ring of the starting quinoline. Quinolines **1o**-**1r**, bearing a methyl group at positions 5 to 8, were tested and afforded comparable yields ranging from 76-79%. The presence of halogen substituents exerted a more pronounced effect. Compounds **2s**-**2u** exhibited reduced yields, with values ranging from 48% to 31% for bromine and fluorine substituents at the 5-position of the indole ring, and 25% for a chlorine substituent at the 6-position. The lowest yield in this series was noted for the quinoline with a methoxycarbonyl substituent at position 6 (**2v**, 18%). We assume that the reduced yields (**2s**-**2v**) arise from radical fragmentations of the intermediates or the decomposition of the products under the reaction conditions. The 3-position of 2,3-disubstituted-quinoline starting materials was diversified and it could be observed that ester, amide and aryl-functionalities were tolerated and the corresponding indoles were isolated with good yields (**2w**-**2z**, 43-62%), except for the secondary amide **2y** that was isolated in 20% yield only along with 18% of 2-phenylindole as a side product. Notably, the good yield for the reductive rearrangement of ethyl ester **1w** was achieved upon using phosphine **P3** (P(*p*-CF₃C₆H₄)₃) in place of **P1**.



We next addressed the 2-position of the quinoline core to determine whether an aryl functionality is required to facilitate the reaction. In this series, the 3-substituent was varied as well. Pleasingly, we could observe that isopropyl (**2aa** and **2ac**), tert-butyl (**2ab**), methyl (**2ad**) and trifluoro (**2ae**) groups were all tolerated as 2-substituents, significantly enlarging the scope of our reaction. The corresponding products were isolated in low to very good yields (23-83%). Furthermore, we found that 2,3-ring-annulated quinolines could be rearranged to give a carbazole **2af** (48%) and indoles **2ag** (72%) and **2ah** (31%), demonstrating that a combined ring-contraction and ring-enlargement of the quinoline starting material is possible.

We scaled up the reaction for substrate **1d** to a 2 mmol scale and were able to isolate the product in 47% yield, compared to 64% at the 0.2 mmol scale (Figure 2B). Further, indole **2a** was successfully prepared in 60% yield at 4 mmol scale. In this case, a lower catalyst loading and reduced solvent volume were used, which led to solubility issues and a slower reaction, explaining the presence of unreacted starting material. Nevertheless, the reaction had to be stopped due to product decomposition observed for this substrate during the reaction. Taking this into account, a yield of 70%, based on the consumed starting material, was obtained.



ARTICLE

A) Scope



B) Scale Up



Figure 2: A) Overview of the scope of the optimized reaction conditions, shown at the top. [a] In this case 7% of defluorinated side-product (3-methyl-2-phenylindole) coeluted and a separation from **2j** was not possible. [b] **P3** (P(C₆H₄CF₃)₃) was used instead of **P1** and the water-amount was reduced to 50 μ L. [c] 2-Phenylindole was isolated as a side product. B) Scale Up experiments. Abbreviations: n.f. – not formed.



ARTICLE

To investigate the reaction mechanism, several control experiments were conducted (Figure 3). Exchange of H₂O with D₂O showed that the incorporated hydrogen atoms originate from the added water (Figure 3A). In our previous work we showed that the Ir-based **PC** used can be oxidatively quenched by the protonated quinoline.³⁰ Further, NMR studies confirmed that quinoline **1a** is protonated by pTsoH, while the phosphine **P1** remains in its neutral form (Figure 3B). Cyclovoltammetry measurement (CV) of **1a** revealed a reduction potential of $E = -2.61$ V vs. Fc/Fc⁺ (-2.21 V vs. SCE) and two signals for the protonated quinoline **1a**-HCl at $E = -0.95$ V vs. Fc/Fc⁺ (-0.55 V vs. SCE) and $E = -2.52$ V vs. Fc/Fc⁺ (-2.12 V vs. SCE) (Figure 3C). Thus, the excited **PC** ($E(\text{Ir}^{4+}/\text{Ir}^{3+}) = -0.89$ V vs. SCE) should be able to SET-reduce the protonated quinoline but not the unprotonated form. According to the Stern-Volmer quenching experiments it can be stated that none of the used components – phosphines **P1** and **P3** as well as the protonated quinoline **1a**-HCl – are quenching with a significant higher rate (Figure 3D). Therefore, an initial reductive quenching of the **PC** by the phosphine or an oxidative quenching by the protonated quinoline is feasible. In our initial work⁶⁹ we proposed a neophyl-type rearrangement for the reductive transformation of 2-monosubstituted quinolines through a radical intermediate in 3-position resulting in the extrusion of the C4 carbon atom. Contrary to this, Chiba and Pratt proposed a 3-exo cyclization of an intermediate radical at the 4-position, followed by ring-opening and reduction based on DFT-calculations and ¹³C-labeling.⁵² Considering the 2,3-disubstituted quinolines our results perfectly align with the proposed mechanism of Chiba and Pratt, as judged from the substitution pattern of the product indoles. Thus, the protonated quinoline is first reduced via SET from the **PC** to form an enamine radical intermediate **I-1**. Protonation leads to **I-2**, which sets the stage for the 3-exo cyclisation to generate **I-3**. Ring-opening then leads to the distonic radical cation **I-4** with upon HAT from the HOPAr₃-

radical and deprotonation eventually gives the rearranged indole **2** (Figure 3F).

To investigate, whether the reductive rearrangement of 2-monosubstituted quinolines may also be explained by the Pratt/Chiba mechanism, we prepared 2-phenylquinoline with D-incorporation in 3- as well as 4-position (Figure 3E). According to the proposed reaction mechanism, deuterium should remain incorporated in the molecule if it was connected to the C3 carbon atom, while the C4 D-labeling should get lost.⁷¹ We first subjected the non-labeled quinoline **1d** to our reaction conditions by using D₂O in place of H₂O. Interestingly we found a higher D-incorporation at the methyl-substituent than expected (78%, 66% expected), indicating that an H/D-exchange might take place under the reaction conditions along with the concluding deuterium atom transfer from the DOPAr₃ phosphoranyl radical to the indolylmethyl radical. This exchange presumably occurs after reduction at the enamine radical intermediate stage, indicating a reversible equilibrium between intermediates **I-1** and **I-2**. Along these lines, when **1d** was reacted with the **PC**, pTsoH, TripSH with D₂O under irradiation, a 33% D-incorporation was found over the positions 3 and 4 (27% mono- and 3% disubstituted) in the non-rearranged quinoline. When substrates **1d-3-D** and **1d-4-D** were subjected to the standard reaction conditions, a remaining deuterium incorporation of 18% was observed for **1d-3-D**, whereas nearly any deuterium was retained in **1d-4-D**, as expected based on the suggested mechanism. The labeling loss for **1d-3-D** is likely a result of the reversible protonation **I-1** in agreement with the studies above. Taken together, we suggest that the 3-exo-cyclization is also occurring for the reductive rearrangement of 2-monosubstituted quinolines. Moreover, these isotope labeling studies gave us the insight that the formed reduced intermediate **I-1** can likely be reversibly protonated under the applied conditions.






Figure 3: Mechanistic investigations including A) Deuteration experiments, B) Chemoselective protonation of the quinoline, C) Redox potentials, D) Stern-Volmer quenching study, E) Deuteration experiments with C2 substituted quinoline and F) Proposed Mechanism.

Conclusions

In summary, we developed a general reductive rearrangement of 2-monosubstituted and 2,3-disubstituted quinolines to indoles via extrusion of the carbon atom at the 3-position, applying a photocatalytic phosphine mediated water activation process. Moreover, the reaction proceeds without the need for

any pre-functionalization of the quinoline substrate, underscoring the practical simplicity of this method. The formed indoles could be isolated in moderate to very good yields and the reaction showed good functional group tolerance. Scalability of the reductive rearrangement was demonstrated. This method provides a valuable addition to the toolbox in the growing field of skeletal editing.



Author contributions

N.S., J.K., J.Z. and A.S. conceived all studies and experiments. N.S., J.K. and J.L. synthesized all compounds. CV measurements were conducted by E.S.H. All authors discussed the results and contributed to the preparation of the manuscript. All authors have given approval to the final version of the manuscript.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data (NMR, HRMS, IR, melting points, CVs) that support the findings of this study have been uploaded as part of the supplementary information.

Acknowledgements

We acknowledge the support from the European Research Council (advanced grant agreement no. 101140173), the Deutsche Forschungsgemeinschaft DFG (GRK 2678-437785492) and the National Research Foundation of Korea (RS-2025-14383014) for supporting this work.

The authors thank the AK Studer Water-Subgroup for scientific discussions, Timon Reckmann for NMR-support and Jan Henrik Deventer for experimental support (all University of Münster).

Notes and references

- 1 C. M. Marshall, J. G. Federice, C. N. Bell, P. B. Cox and J. T. Njardarson, *J. Med. Chem.*, 2024, **67**, 11622–11655.
- 2 E. Vitaku, D. T. Smith and J. T. Njardarson, *J. Med. Chem.*, 2014, **57**, 10257–10274.
- 3 R. D. Taylor, M. MacCoss and A. D. G. Lawson, *J. Med. Chem.*, 2014, **57**, 5845–5859.
- 4 I. J. P. De Esch, D. A. Erlanson, W. Jahnke, C. N. Johnson and L. Walsh, *J. Med. Chem.*, 2022, **65**, 84–99.
- 5 L. D. Pennington and D. T. Moustakas, *J. Med. Chem.*, 2017, **60**, 3552–3579.
- 6 L. D. Pennington, P. N. Collier and E. Comer, *Med. Chem. Res.*, 2023, **32**, 1278–1293.
- 7 In *Alkaloids*, Elsevier, 2015, pp. 63–102.
- 8 I. Kumar, R. Sharma and U. Sharma, *Org. Biomol. Chem.*, 2025, **23**, 2572–2585.
- 9 P. Bellotti, H.-M. Huang, T. Faber and F. Glorius, *Chem. Rev.*, 2023, **123**, 4237–4352.
- 10 C. M. Josephitis, H. M. H. Nguyen and A. McNally, *Chem. Rev.*, 2023, **123**, 7655–7691.
- 11 A. Corio, C. Gravier-Pelletier and P. Busca, *Molecules*, 2021, **26**, 5467.
- 12 H. Cao, Q. Cheng and A. Studer, *Science*, 2022, **378**, 779–785. DOI: 10.1039/D5SC08793G
- 13 B. T. Boyle, J. N. Levy, L. De Lescure, R. S. Paton and A. McNally, *Science*, 2022, **378**, 773–779.
- 14 G.-Q. Sun, P. Yu, W. Zhang, W. Zhang, Y. Wang, L.-L. Liao, Z. Zhang, L. Li, Z. Lu, D.-G. Yu and S. Lin, *Nature*, 2023, **615**, 67–72.
- 15 W. C. Wertjes, E. H. Southgate and D. Sarlah, *Chem. Soc. Rev.*, 2018, **47**, 7996–8017.
- 16 C. J. Huck and D. Sarlah, *Chem*, 2020, **6**, 1589–1603.
- 17 S. P. Roche and J. A. Porco, *Angew. Chem. Int. Ed.*, 2011, **50**, 4068–4093.
- 18 F. Lovering, J. Bikker and C. Humblet, *J. Med. Chem.*, 2009, **52**, 6752–6756.
- 19 F. Lovering, *Med. Chem. Commun.*, 2013, **4**, 515.
- 20 H. Adkins and H. R. Billica, *J. Am. Chem. Soc.*, 1948, **70**, 695–698.
- 21 W.-B. Wang, S.-M. Lu, P.-Y. Yang, X.-W. Han and Y.-G. Zhou, *J. Am. Chem. Soc.*, 2003, **125**, 10536–10537.
- 22 D.-W. Wang, X.-B. Wang, D.-S. Wang, S.-M. Lu, Y.-G. Zhou and Y.-X. Li, *J. Org. Chem.*, 2009, **74**, 2780–2787.
- 23 T. Wang, L.-G. Zhuo, Z. Li, F. Chen, Z. Ding, Y. He, Q.-H. Fan, J. Xiang, Z.-X. Yu and A. S. C. Chan, *J. Am. Chem. Soc.*, 2011, **133**, 9878–9891.
- 24 Z. Zhang and H. Du, *Org. Lett.*, 2015, **17**, 6266–6269.
- 25 X.-H. Hu and X.-P. Hu, *Org. Lett.*, 2019, **21**, 10003–10006.
- 26 S. J. Geier, P. A. Chase and D. W. Stephan, *Chem. Commun.*, 2010, **46**, 4884.
- 27 G. Erős, K. Nagy, H. Mehdi, I. Pápai, P. Nagy, P. Király, G. Tárkányi and T. Soós, *Chem. Eur. J.*, 2012, **18**, 574–585.
- 28 T. McCallum, S. P. Pitre, M. Morin, J. C. Scaiano and L. Barriault, *Chem. Sci.*, 2017, **8**, 7412–7418.
- 29 S. Adak, S. E. Braley and M. K. Brown, *Org. Lett.*, 2024, **26**, 401–405.
- 30 J. Zhang, N. Spreckelmeyer, J. Lammert, M. Wiethoff, M. J. Milner, C. Mück-Lichtenfeld and A. Studer, *Angew. Chem. Int. Ed.*, 2025, **64**, e202502864.
- 31 A. J. Birch, *J. Chem. Soc.*, 1944, 430.
- 32 A. J. Birch, *Nature*, 1946, **158**, 60–60.
- 33 D. Liu and J. Ma, *Angew. Chem. Int. Ed.*, 2024, **63**, e202402819.
- 34 B. K. Peters, K. X. Rodriguez, S. H. Reisberg, S. B. Beil, D. P. Hickey, Y. Kawamata, M. Collins, J. Starr, L. Chen, S. Udyavara, K. Klunder, T. J. Gorey, S. L. Anderson, M. Neurock, S. D. Minter and P. S. Baran, *Science*, 2019, **363**, 838–845.
- 35 J. Burrows, S. Kamo and K. Koide, *Science*, 2021, **374**, 741–746.



- 36 M. Szostak, M. Spain and D. J. Procter, *J. Org. Chem.*, 2014, **79**, 2522–2537.
- 37 M. Yasuda, C. Pac and H. Sakurai, *J. Org. Chem.*, 1981, **46**, 788–792.
- 38 A. Chatterjee and B. König, *Angew. Chem. Int. Ed.*, 2019, **58**, 14289–14294.
- 39 J. P. Cole, D.-F. Chen, M. Kudisch, R. M. Pearson, C.-H. Lim and G. M. Miyake, *J. Am. Chem. Soc.*, 2020, **142**, 13573–13581.
- 40 T. Yuan, L. Sun, Z. Wu, R. Wang, X. Cai, W. Lin, M. Zheng and X. Wang, *Nat. Catal.*, 2022, **5**, 1157–1168.
- 41 E. Y. K. Tan, A. S. Mat Lani, W. Sow, Y. Liu, H. Li and S. Chiba, *Angew. Chem. Int. Ed.*, 2023, **62**, e202309764.
- 42 J. Corpas, E. Rivera-Chao, E. M. Arpa, M. Gomez-Mendoza, Y. Katayama, V. A. De La Peña O'Shea, C. Bouchel, C. Jacob, P.-G. Echeverria, A. Ruffoni and D. Leonori, *Chem*, 2025, **11**, 102342.
- 43 A. K. Bains, A. Sau, B. S. Portela, K. Kajal, A. R. Green, A. M. Wolff, L. F. Patin, R. S. Paton, N. H. Damrauer and G. M. Miyake, *Science*, 2025, **388**, 1294–1300.
- 44 Y.-Z. Cheng, Z. Feng, X. Zhang and S.-L. You, *Chem. Soc. Rev.*, 2022, **51**, 2145–2170.
- 45 A. Shimose and Y. Nagashima, *Synlett*, 2025, **36**, 1111–1122.
- 46 E. H. Southgate, J. Pospech, J. Fu, D. R. Holycross and D. Sarlah, *Nat. Chem.*, 2016, **8**, 922–928.
- 47 J. Ma, S. Chen, P. Bellotti, R. Guo, F. Schäfer, A. Heusler, X. Zhang, C. Daniliuc, M. K. Brown, K. N. Houk and F. Glorius, *Science*, 2021, **371**, 1338–1345.
- 48 J. Ma, S. Chen, P. Bellotti, T. Wagener, C. Daniliuc, K. N. Houk and F. Glorius, *Nat. Catal.*, 2022, **5**, 405–413.
- 49 R. Kleinmans, S. Dutta, K. Ozols, H. Shao, F. Schäfer, R. E. Thielemann, H. T. Chan, C. G. Daniliuc, K. N. Houk and F. Glorius, *J. Am. Chem. Soc.*, 2023, **145**, 12324–12332.
- 50 P. Bellotti, T. Rogge, F. Paulus, R. Laskar, N. Rendel, J. Ma, K. N. Houk and F. Glorius, *J. Am. Chem. Soc.*, 2022, **144**, 15662–15671.
- 51 I. Kumar, S. S. Gupta, R. Kumar, R. Kumar, P. Agrawal, D. Sahal and U. Sharma, *ACS Sustain. Chem. Eng.*, 2020, **8**, 12902–12910.
- 52 E. Y. K. Tan, A. Dehdari, A. S. Mat Lani, D. A. Pratt and S. Chiba, *Chem*, 2024, **10**, 3722–3734.
- 53 F. Lin, T. Song, Y. Liu, X. Wang, S. Guo, Y. Sun and Q. Chen, *Angew. Chem. Int. Ed.*, 2025, **64**, e202513552.
- 54 B. W. Joynson and L. T. Ball, *Helv. Chim. Acta*, 2023, **106**, e202200182.
- 55 J. Jurczyk, J. Woo, S. F. Kim, B. D. Dherange, R. Sarpong and M. D. Levin, *Nat. Synth.*, 2022, **1**, 352–364.
- 56 R. Sharma, M. Arisawa, S. Takizawa and M. S. H. Salem, *Org. Chem. Front.*, 2025, **12**, 1633–1670.
- 57 G. L. Ciamician and M. Dennstedt, *Ber. Dtsch. Chem. Ges.*, 1881, **14**, 1153–1163.
- 58 M. Mortén, M. Hennum and T. Bonge-Hansen, *Beilstein J. Org. Chem.*, 2015, **11**, 1944–1949.
- 59 B. D. Dherange, P. Q. Kelly, J. P. Liles, M. S. Sigman and M. D. Levin, *J. Am. Chem. Soc.*, 2021, **143**, 11337–11344.
- 60 B. W. Joynson, G. R. Cumming and L. T. Ball, *Angew. Chem. Int. Ed.*, 2023, **62**, e202305081.
- 61 C. Kaneko, A. Yamamoto and M. Hashiba, *Chem. Pharm. Bull.*, 1979, **27**, 946–952.
- 62 C. Kaneko, H. Fujii, S. Kawai, A. Yamamoto, K. Hashiba, T. Kimata, R. Hayashi and M. Somei, *Chem. Pharm. Bull.*, 1980, **28**, 1157–1171.
- 63 T. Tsuchiya, S. Okajima, M. Enkaku and J. Kurita, *J. Chem. Soc., Chem. Commun.*, 1981, 211.
- 64 T. Tsuchiya, S. Okajima, M. Enkaku and J. Kurita, *Chem. Pharm. Bull.*, 1982, **30**, 3757–3763.
- 65 J. Woo, A. H. Christian, S. A. Burgess, Y. Jiang, U. F. Mansoor and M. D. Levin, *Science*, 2022, **376**, 527–532.
- 66 J. Woo, T. Zeqiri, A. H. Christian, M. C. Ryan and M. D. Levin, *J. Am. Chem. Soc.*, 2025, **147**, 20120–20131.
- 67 B. Roure, M. Alonso, G. Lonardi, D. B. Yildiz, C. S. Buettner, T. Dos Santos, Y. Xu, M. Bossart, V. Derdau, M. Méndez, J. Llaveria, A. Ruffoni and D. Leonori, *Nature*, 2025, **637**, 860–867.
- 68 Z. Wang, P. Xu, S.-M. Guo, C. G. Daniliuc and A. Studer, *Nature*, 2025, **642**, 92–98.
- 69 J. Zhang, C. Mück-Lichtenfeld and A. Studer, *Nature*, 2023, **619**, 506–513.
- 70 M. Wiethoff, L. Lezius and A. Studer, *Angew. Chem. Int. Ed.*, 2025, **64**, e202501757.
- 71 L.-F. Dai, Y.-X. Jiang, D.-L. Yu, G.-Q. Sun, S.-S. Yan, W. Zhang, J.-H. Ye and D.-G. Yu, *ACS Catal.*, 2025, **15**, 7792–7799.



Data availability The data (NMR, HRMS, IR, melting points, CVs) that support the findings of this study have been uploaded as part of the supplementary information.

