

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. Bera, T. Singha, N. A. Bapat and D. P. Hari, *Chem. Sci.*, 2026, DOI: 10.1039/D6SC00233A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Electrochemical Functionalization of Pyridines

Satabdi Bera,[†] Tushar Singha,[†] Nakul Abhay Bapat,[†] and Durga Prasad Hari*

Pyridines are important nitrogen-containing heteroaromatics with wide applications in pharmaceuticals, agrochemicals, ligands, and materials science. However, their selective C–H functionalization remains challenging due to low nucleophilicity, regioselectivity issues, and catalyst deactivation via nitrogen coordination. Traditional methods such as electrophilic substitution, directed metalation, transition-metal catalysis, and *N*-oxide transformations have enabled significant advances in this field, but often require harsh conditions or pre-functionalized substrates. In contrast, electrochemical synthesis has emerged as a sustainable and versatile alternative, employing electricity as a clean redox agent to enable direct C–H, C–C, and C–heteroatom bond formation under mild, tunable conditions. This review provides a focused overview of how electro-organic synthesis has addressed key challenges in pyridine functionalization associated with traditional methods, tracing its historical development and highlighting recent advances, with particular emphasis on anodic oxidation, cathodic reduction, paired electrolysis, and mediator or metal-catalyzed strategies, thereby offering readers an in-depth understanding of the advancements in this field and its significance in organic synthesis. Key mechanistic insights and representative applications are highlighted to demonstrate the growing potential of electrochemical strategies for scalable, late-stage pyridine diversification. We hope that this review will highlight the salient features of electrochemical strategies for pyridine functionalization in synthetic applications and stimulate the development of new, more advanced methodologies.

1. Introduction

Pyridines are among the essential nitrogen-containing heteroaromatic scaffolds widely found in pharmaceuticals,¹ agrochemicals,² ligands,³ and functional materials⁴ due to their electron-deficient aromatic ring, basic nitrogen atom, and excellent metal-coordination properties. Their prevalence in biologically active compounds¹ and catalytic systems³ has driven considerable interest in methods that allow direct and selective functionalization of the pyridine ring. Classical strategies, including electrophilic substitution,⁵ directed metalation with strong bases,^{6,7} transition-metal catalyzed cross-coupling^{8,9} or C–H activation,^{10,11,12} photoredox catalysis,¹³ and transformations via pyridine *N*-oxides,¹⁴ have enabled valuable transformations but often rely on harsh conditions, stoichiometric reagents, or pre-functionalized substrates, limiting their scalability and sustainability.

In recent years, electrochemical functionalization has emerged as a powerful and environmentally benign alternative to stoichiometric oxidants and reductants, employing electric current as a clean redox reagent to generate reactive species directly at the electrode surface under mild and tunable conditions.¹⁵ This strategy enables selective C–H functionalization of readily oxidizable electron-rich arenes, allowing C–C and C–heteroatom bond formation.^{16,17} However, the efficient C–H or C–X bond modification of electron-deficient pyridines remains significantly more challenging. Owing to the operational simplicity of electro-organic synthesis, broad functional-group tolerance, and precise control over redox

events through constant-current or constant-potential electrolysis, electrochemical methods have emerged as a powerful platform for the late-stage diversification of pyridine-containing molecules in both academic and industrial settings (Scheme 1A).

Although several reviews have discussed the electrochemical functionalization of *N*-heterocycles in general, only a limited number of pyridine examples are included. While these reviews provide valuable insights into broader electrochemical strategies, they primarily focus on other (hetero)arenes (e.g., phenols, indoles, pyrazoles, and imidazoles) and on specific pyridine transformations, such as C2 functionalization via activated pyridinium species and the functionalization of cyano-substituted pyridines.¹⁸ However, numerous recent reports have expanded the scope to enable regioselective functionalization at C2, C4, and C3/C5 positions, highlighting the evolving reactivity patterns of pyridine substrates that are not comprehensively captured in existing reviews. Due to their electron-deficient nature and the presence of a coordinating nitrogen atom, pyridines present unique challenges, including site-selective functionalization (C2 vs C4 vs C3/C5), susceptibility to overreduction, and catalyst deactivation through strong *N*-coordination (Scheme 1B). In electrochemical systems, these factors are closely linked to reduction potentials and pyridine activation modes, which critically influence reaction pathways and method selection. Therefore, a dedicated pyridine-focused perspective enables a more systematic understanding of these features and provides practical guidance for electrochemical functionalization of pyridines.

Building on this perspective, this review surveys recent advances in electrochemical methods for pyridine functionalization, organized according to the underlying

^a Department of Organic Chemistry, Indian Institute of Science, Bangalore, India, 560012.

[†] Authors contributed equally.




electrochemical logic. Specifically, the discussed strategies are classified into three broad categories: net-anodic oxidation, net-cathodic reduction, and paired electrolysis, each exploiting the unique redox properties of pyridine and its derivatives. We discuss both direct electrolysis, where the substrate itself undergoes electron transfer at the electrode surface, and mediator-assisted processes, in which redox-active species facilitate selective transformations. The merging of photocatalysis and transition-metal catalysis with electrochemical approaches is also discussed, wherein the synergy of transition metals with electrochemical conditions is shown to enable challenging C–H, C–C, and C–heteroatom bond formations. Key reaction parameters, including the choice of electrolyte, electrode material, solvent, and the application of either constant current or constant voltage, are analyzed, as these factors drastically influence reaction efficiency, selectivity, and functional group tolerance. Furthermore, selected examples of functionalized pyridine derivatives are presented, illustrating the versatility and synthetic utility of electrochemical approaches for the late-stage diversification of this important heterocyclic scaffold. Finally, the mechanistic aspects of representative transformations are explored to provide insight into electron-transfer pathways, the formation of radical or cationic intermediates, and regioselectivity.

This review is organized to first highlight the early examples of electrochemical pyridine functionalization, followed by sections based on the key electrochemical processes: (i) net oxidation, (ii) net reduction, and (iii) paired electrolysis (Scheme 1C). Within each process-oriented section, transformations are further categorized by the type of bond formed, including C–C,

C–N, C–O, C–B, C–Si, C–S, and annulation reactions. At the end of each major electrochemical process, the choice of reaction parameters, such as electrode materials, electrolyte and electrochemical control mode, etc., is summarised to inform rational reaction design for pyridine activation and regioselectivity control. Finally, the current challenges and future directions in the field are discussed. We hope this review will shed light on the key features of electrochemical pyridine functionalization for synthetic applications and inspire the development of more advanced methodologies.

2. Early Advancements

The first example of electrochemical pyridine functionalization was reported by Inoue and his coworkers in 1973 (Scheme 2a).¹⁹ They demonstrated the electrochemical fluorination of pyridines, which predominantly resulted in perfluorinated derivatives. Subsequently, in 1975, Haszeldine and his coworkers extended this approach to alkyl-substituted pyridines, achieving fluorination that primarily produced mixtures of perfluoro-(*N*-fluoropiperidine) derivatives under constant voltage electrolysis (Scheme 2b).²⁰ A significant milestone was achieved by Ballinger and Teare, who reported the first electrochemical monofluorination of pyridine, affording 2-fluoropyridine in 22% yield.²¹ Performing the electrolysis at a constant voltage of 2.5 V using platinum electrodes was key for the successful monofluorination of pyridine with Me₄NF·2HF serving as both the fluorine source and the supporting electrolyte. The reaction efficiency was highly sensitive to the



Scheme 1: A) Distribution of number of publications year-wise. B) Challenges associated with pyridine functionalization. C) Organization of the review.




applied potential and electrolyte concentration, as higher voltages led to pyridine decomposition (Scheme 2c). These studies collectively laid the groundwork for electrochemical fluorination of pyridines, highlighting the critical influence of reaction parameters on selectivity and product distribution.

Building upon these early fluorination efforts, researchers soon began exploring alkylation strategies to expand the functional diversity of pyridine frameworks. In this context, in 1976, Holman and co-workers reported the electrochemical methylation and trifluoromethylation of pyridine by electrolyzing aqueous pyridine solutions containing acetic acid or trifluoroacetic acid, thereby introducing a sustainable pathway for alkyl-functionalized pyridines (Scheme 2d).²²

The electrochemical coupling of pyridines has evolved remarkably over the past decades, progressing from early homocoupling studies to highly selective cross-coupling methodologies. The first electrochemical homocoupling of pyridines was reported in 1993 by Yanilkin and co-workers using a Ni-catalyst and 2,2'-bipyridine as the ligand (Scheme 2e).²³ In this pioneering study, electrochemically generated Ni(0) undergoes oxidative addition with 2-bromopyridine to produce Ni(II)-Py species. This intermediate is then reduced at the cathode to a Ni(I)-pyridyl species, which undergoes another cycle of oxidative addition followed by reductive elimination, ultimately furnishing the dimerized product. Building upon this foundation, in 1995, Amatore and co-workers reported the first electrochemical cross-coupling of pyridines, utilizing

dihalo-pyridines and 2,6-di-*tert*-butylphenol as the arylating reagent to achieve C–C bond formation under electrochemical conditions (Scheme 2f).²⁴ This study opened a new avenue for expanding the reactivity landscape of pyridines beyond homocoupling. The field advanced further in 2000, when the electrochemical cross-coupling of pyridines with aryl halides was achieved, tolerating a wide variety of pyridine, aryl, and heteroaryl halides to deliver the corresponding arylated pyridines in good to excellent yields (Scheme 2g).²⁵ Continuing this progress, Navarro and co-workers further modified the homocoupling protocol by employing NiBr₂(bpy) as a catalyst and zinc as a sacrificial anode in DMF or acetonitrile, successfully synthesizing 2,2'-bipyridine and dimethyl-2,2'-bipyridines from 2-bromopyridine and 2-bromomethylpyridines, respectively (Scheme 2h).²⁶ Further optimization of nickel catalysts and ligand systems enabled the controlled arylation of polyhalogenated pyridines with improved turnover numbers and selectivity.²⁷ Later, the same group significantly enhanced homocoupling efficiency by introducing an iron anode, which increased yields to 98% by preventing nickel catalyst passivation through iron-ion-mediated trapping of free bipyridine generated during sacrificial anode oxidation.²⁸ Collectively, these advances delineate a clear trajectory from the foundational Ni-catalyzed homocoupling to modern, efficient, and selective cross-coupling methodologies, underscoring the growing maturity and synthetic potential of electrochemical strategies in pyridine chemistry.



Scheme 2: Early examples of electrochemical functionalization of pyridines.



Expanding the scope of electrochemical pyridine functionalization beyond halide coupling, Gosmini and co-workers achieved a landmark advancement by reporting the first electrochemical C–H arylation of pyridines, utilizing in situ-generated organozinc reagents as arylating partners in the presence of a copper catalyst (Scheme 2i).²⁹ The organozinc intermediate coupled with pyridinium salts, which is generated in situ from pyridines and methyl chloroformate. The resulting 1,4-dihydropyridine intermediates were then oxidized to furnish 4-phenylpyridines in moderate to high yields. This study not only demonstrated the feasibility of direct C–H arylation under electrochemical control but also highlighted the synthetic versatility of organozinc intermediates in pyridine derivatization.

Nédélec and co-workers developed a C–P bond-forming strategy by employing chlorodiphenylphosphines as phosphorylating agents in the presence of a nickel catalyst, which opened an avenue for C-heteroatom bond formation in pyridine functionalization (Scheme 2j).³⁰


3. Functionalization via Net Oxidation

Electrochemical net-oxidative strategies have emerged as powerful and sustainable approaches for the formation of C–C and C–heteroatom bonds. These methodologies operate through sequential single-electron transfer (SET) oxidation processes, enabling substrate activation without the need for stoichiometric chemical oxidants. By precisely controlling

anodic potentials, electrochemical oxidation facilitates radical–polar crossover pathways and grants access to valuable molecular architectures that are often difficult to obtain using conventional oxidative methods. Among various net oxidative approaches, electrochemical C–H bond functionalization has gained prominence, as it enables regioselective activation and site-selective modification of heteroaromatic systems such as pyridines, thereby providing an efficient route to complex, functionally diverse compounds.


3.1 C–H Alkylation

Within oxidative C–H functionalization, C–H alkylation of pyridines represents a significant advancement. In this strategy, alkyl radicals generated either directly at the anode or through mediator-assisted oxidation add to pyridinium intermediates to furnish dihydropyridine derivatives. A subsequent anodic oxidation step restores aromaticity, thereby providing the corresponding alkylated pyridines. In 2020, an electrophotocatalytic Minisci-type functionalization of pyridines was demonstrated by Xu and co-workers using a decarboxylative alkylation strategy (Scheme 3).³¹ Two distinct optimized conditions were developed for C–H alkylation and C–H carbamoylation, with $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ and 4CzIPN being employed as photocatalysts, respectively. The alkylation of 2,6-disubstituted pyridine was well tolerated to furnish the product **1**; however, C4-substituted pyridines were found to yield a mixture of mono and bis-alkylated products (**2** and **3**). Notably, pyridine-containing drug molecules such as



Scheme 3: A) C–H alkylation. B) C–H carbamoylation.





Scheme 4: Electrochemical Mn-Catalyzed C-H bond arylation using aryl Grignard reagents.

diflufenican were also compatible, delivering the desired alkylated product **4** in 31% yield. Conversely, the carbamoylation reaction exhibited remarkable selectivity at the C2-position of pyridines and pyridine-containing drug-like molecules, affording mono-substituted products (**5-7**). Mechanistically, the alkylation reaction was initiated *via* anodic oxidation of Ce^{III} to Ce^{IV}, which coordinated with carboxylic acid to form intermediate **I**. The alkyl radical **III** was then proposed to be generated *via* a photoinduced ligand-to-metal charge transfer (LMCT) in the Ce(IV) complex **I**, followed by decarboxylation. After the addition of radical **III** to the activated pyridinium salt, the resulting radical intermediate **IV** was suggested to lose a proton to generate intermediate **V** and subsequently undergo oxidation mediated by Ce(IV) to furnish the alkylated product. However, for carbamoylation, the radical **I** was generated *via* a single-electron transfer (SET) oxidation of oxamate by the excited-state 4CzIPN. The radical **II** generated after the Minisci-type addition to pyridinium salt was first reduced by the photocatalyst to give intermediate **III**, while regenerating the photocatalyst. Subsequent oxidation of **IV** at the anode to yield the carbamoylated product. Finally, the cathodic reaction was taken care of by the reduction of protons to produce hydrogen gas.

3.2 C-H Arylation

Heterobiaryls represent valuable structural motifs found in numerous biologically relevant molecules and advanced functional materials.³² Consequently, the development of efficient and sustainable strategies for their synthesis has attracted considerable attention. In line with recent advances in electrochemical methodologies, the direct oxidative C-H arylation of pyridines has been successfully demonstrated in the synthesis of heterobiaryls.


Ackermann and co-workers reported an electrochemical manganese-catalysed strategy for the oxidative C-H arylation of pyridines bearing weakly coordinating amide directing groups, employing Grignard reagents as aryl sources (Scheme 4).³³ This approach circumvented the need for stoichiometric oxidants and zinc salts, which are typically required in iron-catalyzed systems to prevent Grignard homocoupling. The electrolysis was performed under a constant current of 3 mA with MnCl₂ as the catalyst, neocuproine as the ligand, and

TMEDA as an additive in THF solvent. The versatility of this electrocatalytic protocol was demonstrated through its broad substrate scope, accommodating various substituted pyridines and a range of arylating agents, including heteroaryl counterparts (products **9-11**). Moreover, the strategy was successfully extended to the methylation of pyridine (**12**), underscoring its synthetic utility and generality. Mechanistic investigations, supported by cyclic voltammetry, scanning electron microscopy coupled with energy-dispersive X-ray spectroscopy (SEM-EDS), and DFT calculations, revealed that C-H activation proceeds through a high-spin ligand-to-ligand hydrogen transfer (LLHT) pathway facilitated by single-electron transfer at the anode. The reaction was initiated through the formation of the active Mn(II) intermediate **I** in the presence of ligands and the arylating reagent (ArMgBr). This catalytically active Mn(II) species subsequently underwent C-H bond activation of pyridine to form the cyclometalated intermediate **II** via the LLHT process, followed by transmetalation to generate intermediate **III**. Subsequent anodic oxidation produces intermediate **IV**, which then undergoes reductive elimination to furnish the arylated pyridine. The resulting Mn(I) species was reoxidized at the anode to regenerate the Mn(II) catalyst, demonstrating that manganese electrocatalysis could effectively activate the C-H bond of pyridines with weakly coordinating amides while suppressing overoxidation and homocoupling.

3.3 C-H Borylation

Xu and co-workers recently developed a method for direct C-H borylation of pyridines by merging electrochemistry and photocatalysis, using a borane-trimethylamine (**13**) complex as the borylating agent (Scheme 5).³⁴ The main objective of this transformation was to enable efficient generation of boryl radicals under mild conditions, without the need for stoichiometric chemical oxidants. This was achieved using the photocatalyst anthraquinone-2-sulfonate (AQS) via hydrogen-atom transfer. This strategy proved effective for C4-substituted pyridines, 2,6-disubstituted pyridines, and 2,3,4-trisubstituted pyridines, all of which underwent smooth conversion to the products (**14-16**). Upon photoexcitation, the excited anthraquinone sulfonate (AQS) abstracted a hydrogen






Scheme 5: C-H borylation.

atom from the borane–trimethylamine complex (**13**), generating a boryl radical **I** along with a semiquinone-type radical species (AQS-H $\dot{\text{C}}$). The boryl radical **I** subsequently underwent Minisci-type addition to the pyridinium salt **II** to produce a radical cation intermediate **III**, which underwent deprotonation and further oxidation to yield the borylated product **17**. The AQS-H $\dot{\text{C}}$ species underwent anodic oxidation

followed by deprotonation, regenerating the ground-state AQS and thereby completing the catalytic cycle.


3.4 C-H Amination

An electrochemical direct amination of pyridines has also been developed under mild and sustainable conditions. In 2018, Mei and co-workers reported a copper-catalyzed electrochemical C–H amination of pyridines equipped with a directing group (Scheme 6).³⁵ The use of ¹⁸Bu₄NI as a redox mediator was key to the success of this protocol. Constant-current electrolysis was carried out in an undivided cell employing Cu(OTf)₂ as the catalyst and KOiPr as the base. The substrates bearing picolinamide (NHCO-2-Py) moiety served as an effective bidentate directing group, facilitating coordination of both the amide and *ortho*-pyridyl N atoms to the copper center and enabling site-selective C–H amination under electrochemical conditions. A wide range of pyridine derivatives was compatible with this method, tolerating both electron-donating (**19** and **20**) and electron-withdrawing (**21**) substituents, with selective amination at the C2 position. Mechanistic studies, including kinetic isotope effect measurements, cyclic voltammetry, and radical inhibition experiments, suggested that a high-valent Cu(III) species is involved as a key intermediate. The reaction was believed to be initiated by the coordination of the Cu(II) catalyst with the amine nucleophile **18** and substrate **22**, leading to the formation of a copper(II) complex **I**. This complex was subsequently oxidized by an iodine radical to afford the Cu(III) intermediate **II**, which was proposed to undergo a ligand-to-metal charge transfer (LMCT) process to give radical cation **III**. The intramolecular amine transfer to radical cation **III**, followed by a second LMCT event, yielded intermediate **IV**. The aminated product **19** was then released, and a Cu(I) species was formed. Finally, iodine radical-mediated anodic oxidation of Cu(I) was proposed to regenerate the active Cu(II) catalyst, thereby completing the catalytic cycle.



Scheme 6: C-H amination.






Scheme 7: C-H amination using reduced selectfluor II.

Very recently, Malapit and co-workers reported a directing-group-free electrochemical C2-amination of pyridines using Selectfluor II (**23**) as the aminating reagent (Scheme 7).³⁶ Under optimized electrochemical conditions, the desired aminated products (**24–26**) were obtained in good to excellent yields. The transformation proceeded through a two-step sequence: (i) electrochemical amination to generate DABCONium salts, followed by (ii) neutralization with KCN to

afford the corresponding aminated or piperazine derivatives. Mechanistically, the reaction begins with anodic oxidation of reduced Selectfluor II (**23**), forming the radical dication **I**, which subsequently adds to the pyridine substrate via a charge-transfer complex **II**, generating the intermediate **III**. Further anodic oxidation and proton elimination yield the DABCONium salt **IV**. Finally, nucleophilic substitution (S_N2) of this intermediate with KCN furnishes the piperazine derivative **27**. The high C2-selectivity is attributed to the strong electrophilicity and steric demand of the nitrogen-centered radical dication **I**, favoring regioselective addition at the C2-position of the pyridine ring.

3.5 C-H Sulfonylation

Another noteworthy transformation is the electrochemical sulfonylation of pyridines, which enables direct introduction of sulfonyl groups onto the pyridine core and broadens the repertoire of sulfur-containing derivatives.³⁷ In 2024, Yi and co-workers reported a *meta*-selective electrochemical sulfonylation of pyridines proceeding via a dearomatized oxazino-pyridine species (Scheme 8).³⁸ Electrolysis of the oxazino-pyridine and sulfinate salt was carried out in an undivided cell with a graphite felt (GF) anode and Pt cathode under constant-current conditions using $^n\text{Bu}_4\text{NBF}_4$ as the supporting electrolyte. Subsequent acidic treatment furnished the desired *meta*-sulfonylated pyridines. This method proved broadly applicable to a wide range of sulfonyl groups, including (hetero)aryl sulfinates and alkyl sulfinates (products **28–30**). The protocol was also successfully extended to late-stage functionalization of drug-like molecules containing pyridine motifs, such as vismodegib (product **31**). Mechanistic investigations indicated that preferential anodic oxidation of the sulfinate over the oxazino-pyridine **32** was crucial for generating the sulfonyl radical. The sulfonyl radical **I** was selec-



Scheme 8: C-H sulfonylation.




REVIEW

Chemical Science

tively added at the δ -nitrogen site of the oxazino-pyridine **32**, yielding a key intermediate **II** that undergoes further anodic oxidation to generate intermediate **III**. Subsequent deprotonation and acid-mediated rearomatization afforded the *meta*-sulfonylated product.

3.6 C-H Silylation

Direct C–H silylation represents a powerful strategy in modern organic synthesis, as the resulting silylated products serve as versatile intermediates for further synthetic elaboration.³⁹ In 2023, Wang and co-workers reported an elegant example of site-selective silylation under mild electrophotochemical conditions, employing 9,10-phenanthrenequinone (PQ) as both a photocatalyst and a hydrogen-atom-transfer (HAT) agent (Scheme 9).⁴⁰ The protocol exhibits broad substrate tolerance, accommodating variously monosubstituted pyridines (**34** and **35**). Silylated products (**36** and **37**) were efficiently obtained from 2,6- and 2,4-disubstituted pyridines. Mechanistically, the electrophotochemical C–H silylation begins with photoexcitation of PQ to PQ*, followed by HAT from ^tBuMe₂SiH to generate the silyl radical **I** (^tBuMe₂Si•) and PQH•. The silyl radical **I** is added to the pyridinium salt **II**, forming a key intermediate **III**, which subsequently transfers a hydrogen (HAT) atom to PQH•, followed by deprotonation to yield the silylated pyridine **38**. Finally, PQ is regenerated by anodic oxidation of PQH₂.



Scheme 9: ortho-C-H silylation.

An elegant strategy for the *meta*-C–H silylation of pyridines was recently disclosed by Xu and co-workers (Scheme 10).⁴¹ The transformation proceeded without a photocatalyst or external oxidants, relying instead on *N*-hydroxyphthalimide (NHPI) as a hydrogen-atom-transfer (HAT) catalyst. Constant current electrolysis in acetonitrile at room temperature

efficiently generated silyl radicals in situ, which selectively engaged with the pyridine ring. Notably, the presence of an electron-withdrawing substituent on the pyridine ring was found to enhance *meta*-selectivity, dictating the role of electronic effects in controlling regioselective C–H bond functionalization. While 2,6-dicyanopyridine furnished both mono- and disilylated products (**39** and **40**), 2,4,6-trisubstituted pyridine exclusively delivered the monosilylated product **41**, reflecting the impact of steric effects on second silylation. In addition, the use of bulkier silanes, such as triisopropylsilane was compatible with the reaction, affording the corresponding *meta*-silylated product **42** in 55% yield. A plausible reaction pathway involves single-electron oxidation of NHPI, followed by proton elimination to generate the *N*-oxyl radical **I**, which abstracts a hydrogen atom from silane, generating silyl radical **II**. The silyl radical **II** then preferentially adds to the *meta* position of intermediate **III**, affording intermediate **IV**. Finally, *N*-oxyl radical **I** abstract a hydrogen atom from intermediate **IV**, regenerating NHPI and delivering the *meta*-silylated product **39**. The observed *meta*-selectivity is attributed to a combination of electronic and stabilization effects. Specifically, the strong electron-withdrawing inductive effect of cyano or ester substituents enhances the electrophilic character of the *meta* position relative to the *ortho* and *para* positions.



3.7 C-H Halogenation

Halopyridines are highly important intermediates in organic synthesis and medicinal chemistry due to their versatile reactivity and broad applications. They also serve as key building blocks in the synthesis of pharmaceuticals, functional materials, and agrochemicals, enabling precise modification of the pyridine scaffold to tune biological activity or material properties.^{10, 42} Therefore, the electrochemical halogenation of pyridines opened another avenue for their efficient synthesis. In 2016, Gryaznova and co-workers reported a selective electrochemical fluorination of pyridines and ethylpyridines mediated by transition metal catalysts (Scheme 11).⁴³ The regioselectivity of the transformation was found to depend strongly on the metal catalyst employed: nickel catalysts favored fluorination at the C3-position, whereas silver and cobalt salts preferentially promoted C2-fluorination. The methodology was demonstrated on two representative substrates, affording the corresponding fluorinated products (**44–47**) in good yields. The observed catalyst-dependent regioselectivity likely arises from distinct oxidative fluorination pathways, the mechanistic details of which remain to be fully elucidated.

In 2021, Chen and co-workers reported an electrochemical *meta*-C–H bromination of pyridines via an umpolung strategy (Scheme 12).⁴⁴ Selective *meta*-functionalization was achieved by introducing an electron-donating group at the *ortho* position to enhance nucleophilicity at the *meta* site, and electrophilic bromine was generated in situ from tetrabutylammonium bromide (**48**). A broad substrate scope was displayed by the optimized bromination protocol, with both electron-donating (Me) and halogen (F) substituents being tolerated to deliver the products (**49** and **50**). When the




cyclohexylamino (NHCy) group was used, the corresponding brominated pyridine **51** was obtained in 73% yield. Remarkably, preinstallation of methoxy groups at the 2- and 6-positions of the pyridine led to a highly efficient electrolysis, delivering product **52** in an isolated yield of 95%. CV studies indicated that bromine radical is generated via oxidation of bromide, while the 2-aminopyridine alone was found to be electrochemically inactive. Based on these findings and prior reports, a plausible mechanism was proposed in which the bromide ion was anodically oxidized to Br-radical I, which then forms molecular bromine and undergoes electrophilic addition on the amino-pyridine to yield intermediate II. Subsequent deprotonation furnished the *meta*-brominated product **54**.

Scheme 10: *meta*-C-H silylation.

Scheme 11: Transition-metal catalyzed C-H fluorination.

3.8 Annulation of Pyridine Derivatives

Indolizines are found in biologically active natural products and pharmaceuticals.⁴⁵ They exhibit a broad spectrum of activities, including anti-inflammatory, antimicrobial, anti-tubercular, and anticancer properties^{45b, 46} and have recently attracted attention for applications in dye-sensitized solar cells (DSSCs) and organic light-emitting devices (OLEDs).⁴⁷ In 2022, Sharma and co-workers developed a greener strategy for indolizine synthesis via an electrochemical cascade C–H functionalization and annulation (Scheme 13).⁴⁸ Treatment of pyridines with 2-methylquinazolinone derivatives and chalcone derivatives in an undivided electrochemical cell under constant current (10 mA) efficiently furnished the corresponding annulated indolizine **55**. Notably, the replacement of chalcones with isoxazole-containing styryl-alkene also afforded indolizine **56** in good yield. Based on experimental evidence and prior literature, the mechanism was believed to be initiated by anodic oxidation of iodide to generate an iodine radical, which reacted with quinazolinone **57** to form an intermediate I via HAT, followed by recombination of the heterobenzyl radical with the iodine radical. Pyridine then reacted with I to form intermediate II, which underwent deprotonation followed by an intermolecular cycloaddition with chalcone yield intermediate III. Finally, anodic oxidation of intermediate III restored aromaticity, thereby forming the heterocyclic indolizine product **58**.




Scheme 12: C-H bromination.

Very recently, electrochemical annulation of pyridines was reported by Terent'ev and co-workers for the synthesis of imidazo[1,5-a]pyridines using NH_4SCN as a cyanating agent (Scheme 14).⁴⁹ A wide range of pyridine-2-carboxaldehydes and benzylamines, including chloro-substituted and heteroaryl benzylamines, was compatible and afforded the desired products (**59–63**) in good to moderate yields. Control experiments suggested that cyanide species were generated prior to core formation, with no involvement of free




thiocyanogen. Mechanistically, anodic oxidation of thiocyanate in the presence of water produces the cyanide anion, which attacks the imine intermediate **I** formed from pyridine-2-carboxaldehyde **64** and benzylamine **65**, generating intermediate **II**. Subsequent oxidation of


intermediate **II** with DMSO or Shono-type anodic oxidation pathway, furnishes intermediate **III**, which isomerizes to **IV**. Finally, intramolecular cyclization of intermediate **IV**, followed by oxidation and deprotonation, yields the final product, 1-cyano-imidazo[1,5-a]pyridine **59**.



Scheme 13: Synthesis of indolizines via C-H functionalization of pyridine.



Scheme 14: Oxidative annulation of pyridine.



Scheme 15: Dearomative stereoselective multifunctionalization.

3.9 Dearomative Functionalization

The 'escape from flatland' concept has been applied to pyridines to transform their planar aromatic cores into three-dimensional (3D) dehydropiperidine scaffolds with increased molecular complexity.⁵⁰ Replacing a pyridine core in



pharmaceuticals with an sp^3 -enriched piperidine counterpart can improve solubility and bioavailability.⁵¹ Motivated by this strategy, Ye and co-workers have very recently developed a regio- and diastereoselective dearomative multifunctionalization of pyridines using TMS-halide, TMSCN and alcohol under constant current electrolysis (Scheme 15).⁵² The method exhibits a broad substrate scope across various pyridine derivatives and alcohol nucleophiles; chloro- and bromo-silanes can be employed to facilitate dearomatization with moderate to excellent diastereoselectivity (products **66-68**). The mechanism of this transformation begins with the activation of the pyridine **8** by cyanogen bromide, which is generated electrochemically in situ from TMSBr and TMSCN. The activated pyridinium intermediate **I** is then trapped by methanol to form the dearomative intermediate **II**. Subsequently, an anodically generated bromine radical adds to the nucleophilic enamine double bond in a regio- and diastereoselective manner, affording the radical intermediate **III**. Further anodic oxidation of radical **III** leads to the formation of the iminium ion **IV**, which is subsequently trapped by a nucleophile to afford the final product. The high diastereoselectivity is supported by DFT calculations, which indicate that the addition of the bromine radical to the enamine from the sterically less hindered face is energetically favored.

Collectively, net-oxidative electrochemical strategies for C–H bond functionalization of pyridines offer an attractive and sustainable alternative to conventional methods. In most reported systems, acidic additives are deliberately chosen to generate more electron-deficient pyridinium species, which enhance regioselective radical addition by increasing the electrophilicity of specific positions on the pyridine ring. Polar aprotic solvents (such as MeCN) are typically selected to stabilize charged intermediates and to buffer proton release during the rearomatization step. From a practical standpoint, robust electrode materials such as reticulated vitreous carbon (RVC), platinum, carbon, and graphite felt (GF) are preferred due to their stability under oxidative conditions and corrosion resistance; at the cathode, these electrodes also enable efficient proton reduction, thereby maintaining charge balance throughout the electrochemical process.

4. Functionalization via Net Reduction

In contrast to the extensively studied net oxidative pathways, net reductive electrochemical methods provide an equally versatile and sustainable platform for pyridine functionalization. These approaches proceed via distinct mechanistic pathways that involve radical or anionic intermediates generated by cathodic reduction. Therefore, pyridine functionalization can also be achieved efficiently via cathodic activation of C–X (X = H, Cl, Br, CN) bonds of pyridines, generating reactive intermediates that facilitate efficient C–C and C–heteroatom bond formation with high regioselectivity and broad functional group tolerance.


4.1 C-H Deuteration

Due to the pronounced kinetic isotope effect, deuterated compounds are of great significance in synthetic chemistry,

materials science, and pharmaceutical research.⁵³ Although C–H/C–D exchange provides a straightforward route for deuterium introduction,⁵⁴ traditional methods, such as transition-metal-catalysed exchange⁵⁵ or deuterodehalogenation⁵⁶ typically require noble metals, strong acids/bases, or organometallic reagents, and often suffer from limited regioselectivity, low incorporation efficiency, and a restricted substrate scope. In this context, Xiang and co-workers recently reported an electrocatalytic γ -selective deuteration of pyridines using D_2O as the deuterium source (Scheme 16).⁵⁷ A broad range of substituted pyridines bearing aryl and heteroaryl groups afforded the desired deuterated products (**69-71**) in excellent yields. The practicality of this strategy was further demonstrated through the late-stage deuteration of a pyridine-containing estradiol derivative **71**. Mechanistically, the cathodic reduction of 2-phenyl pyridine **72** produced the pyridyl radical anion **I**, which underwent nucleophilic attack to D_2O at the preferred C4-position to form intermediate **II**. Subsequent oxidation by molecular iodine or molecular oxygen afforded the deuterated pyridine **69**. Molecular iodine is generated in situ from iodide at the anode.

4.2 C-H Carboxylation

Direct carboxylation of organic molecules using CO_2 has emerged as a powerful strategy for the incorporation of carboxylic acid functionalities.⁵⁸ In the context of pyridines, such transformations provide straightforward access to picolinic acid and nicotinic acid derivatives, which serve as valuable building blocks in pharmaceuticals and drug discovery.⁵⁹ The first study in this area was reported by Zare and coworkers, who demonstrated that a nickel complex could efficiently catalyze the electrochemical reduction of CO_2 , generating a carboxylate radical anion as a key reactive intermediate.⁶⁰ This species not only facilitated the indirect electroreduction of pyridines but also coupled directly with



Scheme 16: C-H deuteration.



REVIEW

Chemical Science

pyridyl radical anions, leading to the formation of the corresponding carboxylated pyridine derivatives. However, this reaction is limited to mechanistic studies; its synthetic potential is not explored.


In 2022, Qiu and co-workers made a significant breakthrough by reporting the first site-selective C4-carboxylation of pyridines utilizing CO₂ as an inexpensive and sustainable carbon source (Scheme 17).⁶¹ This protocol is a metal-free, base-free and allows selective C–H carboxylation of electron-deficient pyridines, affording the products (**73–75**) with excellent chemo- and regioselectivity. Notably, 2,6-difluoropyridine was converted to 2,6-difluoronicotinic acid (**73**) in 76% yield rather than to 2,6-difluoroisonicotinic acid, which was likely influenced by the electron-withdrawing effects of the fluorine atoms. Similarly, 2,6-diphenylpyridine was transformed into 2,6-diphenylisonicotinic acid (**74**) under the same conditions. The reaction was initiated by the cathodic reduction of 2-phenyl pyridine **72**, leading to the formation of a pyridyl radical anion **I**, which was subsequently added to CO₂ to generate intermediate **II**. The high regioselectivity observed was attributed to the intrinsic electronic properties of the substrate. Intermediate **II** was then oxidized by iodine to produce **III**, accompanied by the loss of a proton. Finally, protonation afforded the carboxylic acid product **76**. Meanwhile, Et₄N⁺I⁻ was oxidized at the anode, obviating the need for a sacrificial anode.

Concurrently, Lin, Yu and co-workers developed an electrochemical approach that offers complementary site selectivity, in which regioselectivity is governed entirely by the cell configuration: C5-carboxylation is favored in divided cells, whereas C4-carboxylation predominates in undivided cells (Scheme 18).⁶² This strategy was successfully applied to a broad range of pyridines (**77–80**), bipyridine (**81**), and furan-substituted pyridine (**82**), demonstrating its wide applicability. For C5-carboxylation, the reaction is proposed to initiate via single-electron reduction of 2-phenyl pyridine **72**, generating radical anion **I**. DFT calculations indicated that the highest electron density of **I** is located at C5, which undergoes nucleophilic addition to CO₂ to form intermediate **II**. Subsequently, cathodic reduction produces dianion **III**, which is then oxidatively rearomatized by O₂ and methylated in the presence of methyl iodide to furnish the corresponding methyl ester derivative **83**. However, the addition of CO₂ to intermediate **I** is reversible and not energetically favorable (endergonic). This reversibility allows the product ratio to depend on the next irreversible step according to the Curtin–Hammett principle. Additionally, DFT calculations showed that if CO₂ were to add at the C4 position instead, the resulting intermediate would have a weaker C–H bond at that site (by approximately 3.8 kcal/mol), thereby facilitating a hydrogen-atom transfer (HAT) process. So, if a hydrogen-atom acceptor were present, the reaction could be pushed toward C4-carboxylation instead of C5. Under undivided-cell conditions, iodine generated anodically from tetrabutylammonium iodide (tBu₄N⁺I⁻) functioned as an efficient hydrogen-atom acceptor, engaging in either a direct H-atom transfer (HAT) mechanism or a proton-coupled electron-transfer (PCET) process with intermediate **IV** to produce intermediate **V**. Subsequent

treatment with methyl iodide furnished the isomeric nicotinic acid ester **84** as the final product. Despite its high C4-selectivity, the method showed reduced current efficiency and slower reaction rates due to continuous charge consumption in regenerating I₂ instead of driving the desired transformation.


4.2 C–H Silylation

Silylated pyridines are valuable synthetic intermediates, and they can also be efficiently accessed through cathodic reduction of pyridines. Recently, Zhan and co-workers reported an electrochemical C4-silylation of pyridines via a reductive dearomatization strategy in the presence of chlorosilanes (Scheme 19).⁶³ This methodology was demonstrated to be broadly applicable, affording good yields for a wide range of pyridines bearing aryl, heteroaryl and estrone substituents (**85–87**). However, for a bulky silyl group, C5-silylated pyridine **88** was obtained as the major product. The reaction commenced with the formation of a pyridyl radical anion **I** via single electron-reduction, wherein the nitrogen atom, bearing the highest electron density on the pyridine ring, attacked the chlorosilane, followed by cathodic reduction produced the *N*-silylated intermediate **II**, when smaller chlorosilanes were employed. This intermediate **II** subsequently reacted with another chlorosilane at the C4 position to afford intermediate **III**, which, upon hydrolysis and rearomatization by oxidation, yielded the C4-silylated product with excellent regioselectivity. In contrast, when bulky chlorosilanes such as chlorotrisopropylsilane (TIPSCI) were used, steric hindrance disfavored the formation of intermediate **VI**. Instead, silylation occurred at the C5 position via intermediate **IV**, in which the most electron-rich carbon of the pyridine ring participated. The resulting intermediate **V** after hydrolysis and aerobic oxidation furnished the C5-silylated product.



Scheme 17: Electroreductive C–H carboxylation.





Scheme 18: Site-selective carboxylation of pyridines.

4.5 C–H Alkylation

Although the alkylation of pyridines has been extensively explored through oxidative electrochemical Minisci-type reactions employing different alkyl radical precursors, these approaches often suffer from C2/C4 regioselectivity issues and overalkylation. Very recently, Zhan and co-workers furnished a silane-assisted electroreductive C4-alkylation of pyridines (Scheme 20).⁶⁴ The reaction was compatible with a range of pyridines and alkyl halides, affording the corresponding alkylated pyridines (**89–91**) in good yields. The reaction was proposed to begin with the formation of a pyridinium intermediate **I** through the reaction of 2-phenyl pyridine **72** with chlorotrimethylsilane, which was subsequently reduced at the cathode to afford radical intermediate **II**, while magnesium was oxidized to Mg²⁺ at the anode. In parallel, cathodic reduction of cyclohexyl bromide **92** generates a cyclohexyl radical **III**. Radical-radical recombination produced intermediate **IV**. Subsequent hydrolysis of intermediate **IV**, followed by aerobic oxidation, furnished the desired C4-alkylated product **93**.


4.6 Decyanative Alkylation

Following the electroreductive C–H bond functionalization strategy, electrochemical decyanative functionalization has emerged as another efficient approach for the selective bond functionalization of pyridines. Under electrochemical conditions, cyano-pyridines undergo single-electron reduction to form pyridyl radical anions, which can participate in a variety of bond-forming processes. Subsequent cyanide elimination restores aromaticity.


A pioneering advance was achieved with a nickel-catalyzed electroreductive alkylation of pyridines, employing 4-cyanopyridine and electron-deficient alkenes as coupling partners (Scheme 21).⁶⁵ A diverse range of enones was explored, affording the corresponding alkylated pyridines (**95–97**) in moderate to excellent yields. When cinnamide bearing a chiral auxiliary was employed, the alkylated product **97** was obtained in good yield with a 1:1 diastereomeric ratio. Mechanistically, the transformation can proceed through two distinct pathways: In path a, the alkene **98** undergoes cathodic reduction to generate a radical-anion species **I**, which tautomerizes to the radical-enolate intermediate **II**. This intermediate then couples with a nickel-bound pyridine complex **III** at the C4 position, forming intermediate **IV**. Subsequent reduction of **IV** produces the carbanion intermediate **V**, which undergoes cyanide elimination to afford the desired alkylated product **99**. In path b, the enolate intermediate **II** engages in radical coupling with a pyridine radical anion **VI**, generated in situ from 4-cyanopyridine via cathodic reduction, to form adduct **V**, which then follows a similar cyanide elimination and protonation to deliver the final product.

Rovis and co-workers developed an elegant electrochemical decyanative alkylation of pyridines employing iminium salts as alkylating agents (Scheme 22a).⁶⁶ This strategy provided a direct and efficient route to pyridine-containing α -tertiary amines via a proton-coupled electron transfer (PCET) pathway, a transformation of high significance owing to the prevalence of such scaffolds in pharmaceuticals and





Scheme 19: Regioselective C-H silylation.



Scheme 20: C-H bond alkylation.

agrochemicals. The method exhibited broad functional-group tolerance, accommodating pyridines substituted with halides


and alkynes (**100** and **101**), and was equally effective with cyclic iminium salts, affording the corresponding cyclic amines (**102** and **103**). Mechanistically, the reaction proceeds through single-electron cathodic reduction of the iminium salt **104** to generate an α -amino radical **I**, while cyanopyridine **94** undergoes simultaneous proton-coupled electron transfer (PCET) to yield a pyridyl radical **II**. The radical-radical coupling between **I** and **II** affords intermediate **III**, which subsequently undergoes deprotonation followed by cyanide elimination to deliver the sterically hindered pyridyl amine **105**.

Following this, Guan and co-workers developed an electrochemical protocol for the synthesis of benzylic secondary amines via the coupling of imines with 4-cyanopyridine (Scheme 22b).⁶⁷ Under the standard reaction conditions, 4-cyanopyridine **94** afforded the corresponding aminated product **107**, albeit in modest yield (36%). The reaction was initiated via reduction of both the 4-cyanopyridine **94** and the imine **106** via SET to generate the pyridyl radical intermediate and α -amino radical, respectively. Finally, radical-radical cross-coupling followed by cyanide elimination delivered the desired product **107**.

Next, electroreductive alkylation of 4-cyanopyridines using aldehydes and ketones as alkylating agents was reported by Yang and co-workers (Scheme 23).⁶⁸ The reaction exhibited a broad substrate scope, efficiently converting a variety of aliphatic and aromatic aldehydes into the corresponding secondary alcohols (**108-110**), while ketones also participated to afford *tertiary* alcohol **111** in good yield. The reaction mechanism involved a ketyl radical intermediate **I** formed by cathodic reduction of carbonyl compound **112**; concurrently, 4-cyanopyridine **94** was reduced to a persistent radical anion **II**. These radicals underwent radical-radical cross-coupling to form an intermediate **III**, which then eliminated cyanide to afford alcohol **110**. The formation of both transient radical **I** and persistent radical **II** was proposed to occur preferentially at the cathode. At the anode, DABCO undergoes sacrificial oxidation, thereby providing the necessary electron balance to drive the cathodic reduction steps.

In a recent advance, Huang and co-workers developed a decyanative alkylation of cyanopyridines using unactivated alkyl bromides (Scheme 24).⁶⁹ This operationally simple and scalable protocol featured a broad substrate scope including electron-withdrawing, electron-donating groups substituted pyridines, as well as primary, secondary, tertiary and cyclic alkyl bromides (products **113-117**), without the need for a sacrificial anode. The transformation proceeded through single-electron reduction of Cs-coordinated cyanopyridines to pyridyl radical anion intermediate **I**, which subsequently, coupled with alkyl radical intermediate **II** generated from the corresponding alkyl bromides **118** by reduction at the cathode to give intermediate **III**. Finally, intermediate **III** underwent decyanation to furnish product **119**.


Very recently, a complementary decyanative alkylation of 4-cyanopyridines was disclosed by Zeng and co-workers using continuous-flow electrochemical methods (Scheme 25).⁷⁰ Under flow constant current electrolysis, the combination of alkenes, aryl bromides, and cyanopyridines enabled efficient



access to alkylated pyridine frameworks. Aryl bromides bearing electron-withdrawing substituents, such as an ester group, was well tolerated and delivered the corresponding product **120** in 73% yield. Notably, aryl bromides derived from *L*-menthol also participated effectively, providing the desired product **121** in 62% yield, highlighting the functional-group compatibility of the protocol. A plausible mechanistic pathway was proposed for the observed regioselectivity (Scheme 25). The process was initiated by cathodic reduction of the aryl bromide, followed by debromination to generate an aryl radical **I**, which subsequently added to the alkene to form a carbon-centred radical intermediate **II**. In parallel, cyanopyridine underwent cathodic reduction to produce a relatively persistent pyridyl radical anion **III**. Final radical-radical coupling between these two intermediates, followed by decyanation, furnished the alkylated pyridine product.

4.7 Decyanative Carboxylation

Very recently, Li and co-workers employed a decyanative strategy for the carboxylation of cyanopyridines (Scheme 26).⁷¹ A broad range of 4-cyanopyridines bearing various aryl substituents efficiently afforded the corresponding isonicotinic acid derivatives (**122-125**). The proposed mechanism for this transformation began with the cathodic reduction of 4-cyanopyridine **94**, producing a pyridyl radical




Scheme 21: Decyanative alkylation using alkene.

anion intermediate **I** that underwent carboxylation with CO_2 to generate a cyano-1,4-dihydropyridine-4-carboxylate radical anion **II**. Further cathodic reduction of **II** formed the corresponding dianion **III**, which subsequently undergoes decyanation and protonation to afford the desired isonicotinic acid derivative **126**.





4.8 Dehalogenative Alkylation

Electrochemical dehalogenative functionalization has become a powerful method, allowing the otherwise inert C–halogen bond in pyridines to be effectively activated under electrochemical conditions, thereby facilitating a wide variety of C–C and C–heteroatom bond-forming reactions.⁷² In this context, Bio and co-workers reported an electrochemical dehalogenative alkylation of pyridines employing redox-active esters as alkylating reagents, highlighting the potential of electrochemical activation in promoting sustainable and selective pyridine functionalization (Scheme 27).⁷³ Under constant-current electrolysis at 20 mA, this method afforded 3-alkylated pyridines in moderate yields (**127** and **128**). The reaction proceeds *via* the following mechanism: cathodic reduction of the redox-active ester results in decarboxylative fragmentation, generating radical **II**. Simultaneously, Ni(0) underwent oxidative addition with pyridine halides to form intermediate **IV**. The alkyl radical **II** was then captured by the



Scheme 22: Decyanative alkylation using imines.





nickel complex IV, affording intermediate V. Finally, reductive elimination from intermediate V delivered the C3-alkylated pyridine. The resulting Ni(I) species was subsequently reduced at the cathode, completing the catalytic cycle.

4.9 Dehalogenative deuteration

In 2020, Zhang and co-workers reported an elegant electrochemical dehalogenative deuteration of arenes/pyridines using D₂O as the deuterium source (Scheme

28).^{56a} Control experiments with CD₃CN/H₂O established water as the exclusive hydrogen source. The proposed mechanism involves a cathodic reduction of halopyridine to generate pyridyl radicals, which subsequently couple with deuterium radicals produced via D₂O splitting, supported by EPR studies. The suppression of product formation by hydrogen-radical scavenging with ^tBuOH further corroborated this mechanism, although direct deuterium abstraction from D₂O could not be completely excluded. The reaction exhibited




broad scope with aryl halides, and a few examples of pyridine substrates were reported in this study (129–132).


4.10 Dehalogenative phosphorylation

In 2020, Wickens and co-workers reported a strategy for phosphorylation of chloropyridine, where a photocatalyst was first activated electrochemically before photoexcitation, overcoming the energy limitations of visible-light photoredox catalysis (Scheme 29).⁷⁴ This approach enabled radical generation from 2-chloropyridine (133), which are difficult to


reduce owing to their high reduction potentials. The reaction is believed to initiate via cathodic reduction of the photocatalyst NpMI, generating a mild reducing radical anion intermediate I, which is then excited by light to form a highly reducing intermediate II. This excited photocatalyst intermediate II then reduced the pyridine chloride 133, producing pyridyl radical intermediate III. Radical III then added to the phosphorous atom of triethyl phosphite 134 to yield intermediate IV. Finally, the radical intermediate IV undergoes β -scission to give phosphorylated pyridine 135.



Scheme 27: Dehalogenative alkylation using redox-active ester.



Scheme 28: Dehalogenative deuteration.




Scheme 29: Dehalogenative phosphorylation.

4.11 Dearomative carboxylation

Meanwell and co-workers developed a mild, highly chemoselective electrochemical Birch carboxylation of pyridines using CO₂ as a sustainable C1 source, providing a greener and safer alternative to classical Birch reductions (Scheme 30).⁷⁵ Electrolysis conducted in DMF using a magnesium sacrificial anode and a platinum cathode enabled selective reduction and C4-carboxylation of pyridines without affecting other aromatic rings, such as bis(trifluoromethyl)-, methylthio-substituted phenyl, and benzofuran moieties, affording the desired products in good yields (products 136–138). A plausible mechanism involves cathodic reduction of the pyridine substrate 139 via a Birch-type electrochemical process to form a radical anion intermediate I. Delocalization of the resulting benzylic radical into the aryl substituent stabilizes the dearomatized species. Trapping of the nitrogen-centered anion by CO₂ affords intermediate II, which, upon further cathodic reduction, generates a dianionic species III. Capture of a second equivalent of CO₂ produces a dicarboxylated intermediate IV, followed by decarboxylation on work-up, furnishes the corresponding 1,4-dihydropyridine product 140.





Scheme 30: Birch-type dearomative carboxylation.

From a practical standpoint, net-reductive methods enable diverse C–X (X = H, Cl, Br, CN) functionalization of pyridines by generating pyridyl radical anions via controlled single-electron transfer at the cathode. Polar aprotic solvents such as DMF and MeCN are typically selected because they stabilize charged intermediates, efficiently dissolve salts and CO_2 , and provide a wide electrochemical window under reductive conditions. Halide-containing supporting electrolytes are often preferred, as they can participate in the overall redox balance and, in some systems, in situ-generated molecular iodine facilitates rearomatization. Carbon-based electrodes paired with oxidizable electron donors such as amines, or sacrificial metal anodes (e.g., magnesium), are commonly employed, while metal-based electrodes are typically used at the cathode. For instance, the choice of electrochemical cell: a divided versus an undivided cell can be used to control regioselective functionalization.

5. Functionalization via Paired Electrolysis


Paired electrolysis is an electrochemical strategy in which both anodic and cathodic reactions are utilized productively within a single operation. By enabling simultaneous redox events, it maximizes energy efficiency and allows the formation of multiple bonds or products in one step. This approach expands the synthetic scope by integrating complementary oxidation and reduction processes, providing a sustainable and versatile platform for complex molecular transformations.

5.1 C–H Deuteration

A direct, efficient, and regioselective C4–H deuteration strategy for pyridines was reported by Qiu and co-workers (Scheme 31).⁷⁶ Under optimized conditions, a wide range of pyridines and bipyridines were successfully deuterated with excellent yields and high deuterium incorporation (**69** and **141**). Moreover, the late-stage functionalization of pharmaceutically relevant molecules, including carveol (product **142**), further underscored the synthetic utility and practical relevance of this approach. Mechanistic studies supported by cyclic voltammetry suggested the following pathway for deuteration: initially, $n\text{Bu}_4\text{NI}$ dissociates to furnish $n\text{Bu}_3\text{N}$ (**I**) and $n\text{BuI}$ (**II**). The $n\text{Bu}_3\text{N}$ (**I**) undergoes anodic oxidation to generate a radical cation **III**, thereby delivering electrons to the electrochemical cycle. Concurrently, $n\text{BuI}$ and pyridine derivative **72** react to give the pyridinium salt **IV**, which undergoes single-electron reduction at the cathode to form a radical species **V**. The radical **V** is further reduced to an anionic intermediate **VI**. Subsequent reaction of **VI** with D_2O provides a deuterated intermediate **VII**, which undergoes anodic oxidation to furnish the desired deuterated pyridine **69**.


5.2 C–H Alkylation

Paired electrolysis also enabled C–H alkylation of pyridines under mild conditions. Chen and co-workers reported the first electrochemical deaminative alkylation of pyridines, emplo-




Scheme 31: C–H bond deuteration via paired electrolysis.





Scheme 32: Redox-neutral C-H alkylation using Katritzky salts.



Scheme 33: *meta*-C-H bond alkylation via paired electrolysis.

ying Katritzky salts as alkyl radical precursors (Scheme 32).⁷⁷ The methodology exhibited a limited scope with respect to pyridine substrates; however, it is applicable to structurally complex pyridine derivatives, affording the corresponding C2- or C4-alkylated products (**143–145**), albeit in modest yields. A plausible mechanism involved the single-electron reduction of the Katritzky salt **146** at the cathode, leading to the formation of the persistent radical species **I**, which was subsequently exergonically fragmented to produce the alkyl radical **II**. The radical then slowly underwent an energy-demanding addition to the pyridinium salt **IV**, forming intermediate **V**. Simultaneously, a relatively rapid and exergonic radical-radical combination between **I** and **II** generated intermediate **III**. The reduction of intermediate **V** was achieved by the excited state photocatalyst $[\text{Ru}^{\text{II}}(\text{bpy})_3]^*$, resulting in the

formation of intermediate **VI** and $[\text{Ru}^{\text{III}}(\text{bpy})_3]$. The oxidized $[\text{Ru}^{\text{III}}(\text{bpy})_3]$ species could oxidize intermediate **III**, thereby generating the alkyl radical **II** and completing the photocatalytic cycle. Finally, intermediate **VI** was oxidized at the anode to furnish the desired product **147**. Cheng and co-workers further utilized a paired electrolysis strategy for developing a protocol for *meta*-C–H alkylation of 4-cyanopyridines with alkyl halides (Scheme 33).⁷⁸ Using this strategy, the electrochemical alkylation of pyridines with diverse alkyl bromides, including primary, secondary, and tertiary alkyl bromides, was well tolerated, affording the corresponding products **148–151** in moderate to good yields. The transformation proceeds through the cathodic reduction of 4-cyanopyridine **94**, generating the radical anion **I**, which undergoes *meta*-selective $\text{S}_{\text{N}}2$ alkylation with primary alkyl



REVIEW

Chemical Science

bromides to afford intermediate **II**. Subsequent anodic oxidation of intermediate **II** furnishes the desired product. In contrast, secondary and tertiary bromides undergo single-electron reduction to yield the alkyl radical **III**, which couples with the radical anion **IV**, a resonance form of **I**, to produce the anion **V**. Finally, anodic oxidation of **V** delivers the alkylated pyridine. The observed *meta*-selectivity arises from the highest charge and spin density localised at the *meta*-position of radical anion **I**, favouring S_N2 substitution with primary alkyl bromides. For secondary and tertiary bromides, the *meta*-substituted intermediate **V** is energetically more stable, suggesting that post-transition-state bifurcation may contribute to the origin of selectivity.

5.3 C-H Amination


Malapit and co-workers developed a site-selective electrochemical protocol for the direct amination of aromatic C–H bonds, including pyridines (Scheme 34).⁷⁹ Under optimized electrochemical conditions, selective functionalization of the *meta*-C–H bond was observed. Interestingly, substituted pyridine, bipyridine and terpyridine exclusively afforded the corresponding DABCONium salts **153–155** in moderate to good yields (43–68%). Mechanism of the reaction starts with the single-electron reduction of Selectfluor II reagent **152** at the cathode to produce an *N*-radical dication intermediate **I**, which engages in a charge-transfer interaction with pyridine and leads to C–N bond formation. This process generated the radical intermediate **III**, which was then anodically oxidized to give intermediate **IV**. Finally, deprotonation of **IV** by a base afforded the pyridine-DABCONium product **156**. Furthermore, it is proposed that triethylamine may function as a sacrificial reductant through anodic oxidation, thereby promoting the efficient reduction of Selectfluor II.

5.4 C-H Halogenation


Lei and co-workers reported an electrochemical C–H halogenation of 2-aminopyridine **53** by employing a paired electrolysis approach (Scheme 35).⁸⁰ Using CBr_4 (**157**) as the brominating agent, constant-current electrolysis (12 mA) was performed with a carbon rod as the anode, a nickel plate as the cathode, and ${}^n\text{Bu}_4\text{NBF}_4$ as the electrolyte, converting 2-aminopyridine **53** into the corresponding brominated product **54**. Mechanistic studies suggest that under electrolysis, CBr_4 is reduced at the cathode to generate both bromide ions and tribromomethyl radicals. The bromide ions are subsequently oxidized at the anode to form Br_2 , which efficiently brominates 2-aminopyridine via electrophilic bromination.

5.5 Dehalogenative Alkylation

Ye and co-workers reported an electrochemical aminomethylation of halopyridines catalyzed by nickel (Scheme 36).⁸¹ C2 and C4-alkylated pyridines **158** and **159** were obtained from the corresponding bromopyridines **161** and **162**. Mechanistically, Ni(II) was first reduced at the cathode via a sequential two-electron transfer to generate Ni(0), which subsequently underwent a rapid Ni(0)/Ni(II) comproportionation to yield Ni(I) species. The oxidative addition of Ni(I) to the bromo-pyridine **158** produced the



Scheme 34: C-H amination via paired-electrolysis.


Scheme 35: *meta*-Bromination via paired electrolysis.

intermediate **I**, which was further reduced at the cathode to form intermediate **II**. Simultaneously, the α -amino carbon radical **III** was generated via anodic oxidation followed by deprotonation of tertiary-amine **160**, which was captured by intermediate **II** to furnish intermediate **IV**. Finally, reductive elimination from intermediate **IV** affords the coupling product **153**, thereby regenerating the Ni(I) complex.


Given the significance of fluorinated moieties in drug discovery, and their ability to improve pharmacokinetic profiles as well as key physicochemical properties when incorporated into therapeutic or diagnostic molecules, Zhan and co-workers established a nickel-catalyzed electrochemical fluoroalkylation of iodo-pyridines using sodium difluoromethanesulfinate **157** as the fluoroalkylating reagent (Scheme 37).⁸² Under mild electrochemical conditions, 3-iodo-pyridine (**155**) and 2-methoxy-3-iodo-pyridine (**156**) are converted to fluoroalkylated pyridines **158** and **159** in moderate yields. Based on cyclic voltammetry and electron spin resonance (ESR) spectroscopy, two possible pathways were proposed for this electrochemical fluoroalkylative coupling. Firstly, 3-iodo-pyridine **155** undergoes oxidative addition with Ni(I) to form the intermediate **I**, which was subsequently reduced at the cathode to generate the Ni(II)



species **II**. Alternatively, Ni(I) was first cathodically reduced to Ni(0), which then underwent oxidative addition with **155** to generate intermediate **II**. Simultaneously, sodium fluoroalkyl sulfinate **157** was anodically oxidized to produce the fluoroalkyl radical **III**, which was captured by the Ni(II) species **II** to form intermediate **IV**. Finally, reductive elimination from intermediate **IV** affords the fluoroalkylated product **158** and regenerates Ni(I).



Scheme 36: Dehalogenative alkylation via paired electrolysis.




Scheme 37: Dehalogenative fluoroalkylation via paired electrolysis.

5.6 Decyanative Arylation

In 2021, Yang and co-workers reported decyanative arylation of pyridines using quinoxalin-2(1*H*)-ones as a coupling partner (Scheme 38).⁸³ This method demonstrated excellent substrate

scope, enabling the synthesis of the desired products bearing various functional groups, including cyano, bromo, and *N*-benzyl (**168-171**). The reaction mechanism involves protonation of 4-cyanopyridine (**94**) with HFIP, followed by cathodic reduction to generate the radical intermediate **II**. This radical species then adds to substrate **172**, forming the nitrogen-centred radical intermediate **III**, which undergoes anodic oxidation, followed by decyanation to yield the cationic intermediate **IV**. Finally, intermediate **IV** undergoes isomerization and deprotonation, affording the desired product.

Another strategy for the electrochemical arylation of pyridine was developed by Wen and co-workers by employing quinoline as the arylating agent (Scheme 39).⁸⁴ Under constant-current electrolysis in an undivided cell, a variety of heterocycles, including quinoline and phenanthridine, were reacted with 4-cyanopyridines to afford 4-arylated pyridines (**173-175**) in good to excellent yields. Mechanistically, the 4-cyanopyridinium salt **I** was cathodically reduced to generate the radical intermediate **II**, while quinoline **176** was reduced to form the radical **III**. Radicals **II** and **III** subsequently underwent a radical-radical coupling to produce intermediate **IV**, which, upon elimination of HCN, generated intermediate **V**. Finally, the desired arylated pyridine product **177** was obtained via two-electron anodic oxidation and proton elimination from intermediate **V**.



Scheme 38: Decyanative arylation using quinoxalin-2(1*H*)-ones.

5.7 Dehalogenative Etherification


Recent advances in electrochemical etherification have enabled site-selective alkoxylation of arenes, providing a practical and sustainable alternative to conventional ether-



REVIEW

Chemical Science

forming methods. In this context, Baran and co-workers developed a nickel-catalyzed electrochemical etherification of pyridines and arenes with alcohols as an efficient replacement for classical S_NAr strategies (Scheme 40).⁸⁵ The reaction proceeds under mild conditions, using DBU as the base and ${}^n\text{Bu}_4\text{NBr}$ as the electrolyte, affording desired alkoxy-pyridines **178** and **179** in good yields. The mechanism of this C–O cross-coupling is particularly intriguing. The process begins with the cathodic reduction of Ni(II) to generate an active Ni(I) species, which undergoes oxidative addition with 2-bromopyridine (**158**) to form the Ni(III) complex I. Subsequent reduction of intermediate I generates the Ni(II) species II, which then undergoes ligand exchange with the alcohol to afford intermediate III. Because C–O reductive elimination from the Ni(II)-alkoxy complex III is thermodynamically and kinetically unfavourable at room temperature, and therefore anodic oxidation is employed to generate the Ni(III)-alkoxy species IV. From this higher-valent intermediate IV, facile reductive elimination efficiently affords the C–O coupled product **178**.




Scheme 39: Decyanative arylation using quinoline.


5.8 Dehalogenative Phosphorylation

Parallel to the development of C–X bond alkylation and arylation, electrochemical strategies have also been applied to the phosphorylation of pyridines under mild and sustainable conditions. In 2018, Léonel and co-workers reported the first electrochemically assisted Ni-catalyzed phosphorylation of aryl halides (Scheme 41).⁸⁶ When 3-bromopyridine (**180**) was reacted with dimethyl phosphite (**181**) under constant-current electrolysis (0.2 A) using an iron/nickel rod as the anode and nickel foam as the cathode, the corresponding 3-

phosphonylated pyridine (**182**) was obtained in 20% yield. The reaction was proposed to begin with the cathodic reduction of the Ni(II) pre-catalyst to Ni(0), followed by oxidative addition with the halide. The resulting Ni(II)-aryl complex undergoes ligand exchange with dimethyl phosphite to form a phosphoryl-nickel intermediate. Subsequent anodic oxidation, followed by reductive elimination, furnished the desired product **182**, regenerating the Ni(0) catalyst.



Scheme 40: C–O bond formation from bromopyridines.




Scheme 41: Phosphorylation via paired electrolysis.

5.9 Annulation Reactions

In 2021, Li and co-workers developed an electrochemical intermolecular C–H amination strategy for the coupling of pyridines and anilines, providing an efficient route to benzo[4,5]imidazo[1,2-a]pyridines (Scheme 42).⁸⁷ The protocol features a broad substrate scope and excellent tolerance of diverse functional groups, enabling streamlined access to a variety of fused heterocyclic frameworks. A variety of substituted pyridines and anilines underwent smooth annulation, affording the desired products in excellent yields (**183** and **184**). The proposed mechanism begins with anodic oxidation and proton elimination of aniline **185** to generate an N-centered radical I, and a simultaneous cathodic reduction of the 4-cyano pyridine **94**, generating the radical species II. Two radical species, I and II, then recombine to form the intermediate IV, which, on subsequent desilylation via cath-





Scheme 42: Synthesis of benzo[4,5]imidazo[1,2-a]pyridines via paired electrolysis.

odic reduction generates the intermediate **V**. Finally, anodic oxidation of intermediate **V**, followed by cyclization, affords the annulated product **186**. However, this transformation is limited by its requirement for an electron-withdrawing substituent (e.g., CN or CO₂Me) at the C4-position of the pyridine ring, as well as the need for an electron-withdrawing group (e.g., Ts or COMe) on the nitrogen atom of the aniline to ensure successful reaction outcomes.

In paired electrolysis strategies, the choice of electrode is particularly critical, as oxidative and reductive events occur concurrently within the same electrochemical system. Carbon-based materials such as carbon, graphite, and reticulated vitreous carbon (RVC) are typically selected as anodes due to their stability under oxidative conditions, whereas nickel, platinum, or metal foams are commonly employed as cathodes owing to their high conductivity and efficient electron-transfer characteristics under reductive conditions. Polar aprotic solvents such as MeCN, HFIP, DMSO, and DMF are frequently chosen because they stabilize radical and ionic intermediates, ensure effective dissolution of electrolytes and substrates, and provide a broad electrochemical window compatible with simultaneous anodic and cathodic processes.


6. Conclusion and Outlook

In this review, we have charted the development of electrochemical strategies for pyridine functionalization with a focus on challenges and design principles associated with pyridines, from the earliest pioneering studies to the most recent advances, offering researchers an in-depth perspective on the diverse approaches employed. Beyond these foundational methods, the integration of electrochemistry with photocatalysis and transition-metal catalysis has further enhanced efficiency, selectivity, and sustainability, paving the way for innovative strategies in pyridine derivatization. The synthetic power of these methods is exemplified by their successful application in the late-stage functionalization of complex, pyridine-containing molecules, underscoring their growing relevance to pharmaceutical, agrochemical, and materials research. Such developments not only expand the

synthetic utility of electrochemical pyridine functionalization but also promise to unlock entirely new reactivity paradigms. Nevertheless, the studies highlighted in this review reveal that the field of electrochemical pyridine functionalization remains comparatively underdeveloped, with several important areas still requiring systematic exploration.

1. A significant number of studies summarized in this review utilize pre-functionalized pyridines (e.g., cyano- or halo-substituted derivatives) as starting materials. To further broaden the synthetic versatility and practical applicability of these electrochemical strategies, the direct C–H functionalization of pyridines remains a highly desirable and impactful goal.
2. Several valuable functional groups, such as trifluoromethyl, thiotrifluoromethyl, nitro and azide have not yet been introduced into pyridine frameworks via electrochemical methodologies. Therefore, developing efficient and general electrochemical protocols for incorporating these functionalities is an important direction for future research.
3. Despite significant progress in alkylation and arylation cross-coupling reactions of pyridines, alkenylation, alkynylation, and allenylation processes remain unexplored. The development of such transformations would enable greater molecular complexity and structural diversity, thereby expanding the synthetic utility of electrochemical pyridine functionalization.
4. To date, there are no reported examples of asymmetric electrochemical functionalization of pyridines employing chiral ligands or catalysts, highlighting a significant opportunity for advancement toward enantioselective electrochemical methodologies.

Taken together, these observations highlight the pressing need for continued innovation in catalyst design, reaction development, and mechanistic investigation to fully realize the potential of electrochemical strategies in pyridine functionalization. Overall, we anticipate that bringing together these key historical and recent advances will inspire researchers in both organic synthesis and medicinal chemistry to adopt electrochemical approaches for accessing functionalized pyridines.



Acknowledgements

Financial support from the Anusandhan National Research Foundation (ANRF), Government of India (File ANRF/ARG/2025/006468/CS) is gratefully acknowledged. D. P. H. thanks IISc Bangalore for the infrastructure. S.B. thanks CSIR (File 321-4245-9407/2K24/1) for the post-doctoral fellowship. T.S. thanks the Ministry of Education, Government of India, for Prime Minister's Research Fellowship.

Conflict of Interest

The authors declare no conflict of interest.

Keywords

Electrochemistry, Pyridines, C–H Functionalization, Decyanation.

References

- (1) Altaf, A. A.; Shahzad, A.; Gul, Z.; Rasool, N.; Badshah, A.; Lal, B.; Khan, E. *J. Drug Des. Med. Chem.* **2015**, *1*, 1-11, DOI: 10.11648/j.jddmc.20150101.11.
- (2) Vitaku, E.; Smith, D. T.; Njardarson, J. T. *J. Med. Chem.* **2014**, *57*, 10257-10274, DOI: 10.1021/jm501100b.
- (3) Pal, S. Pyridine: A Useful Ligand in Transition Metal Complexes. In *Pyridine*, Parashar Pandey, P. Ed.; IntechOpen, 2018; DOI: 10.5772/intechopen.76986.
- (4) (a) Kallitsis, J. K.; Geormezi, M.; Neophytides, S. G. *Polym. Int.* **2009**, *58*, 1226-1233, DOI: 10.1002/pi.2661. (b) Zafar, M. N.; Atif, A. H.; Nazar, M. F.; Sumrra, S. H.; Gul, E. S.; Paracha, R. *Russ. J. Coord. Chem.* **2016**, *42*, 1-18, DOI: 10.1134/S1070328416010097.
- (5) (a) Klapars, A.; Vedejs, E. *Chem. Heterocycl. Compd.* **2004**, *40*, 759-766, DOI: 10.1023/B:COHC.0000040772.13427.0b. (b) Katritzky, A. R.; Scriven, E. F. V.; Majumder, S.; Akhmedova, R. G.; Vakulenko, A. V.; Akhmedov, N. G.; Murugan, R.; Abboud, K. A. *Org. Biomol. Chem.* **2005**, *3*, 538-541, DOI: 10.1039/B413285H. (c) Levy, J. N.; Alegre-Requena, J. V.; Liu, R.; Paton, R. S.; McNally, A. *J. Am. Chem. Soc.* **2020**, *142*, 11295-11305, DOI: 10.1021/jacs.0c04674. (d) Boyle, B. T.; Levy, J. N.; de Lescure, L.; Paton, R. S.; McNally, A. *Science* **2022**, *378*, 773-779, DOI: 10.1126/science.add8980. (e) Domingo, L. R.; Aurell, M. J. *New J. Chem.* **2023**, *47*, 5193-5202, DOI: 10.1039/D2NJ05845F.
- (6) Turck, A.; Plé, N.; Mongin, F.; Quéguiner, G. *Tetrahedron* **2001**, *57*, 4489-4505, DOI: 10.1016/S0040-4020(01)00225-3.
- (7) Marsais, F.; Queguiner, G. *Tetrahedron* **1983**, *39*, 2009-2021, DOI: 10.1016/S0040-4020(01)91919-2.
- (8) Dias Pires, M. J.; Poeira, D. L.; Marques, M. M. B. *Eur. J. Org. Chem.* **2015**, *2015*, 7197-7234, DOI: 10.1002/ejoc.201500952.
- (9) Zhou, F. Y.; Jiao, L. *Synlett* **2020**, *32*, 159-178, DOI: 10.1055/s-0040-1706552.
- (10) Murakami, K.; Yamada, S.; Kaneda, T.; Itami, K. *Chem. Rev.* **2017**, *117*, 9302-9332, DOI: 10.1021/acs.chemrev.7b00021.
- (11) Sindhe, H.; Reddy, M. M.; Rajkumar, K.; Kamble, A.; Singh, A.; Kumar, A.; Sharma, S. *Beilstein J. Org. Chem.* **2023**, *19*, 820-863, DOI: 10.3762/bjoc.19.62.
- (12) Manolikakes, S. M.; Barl, N. M.; Sämman, C.; Knochel, P. Z. *Naturforsch. B* **2013**, *68*, 411-422, DOI: doi:10.5560/znb.2013-3061.
- (13) Zhang, L.; Wu, Z. Q.; Jiao, L. *Angew. Chem. Int. Ed.* **2020**, *59*, 2095-2099, DOI: 10.1002/anie.201912564.
- (14) Habib, I.; Singha, K.; Hossain, M. *ChemistrySelect* **2023**, *8*, e202204099, DOI: 10.1002/slct.202204099.
- (15) (a) Horn, E. J.; Rosen, B. R.; Baran, P. S. *ACS Cent. Sci.* **2016**, *2*, 302-308, DOI: 10.1021/acscentsci.6b00091. (b) Yan, M.; Kawamata, Y.; Baran, P. S. *Chem. Rev.* **2017**, *117*, 13230-13319, DOI: 10.1021/acs.chemrev.7b00397. (c) Möhle, S.; Zirbes, M.; Rodrigo, E.; Gieshoff, T.; Wiebe, A.; Waldvogel, S. R. *Angew. Chem. Int. Ed.* **2018**, *57*, 6018-6041, DOI: 10.1002/anie.201712732. (d) Wiebe, A.; Gieshoff, T.; Möhle, S.; Rodrigo, E.; Zirbes, M.; Waldvogel, S. R. *Angew. Chem. Int. Ed.* **2018**, *57*, 5594-5619, DOI: 10.1002/anie.201709766. (e) Yuan, Y.; Lei, A. *Nat Commun* **2020**, *11*, 802, DOI: 10.1038/s41467-020-14322-z.
- (16) Waldvogel, S. R.; Lips, S.; Selt, M.; Riehl, B.; Kampf, C. J. *Chem. Rev.* **2018**, *118*, 6706-6765, DOI: 10.1021/acs.chemrev.8b00233.
- (17) Pollok, D.; Waldvogel, S. R. *Chem. Sci.* **2020**, *11*, 12386-12400, DOI: 10.1039/D0SC01848A.
- (18) (a) Wiebe, A.; Gieshoff, T.; Möhle, S.; Rodrigo, E.; Zirbes, M.; Waldvogel, S. R. *Angew. Chem. Int. Ed.* **2018**, *57*, 5594-5619, DOI: 10.1002/anie.201711060. (b) Li, N.; Sitdikov, R.; Kale, A. P.; Steverlyncck, J.; Li, B.; Rueping, M. *Beilstein J. Org. Chem.* **2024**, *20*, 2500-2566, DOI: 10.3762/bjoc.20.214. (c) Maria, F. A. M.; Guilherme, B. S.; Eliakin Sato de, B.; Timothy John, B.; Guilherme, M. M.; Kleber, T. d. O. *Chem. Synth.* **2025**, *5*, 41, DOI: 10.20517/cs.2024.177. (d) Yadav, M. K.; Chowdhury, S. *Org. Biomol. Chem.* **2025**, *23*, 506-545, DOI: 10.1039/D4OB01187B.
- (19) Inoue, Y.; Nagase, S.; Kodaira, K.; Baba, H.; Abe, T. *Bull. Chem. Soc. Jpn.* **1973**, *46*, 2204-2207, DOI: 10.1246/bcsj.46.2204.
- (20) Davis, V. J.; Haszeldine, R. N.; Tipping, A. E. *J. Chem. Soc., Perkin Trans. 1* **1975**, 1263-1267, DOI: 10.1039/P19750001263.
- (21) Ballinger, J. R.; Teare, F. W.; Bowen, B. M.; Garnett, E. S. *Electrochim. Acta* **1985**, *30*, 1075-1077, DOI: 10.1016/0013-4686(85)80175-4.
- (22) Utlej, J. H. P.; Holman, R. J. *Electrochim. Acta* **1976**, *21*, 987-989, DOI: 10.1016/0013-4686(76)85075-X.
- (23) Budnikova, Y. G.; Kargin, Y. M.; Yanilkin, V. V. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **1992**, *41*, 1299-1300, DOI: 10.1007/BF00864205.
- (24) Amatore, C.; Combellas, C.; Lebbar, N. E.; Thiébaud, A.; Verpeaux, J.-N. *J. Org. Chem.* **1995**, *60*, 18-26.
- (25) Gosmini, C.; Lasry, S.; Nédélec, J. Y.; Périchon, J. *Tetrahedron* **1998**, *54*, 1289-1298, DOI: 10.1016/S0040-4020(97)10225-3.
- (26) de França, K. W. R.; Navarro, M.; Léonel, É.; Durandetti, M.; Nédélec, J. Y. *J. Org. Chem.* **2002**, *67*, 1838-1842, DOI: 10.1021/jo016280y.
- (27) de França, K. W. R.; de Lira Oliveira, J.; Florêncio, T.; da Silva, A. P.; Navarro, M.; Léonel, E.; Nédélec, J. Y. *J. Org. Chem.* **2005**, *70*, 10778-10781, DOI: 10.1021/jo0517491.
- (28) Oliveira, J. L.; Silva, M. J.; Florêncio, T.; Urgin, K.; Sengmany, S.; Léonel, E.; Nédélec, J. Y.; Navarro, M. *Tetrahedron* **2012**, *68*, 2383-2390, DOI: 10.1016/j.tet.2012.01.017.
- (29) Gall, E. L.; Gosmini, C.; Nédélec, J. Y.; Périchon, J. *Tetrahedron* **2001**, *57*, 1923-1927, DOI: 10.1016/S0040-4020(01)00015-1.
- (30) Budnikova, Y.; Kargin, Y.; Nédélec, J.-Y.; Périchon, J. *J. Organomet. Chem.* **1999**, *575*, 63-66, DOI: 10.1016/S0022-328X(98)00963-2.
- (31) Lai, X.-L.; Shu, X.-M.; Song, J.; Xu, H.-C. *Angew. Chem. Int. Ed.* **2020**, *59*, 10626-10632, DOI: 10.1002/anie.202002900.
- (32) (a) Billingsley, K. L.; Anderson, K. W.; Buchwald, S. L. *Angew. Chem. Int. Ed.* **2006**, *45*, 3484-3488, DOI: 10.1002/anie.200600493. (b) Guram, A. S.; Wang, X.; Bunel, E. E.; Faul, M. M.; Larsen, R. D.; Martinelli, M. J. *J. Org. Chem.* **2007**, *72*, 5104-5112, DOI:



- 10.1021/jo070341w. (c) Kudo, N.; Perseghini, M.; Fu, G. C. *Angew. Chem. Int. Ed.* **2006**, *45*, 1282-1284, DOI: 10.1002/anie.200503479. (d) Campeau, L.-C.; Fagnou, K. *Chem. Soc. Rev.* **2007**, *36*, 1058-1068, DOI: 10.1039/B616082D. (e) Kiehne, U.; Bunzen, J.; Staats, H.; Lützen, A. *Synthesis* **2007**, *2007*, 1061-1069, DOI: 10.1055/s-2007-965952. (f) Zhao, D.; You, J.; Hu, C. *Chem. Eur. J.* **2011**, *17*, 5466-5492, DOI: 10.1002/chem.201003039. (g) Markovic, T.; Rocke, B. N.; Blakemore, D. C.; Mascitti, V.; Willis, M. C. *Chem. Sci.* **2017**, *8*, 4437-4442, DOI: 10.1039/C7SC00675F.
- (33) Massignan, L.; Zhu, C.; Hou, X.; Oliveira, J. C. A.; Salamé, A.; Ackermann, L. *ACS Catal.* **2021**, *11*, 11639-11649, DOI: 10.1021/acscatal.1c02516.
- (34) Song, L.; Zhuang, J.-L.; Xiong, P.; Xu, H.-C. *Green Chem.* **2025**, *27*, 10556-10561, DOI: 10.1039/D5GC03231H.
- (35) Yang, Q.-L.; Wang, X.-Y.; Lu, J.-Y.; Zhang, L.-P.; Fang, P.; Mei, T.-S. *J. Am. Chem. Soc.* **2018**, *140*, 11487-11494, DOI: 10.1021/jacs.8b07380.
- (36) Stewart, G.; Alvarez, E. M.; Rapala, C.; Sklar, J. H.; Kalow, J. A.; Malapit, C. A. *Nat. Synth.* **2025**, DOI: 10.1038/s44160-025-00890-9.
- (37) Stevens, T.; Ekholm, K.; Gränse, M.; Lindahl, M.; Kozma, V.; Jungar, C.; Ottosson, T.; Falk-Håkansson, H.; Churg, A.; Wright, J. L.; Lal, H.; Sanfridson, A. *J. Pharmacol. Exp. Ther.* **2011**, *339*, 313-320, DOI: 10.1124/jpet.111.182139.
- (38) Qin, S.; Yang, M.; Xu, M.; Peng, Z.-H.; Cai, J.; Wang, S.; Gao, H.; Zhou, Z.; Hashmi, A. S. K.; Yi, W.; Zeng, Z. *Nat. Commun.* **2024**, *15*, 7428, DOI: 10.1038/s41467-024-50644-y.
- (39) (a) Nakao, Y.; Hiyama, T. *Chem. Soc. Rev.* **2011**, *40*, 4893-4901, DOI: 10.1039/C1CS15122C. (b) Franz, A. K.; Wilson, S. O. *J. Med. Chem.* **2013**, *56*, 388-405, DOI: 10.1021/jm3010114. (c) Cheng, C.; Hartwig, J. F. *Chem. Rev.* **2015**, *115*, 8946-8975, DOI: 10.1021/cr5006414.
- (40) Wan, Q.; Hou, Z.-W.; Zhao, X.-R.; Xie, X.; Wang, L. *Org. Lett.* **2023**, *25*, 1008-1013, DOI: 10.1021/acs.orglett.3c00144.
- (41) Wang, Z.-X.; Yang, D.-Y.; Zhang, Z.; Xu, P.; Zhou, F.; Yang, S.; Zhou, J. *J. Org. Chem.* **2025**, DOI: 10.1021/acs.joc.5c02473.
- (42) (a) Bunnett, J. F.; Zahler, R. E. *Chem. Rev.* **1951**, *49*, 273-412, DOI: 10.1021/cr60153a002. (b) Grimmett, M. R. Halogenation of Heterocycles: II. Six- and Seven-Membered Rings. In *Advances in Heterocyclic Chemistry*, Katritzky, A. R. Ed.; Vol. 58; Academic Press, 1993; pp 271-345 DOI: 10.1016/S0065-2725(08)60288-3.
- (43) Gryaznova, T. V.; Khrizanforova, V. V.; Kholin, K. V.; Khrizanforov, M. N.; Budnikova, Y. H. *Russ. Chem. Bull.* **2016**, *65*, 1798-1804, DOI: 10.1007/s11172-016-1513-x.
- (44) Wu, Y.; Xu, S.; Wang, H.; Shao, D.; Qi, Q.; Lu, Y.; Ma, L.; Zhou, J.; Hu, W.; Gao, W.; Chen, J. *J. Org. Chem.* **2021**, *86*, 16144-16150, DOI: 10.1021/acs.joc.1c00923.
- (45) (a) Leboeuf, M.; Cavé, A.; Ranaivo, A.; Moskowitz, H. *Can. J. Chem.* **1989**, *67*, 947-952, DOI: 10.1139/v89-145. (b) Gundersen, L.-L.; Charnock, C.; Negussie, A. H.; Rise, F.; Teklu, S. *Eur. J. Pharm. Sci.* **2007**, *30*, 26-35, DOI: 10.1016/j.ejps.2006.09.006. (c) Michael, J. P. *Nat. Prod. Rep.* **2008**, *25*, 139-165, DOI: 10.1039/B612166G.
- (46) Bedjeguel, K.; Bienaymé, H.; Dumoulin, A.; Poigny, S.; Schmitt, P.; Tam, E. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3998-4001, DOI: 10.1016/j.bmcl.2006.05.014.
- (47) (a) Wan, J.; Zheng, C.-J.; Fung, M.-K.; Liu, X.-K.; Lee, C.-S.; Zhang, X.-H. *J. Mater. Chem.* **2012**, *22*, 4502-4510, DOI: 10.1039/C2JM14904D. (b) Delcamp, J. H.; Yella, A.; Holcombe, T. W.; Nazeeruddin, M. K.; Grätzel, M. *Angew. Chem. Int. Ed.* **2013**, *52*, 376-380, DOI: 10.1002/anie.201205007. (c) Song, Y. R.; Lim, C. W.; Kim, T. W. *Luminescence* **2016**, *31*, 364-371, DOI: 10.1002/bio.2968.
- (48) Malviya, B. K.; Jassal, A. K.; Karnatak, M.; Verma, V. P.; Sharma, S. *J. Org. Chem.* **2022**, *87*, 2898-2911, DOI: 10.1021/acs.joc.1c02773.
- (49) Grishin, S. S.; Mulina, O. M.; Vil', V. A.; Terent'ev, A. O. *Org. Chem. Front.* **2024**, *11*, 327-335, DOI: 10.1039/D3QO01690K.
- (50) (a) Lovering, F.; Bikker, J.; Humblet, C. *J. Med. Chem.* **2009**, *52*, 6752-6756, DOI: 10.1021/jm901241e. (b) Ling, Y.; Hao, Z.-Y.; Liang, D.; Zhang, C.-L.; Liu, Y.-F.; Wang, Y. *Drug Des. Dev. Ther.* **2021**, *4289*-4338, DOI: 10.2147/DDDT.S329547.
- (51) Chen, Q.-S.; Li, J.-Q.; Zhang, Q.-W. *Pharm. Fronts* **2023**, *05*, e1-e14, DOI: 10.1055/s-0043-1764218.
- (52) Wang, Z.; Liang, W.-F.; Gong, C.-X.; He, X.; Li, J.-H.; Lin, Y.; Ye, K.-Y. *CCS Chem.* **2026**, *0*, 1-19, DOI: 10.31635/ccschem.025.202507069.
- (53) (a) Katsnelson, A. *Nat. Med.* **2013**, *19*, 656, DOI: 10.1038/nm0613-656. (b) Allais, C.; Grassot, J. M.; Rodriguez, J.; Constantieux, T. *Chem. Rev.* **2014**, *114*, 10829-10868, DOI: 10.1021/cr500099b. (c) Mullard, A. *Nat. Rev. Drug Discov.* **2016**, *15*, 219-221,
- (54) Junk, T.; Catallo, W. *J. Chem. Soc. Rev.* **1997**, *26*, 401-406, DOI: 10.1039/CS9972600401.
- (55) (a) Alexakis, E.; Jones, J. R.; Lockley, W. J. S. *Tetrahedron Lett.* **2006**, *47*, 5025-5028, DOI: 10.1016/j.tetlet.2006.05.106. (b) Yang, H.; Zarate, C.; Palmer, W. N.; Rivera, N.; Hesk, D.; Chirik, P. J. *ACS Catal.* **2018**, *8*, 10210-10218, DOI: 10.1021/acscatal.8b03717.
- (56) (a) Liu, C.; Han, S.; Li, M.; Chong, X.; Zhang, B. *Angew. Chem. Int. Ed.* **2020**, *59*, 18527-18531, DOI: 10.1002/anie.202009155. (b) Li, Y.; Ye, Z.; Lin, Y.-M.; Liu, Y.; Zhang, Y.; Gong, L. *Nat. Commun.* **2021**, *12*, 2894, DOI: 10.1038/s41467-021-23255-0.
- (57) Shi, L.; Liu, M.; Zheng, L.; Gao, Q.; Wang, M.; Wang, X.; Xiang, J. *Org. Lett.* **2024**, *26*, 4318-4322, DOI: 10.1021/acs.orglett.4c01296.
- (58) (a) Huang, K.; Sun, C. L.; Shi, Z. *J. Chem. Soc. Rev.* **2011**, *40*, 2435-2452, DOI: 10.1039/C0CS00129E. (b) Weaver, J. D.; Recio, A., III; Grenning, J. A.; Tunge, J. A. *Chem. Rev.* **2011**, *111*, 1846-1913, DOI: 10.1021/cr1002744. (c) Wang, S.; Xi, C. *Chem. Soc. Rev.* **2019**, *48*, 382-404, DOI: 10.1039/C8CS00281A. (d) Ye, J. H.; Ju, T.; Huang, H.; Liao, L. L.; Yu, D. G. *Acc. Chem. Res.* **2021**, *54*, 2518-2531, DOI: 10.1021/acs.accounts.1c00135. (e) Sun, G.-Q.; Liao, L.-L.; Ran, C.-K.; Ye, J.-H.; Yu, D.-G. *Acc. Chem. Res.* **2024**, *57*, 2728-2745, DOI: 10.1021/acs.accounts.4c00417.
- (59) (a) Jew, S.-s.; Park, B.-s.; Lim, D.-y.; Kim, M. G.; Chung, I. K.; Kim, J. H.; Hong, C. I.; Kim, J.-K.; Park, H.-J.; Lee, J.-H.; Park, H.-g. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 609-612, DOI: 10.1016/S0960-894X(02)01041-7. (b) Cokoja, M.; Bruckmeier, C.; Rieger, B.; Herrmann, W. A.; Kühn, F. E. *Angew. Chem. Int. Ed.* **2011**, *50*, 8510-8537, DOI: 10.1002/anie.201102010.
- (60) Khoshro, H.; Zare, H. R.; Jafari, A. A.; Gorji, A. *Electrochem. Commun.* **2015**, *51*, 69-71, DOI: 10.1016/j.elecom.2014.11.025.
- (61) Zhao, Z.; Liu, Y.; Wang, S.; Tang, S.; Ma, D.; Zhu, Z.; Guo, C.; Qiu, Y. *Angew. Chem. Int. Ed.* **2023**, *62*, e202214710, DOI: 10.1002/anie.202214710.
- (62) (a) Sun, G.-Q.; Yu, P.; Zhang, W.; Zhang, W.; Wang, Y.; Liao, L.-L.; Zhang, Z.; Li, L.; Lu, Z.; Yu, D.-G.; Lin, S. *Nature* **2023**, *615*, 67-72, DOI: 10.1038/s41586-022-05667-0. (b) Zhong, G.; Huang, Y.; He, L. *Chem. Synth.* **2023**, *3*, 19, DOI: 10.20517/cs.2023.03. (c) Surke, M.; Zhao, R.; Ackermann, L. *Sci. China Chem.* **2023**, *66*, 1549-1550, DOI: 10.1007/s11426-023-1585-6.
- (63) Yang, Z.; Liu, X.; Zhang, T.; Maekawa, H.; Hao, X.-Q.; Song, M.-P. *Org. Chem. Front.* **2024**, *11*, 5545-5552, DOI: 10.1039/D4QO01066C.



REVIEW

Chemical Science

- (64) Yang, Z.; Liu, X.; Chen, Y.-X.; Gao, Y.; Xiao, X.; Maekawa, H.; Zhang, T.; Hao, X.-Q.; Song, M.-P. *Sci. Adv.* **2025**, *11*, eadz3401, DOI: 10.1126/sciadv.adz3401.
- (65) Zhang, S.; Li, L.; Li, X.; Zhang, J.; Xu, K.; Li, G.; Findlater, M. *Org. Lett.* **2020**, *22*, 3570-3575, DOI: 10.1021/acs.orglett.0c01014.
- (66) Lehnerr, D.; Lam, Y.-h.; Nicastrì, M. C.; Liu, J.; Newman, J. A.; Regalado, E. L.; DiRocco, D. A.; Rovis, T. *J. Am. Chem. Soc.* **2020**, *142*, 468-478, DOI: 10.1021/jacs.9b10870.
- (67) Zeng, W.-M.; He, Y.-H.; Guan, Z. *Org. Lett.* **2022**, *24*, 7178-7182, DOI: 10.1021/acs.orglett.2c02900.
- (68) Zhang, X.; Yang, C.; Gao, H.; Wang, L.; Guo, L.; Xia, W. *Org. Lett.* **2021**, *23*, 3472-3476, DOI: 10.1021/acs.orglett.1c00920.
- (69) Deng, W.; Li, X.; Li, Z.; Wen, Y.; Wang, Z.; Lin, Z.; Li, Y.; Hu, J.; Huang, Y. *Org. Lett.* **2023**, *25*, 9237-9242, DOI: 10.1021/acs.orglett.3c03984.
- (70) Sun, X.; Meng, F.; Xu, K.; Zeng, C. *Org. Lett.* **2025**, *27*, 12784-12789, DOI: 10.1021/acs.orglett.5c04052.
- (71) Sui, J.-L.; Zeng, L.; Hu, M.; Sun, F.-J.; Cui, P.-Z.; Li, J.-H. *Org. Lett.* **2025**, *27*, 11510-11514, DOI: 10.1021/acs.orglett.5c03529.
- (72) Li, P. F.; Kou, G. S.; Qi, L. P.; Qiu, Y. A. *J. Electrochem.* **2024**, *30*, 2313005, DOI: 10.61558/2993-074X.3442.
- (73) Li, H.; Breen, C. P.; Seo, H.; Jamison, T. F.; Fang, Y.-Q.; Bio, M. M. *Org. Lett.* **2018**, *20*, 1338-1341, DOI: 10.1021/acs.orglett.8b00070.
- (74) Cowper, N. G. W.; Chernowsky, C. P.; Williams, O. P.; Wickens, Z. K. *J. Am. Chem. Soc.* **2020**, *142*, 2093-2099, DOI: 10.1021/jacs.9b12328.
- (75) Sarkar, S.; Rohit; Meanwell, M. W. *Green Chem.* **2025**, *27*, 2900-2906, DOI: 10.1039/D4GC05976J.
- (76) Zhao, Z.; Zhang, R.; Liu, Y.; Zhu, Z.; Wang, Q.; Qiu, Y. *Nat. Commun.* **2024**, *15*, 3832, DOI: 10.1038/s41467-024-48262-9.
- (77) Wang, K.; Liu, X.; Yang, S.; Tian, Y.; Zhou, M.; Zhou, J.; Jia, X.; Li, B.; Liu, S.; Chen, J. *Org. Lett.* **2022**, *24*, 3471-3476, DOI: 10.1021/acs.orglett.2c01022.
- (78) Pan, J.; Cheng, X. *ChemCatChem* **2025**, *17*, e202402139, DOI: 10.1002/cctc.202402139.
- (79) Alvarez, E. M.; Stewart, G.; Ullah, M.; Lalisce, R.; Gutierrez, O.; Malapit, C. A. *J. Am. Chem. Soc.* **2024**, *146*, 3591-3597, DOI: 10.1021/jacs.3c11506.
- (80) Zhou, Z.; Yuan, Y.; Cao, Y.; Qiao, J.; Yao, A.; Zhao, J.; Zuo, W.; Chen, W.; Lei, A. *Chin. J. Chem.* **2019**, *37*, 611-615, DOI: 10.1002/cjoc.201900091.
- (81) Ma, Y.; Hong, J.; Yao, X.; Liu, C.; Zhang, L.; Fu, Y.; Sun, M.; Cheng, R.; Li, Z.; Ye, J. *Org. Lett.* **2021**, *23*, 9387-9392, DOI: 10.1021/acs.orglett.1c03500.
- (82) Zou, Z.; Li, H.; Huang, M.; Zhang, W.; Zhi, S.; Wang, Y.; Pan, Y. *Org. Lett.* **2021**, *23*, 8252-8256, DOI: 10.1021/acs.orglett.1c02997.
- (83) Wen, J.; Yang, X.; Yan, K.; Qin, H.; Ma, J.; Sun, X.; Yang, J.; Wang, H. *Org. Lett.* **2021**, *23*, 1081-1085, DOI: 10.1021/acs.orglett.0c04296.
- (84) Niu, C.; Yang, J.; Yan, K.; Xie, J.; Jiang, W.; Li, B.; Wen, J. *iScience* **2022**, *25*, 104253, DOI: 10.1016/j.isci.2022.104253.
- (85) Zhang, H. J.; Chen, L.; Oderinde, M. S.; Edwards, J. T.; Kawamata, Y.; Baran, P. S. *Angew. Chem. Int. Ed.* **2021**, *60*, 20700-20705, DOI: 10.1002/anie.202107820.
- (86) Sengmany, S.; Ollivier, A.; Le Gall, E.; Léonel, E. *Org. Biomol. Chem.* **2018**, *16*, 4495-4500, DOI: 10.1039/C8OB00500A.
- (87) Luo, M. J.; Ouyang, X. H.; Zhu, Y. P.; Li, Y.; Li, J. H. *Green Chem.* **2021**, *23*, 9024-9029, DOI: 10.1039/D1GC02922C.

View Article Online
DOI: 10.1039/D6SC00233A

Chemical Science Accepted Manuscript



No new primary research results, software, or code were included, and no data were generated or analyzed as part of this review.

View Article Online
DOI: 10.1039/D6SC00233A

