



Cite this: DOI: 10.1039/d6sc00382f

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 14th January 2026  
Accepted 18th February 2026

DOI: 10.1039/d6sc00382f

rsc.li/chemical-science

# Photocatalytic multicomponent alkene dicarbofunctionalization *via* PCET/nickel dual catalysis

Yeersen Patehebieke, Victoria Jansson, Jakob Öberg and Carl-Johan Wallentin \*

Herein we report a synergistic photoredox PCET/nickel dual catalytic strategy that enables three-component alkene dicarbofunctionalization (DCF) using unprotected alcohols as alkyl radical precursors. This transformation merges concerted proton-coupled electron transfer (PCET) activation of alcohols with nickel-catalyzed cross-coupling, achieving sequential Giese addition and C(sp<sup>3</sup>)-C(sp<sup>3</sup>), C(sp<sup>3</sup>)-C(sp<sup>2</sup>) bond formation under visible light and redox-neutral conditions. The method proceeds directly from secondary and tertiary alcohols, exhibits broad functional-group tolerance. Notably, employing a chiral Ni catalyst enables the first enantioselective variant of this type of PCET mediated transformation. This work establishes a general platform for deconstructive alkene difunctionalization from simple alcohol feedstocks, expanding the synthetic reach of PCET in dual catalytic radical chemistry.

## Introduction

Multicomponent reactions (MCRs) provide a powerful platform for the rapid assembly of molecular complexity from simple precursors in a single operation.<sup>1,2</sup> Their inherent convergence, efficiency, and atom economy have made them indispensable tools for the synthesis of pharmaceuticals, agrochemicals, and functional materials.<sup>3</sup> By forming multiple bonds in a single step, MCRs minimize purification steps, reduce waste, and enable modular access to structurally diverse scaffolds. These features align MCRs with the principles of sustainable chemistry<sup>4</sup> and make them highly attractive for late-stage diversification in medicinal chemistry.<sup>5</sup> The challenge, however, lies in controlling selectivity when multiple reactive intermediates are generated simultaneously. Recent advances in catalysis, particularly visible-light photoredox catalysis, have opened new avenues to address this challenge by enabling the controlled generation of reactive radical intermediates under exceptionally mild and tunable conditions.<sup>6–8</sup>

Photoredox catalysis relies on single-electron transfer (SET) processes from photoexcited catalysts to transform simple precursors into radicals with excellent control over reactivity and selectivity.<sup>9</sup> As a result, photoredox catalysis has greatly expanded the scope of multicomponent transformations, enabling the direct incorporation of diverse alkyl fragments into complex molecular frameworks. Precursors such as carboxylic acids,<sup>10–15</sup> alkyltrifluoroborates,<sup>16–28</sup> oxalates,<sup>29–32</sup> *N*-(acyloxy)phthalimides (NHPI esters),<sup>33–44</sup> alkyl halides,<sup>45–53</sup> silicates,<sup>54–58</sup> Katritzky pyridinium salts,<sup>59–61</sup> aldehydes,<sup>62–65</sup> alcohols,<sup>66–70</sup> and




Fig. 1 (a) Representative alkyl radical precursors in photocatalytic multicomponent reactions. (b) Photocatalytic Ni-catalyzed multicomponent reactions using alcohols as alkyl radical precursors.

Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg SE 41390, Sweden. E-mail: carl.wallentin@chem.gu.se

even unactivated C(sp<sup>3</sup>)-H bonds<sup>66,68,71-75</sup> have been converted into carbon-centered radicals and engaged in sequential bond-forming events with high functional-group tolerance (Fig. 1a).

These developments have transformed MCRs into versatile tools for step-economic construction of C-C and C-heteroatom bonds and have set the stage for increasingly sophisticated radical cascades. Among the various strategies for radical generation, proton-coupled electron transfer (PCET) has emerged as a particularly powerful approach for activating strong X-H bonds under mild, redox-neutral conditions.<sup>76,77</sup> By orchestrating electron transfer and proton transfer in a single concerted event, PCET overcomes the high bond dissociation energies of substrates such as alcohols (BDE ≈ 105 kcal mol<sup>-1</sup>), enabling direct access to alkoxy or carbon-centered radicals without prefunctionalization.<sup>76,78</sup> Pioneering studies by Knowles,<sup>79-83</sup> and others<sup>84-91</sup> established the foundation for PCET activation of alkanols, enabling ring-opening reactions and Giese-type additions that rapidly build molecular complexity. These breakthroughs demonstrated that alcohols can serve as practical radical precursors, circumventing the need for laborious derivatization steps. More recently, PCET also has been integrated into dual catalytic systems, where the radicals generated from alcohols are intercepted by nickel, unlocking cross-coupling pathways that were previously inaccessible.<sup>92-95</sup>

Recently, our group developed a PCET based metal-free photocatalytic strategy for the β-scission of secondary and tertiary alcohols, enabling their direct conversion into alkyl radicals under visible light irradiation.<sup>96</sup> This protocol allowed the efficient alkylation of diverse electron-deficient alkenes and demonstrated late-stage functionalization of pharmaceutically relevant scaffolds under mild, redox-neutral conditions (Fig. 2a). In subsequent work, we merged concerted PCET-mediated radical generation with nickel cross-coupling to achieve a practical deconstructive arylation of aliphatic alcohols (Fig. 2a).<sup>97</sup> Together, these studies established that free alcohols

can serve as versatile radical precursors for both Giese-type additions and Ni-catalyzed cross-couplings, thereby laying the foundation for combining these two manifolds in a single multicomponent transformation.

A few photocatalytic three-component reactions that combine radical 1,4-addition with nickel-catalyzed cross-coupling have been reported as powerful methods for vicinal difunctionalization of alkenes under mild conditions.<sup>98,99</sup> In these processes carbon-centered radicals are derived from pre-activated species like NHPI esters,<sup>37-39,42,43</sup> oxalates,<sup>29-31</sup> or trifluoroborates<sup>16,18-20,24-28</sup> which require multistep synthesis and limit late-stage applicability. Direct use of free alcohol in such three-component transformations remains rare (Fig. 1b).

In 2021, Molander and Gutierrez presented photoredox/nickel dual catalytic utilization of alcohols (among other feedstocks like ethers and amides) by performing photocatalyzed hydrogen atom transfer (photo-HAT) to activate aliphatic C-H bonds giving alkyl radicals. The subsequent steps involve Giese addition of the alkyl radicals and nickel-catalyzed coupling.<sup>66</sup> Later in 2024, Nevado realized a similar alkene dicarbofunctionalization *via* an enantioselective photo-HAT/DCF process.<sup>68</sup> In the same year, the MacMillan group utilized benzoxazolium activated primary alcohols as alkyl radical precursors in photoredox/nickel dual catalytic DCF of alkenes. The reaction proceeds through a triple radical sorting mechanism that merges photoredox and nickel-catalyzed bimolecular homolytic substitution (S<sub>H2</sub>) catalysis to achieve alkene dialkylation.<sup>69</sup>

There has not been any report that utilize PCET in photoredox/nickel dual catalytic DCF of alkenes. Combining PCET-mediated radical generation from unactivated alcohols in DCF type transformations. Consequently, a PCET based approach would deliver a more general and modular three-component strategy towards molecular construction.

Herein, we report a synergistic photoredox PCET/nickel dual catalytic system that accomplishes the three-component dicarbofunctionalization of electron-deficient alkenes using unprotected alcohol as the radical source (Fig. 2b). The method features broad substrate scope, including challenging secondary and tertiary alcohols, and demonstrates excellent functional-group tolerance. We also demonstrated the first enantioselective variant of a PCET initiated DCF process.

## Results and discussion

To evaluate whether our previously developed PCET-mediated alkyl radical generation from alcohols is compatible with three-component alkene dicarbofunctionalization, we selected tertiary alcohol **1**, phenyl acrylate **2**, and 4-bromoacetophenone **3** as model substrates. The reaction conditions were adapted from our previously reported nickel-catalyzed deconstructive cross-coupling methods.<sup>97</sup> Using the Fukuzumi catalyst [Mes-Acr-Me]<sup>+</sup>ClO<sub>4</sub><sup>-</sup> as the photocatalyst, NiBr<sub>2</sub>·dtbbpy (4,4'-di-*tert*-butyl-2,2'-dipyridyl) as cross-coupling catalyst, 2,4,6-collidine as the base, in a solvent mixture of 1,2-dichloroethane (DCE) + 5% MeCN at 35 °C under 450 nm blue light irradiation for 48 h. To




Fig. 2 (a) Our previous work on PCET mediated alkyl radical generation from alcohol and its utilization. (b) This work: photoredox PCET/nickel dual catalytic three-component alkene dicarbofunctionalization by alcohols.




our delight, the desired dicarbofunctionalized product **4** was obtained in a very good yield of 82% (Table 1, entry 1).

Encouraged by this result, we performed additional screening to refine the conditions. Using DCE alone reduced the yield to 73% (entry 2), while MeCN as the sole solvent failed to produce the desired product (entry 3). Interestingly, employing MeCN as a co-solvent improved the yield, possibly due to an optimal balance of polarity and Brønsted basicity that promotes both precursor complex formation and disaggregation of the successor complex, thereby mitigating non-productive back electron transfer.<sup>100</sup> Other solvents gave significantly lower or trace yields (entries 4–5). Altering the reaction stoichiometry negatively impacted the outcome (see SI for details). Changing the photocatalyst counterion from ClO<sub>4</sub><sup>−</sup> to BF<sub>4</sub><sup>−</sup> had little effect (entry 6), and screening other acridinium photocatalysts resulted in decreased yields (see SI for details).

Shortening the reaction time to 6 h slightly reduced the yield to 73% (entry 7), with <sup>1</sup>H NMR indicating residual starting alcohol. At 12 h, the alcohol was nearly consumed, and the yield improved to 81% (entry 8). Increasing light intensity to 100% accelerated alcohol consumption within 12 h but lowered the product yield to 74% (entry 9), likely due to increased side-product formation and photocatalyst degradation. Reducing light intensity to 75% and maintaining a 12 h reaction time afforded the highest yield of 84% (entry 11). Finally, control experiments confirmed the essential role of all components—photocatalyst, nickel catalyst, base, and light—as omission of any resulted in no product formation (entries 12–15).

Table 1 Optimization of reaction conditions<sup>a</sup>



Entry	Deviation from standard conditions	Yield <sup>b</sup>
1	No deviation	82%
2	DCE	73%
3	MeCN	n.d.
4	DCM	57%
5	DMF	Trace
6	[Mes-Acr-Me] <sup>+</sup> BF <sub>4</sub> <sup>−</sup>	81%
7	6 h	73%
8	12 h	81%
9	12 h, 100% intensity	74%
10	6 h, 75% intensity	74%
11	12 h, 75% intensity	84%
12	12 h, no photocatalyst	n.d.
13	12 h, no nickel catalyst	n.d.
14	12 h, no base	n.d.
15	12 h, no light	n.d.

<sup>a</sup> Reaction conditions: alcohol (0.3 mmol), alkene (0.2 mmol), aryl bromide (0.1 mmol), 2,4,6-collidine (0.3 mmol), [Mes-Acr-Me]<sup>+</sup>ClO<sub>4</sub><sup>−</sup> (0.01 mmol), NiBr<sub>2</sub>·dtbbpy (0.02 mmol), DCE + 5% MeCN (3 mL), time (48 h), Lucent 360 photo reactor (450 nm, 50% light intensity), temperature (35 °C). <sup>b</sup> Yields were determined by <sup>1</sup>H NMR analysis of the crude reaction mixtures using ethylene carbonate as an internal standard.

With the optimal reaction conditions established, we next explored the substrate scope of this transformation. We began by examining the aryl bromide scope. A wide range of aryl bromides bearing diverse functional groups were well tolerated under optimized conditions (Scheme 1). Substrates containing nitrile **5**, sulfone **6** & **10**, trifluoromethyl **7**, ketone **8**, ester **9**, aldehyde **11**, and boronic ester groups **12** afforded the desired products in good to excellent yields (92–63%). Notably, aryl bromides bearing aldehyde **11** or Bpin **12** groups posed purification challenges, as the target products co-eluted with the ketone byproduct and trace amounts of unreacted alcohol.

To address this, we replaced *tert*-butyl acrylate with phenyl acrylate, anticipating improved separation. Phenyl acrylate delivered good yields (84–63%) with electron-withdrawing substituents on the aryl halide **4**, **13–15**, however, electron-neutral **16** and electron-donating substituents **17** & **18** significantly diminished the yields. This limitation is consistent with known trends in photoredox/nickel cross-coupling reactions, where efficient oxidative addition to low-valent nickel species requires a polarized C–X bond and an electrophilic aryl carbon. Electron-donating groups reduce electrophilicity, slowing oxidative addition and increasing decomposition pathways.<sup>101</sup> Unfortunately, switching to phenyl acrylate did not resolve the purification issue, so we reverted to *tert*-butyl acrylate as the alkene partner.


Extended aromatic systems such as biphenyl **19**, naphthalene **20**, and phthalide derivatives **21** provided decent yields (51–46%). Heteroaryl bromides, including pyridine **22**, quinoline **23** & **25**, and pyrimidine derivatives **24**, also furnished good to moderate yields (76–50%). Attempts to employ alkyl bromides **26** or alkynyl bromides **27** as coupling partners were unsuccessful, with reactions failing under the optimized conditions.

After achieving satisfactory results with aryl halides, we investigated the scope of alkenes and alkynes. Electron-deficient olefins such as vinylboronic acid pinacol ester **28** and *N,N*-dimethylacrylamide (DMAA) **29** exhibited reactivity comparable to acrylates, delivering good to synthetically useful yields. Ethynylbenzene derivatives bearing electron-neutral **30**, electron-donating **31**, and electron-withdrawing substituents **32** also afforded reasonable yields.


Finally, we examined the alcohol scope for alkyl radical generation (Scheme 2). Tertiary alcohols bearing alkyl, cycloalkyl, and heterocyclic groups, including *tert*-butyl **22**, cyclohexyl **33**, cyclopentyl **34**, piperidine **35**, tetrahydropyran **36**, and isopropyl **37**, provided dicarbofunctionalized products in yields ranging from 76% to 23%. Secondary alcohols with tertiary **22**, secondary **33** & **38**, benzylic **39**, and primary **40** substituents exhibited similar trends, affording products in 78% to trace yield. The observed reactivity correlates with radical stability, following the order 3° > 2° > benzylic > 1°. More stable and sterically hindered radicals favor Giese-type addition to the olefin, after which the resulting radical engages the nickel catalytic cycle to furnish the difunctionalized product.

In contrast, less stable and less hindered radicals are more prone to nickel-mediated cross-coupling, competing with Giese addition and thereby reducing overall yields. These trends are





**Scheme 1** Scope of aryl halides, alkenes and alkynes. Reaction conditions: alcohol (0.3 mmol), alkene (0.2 mmol), aryl bromide (0.1 mmol), 2,4,6-collidine (0.3 mmol), [Mes-Acr-Me]<sup>+</sup>ClO<sub>4</sub><sup>-</sup> (0.01 mmol), NiBr<sub>2</sub>·dtbbpy (0.02 mmol), DCE + 5% MeCN (3 mL), time (12 h), Lucent 360 photo reactor (450 nm, 75% light intensity), temperature (35 °C). <sup>a</sup>Isolated yield. <sup>b</sup><sup>1</sup>H NMR yield.



**Scheme 2** Scope of alcohols. <sup>a</sup>Isolated yield. <sup>b</sup><sup>1</sup>H NMR yield.

clearly reflected in quantitative <sup>1</sup>H NMR analysis of the crude reaction mixtures of **4**, **33**, and **40** (see SI). For the tertiary radical involved in the synthesis of **4**, no cross-coupling product is detected; instead, the Giese-type addition product is formed in a 1.2 : 1.0 ratio relative to the targeted difunctionalized product **4**. In contrast, secondary radicals predominantly yield the cross-coupled product, while the Giese-type and desired difunctionalized products are formed in comparable amounts (1.15 : 1.00 for **33**). The least stable and least sterically demanding primary radical likewise furnishes the cross-coupled product as the major component, with only trace amounts of the difunctionalized product observed, as exemplified by **40**.

Encouraged by the work of Chu and Gutierrez<sup>18</sup> and other groups<sup>24,47,50,68,102</sup> we wanted to explore the potential to install the aromatic moiety in an enantioenriched fashion. Consequently, we engaged the synthesis of **22** utilizing a chiral Ni catalyst employing the standard conditions. To our delight **22** was generated in 66% yield with an ee of >94% (Scheme 2). This illustrates that this dicarbofunctionalization strategy indeed is compatible with enantioselective multicomponent based diversification of electron deficient alkenes. Studies focused on further optimization of the chiral ligand and reaction parameters, as well as expansion of the enantioselective scope are ongoing in our lab.

In our previous investigation of the reaction mechanism, we have concluded the concerted PCET is the main mechanism of






Fig. 3 Proposed reaction mechanism.

alkyl radical generation from these aliphatic alcohols. So based on our previous mechanistic studies<sup>96,97</sup> and the reported literature on photoredox/nickel dual catalytic cross-coupling reactions,<sup>103–105</sup> we propose the following plausible mechanism for the alkene dicarbofunctionalization reaction (Fig. 3). Upon visible-light irradiation, the photocatalyst (41) in its excited state (42) oxidizes the base complex substrate (43) to its corresponding oxidized form (44). This oxidized species then undergoes a concerted proton-coupled electron transfer (PCET), generating the key alkyl radical (48) along with a protonated base (46) and a ketone byproduct (47). The alkyl radical subsequently adds to an electron-deficient olefin (49), forming a new radical intermediate (50). This intermediate then combines with a Ni(I)–Br complex (51) to yield a Ni(II)–alkyl–bromo species (52). Next, a single-electron transfer (SET) from the reduced photocatalyst (45) to the nickel complex (52) regenerates the ground-state photocatalyst and produces a Ni(I)–alkyl species (53). Oxidative addition of the aryl halide (54) to this Ni(I) complex forms a Ni(III)–alkyl–aryl–bromo intermediate (55), which undergoes reductive elimination to deliver the desired difunctionalized product (56) and a Ni(I)–bromo species (51), thereby completing the nickel catalytic cycle.

## Conclusions

In conclusion, we have developed an efficient PCET mediated photoredox/nickel dual catalytic strategy that achieves three-component alkene dicarbofunctionalization directly from unprotected alcohols. This redox-neutral protocol merges

concerted PCET-mediated alkyl radical generation with nickel-catalyzed cross-coupling, enabling the direct use of secondary and tertiary alcohols as alkyl radical precursors under visible light. The method proceeds under mild conditions, displays broad functional-group tolerance, and provides access to diverse vicinal aryl–alkyl disubstituted products. The observed reactivity trend across different alcohol classes correlates with radical stability and underlines the mechanistic synergy between photoredox and nickel catalysis. Product distribution analysis further reveals a radical stability dependent competition between conjugate addition and direct nickel capture, rationalizing diminished efficiency for less stabilized radicals. Moreover, the successful realization of an enantioselective example highlights the potential of this approach for asymmetric multicomponent synthesis. Ongoing efforts are directed toward developing broadly applicable enantioselective variants of this transformation through catalyst and ligand design. This work establishes a general and practical platform for deconstructive difunctionalization of alkenes from readily available alcohols and paves the way toward sustainable, atom-economical radical transformations driven by PCET.

## Author contributions

Y. P.: conceptualization (lead); methodology; investigation; validation; formal analysis; data curation; writing – original draft. V. J.: investigation; validation; formal analysis; data curation. J. Ö.: investigation; validation; formal analysis; data



curation. C.-J. W.: conceptualization (lead); supervision; project administration; funding acquisition; writing – review & editing.

## Conflicts of interest

There are no conflicts to declare.

## Data availability

Supplementary information (SI): all experimental procedures, characterization data, and copies of NMR spectra for new compounds. See DOI: <https://doi.org/10.1039/d6sc00382f>.

## Acknowledgements

Financial support from the Knut and Alice Wallenberg Foundation (Grant 2019.0071) and The Swedish Research Council are gratefully acknowledged.

## Notes and references

- J. Zhu, Q. Wang and M. Wang, *Multicomponent reactions in organic synthesis*, John Wiley & Sons, 2015.
- R. P. Herrera and E. Marqués-López, *Multicomponent reactions: concepts and applications for design and synthesis*, John Wiley & Sons, 2015.
- A. Dömling, W. Wang and K. Wang, Chemistry and Biology Of Multicomponent Reactions, *Chem. Rev.*, 2012, **112**, 3083–3135.
- B. Martinengo, E. Diamanti, E. Uliassi and M. L. Bolognesi, Harnessing the 12 Green Chemistry Principles for Sustainable Antiparasitic Drugs: Toward the One Health Approach, *ACS Infect. Dis.*, 2024, **10**, 1856–1870.
- E. Ruijter and R. V. A. Orru, Multicomponent reactions – opportunities for the pharmaceutical industry, *Drug Discov. Today Technol.*, 2013, **10**, e15–e20.
- L. Marzo, S. K. Pagire, O. Reiser and B. König, Visible-Light Photocatalysis: Does It Make a Difference in Organic Synthesis?, *Angew. Chem., Int. Ed.*, 2018, **57**, 10034–10072.
- J. Twilton, C. Le, P. Zhang, M. H. Shaw, R. W. Evans and D. W. C. MacMillan, The merger of transition metal and photocatalysis, *Nat. Rev. Chem.*, 2017, **1**, 0052.
- C. K. Prier, D. A. Rankic and D. W. C. MacMillan, Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis, *Chem. Rev.*, 2013, **113**, 5322–5363.
- J. M. R. Narayanam and C. R. J. Stephenson, Visible light photoredox catalysis: applications in organic synthesis, *Chem. Soc. Rev.*, 2011, **40**, 102–113.
- C. Zhu, H. Yue, B. Maity, I. Atodiresei, L. Cavallo and M. Rueping, A multicomponent synthesis of stereodefined olefins via nickel catalysis and single electron/triplet energy transfer, *Nat. Catal.*, 2019, **2**, 678–687.
- R. S. Mega, V. K. Duong, A. Noble and V. K. Aggarwal, Decarboxylative Conjunctive Cross-coupling of Vinyl Boronic Esters using Metallaphotoredox Catalysis, *Angew. Chem., Int. Ed.*, 2020, **59**, 4375–4379.
- H. Jiang, X. Yu, C. G. Daniliuc and A. Studer, Three-Component Aminoarylation of Electron-Rich Alkenes by Merging Photoredox with Nickel Catalysis, *Angew. Chem., Int. Ed.*, 2021, **60**, 14399–14404.
- V. T. Nguyen, G. C. Haug, V. D. Nguyen, N. T. H. Vuong, G. B. Karki, H. D. Arman and O. V. Larionov, Functional group divergence and the structural basis of acridine photocatalysis revealed by direct decarboxysulfonylation, *Chem. Sci.*, 2022, **13**, 4170–4179.
- N. J. Venditto and J. A. Boerth, Photoredox-Catalyzed Multicomponent Synthesis of Functionalized  $\gamma$ -Amino Butyric Acids via Reductive Radical Polar Crossover, *Org. Lett.*, 2023, **25**, 3429–3434.
- F. Pasca, Y. Gelato, M. Andresini, D. Serbetci, P. Natho, G. Romanazzi, L. Degennaro, M. Colella and R. Luisi, Continuous Flow Decarboxylative Monofluoroalkylation Enabled by Photoredox Catalysis, *JACS Au*, 2025, **5**, 684–692.
- M. W. Campbell, J. S. Compton, C. B. Kelly and G. A. Molander, Three-Component Olefin Dicarbofunctionalization Enabled by Nickel/Photoredox Dual Catalysis, *J. Am. Chem. Soc.*, 2019, **141**, 20069–20078.
- J. Yi, S. O. Badir, R. Alam and G. A. Molander, Photoredox-Catalyzed Multicomponent Petasis Reaction with Alkyltrifluoroborates, *Org. Lett.*, 2019, **21**, 4853–4858.
- L. Guo, M. Yuan, Y. Zhang, F. Wang, S. Zhu, O. Gutierrez and L. Chu, General Method for Enantioselective Three-Component Carboarylation of Alkenes Enabled by Visible-Light Dual Photoredox/Nickel Catalysis, *J. Am. Chem. Soc.*, 2020, **142**, 20390–20399.
- W. C. Huang, S. Keess and G. A. Molander, Dicarbofunctionalization of 1.1.1 Propellane Enabled by Nickel/Photoredox Dual Catalysis: One-Step Multicomponent Strategy for the Synthesis of BCP-Aryl Derivatives, *J. Am. Chem. Soc.*, 2022, **144**, 12961–12969.
- W. C. Huang, S. Keess and G. A. Molander, One step synthesis of unsymmetrical 1,3-disubstituted BCP ketones via nickel/photoredox-catalyzed 1.1.1 propellane multicomponent dicarbofunctionalization, *Chem. Sci.*, 2022, **13**, 11936–11942.
- S. Li, P. W. Davies and W. Shu, Modular synthesis of  $\alpha$ -arylated carboxylic acids, esters and amides via photocatalyzed triple C–F bond cleavage of methyltrifluorides, *Chem. Sci.*, 2022, **13**, 6636–6641.
- C. Zhang, Z. M. Tang, Y. J. Qiu, J. Tang, S. Q. Ye, Z. M. Li and J. Wu, Access to axially chiral styrenes via a photoinduced asymmetric radical reaction involving a sulfur dioxide insertion, *Chem Catal.*, 2022, **2**, 164–177.
- Q. Li, H. H. Sun, F. Y. Yan, Y. Y. Zhao, Y. C. Zhang, C. Zhou, M. Y. Han, H. J. Li and X. W. Sui, Modular construction of N-arylated amino acid esters enabled by a photoredox-catalyzed multicomponent reaction, *Green Chem.*, 2023, **25**, 6226–6230.
- X. Li, M. Yuan, F. Chen, Z. Huang, F.-L. Qing, O. Gutierrez and L. Chu, Three-component enantioselective alkenylation of organophosphonates via nickel metallaphotoredox catalysis, *Chem*, 2023, **9**, 154–169.



- 25 Z.-K. Wang, Y.-P. Wang, Z.-W. Rao, C.-Y. Liu, X.-H. Pan and L. Guo, General Method for Selective Three-Component Carboacylation of Alkenes via Visible-Light Dual Photoredox/Nickel Catalysis, *Org. Lett.*, 2023, **25**, 1673–1677.
- 26 Q. Zeng, F. Y. Gao, J. Benet-Buchholz and A. W. Kleij, Stereoselective Three-Component Allylic Alkylation Enabled by Dual Photoredox/Ni Catalysis, *ACS Catal.*, 2023, **13**, 7514–7522.
- 27 X. He, E. X. Shi and J. H. Xiao, Enantioselective Synthesis of Benzylic Boronic Esters via Photoredox/Nickel-Catalyzed Multicomponent Cross-Couplings, *Org. Lett.*, 2024, **26**, 9728–9734.
- 28 L. J. Niu, S. Z. Jin, M. Q. Zhu, L. F. Wang and D. Y. Wang, Three-Component 1,2-Carboalkynylation of Alkenes Via Cooperative Nickel/Photoredox Catalysis, *Org. Lett.*, 2025, **27**, 3183–3187.
- 29 L. Guo, F. Song, S. Zhu, H. Li and L. Chu, *syn*-Selective alkyarylation of terminal alkynes via the combination of photoredox and nickel catalysis, *Nat. Commun.*, 2018, **9**, 4543.
- 30 L. Guo, H.-Y. Tu, S. Zhu and L. Chu, Selective, Intermolecular Alkyarylation of Alkenes via Photoredox/Nickel Dual Catalysis, *Org. Lett.*, 2019, **21**, 4771–4776.
- 31 H. Li, L. Guo, X. Feng, L. Huo, S. Zhu and L. Chu, Sequential C–O decarboxylative vinylation/C–H arylation of cyclic oxalates via a nickel-catalyzed multicomponent radical cascade, *Chem. Sci.*, 2020, **11**, 4904–4910.
- 32 W.-H. Sun, J.-Y. Zou, X.-J. Xu, J.-L. Wang, M.-L. Liu and X.-Y. Liu, Photo-Catalyzed Redox-Neutral 1,2-Dialkylation of Alkenes, *Adv. Synth. Catal.*, 2022, **364**, 2260–2265.
- 33 A. Tlahuext-Aca, R. A. Garza-Sanchez and F. Glorius, Multicomponent Oxyalkylation of Styrenes Enabled by Hydrogen-Bond-Assisted Photoinduced Electron Transfer, *Angew. Chem., Int. Ed.*, 2017, **56**, 3708–3711.
- 34 Y. Shen, M.-L. Shen and P.-S. Wang, Light-Mediated Chiral Phosphate Catalysis for Asymmetric Dicarbofunctionalization of Enamides, *ACS Catal.*, 2020, **10**, 8247–8253.
- 35 M. J. Cabrera-Afonso, A. Sookezian, S. O. Badir, M. El Khatib and G. A. Molander, Photoinduced 1,2-dicarbofunctionalization of alkenes with organotrifluoroborate nucleophiles via radical/polar crossover, *Chem. Sci.*, 2021, **12**, 9189–9195.
- 36 W. Dong, E. Yen-Pon, L. Li, A. Bhattacharjee, A. Jolit and G. A. Molander, Exploiting the  $sp^2$  character of bicyclo [1.1.1]pentyl radicals in the transition-metal-free multicomponent difunctionalization of [1.1.1]propellane, *Nat. Chem.*, 2022, **14**, 1068–1077.
- 37 S. Z. Duan, Y. J. Zi, L. L. Wang, J. L. Cong, W. Chen, M. Y. Li, H. B. Zhang, X. D. Yang and P. J. Walsh,  $\alpha$ -Branched amines through radical coupling with 2-azaallyl anions, redox active esters and alkenes, *Chem. Sci.*, 2022, **13**, 3740–3747.
- 38 Y. Li, J. Yang, X. Geng, P. Tao, Y. Shen, Z. Su and K. Zheng, Modular Construction of Unnatural  $\alpha$ -Tertiary Amino Acid Derivatives by Multicomponent Radical Cross-Couplings, *Angew. Chem., Int. Ed.*, 2022, **61**, e202210755.
- 39 X. Xi, Y. Chen and W. Yuan, Nickel-Catalyzed Three-Component Alkylacylation of Alkenes Enabled by a Photoactive Electron Donor-Acceptor Complex, *Org. Lett.*, 2022, **24**, 3938–3943.
- 40 W. Z. Dong, S. Keess and G. A. Molander, Nickel-mediated alkyl-, acyl-, and sulfonylcyanation of 1.1.1 propellane, *Chem Catal.*, 2023, **3**, 100608.
- 41 J. Xu, P. F. Zhang and W. M. Li, Synthesis of BCP nitriles enabled by a metallaphotoredox-based multi-component reaction, *Chem Catal.*, 2023, **3**, 100618.
- 42 J. Z. Wang, E. Mao, J. A. Nguyen, W. L. Lyon and D. W. C. MacMillan, Triple Radical Sorting: Aryl-Alkylation of Alkenes, *J. Am. Chem. Soc.*, 2024, **146**, 15693–15700.
- 43 T. Zhou, Z.-W. Zhang, J. Nie, F. Y. Kwong, J.-A. Ma and C. W. Cheung, Metallaphotocatalytic triple couplings for modular synthesis of elaborate N-trifluoroalkyl anilines, *Nat. Commun.*, 2024, **15**, 9926.
- 44 P. Cao, W. Xiong and T. Yang, Modular Access to Chiral Benzylamines via Ni/Photoredox-Catalyzed Multicomponent Cross-Electrophile Coupling, *Org. Lett.*, 2025, **27**, 1475–1480.
- 45 H.-M. Huang, P. Bellotti, P. M. Pflüger, J. L. Schwarz, B. Heidrich and F. Glorius, Three-Component, Interrupted Radical Heck/Allylic Substitution Cascade Involving Unactivated Alkyl Bromides, *J. Am. Chem. Soc.*, 2020, **142**, 10173–10183.
- 46 S.-Z. Sun, Y. Duan, R. S. Mega, R. J. Somerville and R. Martin, Site-Selective 1,2-Dicarbofunctionalization of Vinyl Boronates through Dual Catalysis, *Angew. Chem., Int. Ed.*, 2020, **59**, 4370–4374.
- 47 P. C. Qian, H. X. Guan, Y. E. Wang, Q. Q. Lu, F. Zhang, D. Xiong, P. J. Walsh and J. Y. Mao, Catalytic enantioselective reductive domino alkyl arylation of acrylates via nickel/photoredox catalysis, *Nat. Commun.*, 2021, **12**, 6613.
- 48 Y.-F. Han, Y. Huang, H. Liu, Z.-H. Gao, C.-L. Zhang and S. Ye, Photoredox cooperative N-heterocyclic carbene/palladium-catalysed alkylacylation of alkenes, *Nat. Commun.*, 2022, **13**, 5754.
- 49 C. Zhu, H. Yue and M. Rueping, Nickel catalyzed multicomponent stereodivergent synthesis of olefins enabled by electrochemistry, photocatalysis and photo-electrochemistry, *Nat. Commun.*, 2022, **13**, 3240.
- 50 M. S. Liu and W. Shu, Rapid Synthesis of  $\beta$ -Chiral Sulfones by Ni-Organophotocatalyzed Enantioselective Sulfonylalkenylation of Alkenes, *JACS Au*, 2023, **3**, 1321–1327.
- 51 M. A. Smith, R. J. D. Kang, R. Kumar, B. Roy and M. J. Gaunt, Modular synthesis of  $\alpha$ -branched secondary alkylamines via visible-light-mediated carbonyl alkylative amination, *Chem. Sci.*, 2024, **15**, 14888–14898.
- 52 J. Zhu, Y. Hong, Y. Wang, Y. Guo, Y. Zhang, Z. Ni, W. Li and J. Xu, Synthesis of 1-(Halo)alkyl-3-heteroaryl Bicyclo[1.1.1]pentanes Enabled by a Photocatalytic Minisci-type Multicomponent Reaction, *ACS Catal.*, 2024, **14**, 6247–6258.



- 53 A. Gimn Khan, R. Kancherla, K. Muralirajan and M. Rueping, Selective Alkylarylation Difunctionalization of 1,3-Butadienes via Nickel/Photoredox Dual Catalysis, *Org. Lett.*, 2025, **27**, 5509–5514.
- 54 A. García-Domínguez, R. Mondal and C. Nevado, Dual Photoredox/Nickel-Catalyzed Three-Component Carbonylation of Alkenes, *Angew. Chem., Int. Ed.*, 2019, **58**, 12286–12290.
- 55 S. Zheng, Z. Chen, Y. Hu, X. Xi, Z. Liao, W. Li and W. Yuan, Selective 1,2-Aryl-Aminoalkylation of Alkenes Enabled by Metallaphotoredox Catalysis, *Angew. Chem., Int. Ed.*, 2020, **59**, 17910–17916.
- 56 X. Yu, C. G. Daniliuc, F. A. Alasmay and A. Studer, Direct Access to  $\alpha$ -Aminosilanes Enabled by Visible-Light-Mediated Multicomponent Radical Cross-Coupling, *Angew. Chem., Int. Ed.*, 2021, **60**, 23335–23341.
- 57 F. Ye, S. L. Zheng, Y. X. Luo, X. T. Qi and W. M. Yuan, Ligand-Controlled Regioreversed 1,2-Aryl-Aminoalkylation of Alkenes Enabled by Photoredox/Nickel Catalysis, *ACS Catal.*, 2024, **14**, 8505–8517.
- 58 W. H. Xiong, H. Y. Xiao, G. Y. Sun, P. T. Cao, Y. Chen, W. R. Yang, M. J. Koh and T. Yang, Enantioselective multicomponent 1,2-arylaminoalkylation of acrylates by photoredox/nickel dual catalysis, *Cell Rep. Phys. Sci.*, 2025, **6**(2), 102435.
- 59 F. J. R. Klauck, H. Yoon, M. J. James, M. Lautens and F. Glorius, Visible-Light-Mediated Deaminative Three-Component Dicarbofunctionalization of Styrenes with Benzylic Radicals, *ACS Catal.*, 2019, **9**, 236–241.
- 60 S. Nandi, P. Das, S. Das, S. Mondal and R. Jana, Visible-light-mediated  $\beta$ -acylative divergent alkene difunctionalization with Katritzky salt/CO<sub>2</sub>, *Green Chem.*, 2023, **25**, 3633–3643.
- 61 S. Das, S. A. Rahaman, K. Pradhan and R. Jana, Organophotoredox-Catalyzed Synthesis of Unnatural  $\alpha/\beta$  Amino Acids and Peptides via Deaminative Three-Component Coupling, *Org. Lett.*, 2024, **26**, 6955–6960.
- 62 A. Trowbridge, D. Reich and M. J. Gaunt, Multicomponent synthesis of tertiary alkylamines by photocatalytic olefin-hydroaminoalkylation, *Nature*, 2018, **561**, 522–527.
- 63 H.-M. Huang, P. Bellotti, S. Kim, X. Zhang and F. Glorius, Catalytic multicomponent reaction involving a ketyl-type radical, *Nat. Synth.*, 2022, **1**, 464–474.
- 64 D. Wang and L. Ackermann, Three-component carboacylation of alkenes via cooperative nickelaphotoredox catalysis, *Chem. Sci.*, 2022, **13**, 7256–7263.
- 65 T. Yang, W. H. Xiong, G. Y. Sun, W. R. Yang, M. D. Lu and M. J. Koh, Multicomponent Construction of Tertiary Alkylamines by Photoredox/Nickel-Catalyzed Aminoalkylation of Organohalides, *J. Am. Chem. Soc.*, 2024, **146**, 29177–29188.
- 66 M. W. Campbell, M. B. Yuan, V. C. Polites, O. Gutierrez and G. A. Molander, Photochemical C-H Activation Enables Nickel-Catalyzed Olefin Dicarbofunctionalization, *J. Am. Chem. Soc.*, 2021, **143**, 3901–3910.
- 67 L.-M. Feng, S. Liu, Y.-H. Tu, P.-X. Rui and X.-G. Hu, Radical Deoxygenative Three-Component Reaction of Alcohols, Aryl Alkenes, and Cyanopyridines, *Org. Lett.*, 2024, **26**, 6225–6229.
- 68 X. Hu, I. Cheng-Sánchez, W. Q. Kong, G. A. Molander and C. Nevado, Nickel-catalysed enantioselective alkene dicarbofunctionalization enabled by photochemical aliphatic C-H bond activation, *Nat. Catal.*, 2024, **7**, 655–665.
- 69 J. Z. Wang, W. L. Lyon and D. W. C. MacMillan, Alkene dialkylation by triple radical sorting, *Nature*, 2024, **628**, 104–109.
- 70 B.-Q. He, L. Zhao, J. Zhang, W.-H. Bao, M. Yang and X. Wu, Alcohol Activation by Benzodithiolylium for Deoxygenative Alkylation Driven by Photocatalytic Energy Transfer, *Angew. Chem., Int. Ed.*, 2025, **64**, e202423795.
- 71 X.-H. Ouyang, Y. Li, R.-J. Song, M. Hu, S. Luo and J.-H. Li, Intermolecular dialkylation of alkenes with two distinct C(sp<sup>3</sup>)-H bonds enabled by synergistic photoredox catalysis and iron catalysis, *Sci. Adv.*, 2019, **5**, eaav9839.
- 72 S. Xu, H. Chen, Z. Zhou and W. Kong, Three-Component Alkene Difunctionalization by Direct and Selective Activation of Aliphatic C-H Bonds, *Angew. Chem., Int. Ed.*, 2021, **60**, 7405–7411.
- 73 Y. Hong, M.-Y. Dong, D.-S. Li and H.-P. Deng, Photoinduced Three-Component Carboarylation of Unactivated Alkenes with Protic C(sp<sup>3</sup>)-H Feedstocks, *Org. Lett.*, 2022, **24**, 7677–7684.
- 74 M. Kim, E. You, J. Kim and S. Hong, Site-Selective Pyridylic C-H Functionalization by Photocatalytic Radical Cascades, *Angew. Chem., Int. Ed.*, 2022, **61**, e202204217.
- 75 W. Liu, C. Liu, M. Wang and W. Kong, Modular Synthesis of Multifunctionalized CF<sub>3</sub>-Allenenes through Selective Activation of Saturated Hydrocarbons, *ACS Catal.*, 2022, **12**, 10207–10221.
- 76 E. C. Gentry and R. R. Knowles, Synthetic Applications of Proton-Coupled Electron Transfer, *Acc. Chem. Res.*, 2016, **49**, 1546–1556.
- 77 P. R. D. Murray, J. H. Cox, N. D. Chiappini, C. B. Roos, E. A. McLoughlin, B. G. Hejna, S. T. Nguyen, H. H. Ripberger, J. M. Ganley, E. Tsui, N. Y. Shin, B. Koronkiewicz, G. Qiu and R. R. Knowles, Photochemical and Electrochemical Applications of Proton-Coupled Electron Transfer in Organic Synthesis, *Chem. Rev.*, 2022, **122**, 2017–2291.
- 78 R. Tyburski, T. Liu, S. D. Glover and L. Hammarström, Proton-Coupled Electron Transfer Guidelines, Fair and Square, *J. Am. Chem. Soc.*, 2021, **143**, 560–576.
- 79 H. G. Yayla, H. Wang, K. T. Tarantino, H. S. Orbe and R. R. Knowles, Catalytic Ring-Opening of Cyclic Alcohols Enabled by PCET Activation of Strong O-H Bonds, *J. Am. Chem. Soc.*, 2016, **138**, 10794–10797.
- 80 K. Zhao, G. Seidler and R. R. Knowles, 1,3-Alkyl Transposition in Allylic Alcohols Enabled by Proton-Coupled Electron Transfer, *Angew. Chem., Int. Ed.*, 2021, **60**, 20190–20195.
- 81 K. Zhao, K. Yamashita, J. E. Carpenter, T. C. Sherwood, W. R. Ewing, P. T. W. Cheng and R. R. Knowles, Catalytic



- Ring Expansions of Cyclic Alcohols Enabled by Proton-Coupled Electron Transfer, *J. Am. Chem. Soc.*, 2019, **141**, 8752–8757.
- 82 E. Ota, H. Wang, N. L. Frye and R. R. Knowles, A Redox Strategy for Light-Driven, Out-of-Equilibrium Isomerizations and Application to Catalytic C–C Bond Cleavage Reactions, *J. Am. Chem. Soc.*, 2019, **141**, 1457–1462.
- 83 D. Q. Thach and R. R. Knowles, Ring Expansions of 1,2-Amino Alcohols to Amine Heterocycles Enabled by Proton-Coupled Electron Transfer, *J. Am. Chem. Soc.*, 2025, **147**, 21818–21823.
- 84 T. Ji, X.-Y. Chen, L. Huang and M. Rueping, Remote Trifluoromethylthiolation Enabled by Organophotocatalytic C–C Bond Cleavage, *Org. Lett.*, 2020, **22**, 2579–2583.
- 85 T. Kikuchi, K. Yamada, T. Yasui and Y. Yamamoto, Synthesis of Benzo-Fused Cyclic Ketones via Metal-Free Ring Expansion of Cyclopropanols Enabled by Proton-Coupled Electron Transfer, *Org. Lett.*, 2021, **23**, 4710–4714.
- 86 X. Wang, Y. Li and X. Wu, Photoredox/Cobalt Dual Catalysis Enabled Regiospecific Synthesis of Distally Unsaturated Ketones with Hydrogen Evolution, *ACS Catal.*, 2022, **12**, 3710–3718.
- 87 K. Liao, F. Wu, J. Chen and Y. Huang, Catalytic cleavage and functionalization of bulky and inert Csp<sup>3</sup>–Csp<sup>3</sup> bonds via a relayed proton-coupled electron transfer strategy, *Cell Rep. Phys. Sci.*, 2022, **3**, 100763.
- 88 K. Liao, C. Y. Chan, S. Liu, X. Zhang, J. Chen and Y. Huang, Photoredox Cleavage of a Csp<sup>3</sup>–Csp<sup>3</sup> Bond in Aromatic Hydrocarbons, *J. Am. Chem. Soc.*, 2023, **145**, 12284–12292.
- 89 S. Horiuchi, M. Oishi, A. Mizutani, H. Takamura, I. Kadota and K. Tanaka, Generation of alkyl radicals via C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bond cleavage of xanthene-based precursors for photocatalytic Giese-type reaction, *Chem. Commun.*, 2025, **61**, 16376–16379.
- 90 X. Hu, L. Ribadeau-Dumas and C. Nevado, Redox-neutral photocatalytic cleavage and *gem*-difluoroalkenylation of lignin linkages, *Sci. Adv.*, 2025, **11**, eady2227.
- 91 T. Matsuo, M. Sano, Y. Sumida and H. Ohmiya, Organic photoredox-catalyzed unimolecular PCET of benzylic alcohols, *Chem. Sci.*, 2025, **16**, 3150–3156.
- 92 L. Huang, T. Ji and M. Rueping, Remote Nickel-Catalyzed Cross-Coupling Arylation via Proton-Coupled Electron Transfer-Enabled C–C Bond Cleavage, *J. Am. Chem. Soc.*, 2020, **142**, 3532–3539.
- 93 L. Huang, T. Ji, C. Zhu, H. Yue, N. Zhumabay and M. Rueping, Bioinspired desaturation of alcohols enabled by photoredox proton-coupled electron transfer and cobalt dual catalysis, *Nat. Commun.*, 2022, **13**, 809.
- 94 N. Varabyeva, M. Barysevich, Y. Aniskevich and A. Hurski, Ti(O<sup>i</sup>Pr)<sub>4</sub>-Enabled Dual Photoredox and Nickel-Catalyzed Arylation and Alkenylation of Cyclopropanols, *Org. Lett.*, 2021, **23**, 5452–5456.
- 95 J. Wang and X. Li, Asymmetric β-arylation of cyclopropanols enabled by photoredox and nickel dual catalysis, *Chem. Sci.*, 2022, **13**, 3020–3026.
- 96 Y. Patehebieke, R. Charaf, H. P. Bryce-Rogers, K. Ye, M. Ahlquist, L. Hammarström and C.-J. Wallentin, β-Scission of Secondary Alcohols via Photosensitization: Synthetic Utilization and Mechanistic Insights, *ACS Catal.*, 2024, **14**, 585–593.
- 97 Y. Patehebieke, R. Charaf, K. B. Pal, B. M. Baamonde, A. Brnovic, L. Hammarström and C.-J. Wallentin, PCET-mediated deconstructive cross-coupling of aliphatic alcohols, *Chem. Sci.*, 2025, **16**, 7720–7729.
- 98 S. Q. Zhu, X. Zhao, H. Li and L. L. Chu, Catalytic three-component dicarbofunctionalization reactions involving radical capture by nickel, *Chem. Soc. Rev.*, 2021, **50**, 10836–10856.
- 99 P. Gao, Y. J. Niu, F. Yang, L. N. Guo and X. H. Duan, Three-component 1,2-dicarbofunctionalization of alkenes involving alkyl radicals, *Chem. Commun.*, 2022, **58**, 730–746.
- 100 J. W. Tucker and C. R. J. Stephenson, Shining Light on Photoredox Catalysis: Theory and Synthetic Applications, *J. Org. Chem.*, 2012, **77**, 1617–1622.
- 101 S. Ni, R. Halder, D. Ahmadli, E. J. Reijerse, J. Cornella and T. Ritter, C–heteroatom coupling with electron-rich aryls enabled by nickel catalysis and light, *Nat. Catal.*, 2024, **7**, 733–741.
- 102 X. Du, I. Cheng-Sánchez and C. Nevado, Dual Nickel/Photoredox-Catalyzed Asymmetric Carbosulfonylation of Alkenes, *J. Am. Chem. Soc.*, 2023, **145**, 12532–12540.
- 103 L. Anghileri, H. Baunis, A. R. Bena, C. Giannoudis, J. H. Burke, S. Reischauer, C. Merschjann, R. F. Wallick, T. Al Said, C. E. Adams, G. Simionato, S. Kovalenko, L. Dell'Amico, R. M. van der Veen and B. Pieber, Evidence for a Unifying Ni<sup>I</sup>/Ni<sup>III</sup> Mechanism in Light-Mediated Cross-Coupling Catalysis, *J. Am. Chem. Soc.*, 2025, **147**, 13169–13179.
- 104 N. A. Till, L. Tian, Z. Dong, G. D. Scholes and D. W. C. MacMillan, Mechanistic Analysis of Metallaphotoredox C–N Coupling: Photocatalysis Initiates and Perpetuates Ni(I)/Ni(III) Coupling Activity, *J. Am. Chem. Soc.*, 2020, **142**, 15830–15841.
- 105 N. A. Till, S. Oh, D. W. C. MacMillan and M. J. Bird, The Application of Pulse Radiolysis to the Study of Ni(I) Intermediates in Ni-Catalyzed Cross-Coupling Reactions, *J. Am. Chem. Soc.*, 2021, **143**, 9332–9337.

