

# Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. Z. Bacho, S. Wu, T. Muramatsu, C. Nagababu, D. Harano, S. Ochiai and N. Shibata, *Chem. Sci.*, 2026, DOI: 10.1039/D6SC01441K.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## ARTICLE

**Bidirectional skeletal remodelling of SF<sub>5</sub>-nitrobenzenes into azepine, bicyclic, and benzimidazole frameworks****Muhamad Zulfaqar Bacho,<sup>a</sup> Shiwei Wu,<sup>b</sup> Takuya Muramatsu,<sup>b</sup> Chavakula Nagababu,<sup>a</sup> Daiki Harano,<sup>b</sup> Seishu Ochiai<sup>b</sup> and Norio Shibata<sup>\* a,b</sup>**Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

**Abstract:** The development of PFAS-free fluorinated scaffolds that preserve the desirable physicochemical attributes of perfluoroalkyl groups remains a central challenge in contemporary fluorine chemistry. Herein, we report a rapid and bidirectional skeletal-remodelling platform that enables controlled interconversion between aromatic, medium-sized, and bicyclic SF<sub>5</sub>-containing heterocycles from readily accessible SF<sub>5</sub>-nitrobenzenes. Phosphorus-catalyzed iterative deoxygenation of SF<sub>5</sub>-nitrobenzenes generates aryl nitrene intermediates that undergo remarkably accelerated dearomative ring expansion, furnishing seven-membered SF<sub>5</sub>-azepines within dramatically shortened reaction times compared to non-SF<sub>5</sub> analogues. These azepines function as versatile skeletal nodes, enabling divergent downstream transformations: photoinduced 4π-electrocyclization provides access to previously unexplored SF<sub>5</sub>-azabicyclo[3.2.0]hepta-2,6-diene frameworks, while selective fluoroacylative activation promotes reverse skeletal reconstruction to restore aromaticity and deliver SF<sub>5</sub>-substituted benzimidazoles. Collectively, this work demonstrates that strategic incorporation of the SF<sub>5</sub> group not only expands accessible heterocyclic architectures but also fundamentally alters skeletal rearrangement kinetics, enabling rapid and controllable skeletal editing from a common, practical precursor. Given the OECD classification of SF<sub>5</sub>-containing molecules as non-PFAS, this unified skeletal-remodelling approach substantially broadens the design space of fluorinated scaffolds for applications in pharmaceuticals, agrochemicals, and functional materials, advancing the principles of sustainable fluorine chemistry.

**Introduction**

Fluorinated heterocycles are indispensable motifs in agrochemicals, pharmaceuticals, and functional materials, as fluorine substitution profoundly enhances lipophilicity, metabolic stability, and target engagement.<sup>1</sup> In agrochemical design, CF<sub>3</sub>-substituted heterocycles are particularly dominant, accounting for more than 40% of all fluorinated agrochemicals.<sup>1b</sup> However, CF<sub>3</sub> groups are classified as per- and polyfluoroalkyl substances (PFAS) under the OECD definition, and their extreme environmental persistence has become a serious global concern.<sup>2</sup> The exceptional stability of C–F bonds leads to long-term accumulation and the formation of highly mobile degradation products such as trifluoroacetic acid (TFA, CF<sub>3</sub>CO<sub>2</sub>H).<sup>3</sup> While CF<sub>3</sub> groups remain manageable in pharmaceuticals,<sup>1a</sup> their intentional environmental release via agrochemicals<sup>1b,3</sup> is increasingly unsustainable. Consequently,

the development of PFAS-free fluorinated alternatives that retain the functional advantages of CF<sub>3</sub> substituents has emerged as a critical challenge.

The pentafluorosulfanyl (SF<sub>5</sub>) group represents a compelling solution.<sup>4,5</sup> As a hypervalent sulfur–fluorine functionality, SF<sub>5</sub> combines strong electron-withdrawing character, high lipophilicity, and substantial steric bulk, and is therefore widely regarded as a “super-CF<sub>3</sub>” substituent. Importantly, unlike CF<sub>3</sub>, SF<sub>5</sub> does not rely on persistent C–F bonds and exhibits the potential for environmentally benign mineralization, positioning SF<sub>5</sub>-containing molecules as classified as PFAS-free.<sup>2</sup> In molecular design, SF<sub>5</sub> functions as a bioisostere for CF<sub>3</sub>, *tert*-butyl, and nitro groups, with comparable hydrophobicity and intrinsic volume (Figure 1a). Despite these advantages, the structural diversity of SF<sub>5</sub>-containing molecules—particularly SF<sub>5</sub>-heterocycles—remains severely limited.<sup>5c</sup>


This limitation arises from a fundamental synthetic bottleneck. Traditional SF<sub>5</sub> installation typically relies on radical reactions with alkenes or alkynes using SF<sub>5</sub>Cl or related reagents, which significantly limits substrate scope.<sup>5h</sup> In contrast, oxidative chloro-fluorination of aryl sulfides and related protocols<sup>4b</sup> require harsh reaction conditions, specialized equipment, and stringent moisture control, rendering their application to the synthesis of SF<sub>5</sub>-heterocycles highly limited.<sup>6</sup> As a result, SF<sub>5</sub> chemistry has remained largely limited to simple alkyl, alkenyl,

<sup>a</sup> Department of Nanopharmaceutical Sciences, Nagoya Institute of Technology, Gokiso, Showa-ku, Nagoya 466-8555, Japan

<sup>b</sup> Department of Engineering, Nagoya Institute of Technology, Gokiso, Showa-ku, Nagoya 466-8555, Japan

<sup>\*</sup> Dedicated to Professor David O'Hagan to mark his retirement from the University of St Andrews (UK) and his significant contributions to the field of organofluorine chemistry.





and alkynyl motifs. Only recently have new classes of SF<sub>5</sub> reagents or new protocols been developed,<sup>7</sup> expanding the diversity of accessible substrates and reactivity profiles. Nevertheless, structurally complex heterocycles—particularly those incorporating SF<sub>5</sub> or SF<sub>4</sub><sup>8</sup>—remain largely unexplored and synthetically inaccessible.

Concurrently, skeletal rearrangements have emerged as a powerful strategy for heterocycle synthesis, enabling rapid access to strained and unconventional architectures that are difficult to construct using classical cyclization methods.<sup>9</sup> We recognized that this framework-reorganization approach is uniquely suited to SF<sub>5</sub> chemistry, where conventional ring-construction strategies are intrinsically constrained. Consistent with this concept, we recently reported the photo-induced synthesis of SF<sub>5</sub>-azepine aryl ethers from SF<sub>5</sub>-azido benzene with phenols via skeletal rearrangement, thereby establishing this approach as a viable entry to medium-sized SF<sub>5</sub>-heterocycles (Figure 1b).<sup>10</sup> Notably, the strong electron-withdrawing nature of the SF<sub>5</sub> substituent was found to facilitate the key ring-expansion step from benzene to azepine frameworks. Encouraged by these findings, we herein extend this concept to sequential skeletal remodelling, enabling access to a range of previously inaccessible SF<sub>5</sub>-containing heterocycles, including SF<sub>5</sub>-azepin-2-amines **3**, SF<sub>5</sub>-azabicyclo[3.2.0]hepta-2,6-dienes **4**, and SF<sub>5</sub>-benzimidazoles **5**. In particular, the SF<sub>5</sub>-azabicyclo[3.2.0]hepta-2,6-diene derivatives represent the first examples of this rigid bicyclic scaffold bearing an SF<sub>5</sub> substituent (Figure 1c). Importantly, the synthesis originates from readily available SF<sub>5</sub>-nitrobenzenes **1**, rather than inherently energetic SF<sub>5</sub>-azido benzenes, and proceeds through consecutive skeletal rearrangements. These transformations establish a unified skeletal-editing manifold


that interconverts aromatic, medium-sized, and bicyclic SF<sub>5</sub>-containing architectures.

DOI: 10.1039/D6SC01441K

## Results and discussion

### Development of a rapid skeletal rearrangement to amino-SF<sub>5</sub>-azepines


Seven-membered SF<sub>5</sub>-heterocycles are essentially unexplored, despite the prevalence of azepine frameworks in biologically active molecules and marketed drugs.<sup>11</sup> Given our interest in 2-amino-SF<sub>5</sub>-azepines, we first examined the skeletal remodelling of SF<sub>5</sub>-azido benzene **6** using diethylamine **2a** as the nucleophile in the presence of DABCO, following our reported LED conditions (Figure 2).<sup>10</sup> Under these conditions, the desired 2-diethylamino-SF<sub>5</sub>-azepine **3a** was obtained in 45% yield after 48 h at room temperature, confirming the feasibility of azepine formation from SF<sub>5</sub>-azido benzene, albeit with limited efficiency and prolonged reaction times. We next turned to an alternative nitrene-generation strategy based on phosphorus-catalyzed deoxygenation of SF<sub>5</sub>-nitrobenzene, as developed by Radosevich and co-workers.<sup>12</sup> Under these conditions, phenylnitrene are generated thermally via P(III)/P(V)=O redox cycling from P(V)=O-catalyst with phenylsilane. Pleasingly, application of this protocol to SF<sub>5</sub>-nitrobenzene **1a** proved markedly more effective, delivering the corresponding SF<sub>5</sub>-azepine **3a** in 72% yield within only 1 h at 120 °C. Notably, in non-SF<sub>5</sub> systems, analogous nitrobenzene-to-azepine conversions under Radosevich conditions typically require approximately 12 h to reach completion.<sup>12</sup> To further probe the origin of this pronounced rate enhancement, we evaluated nitrobenzene **7** and fluoro-substituted nitrobenzene **8** under identical conditions. In both cases, the corresponding azepin-2-



Scheme 1. Substrate scope for skeletal rearrangement to amino-SF<sub>5</sub>-azepines

View Article Online

DOI: 10.1039/D6SC01441K



<sup>a</sup>Unless otherwise noted, the reactions were carried out with **1a** (0.2 mmol, 1.0 equiv), **2** (0.4 mmol, 2.0 equiv), **P(V)=O-catalyst** (15 mol%), PhSiH<sub>3</sub> (0.4 mmol, 2.0 equiv), in toluene (0.4 mL, 0.5 M) stirred at 120 °C for 1 hour. Isolated yields are given. <sup>b</sup>A mixture of **1a** (0.1 mmol, 1.0 equiv), **2a** (0.2 mmol, 2.0 equiv), **P(V)=O-catalyst** (15 mol%), PhSiH<sub>3</sub> (0.2 mmol, 2.0 equiv), in toluene (0.2 mL, 0.5 M) stirred at 120 °C for 1 hour. Isolated yields are given. <sup>c</sup>A mixture of **1a** (4.0 mmol, 1.0 equiv), **2a** (8.0 mmol, 2.0 equiv), **P(V)=O-catalyst** (15 mol%), PhSiH<sub>3</sub> (8.0 mmol, 2.0 equiv), in toluene (8.0 mL, 0.5 M) stirred at 120 °C for 1 hour. Isolated yields are given. <sup>d</sup>Yield was determined by <sup>19</sup>F-NMR spectroscopy (fluorobenzene as standard).

amines (**10** and **11**) were obtained in only 16–17% yield after 1 h, with yields increasing gradually to 43–45% after 12 h. In sharp contrast, the SF<sub>5</sub>-substituted nitrobenzene underwent rapid and efficient skeletal rearrangement, suggesting that the strongly electron-withdrawing SF<sub>5</sub> substituent significantly accelerates the ring-expansion process, likely by facilitating aryl nitrene insertion and subsequent electrocyclic rearrangement.<sup>10</sup> Given this substantial rate enhancement and the improved safety and practicality of using SF<sub>5</sub>-nitrobenzenes instead of SF<sub>5</sub>-azido benzenes, we undertook a systematic investigation of the phosphorus-catalyzed conversion of SF<sub>5</sub>-nitrobenzenes into SF<sub>5</sub>-azepines. A similar acceleration in the

ring-expansion was observed for trifluoromethyl-substituted nitrobenzene **9** under standard conditions, affording azepine-2-amine **12** in 83% yield within 1 h, compared to 12 h in the reported method.<sup>12</sup>




Substrate scope for skeletal rearrangement to amino-SF<sub>5</sub>-azepines

The generality of the phosphorus-catalyzed ring-expansion reaction was first evaluated with a broad range of acyclic secondary amines (Scheme 1, top). Straight-chain dialkylamines such as diethylamine (**2a**), dibutylamine (**2b**), and didecylamine (**2c**) were smoothly converted into the corresponding azepines **3aa–3ac** in 57–66% yields. Branched secondary amines were



Scheme 2. Skeletal editing of azepines: transformations to bicyclic SF<sub>5</sub>-azabicyclo[3.2.0]hepta-2,6-dienes and SF<sub>5</sub>-benzimidazoles

DOI: 10.1039/D6SC01441K

a) Skeletal rearrangement to bicyclic SF<sub>5</sub>-azabicyclo[3.2.0]hepta-2,6-dienes<sup>a</sup>b) Skeletal rearrangement to SF<sub>5</sub>-benzimidazoles<sup>b</sup>c) Plausible mechanism for SF<sub>5</sub>-benzimidazoles

<sup>a</sup>Reaction conditions: **3** (0.1 mmol, 1.0 equiv) in MeCN (3.0 mL, 0.033M) irradiated under (365 nm) at rt for 40 hours. Isolated yields are given. <sup>b</sup>Reaction conditions: **3** (0.1 mmol, 1.0 equiv), DABCO (0.11 mmol, 1.1 equiv), TFAA or DFAA (0.33 mmol, 3.3 equiv) in toluene (2.0 mL, 0.05 M) at 120 °C for 24 hours. Isolated yields are given.

also compatible, although increased steric demand led to somewhat diminished efficiencies, affording products **3ad** and **3ae** in 29–51% yields. An electron-rich ether-substituted amine (**2f**) was well tolerated, delivering **3af** in 74% yield. Aromatic and alicyclic secondary amines, including dibenzylamine (**2g**), dicyclohexylamine (**2h**), and unsymmetrical amines (**2i–2j**), furnished azepines **3ag–3aj** in 30–72% yields. The influence of cyclic amine ring size was next evaluated. Four-, five-, six-, and seven-membered cyclic amines (**2k–2n**) participated successfully, affording azepines **3ak–3an** in moderate yields

(38–59%). Notably, heteroatom-containing cyclic amines also proved to be competent nucleophiles: morpholine (**2o**), thiomorpholine (**2p**), and difluorinated piperidine (**2q**) delivered products **3ao–3aq** in good to excellent yields (47–76%). In addition, a complex aryloxy-substituted piperidine (**2r**) underwent smooth coupling to afford **3ar** in 75% yield, highlighting the functional-group tolerance of the process.

To further assess the generality of the ketenimine intermediate derived from **1a**, we investigated nucleophiles bearing diverse functional groups. Esters (**2s**), alcohols (**2t**),



amides (**2u**), and bicyclic amines (**2v**) were all successfully incorporated, furnishing products **3as–3av** in 48–83% yields. The structure of **3av** was unambiguously confirmed by single-crystal X-ray diffraction (CCDC 2501744). Notably, this transformation is applicable not only to *p*-substituted SF<sub>5</sub>-nitrobenzene **1a** but also to *m*-substituted analogue **1b**, which afforded the regioselective product **3ba** in 40% isolated yield.

Encouraged by the breadth of secondary amine compatibility, we extended the methodology to primary amines (Scheme 1, middle). Benzylamines bearing either electron-donating or electron-withdrawing substituents at the para position (**2w–2aa**) underwent efficient ring expansion to give products **3aw–3aaa** in 36–66% yields. Primary aliphatic amines (**2ab** and **2ac**) were likewise competent substrates, affording azepines **3aab** and **3aac** in 39–57% yields.

Finally, the synthetic utility of this transformation was demonstrated through late-stage functionalization of biologically relevant amines (Scheme 1, bottom). Complex drug-derived amines, including fluoxetine (**2ad**), betahistine (**2ae**), duloxetine (**2af**), and sertraline (**2ag**), were successfully converted into the corresponding SF<sub>5</sub>-azepine derivatives **3aad–3aag** in good yields (60–69%). These results underscore the robustness of the method and its potential for late-stage diversification in medicinal and agrochemical discovery.

Anilines were not suitable as amine nucleophiles; instead of azepines **3**, N–N bond coupling products were formed, consistent with previous reports.<sup>13</sup>

### Second skeletal rearrangement to bicyclic SF<sub>5</sub>-azabicyclo[3.2.0]hepta-2,6-dienes

With gram-scale quantities of azepine **3aa** available, we next investigated the photochemical 4π-electrocyclization developed by Leonori<sup>14</sup> for the synthesis of bicyclic SF<sub>5</sub>-attached azabicyclo[3.2.0]hepta-2,6-dienes (Scheme 2a, top). Under simple irradiation of azepines **3** at 365 nm in acetonitrile at room temperature cleanly delivered the corresponding bicyclic products, SF<sub>5</sub>-attached azabicyclo[3.2.0]hepta-2,6-dienes **4** in moderate to good yields. SF<sub>5</sub>-azepines **3** bearing electron-donating alkyl substituents, including ethyl (**3aa**), *n*-butyl (**3ab**), *n*-decyl (**3ac**), and 2-methoxyethyl (**3af**) groups, were well tolerated, affording bicyclic derivatives **4aa**, **4ab**, **4ac**, and **4af** in yields of up to 67%. Azepines containing unsymmetrical secondary amines (**3ai** and **3aj**) also underwent smooth skeletal rearrangement to furnish bicyclic products **4ai** and **4aj** in 40–55% yields. Notably, this photochemical transformation proved compatible with cyclic amine-substituted azepines: substrates **3ao–3as** were efficiently converted into the corresponding bicyclic compounds **4ao–4as** in consistently good yields (53–62%). Furthermore, SF<sub>5</sub>-azepines bearing primary amine substituents, exemplified by 4-bromobenzylamine-derived azepine **3az**, participated smoothly to afford bicyclic product **4az** in 58% yield. The relative stereochemistry of the bicyclic products **4** was established by detailed 2D NMR analysis of compound **4az** (see Supporting Information) and was found to be fully consistent with previously reported azabicyclo[3.2.0] frameworks.

### Third skeletal rearrangement to SF<sub>5</sub>-benzimidazoles

View Article Online

DOI: 10.1039/D6SC01441K

To further probe the structural plasticity of the SF<sub>5</sub>-azepine framework and to demonstrate bidirectional skeletal editing from a common intermediate, we investigated the re-aromatization of SF<sub>5</sub>-azepines to SF<sub>5</sub>-benzimidazoles. Following a protocol related to that reported by Radosevich and co-workers,<sup>12</sup> a primary amine-substituted azepine (**3aw**, derived from benzylamine) was treated with trifluoroacetic anhydride (TFAA) in the presence of DABCO in toluene at 120 °C for 24 h. Under these conditions, re-aromatization of the azepine core was accompanied by intramolecular cyclodehydration, affording the corresponding SF<sub>5</sub>-benzimidazole **5a** in 53% yield (Scheme 2b, middle). This cyclodehydration is plausibly driven by electrophilic activation of the azepine nitrogen by the acyl anhydride, followed by intramolecular nucleophilic attack of the pendant amine and subsequent loss of water to restore aromaticity (Scheme 2c, bottom)

Because the CF<sub>3</sub> group is classified as a PFAS motif, we next examined the transformation using difluoroacetic anhydride (DFAA) under otherwise identical conditions. As anticipated, the reaction proceeded efficiently to deliver the CF<sub>2</sub>H-substituted SF<sub>5</sub>-benzimidazole **5b** in 80% yield. Notably, this skeletal reconstruction pathway proved tolerant of diverse substituents on the aromatic ring, including OMe, F, Br, and Cl, furnishing a series of CF<sub>2</sub>H-substituted SF<sub>5</sub>-benzimidazoles (**5c–5f**) in good to excellent yields.

### Conclusions

This study demonstrates that skeletal rearrangement of SF<sub>5</sub>-nitrobenzenes provides a general and efficient strategy for constructing diverse SF<sub>5</sub>-heterocycles, including SF<sub>5</sub>-azepines, SF<sub>5</sub>-azabicyclo[3.2.0] frameworks, and SF<sub>5</sub>-benzimidazoles. Importantly, these motifs represent PFAS-free alternatives to CF<sub>3</sub>-based scaffolds, offering structurally rich and functionally attractive building blocks for pharmaceuticals, agrochemicals, and materials science. In light of the OECD classification of SF<sub>5</sub>-containing molecules as non-PFAS, this work contributes to the advancement of sustainable fluorine chemistry by demonstrating that strategic skeletal remodelling can unlock both reactivity and chemical space unique to SF<sub>5</sub> substitution.

### Author contributions

MZB optimized the reaction conditions, surveyed the substrate scope, analyzed the data and discussed the results with NS. SW, TM, and CN prepared starting materials and attempted reactions. HD helped to prepare the starting materials. SO took the X-ray crystallography of **3av**. MZB wrote the initial draft and NS wrote the manuscript. NS supervised the study. All authors contributed to the manuscript and approved the final version of the manuscript.

### Conflicts of interest

There are no conflicts to declare.



## Data availability

CCDC 2501744 contains the supplementary crystallographic data for this paper.

The data that support the findings of this study are available within the article and the supplementary information (SI). Supplementary information: materials and methods, experimental procedures, characterization data, and NMR spectra. See DOI:

## Acknowledgements

This study was supported by the CREST program of the Japan Science and Technology Agency, entitled "Precise Material Science for Degradation and Stability" (grant number: JPMJCR21L1), by Dr Seiji Motojima (CMC Research Institute, Japan), and Kenji Koike (Tosoh Corporation).

## References

- (a) M. Inoue, Y. Sumii and N. Shibata, *ACS Omega*, 2020, **5**, 10633–10640. (b) Y. Ogawa, E. Tokunaga, O. Kobayashi, K. Hirai and N. Shibata, *iScience*, 2020, **23**, 101467. (c) R. Berger, G. Resnati, P. Metrangolo, E. Weber and J. Hulliger, *Chem. Soc. Rev.*, 2011, **40**, 3496–3508.
- (a) Z. Wang, A. M. Buser, I. T. Cousins, S. Demattio, W. Drost, O. Johansson, K. Ohno, G. Patlewicz, A. M. Richard, G. W. Walker, G. S. White and E. Leinala, *Environ. Sci. Technol.*, 2021, **55**, 15575–15578. (b) E. L. Schymanski, J. Zhang, P. A. Thiessen, P. Chirsir, T. Kondic and E. E. Bolton, *Environ. Sci. Technol.* 2023, **57**, 16918–16928.
- (a) H. Joerss, F. Freeling, S. Van Leeuwen, J. Hollender, X. Liu, K. Nödler, Z. Wang, B. Yu, D. Zahn and G. Sigmund, *Environ. Int.* 2024, **193**, 109061. (b) A. Alinezhad, H. Shao, K. Litvanova, R. Sun, A. Kubatova, W. Zhang, Y. Li and F. Xiao, *Environ. Sci. Technol.*, 2023, **57**, 8796–8807. (c) H. P. H. Arp, A. Gredelj, J. Glüge, M. Scheringer and I. T. Cousins, *Environ. Sci. Technol.*, 2024, **58**, 19925–19935. (d) D. A. Jackson and S. A. Mabury, *Environ. Toxicol. Chem.*, 2009, **28**, 1866–1873.
- (a) W. A. Sheppard, *J. Am. Chem. Soc.*, 1960, **82**, 4751–4752. (b) T. Umemoto, L. M. Garrick and N. Saito, *Beilstein J. Org. Chem.*, 2012, **8**, 461–471.
- (a) P. R. Savoie and J. T. Welch, *Chem. Rev.*, 2015, **115**, 1130–1190 (b) G. Haufe *Tetrahedron*, 2022, **109**, 132656. (c) P. Das, E. Tokunaga and N. Shibata, *Tetrahedron Lett.*, 2017, **58**, 4803–4815. (d) M. F. Sowailah, R. A. Hazlitt and D. A. Colby, *Chem. Med. Chem.*, 2017, **12**, 1481–1490. (e) J. M. W. Chan, *J. Mater. Chem. C*, 2019, **7**, 12822–12834. (f) M. Sani and M. Zanda, *Synthesis*, 2022, **54**, 4184–4209. (g) Y. Yang, L. Han, L. Brettnacher, L. Canavero, A. Tlili, *Chem. Rev.* 2025, **125**, 8426–8476. (h) J. Shou and F. Qing, *Chin. J. Chem.*, 2025, **43**, 3033–3042.
- (a) O. S. Kanishchev and W. R. Dolbier, *Angew. Chem. Int. Ed.*, 2015, **54**, 280–284. (b) M. Kosobokov, B. Cui, A. Balia, K. Matsuzaki, E. Tokunaga, N. Saito and Shibata, *Angew. Chem. Int. Ed.*, 2016, **55**, 10781–10785. (c) B. Cui, M. Kosobokov, K. Matsuzaki, E. Tokunaga, and N. Shibata, *Chem. Comm.* 2017, **53**, 5997–6000. (d) B. Cui, S. Jia, E. Tokunaga, N. Saito, and Shibata, *Chem. Commun.* 2017, **53**, 12738–12741. (e) Saidalimu, Y. Liang, K. Niina, K. Tanagawa, N. Saito and N. Shibata, *Org. Chem. Front.*, 2019, **6**, 1157–1161. (f) K. Tanagawa, Z. Zhao, N. Saito and N. Shibata, *Bull. Chem. Soc. Jpn.*, 2021, **94**, 1682–1684.
- Selected recent papers. (a) J. Shou, X. Xu and F. Qing, *Angew. Chem. Int. Ed.*, 2021, **60**, 15271–15275. (b) Y. Jiang, X. Meng, J. Zhang, G. Wu, X. Lin and S. Guo, *Nat. Commun.*, 2024, **15**, 9705. (c) M. Birepinte, P. A. Champagne and J. Paquin, *Angew. Chem. Int. Ed.*, 2021, **61**, e202112575. (d) L. Verret, K. Burchell-Reyes, J. F. Morin and J. F. Paquin, *J. Fluor. Chem.* 2025, **282**, 110387. (e) J. Y. Shou, F. L. Qing, *Angew. Chem., Int. Ed.* 2022, **61**, e202208860. (f) Z. Fang, J. Y. Shou, M. N. Khan, X. B. He, X. Jia and F. L. Qing, *J. Org. Chem.* 2025, **90**, 7841–7848. (g) Y. Kraemer, J. A. Buldt, W. Kong, A. M. Stephens, A. N. Ragan, S. Park, Z. C. Haidar, A. H. Patel, R. Shey, R. Dagan, C. P. McLoughlin, J. C. Fettinger, D. J. Tantillo and C. R. Pitts, *Angew. Chem. Int. Ed.*, 2024, **136**, e202319930. (h) Y. Y. Jiang, J. S. Zhang, D. Sun, G. Wu, Z.-C. Kuang, N. Zhang, X. Y. Shen, X. L. Zhang, J. Jiang, H. Y. Yang, W. Zhang, X. S. Xue and S. Guo, *J. Am. Chem. Soc.*, 2025, **147**, 43785–43797. (i) R. Li, C. Liu, C. Hu, J. Tsien, M. Hayes, S.-J. Chen, Y. Kanda, R. R. Merchant, B. S. Matsuura and T. Qin, *Nat. Commun.*, 2025, **16**, 10185. (j) Y. Jiang, X. Meng, J. Zhang, G. Wu, X. Lin and S. Guo, *Nat. Commun.*, 2024, **15**, 9705. (k) J. Y. Li, Y. Liu, D. Sun, G. Wu, S. S. Yue, H. Y. Yang, H. Zhang, W. Zhang, M. M. Zheng, X. S. Xue and S. Guo, *J. Am. Chem. Soc.*, 2025, **147**, 26124–26132. (l) R. Li, C. Hu, C. Liu, T. Lyness, W. Li, C. Y. Cai, E. Crossley, Y. Kanda, R. R. Merchant, B. S. Matsuura, N. S. Williams and T. Qin, *J. Am. Chem. Soc.*, 2025, **147**, 34218–34224. (m) Y. Kraemer, S. Park, W. Y. Kong, Y. Chen, A. J. Witt, J. A. Buldt, A. N. Ragan, L. M. Holder, D. J. Tantillo and C. R. Pitts, *J. Am. Chem. Soc.*, 2025, **147**, 27892–27904. (n) Y. Yang, L. Han, L. Canavero, L. Brettnacher, L. Khrouz, C. Monnereau, M. Médebielle and A. Tlili, *J. Am. Chem. Soc.*, 2025, **147**, 27905–27911. (o) H. Y. Yang, H. B. Du, Y. P. Bai, Y. H. Ding, Y.-F. Ma, G. Wu, N. Zhang, N. K. Szymczak and S. Guo, *Sci. Adv.*, 2025, **11**, eadw8408.
- (a) P. Das, K. Niina, T. Hiromura, E. Tokunaga, N. Saito and N. Shibata, *Chem. Sci.*, 2018, **9**, 4931. (b) T. Aggarwal, K. Hada, Y. Murata, Y. Sumii, K. Tanagawa, K. Niina, S. Mori, J. Escorihuela and N. Shibata, *Angew. Chem., Int. Ed.*, 2023, **62**, e202307090. (c) Y. Murata, K. Hada, T. Aggarwal, J. Escorihuela and N. Shibata, *Angew. Chem., Int. Ed.*, 2024, **63**, e202318086. (d) S. R. Narra, M. Z. Bacho, M. Hattori and N. Shibata, *Adv. Sci.*, 2024, **11**, 2306554.
- (a) J. Li, P. Tang, Y. Fan and H. Lu, *Science*, 2025, **389**, 275–281. (b) M. Puriş, H. Nakahara and M. D. Levin, *Science*, 2025, **389**, 295–298. (c) B. Ghosh, P. Kafle, R. Mukherjee, R. Welles, D. Herndon, K. M. Nicholas, Y. Shao and I. Sharma, *Science*, 2025, **387**, 102–107. (d) Z. Wang, P. Xu, S. M. Guo, C. G. Daniliuc and A. Studer, *Nature*, 2025, **642**, 92–98. (e) J. Qi, C. Wang, G. Wang, P. O'Neill, S. R. Dubbaka, H. T. Ang, X. Chen and J. Wu, *Angew. Chem. Int. Ed.*, 2024, **64**, e202413723. (f) R. T. Steele, M. Fujiu and R. Sarpong, *Science*, 2025, **388**, 631–638. (g) P. Rai, S. Naik, K. Gupta, K. Maji, G. Jindal and B. Maji, *Nat. Commun.*, 2025, **16**, 2991.
- C. Nagababu, T. Muramatsu, M. Z. Bacho, S. Wu, S. Ochiai, J. Escorihuela and N. Shibata, *Chem. Sci.*, 2025, **17**, 1840–1849.
- (a) M. Kaur, S. Garg, D. S. Malhi, S. H. Sohal, *Curr. Org. Chem.*, 2021, **25**, 449–506. (b) A. Dorababu, *Polycycl. Aromat. Comp.*, 2022, **43**, 2250–2268. (c) D. McTavish and P. Benfield, *Drugs*, 1990, **39**, 136–153. (d) T. Miyazaki, H. Fujiki, Y. Yamamura, S. Nakamura and T. Mori, *Cardiovasc. Drug Rev.*, 2007, **25**, 1–13. (e) J. A. Balfour and K. L. Goa, *Drugs*, 1991, **42**, 511–539. (f) M. Seto, Y. Aramaki, T. Okawa, N. Miyamoto, K. Aikawa, N. Kanzaki, S. Niwa, Y. Iizawa and M. Shiraiishi, *Chem. Pharm. Bull.*, 2004, **52**, 577–590.
- G. Li, M. N. Lavagnino, S. Z. Ali, S. Hu and A. T. Radosevich, *J. Am. Chem. Soc.*, 2022, **145**, 41–46.
- G. Li, S. P. Miller and A. T. Radosevich, *J. Am. Chem. Soc.* 2021, **143**, 14464–14469.



- 14 E. Matador, M. J. Tilby, I. Saridakis, M. Pedrón, D. Tomczak, J. Llaveria, I. Atodiressei, P. Merino, A. Ruffoni and D. Leonori, *J. Am. Chem. Soc.*, 2023, **145**, 27810–27820.

View Article Online  
DOI: 10.1039/D6SC01441K

Open Access Article. Published on 20 March 2026. Downloaded on 4/10/2026 4:16:06 AM.  
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



### Data availability

CCDC 2501744 contains the supplementary crystallographic data for this paper. The data that support the findings of this study are available within the article and the supplementary information (SI).

Supplementary information: materials and methods, experimental procedures, characterization data, and NMR spectra. See DOI:

