

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. Poryvai, A. Vasievska, K. Bangievska, J. Tarábek, S. Gerber-Lemaire and T. Slanina, *Chem. Sci.*, 2026, DOI: 10.1039/D6SC01848C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Light-Driven Radical Catch-and-Release with BODIPY Photocages

Anna Poryvai,^a Anna Vasiļevska,^{b,c} Karolína Bangievská,^b Ján Tarábek,^b Sandrine Gerber-Lemaire^a and Tomáš Slanina^bReceived 00th January 20xx,
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

Photocages release payloads upon light irradiation and are widely used for spatiotemporal control in chemical biology and materials science. Although payload release is almost universally described as a heterolytic process, homolytic pathways that generate radicals can interfere and produce unintended off-target effects. If controlled, radical photorelease would open new avenues for applications such as polymerization, however the molecular factors that govern this process in photocages remain unknown. Here, we investigate how photophysics and payload identity influence heterolytic vs homolytic reactivity in BODIPY photocages. We find that high fluorescence quantum yields correlate with efficient homolytic cleavage, enabling reversible radical *catch-and-release*: light-assisted capture of radical payloads followed by clean photorelease under green light. This radical release can be suppressed by the introduction of iodide or boron-methyl substituents which promote intersystem crossing. We achieve the highest photorelease quantum yield reported to date for green-light-driven radical generation ($\Phi_r = 0.5\%$), surpassing heterolytic carboxylate uncaging. We further exploit this reactivity in Type I photoinitiation of RAFT polymerization, yielding fluorescently labelled polymers with defined dispersity. This work establishes a structure–reactivity framework for predictable light-controlled radical generation, enabling mitigation of off-target radical effects and opening new avenues for late-stage photochemical payload installation.

Photocages are photoremovable protecting groups that temporarily mask active substrates through covalent bonds and release them upon irradiation.^[1,2] This capability enables precise spatial and temporal control over chemical and biological processes. In the majority of organic photocages, payload release is thought to proceed *via* heterolytic bond cleavage, yielding an anionic payload and a cationic cage (Scheme 1).^[1,2] In contrast, homolytic bond cleavage leading to radical payloads remains rare. Critically, the parameters that govern pathway selection—whether a given photocage produces ions or radicals upon irradiation—remain unknown. The absence of photocage design principles restricts predictive control over photocage reactivity, hinders rational design for radical-sensitive chemical and biological environments, and hinders the mitigation of undesired radical side effects.

Only a limited number of photocages have been shown to release radical payloads, namely nitric oxide,^[3–7] nitroxyl radicals,^[8–19] bromine- and sulfur-based groups.^[20–23] Homolytic release, however, interferes with heterolytic pathways, preventing clean and predictable radical generation.^[8,15,20–23] Recently, coumarin-based photocages were shown to favor homolytic release of bromine- and sulfur-centered payloads,^[20,21] while our prior work on BODIPY photocages revealed partial radical release of acetate and thiodansyl groups.^[22,23] Collectively, these studies underscore the need for

rational photocage design principles that enable predictable pathway selection. There is also a need for strategies that enable easy, efficient late-stage payload installation on the photocage. Current approaches often rely on multi-step organic synthesis to install payloads, resulting in low yields and restricting access to sensitive compounds. A versatile *catch-and-release* approach, where a payload can be photochemically installed at a late stage in the synthesis (“*catch*”) and selectively photocleaved (“*release*”) would address this gap directly.



BODIPY: X = H, R = F; CH₃-BODIPY: X = H, R = CH₃; I-BODIPY: X = I, R = F; R₁: TEMPO or RAFT agent
Scheme 1. *Catch-and-release* photoreactivity of BODIPY-Cl.

Here, we report that green-light-activatable BODIPY photocages can be directed to highly controllable reversible homolytic bond cleavage, enabling a radical *catch-and-release* process (Scheme 1). Using Cl and a persistent radical TEMPO payloads, we demonstrate (i) homolytic BODIPY–Cl bond cleavage to generate a BODIPY radical, (ii) *catch*: capture of this radical by TEMPO or RAFT agent, and (iii) *release*: efficient photorelease of the captured radical. Furthermore, we exploit this reactivity for living reversible addition–fragmentation chain transfer (RAFT) and free radical (FR) chain polymerizations of isobornylacrylate (IBOA). Together, these results establish that

^a École Polytechnique Fédérale de Lausanne, SB ISIC SCI-SB-SG, Station 6, CH-1015 Lausanne, Switzerland.

^b Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, Prague 6, 160 00, Czech Republic.

^c Department of Organic Chemistry, Faculty of Science, Charles University, Albertov 6, 128 00 Prague 2



BODIPY photocage reactivity can be directed toward efficient homolytic pathways and provide design principles for predictable light-controlled radical *catch-and-release*.

Heterolytic vs homolytic reactivity. To investigate the interplay between ionic and radical photoreactivity, we examined BODIPY-Cl in the presence or absence of nucleophiles (methanol or *t*-butanol), which trap the reactive species formed through the heterolytic (photo-S_N1) pathway, or TEMPO, which traps the intermediates of the homolytic (radical) pathway (Chart 1, Chapter S2-2). UV-Vis and LC-MS analysis of the reaction mixtures revealed that the irradiation of BODIPY-Cl in aerated anhydrous MeOH leads predominantly to the photo-S_N1 product, BODIPY-OMe (Tab. S2-3: A), accompanied by traces of BODIPY-OH and an oxidation product, BODIPY-CHO²² (~4% and 1%, respectively, Chart 1). Reaction in *t*-BuOH (as a bulky nucleophile) resulted in multiple products (Tab. S2-3: C, D, Fig. S2-9: C, D), while irradiation in PhH (a non-nucleophilic solvent) led solely to BODIPY-OH and BODIPY-CHO (Tab. S2-3: E, Fig. S2-10). Notably, the addition of 8 eq. of TEMPO shifted the product distribution markedly, increasing the ratios of BODIPY-OH and BODIPY-CHO, to 19 and 8% in MeOH, respectively (Chart 1, Tab. S2-3: B, H). By contrast, a large excess of MeOH (180 eq.) in PhH did not produce BODIPY-OMe; the photo-S_N1 product was only observed when MeOH was present in large excess as co-solvent with PhH (280 000 eq., 1/1 v/v, Tab. S2-3: G, H, I). Overall, these results demonstrate that the presence TEMPO efficiently diverts the photoreaction course from the heterolytic photo-S_N1 pathway toward the radical pathway. We therefore hypothesised that irradiation of BODIPY-Cl generates a BODIPY-centred radical intermediate that is subsequently intercepted by TEMPO in a *catch-and-release* mechanism, with BODIPY-OH and BODIPY-CHO arising as by-products of the radical pathway. To probe this hypothesis, we next investigated the catch process directly.




Chart 1. Reactivity of BODIPY-Cl in MeOH and corresponding traces from LC-MS analysis of solutions of BODIPY-Cl in MeOH with or without TEMPO (8 eq.) before and after illumination with 525 nm light (8 min). DOI: 10.1039/D6SC01848C

Catch. To confirm this hypothesis, we attempted to *catch* a TEMPO payload onto the BODIPY-methyl scaffold (Tab. 1). A concentrated solution of BODIPY-Cl with TEMPO was initially irradiated in a flask, providing BODIPY-TEMPO in a low yield, which was improved by performing the photoreaction in a flow reactor^[24] with the more uniform illumination of the reaction mixture (Tab. 1, entries 1, 2). Moving from benzene to toluene (PhCH₃, Tab. 1, entry 3) and optimizing the reagents ratio and flow rate led to a significant yield increase (up to 87%, Tab. 1, entry 4).

Table 1. Optimization of the *catch* process of TEMPO by BODIPY-Cl.

Entry	c(BODIPY) [mM]	TEMPO [eq.]	Solv.	Isol. yield [%]	Setup, Flowrate [mL/min]
1	20	2	PhH	9	Batch
2	4	9	PhH	52	Flow, 0.15
3	4	9	PhCH ₃	43	Flow, 0.05
4	4	11	PhCH ₃	87	Flow, 0.5

Boron-methylation and iodination of the BODIPY scaffold have previously been shown to significantly increase the efficiency of heterolytic photo-S_N1 release in MeOH ($\Phi_r \sim 2\%$ for BODIPY-Cl vs $\Phi_r \sim 35\%$ and 45% for CH₃-BODIPY-Cl and I-BODIPY-Cl, respectively).^[25] We therefore investigated whether these structural modifications also enhance radical reactivity. Therefore, we studied the abovementioned chlorides as substrates for TEMPO *catch*. Addition of TEMPO to a methanolic solution of CH₃-BODIPY-Cl resulted in higher conversion to the homolytic bond cleavage products (Fig. S2-14). Irradiation in PhH, however, led to a rapid decomposition of CH₃-BODIPY-Cl with no chromophoric products (Fig. S2-14). I-BODIPY-Cl remained unreactive in PhH. High photo-S_N1 reactivity was observed in MeOH with and without TEMPO additive (Fig. S2-16). Addition of a larger excess of TEMPO (34 eq.) resulted in the negligible formation of I-BODIPY-OH alongside very efficient conversion to the photo-S_N1 product I-BODIPY-OMe (Fig. S2-16).

The intermediacy of the BODIPY-methyl radical was probed by electron paramagnetic resonance (EPR) spectroscopy, monitoring the appearance of the aminoxyl radical product of its addition to a *N*-tert-butyl- α -phenylnitron spin-trap (Chart 2, Fig. S2-11). As a complementary experiment we studied the *catch* process upon irradiation of excess BODIPY-Cl and TEMPO in PhH, leading to the BODIPY-TEMPO conjugate, by monitoring the decrease of TEMPO concentration (Fig. S2-12). The EPR measurements of I-BODIPY-Cl with TEMPO showed no decrease in TEMPO concentration, further supporting the lack of photoreactivity observed by UV-Vis and LC-MS techniques (Fig.



S2-17). These results corroborate our hypothesis of BODIPY-Cl undergoing homolytic bond cleavage upon irradiation, generating a BODIPY-methyl radical that is efficiently intercepted by TEMPO, thereby diverting the reaction pathway from photo-S_N1 reactivity to productive radical trapping (Scheme 1).




Chart 2. Evolution of EPR spectra recorded in time of solutions of BODIPY-Cl in PhH in the presence of *N-tert-butyl-α-phenylnitron* spin trap, measured in dark (left), upon illumination with 525 nm light (middle), and after irradiation (right).

Release. Photoinduced *release* of TEMPO was investigated on three BODIPY scaffolds BODIPY-TEMPO, CH₃-BODIPY-TEMPO, and I-BODIPY-TEMPO conjugates (Chart 3, Ch. S1-2, S2-3). The spectral properties of the X-BODIPY-TEMPO conjugates closely matched those of the corresponding X-BODIPY-Cl precursors (Tab. 2, S2-1, S2-2), confirming that the TEMPO substituent does not significantly perturb the BODIPY chromophore. Introduction of substituents resulted in decrease in fluorescence quantum yields (Φ_f) and fluorescence lifetimes (τ) and in increase of triplet state quantum yields (Φ_Δ) (Fig. S2-5, S2-6, S2-7). The LC-MS analysis and ¹H NMR measurements gave evidence for the formation of three products as a result of photorelease from BODIPY-TEMPO: BODIPY-OH, BODIPY-CHO, and TEMPO radical (Chart 3B, S2-18, S2-19, S2-20). Neither *N-O* bond cleavage products^[8] nor S_N1 substitution products were observed, demonstrating that TEMPO *release* proceeds exclusively *via* homolytic bond cleavage. This establishes BODIPY photocages as a scaffold for controlled light-driven radical generation.




Chart 3. (A) Reactivity of BODIPY-TEMPO upon illumination with 525 nm light in MeOH and corresponding: (B) traces from LC-MS analysis; (C) evolution of EPR spectra (before (blue), in time (grey), and after (red) illumination). (D) Concentration (blue axis) and yield (red axis) of TEMPO released over time from CH₃-BODIPY-TEMPO (blue stars, PhH), I-BODIPY-TEMPO (green squares, PhH), BODIPY-TEMPO (black circles, PhH; red triangles, MeOH).

The photorelease of TEMPO was further supported by the EPR kinetic measurements (Chart 3C, D). Under green light illumination, BODIPY-TEMPO showed high reactivity with 95% of TEMPO released in PhH (Tab. 2), exceeding the values reported for quinone methide^[18] and 2-(4-nitrophenyl)benzofuran^[17] photocages (~80%), and similar to the performance of UV-absorbing pyrene-based photocage (95%).^[19] Our system achieved an excellent uncaging cross-section value ($\delta_r \sim 390 \text{ M}^{-1}\text{cm}^{-1}$) and the highest photorelease quantum yield ($\Phi_r = 0.5\%$) reported so far for green-light-induced radical release (compared to the 0.047% for a QM photocage^[18]). Contrary to BODIPY-Cl, which underwent partial photo-S_N1 reaction in MeOH, BODIPY-TEMPO afforded only the products of homolysis with 85% yield in MeOH, demonstrating that the nucleophilic solvent does not affect the photoinduced homolysis. Boron methylation and iodination of the BODIPY-core, however, resulted in a up to ~7-fold decrease of the photorelease efficiency and a loss of ~30% in the yield of released TEMPO.




Table 2. Efficiency of the release of TEMPO from X-BODIPY-TEMPO.

Code	Solv.	Φ_{fl} [%]	Φ_{Δ} [%]	τ , ns	Φ_r [%]	Yield [%]
BODIPY-TEMPO	PhH	38	3	5	0.5	95
	MeOH	n. d.	n. d.	n. d.	0.6	85
CH ₃ -BODIPY-TEMPO	PhH	15	9	4	0.07	65
I-BODIPY-TEMPO	PhH	2	80	b.l.d.	0.09	62

These results are consistent with the reduced radical reactivity observed for I-BODIPY-Cl and CH₃-BODIPY-Cl in the *catch* experiments. The reduced reactivity may be caused by nitroxyl radicals released during the reaction progress, which might promote the intersystem crossing (ISC) from the singlet and triplet excited states,^[26–31] as demonstrated on a series of BODIPY derivatives.^[32–34] Therefore, TEMPO released during the reaction may influence the photodynamics and efficiency of the system by promoting ISC and other deactivation pathways. Altogether, the photoreactivity of X-BODIPY-Cl/TEMPO demonstrated that the homolytic BODIPY-payload bond cleavage is more efficient for compounds with the high Φ_{fl} , and therefore could originate from the excited singlet state.

Polymerization. Taking advantage of the facile radical reactivity of BODIPY-Cl, we then explored the possibility of using BODIPY-Cl as a Type-I radical photoinitiator for the synthesis of fluorescent isobornyl acrylate (IBOA) polymers through a Giese-type radical addition to the activated C-C double bond of IBOA (Scheme 2).



Scheme 2. FR (i) and RAFT (ii, DDMAT RAFT agent) polymerization of IBOA upon irradiation with 525 nm light using BODIPY-Cl photoinitiator (Tab S3-1, e-1-5) and overview of the polymers' properties (λ_{abs} – absorbance and fluorescence maxima, Φ_{fl} , M_n – molecular weight, and \bar{M}_w/\bar{M}_n – dispersity).

Free radical (FR) polymerization of BODIPY-Cl (0.08 mol%) led to low yields of polyacrylates (poly IBOA, 11-19%), with high molecular weight ($M_n = 295\text{--}372$ kDa) and high dispersity ($\bar{M}_w/\bar{M}_n \sim 2.3$, Tab. S3-1). To improve the dispersity of prepared fluorescent polymers, BODIPY-Cl was further exploited as photoinitiator in a RAFT polymerization, leading to poly-IBOAs of lower molecular weights ($M_n = 11.6\text{--}34.1$ kDa) and in higher yields (up to 69%) with excellent \bar{M}_w/\bar{M}_n (~ 1.5) at the lower initiator concentrations (Fig. S3-9A: e-4c, Tab. S3-1). Solutions of poly IBOAs prepared under the same conditions using both FR and RAFT approaches (e-1a,b, Tab. S3-1) have identical absorption and emission spectra ($\lambda_{abs} = 502$ nm, $\lambda_{em} = 511$ nm) with $\Phi_{fl} \sim 42\%$, comparable to the values measured for the standard compound BODIPY-OAc ($\sim 48\%$, Fig. S3-7) indicating that attaching the BODIPY fluorophore onto a polymer chain does not significantly decrease its brightness. Kinetic studies showed

that the conversion rate and properties of the resulting poly-IBOAs (Φ_{fl} , M_n , \bar{M}_w/\bar{M}_n) can be efficiently controlled by the irradiation time: longer irradiation time led to higher conversions and a decrease in Φ_{fl} associated with the dye photobleaching (Fig. S3-6, Fig. S3-7, Fig. S3-9B, e-5, Tab. S3-1). While studying the polymerization mechanism, we detected a BODIPY-RAFT intermediate analogous to the BODIPY-TEMPO conjugate (Fig. S3-10), indicating that the *catch-and-release* mechanism extends to RAFT agents and that light-induced radical generation is not limited to the initiation phase.

Conclusions

In conclusion, we demonstrated that BODIPY photocages can efficiently release radical payloads *via* homolytic cleavage of the photocage–payload bond, challenging the prevailing evidence that BODIPY uncaging proceeds exclusively through heterolytic pathways. This insight enabled us to design a reversible radical *catch-and-release* system based on homolytic bond formation and cleavage, allowing straightforward caging of radical payloads and their efficient photorelease. Using TEMPO as a model payload, radical release was found to be four times more efficient than heterolytic acetate uncaging from the analogous BODIPY-OAc compound. Structure–reactivity analysis of X-BODIPY-TEMPO revealed that radical photorelease is favoured for cages with high fluorescence quantum yields ($X = H$) and diminishes upon boron methylation ($X = CH_3$) or iodination ($X = I$). As a representative application, we exploited the radical reactivity of BODIPY-Cl in free radical and RAFT polymerizations of isobornyl acrylate, yielding polymers covalently linked to the BODIPY core. Detection of a BODIPY-RAFT intermediate analogous to the BODIPY-TEMPO conjugate confirmed that the *catch-and-release* mechanism participates in the RAFT process beyond the initiation phase. This work establishes that BODIPY photocage reactivity is not intrinsically heterolytic but is governed by payload character, reaction conditions, and photocage photophysics. Optimizing these factors enables predictable light-controlled radical generation under mild conditions, and opens opportunities for late-stage photochemical payload installation.

Author contributions

Conceptualization: A. P., S. G.-L., T. S. **Data curation:** A. P., A. V., K. B., J. T. **Formal analysis:** A. P., A. V., J. T. **Funding acquisition:** A. P., S. G.-L., T. S. **Investigation:** A. P., A. V., K. B., J. T. **Methodology:** A. P., J. T. **Project administration:** A. P. **Resources:** A. P. **Supervision:** A. P., S. G.-L., T. S. **Validation:** A. P. **Visualization:** A. P. **Writing - original draft:** A. P. **Writing - reviewing and editing:** A. P., A. V., K. B., J. T., S. G.-L., T. S.

Conflicts of interest

There are no conflicts to declare.



Data availability

The Supporting Information is available as a PDF file, including general procedures, synthetic procedures, structure characterization, photophysical and photochemical properties of the BODIPY derivatives, polymerization procedures, and characterization of polymers. The raw data that support the findings of this study are openly available in ZENODO at 10.5281/zenodo.17405857. A preprint version of this work has been posted on ChemRxiv (10.26434/chemrxiv-2025-96t6r/v2).

Acknowledgements

The authors acknowledge the Swiss National Science Foundation (grant N° TMPFP2_217233) and the Czech Science Foundation (grant N° 22-20319S) for financial support. The authors acknowledge Dr. Alireza Kavand for the introduction to the polymerization techniques, Dr. Tomáš Mašek for discussions on radical chemistry, and Dr. Milena Schuhmacher for mentoring and Hannah E. Pfeifer for her help with editing.

Notes and references

- P. Klan, T. Solomek, C. G. Bochet, A. Blanc, R. Givens, M. Rubina, V. Popik, A. Kostikov, J. Wirz, *Chem. Rev.* 2013, **113**, 119–191.
- R. Weinstain, T. Slanina, D. Kand, P. Klán, *Chem. Rev.* 2020, **120**, 13135–13272.
- C. Parisi, F. Laneri, A. Fraix, S. Sortino, *J. Med. Chem.* 2024, **67**, 16932–16950.
- M. Blangetti, A. Fraix, L. Lazzarato, E. Marini, B. Rolando, F. Sodano, R. Fruttero, A. Gasco, S. Sortino, *Chem. Eur. J.* 2017, **23**, 9026–9029.
- E. Vittorino, E. Ciciarella, S. Sortino, *Chem. Eur. J.* 2009, **15**, 6802–6806.
- E. Y. Zhou, H. J. Knox, C. J.; Reinhardt, G. Partipilo, M. J. Nilges, J. Chan, *J. Am. Chem. Soc.* 2018, **140**, 11686–11697.
- K. Kitamura, N. Ieda, K. Hishikawa, T. Suzuki, N. Miyata, K. Fukuhara, H. Nakagawa, *Bioorg. Med. Chem. Lett.* 2014, **24**, 5660–5662.
- A. Kugele, B. Silkenath, J. Langer, V. Wittmann, M. Drescher, *ChemBioChem* 2019, **20**, 2479–2484.
- D.-L. Versace, J. Lalevéé, J.-P. Fouassier, D. Gigmes, Y.; Guillaneuf, D.; Bertin, Photosensitized alkoxyamines as bicomponent radical photoinitiators. *J. Polym. Sci. A Polym. Chem.* 2010, **48**, 2910–2915.
- J. C. Scaiano, T. J. Connolly, N. Mohtat, C. N. Pliva, *Can. J. Chem.* 1997, **92**–97.
- D.-L. Versace, Y. Guillaneuf, D. Bertin, J. P. Fouassier, J. Lalevéé, D. Gigmes, *Org. Biomol. Chem.* 2011, **9**, 2892–2898.
- S. E. Bottle, J.-L. Clement, M. Fleige, E. M. Simpson, Y. Guillaneuf, K. E. Fairfull-Smith, D. Gigmes, J. P. Blinco, *RSC Adv.* 2016, **6**, 80328–80333.
- A. Goto, J. C. Scaiano, L. Maretti, *Photochem. Photobiol. Sci.* 2007, **833**–835.
- S. Hu, J. H. Malpert, X. Yang, D. C. Neckers, *Polymer* 2000, **41**, 445–452.
- M. Herder, J.-M. Lehn, *J. Am. Chem. Soc.* 2018, **140**, 7647–7657.
- Y. Guillaneuf, D.-L. Versace, D. Bertin, J. Lalevéé, D. Gigmes, J.-P. Fouassier, *Macromol. Rapid Commun.* 2010, **31**, 1909–1913.
- A. Yamada, M. Abe, Y. Nishimura, S. Ishizaka, M. Namba, T. Nakashima, K. Shimoji, N. Hattori, *Beilstein J. Org. Chem.* 2019, **15**, 863–873.
- P. Kieley, P. Farràs, P. McArdle, D. A. Smith, F. Aldabbagh, *Chem. Commun.* 2019, **55**, 14665–14668.
- M. Baron, J. C. Morris, S. Telitel, J.-L. Clément, J. Lalevéé, F. Morlet-Savary, A. Spangenberg, J.-P. Malval, O. Soppera, D. Gigmes, Y. Guillaneuf, *J. Am. Chem. Soc.* 2018, **140**, 3339–3344.
- H. D. Nguyen, M. Abe, *J. Am. Chem. Soc.* 2024, **146**, 10993–11001.
- M. Takano, M. Abe, *Org. Lett.* 2022, **24**, 2804–2808.
- A. Poryvai, M. Galkin, V. Shvadchak, T. Slanina, *Angew. Chem. Int. Ed.* 2022, **61**, e202205855.
- L. Wohlrábová, J. Okoročenkova, E. Palao, E. Kužmová, K. Chalupský, P. Klán, T. Slanina, *Org. Lett.* 2023, **25**, 6705–6709.
- L. Buglioni, F. Raymenants, A. Slattery, S. D. A. Zondag, T. Noël, *Chem. Rev.* 2022, **122**, 2752–2906.
- T. Slanina, P. Shrestha, E. Palao, D. Kand, J. A. Peterson, A. S. Dutton, N. Rubinstein, R. Weinstain, A. H. Winter, P. Klán, P. J. *Am. Chem. Soc.* 2017, **139**, 15168–15175.
- A. R. Watkins, *Chem. Phys. Lett.* 1974, **29**, 526–528.
- A. R. Watkins, *Chem. Phys. Lett.* 1980, **70**, 262–265.
- R. E. Schwerzel, R. A. Caldwell, *J. Am. Chem. Soc.* 1973, **95**, 1382–1389.
- E. Chernikova, A. Morozov, E. Leonova, E. Garina, V. Golubev, C. Bui, B. Charleux, *Macromolecules* 2004, **37**, 6329–6339.
- A. Ito, A. Shimizu, N. Kishida, Y. Kawanaka, D. Kosumi, H. Hashimoto, Y. Teki, *Angew. Chem. Int. Ed.* 2014, **53**, 6715–6719.
- V. Rane, R. Das, *J. Phys. Chem. A* 2015, **119**, 5515–5523.
- Z. Wang, J. Zhao, A. Barbon, A. Toffoletti, Y. Liu, Y. An, L. Xu, A. Karatay, H.-G. Yaglioglu, E. A. Yildiz, M. Hayvali, *J. Am. Chem. Soc.* 2017, **139**, 7831–7842.
- P. K. Pachpatil, S. V. Kanojia, A. Ghosh, A. G. Majumdar, A. Wadawale, M. Mohapatra, B. S. Patro, T. K. Ghanty, D. Goswami, *Sens. Actuators B Chem.* 2022, **370**, 132474.
- M. Mayländer, T. Quintes, M. Franz, X. Allona, A. V. Jentsch, S. Richert, *Chem. Sci.* 2023, **14**, 5361–5368.



Lausanne, March 4th 2026The Editorial Board
*Chemical Science***Data availability statement for a manuscript entitled “Light-Driven Radical Catch-and-Release with BODIPY Photocages”**

Dear Editor,

The Supporting Information for this manuscript is available as a PDF file.
The raw data supporting the findings of this study are openly available on ZENODO at <https://zenodo.org/records/17405857>.

The data include: general procedures, synthetic procedures, structural characterization, photophysical and photochemical properties of the BODIPY derivatives, polymerization procedures, and characterization of polymers.

A preprint version of this work has been posted on ChemRxiv (<https://chemrxiv.org/doi/full/10.26434/chemrxiv-2025-96t6r/v2>).

Thank you for your time and consideration,

Sincerely,

Dr. Anna Poryvai

