



Cite this: DOI: 10.1039/d6sc02383e

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 23rd March 2026
Accepted 13th May 2026

DOI: 10.1039/d6sc02383e

rsc.li/chemical-science

Fluorinated carbohydrate-based vaccines†

Elena Chikunova, James Suri, Kathrin Siebold and Ryan Gilmour*

The structural virtuosity of carbohydrates forms the basis of a molecular recognition language that is ubiquitous in immunomodulatory processes. This diversity generates the complexity required to accommodate the bandwidth of information generated by immune interactions: it logically follows that this capacity underpins the evolutionary success of oligosaccharide permutations in this endeavour. Placing glycan function on a structural level is therefore a core research endeavour in understanding the non-covalent interactions that elicit an immune response and, by extension, leveraging these data to identify and validate new vaccine leads. Innovations to expedite the construction of well-defined, immuno-relevant carbohydrates has revolutionized the field, and focused attention on further structural refinements that are new to biology. Echoing the success of site-selective fluorination in pharmaceutical design, the precision introduction of C(sp³)-F bonds to tailor the physicochemical properties of key targets is an exciting new frontier in targeted immunology. In this short Perspective, the key milestones in the design and validation of carbohydrate-based vaccine leads are discussed through the lense of fluorination.

1. Introduction

Vaccination is a fundamental pillar of public health and the legacy of Salk's totemic introduction of a safe and effective polio vaccine in the mid-1950s.¹⁻³ Its impact is reflected in the World Health Organization's (WHO) estimate that vaccination programmes currently prevent some 3.5-5 million deaths per year worldwide.⁴ This approach to disease prevention and management has played a significant role in reducing the global under-five mortality rate,⁵⁻⁷ thereby lessening the risk posed by numerous fatal infections. Success has been grounded in centuries of observation, experimentation, failure, and persistence, resulting in the aforementioned (r)evolution.⁸⁻¹¹ In response to these efforts, pathogens also continually evolve, developing resistance to vaccines and antibiotics.¹²⁻¹⁴ Indeed, while some vaccines have proved to be remarkably robust over time,¹⁵⁻¹⁹ others require continuous innovation.^{12,20-27} It logically follows that the scientific community must continually innovate to ensure that pathogens do not gain the upper hand. Vaccines efficiently exploit the immune system's natural capacity to recognize and establish long-term defenses against the unique markers of disease-causing agents.²⁸ To successfully achieve protective immunity, a vaccine must include antigens that are either derived directly from the pathogen or introduced in the form of synthetic mimics.^{29,30} Generally, antigens include any substances able to induce a specific immune response or to be recognized by products of an immune response (such as

antibodies).³¹ They include proteins, carbohydrates, nucleic acids, lipids, and haptens.³² The expansive natural diversity of antigens provides researchers who are actively searching for new leads with a vast pool of perspective immunostimulating agents. Carbohydrates are particularly attractive in this regard owing to the multitude of constituent monosaccharide building blocks, their diversity and complexity, and finally their ubiquity in nature.^{33,34} These biopolymers decorate the surface of almost³⁵⁻³⁷ every natural cell (prokaryotic and eukaryotic),³⁸ where they are key mediators in essential recognition processes.^{39,40} The following cases are instructive to illuminate the societal benefits that may be gleaned from pursuing carbohydrate-based approaches: In bacterial cells, unique capsular polysaccharides (CPS) are common primary virulence factors, but they do not have easily targetable proteins on their surfaces (Fig. 1).⁴⁰

Designing bespoke vaccines to emulate bacterial CPS⁴¹ can therefore lead to highly serotype-specific antibodies that precisely neutralize the desired pathogen.⁴²⁻⁴⁴ Moreover, numerous types of tumor cells display aberrant glycosylation patterns (tumor-associated carbohydrate antigens or TACAs) on their surfaces that aid detection.⁴⁵⁻⁴⁷ These specific patterns have inspired the development of carbohydrate-based therapeutic cancer vaccines that are designed to activate the immune system towards these unique signatures.⁴⁸⁻⁵⁰

Importantly, carbohydrate cellular fingerprints extend beyond these scenarios to encompass other chronic conditions such as neurodegenerative diseases⁵¹ and autoimmune disorders,^{52,53} among others.^{54,55} That said, the design and conception of functional carbohydrate-based vaccines remains a multifaceted challenge and general solutions remain elusive.^{56,57} As well as masking pathogenic proteins to enable

Institute for Organic Chemistry, University of Münster, Corrensstraße 36, 48149 Münster, Germany. E-mail: ryan.gilmour@uni-muenster.de

† This manuscript is dedicated to Prof. David O'Hagan (University of St Andrews) for his inspiration and example in organofluorine chemistry.






Fig. 1 Top: General principles of carbohydrate-based vaccine design. Identification and isolation or synthesis of the target glycans from the target pathogen. Glycan conjugation to an immunogenic carrier protein. Creation of a vaccine formulation that may also include an adjuvant. Following administration, the recipient produces antibodies specific for the target pathogen; bottom: a key for symbol nomenclature used throughout this perspective.

immune evasion/infection, glycans are widely presented on human cells and engage in a multitude of functions.^{58–60} Consequently, eliciting sufficient immunogenicity poses a significant obstacle for glycan-based vaccines.^{61–64} The target glycan epitope is required to be distinguishable from sugars on healthy human cells to minimise the risk of autoimmune reactions.^{56,65} Simultaneously, the antigen must be sufficiently similar to the natural epitope to trigger the production of antibodies capable of recognizing the original structure,^{60,66–68} *i.e.* cross-reactive antibodies.⁶⁹

This requirement for structural similarity renders site selective fluorination an appealing molecular editing strategy.^{70,71} A combination of modest steric footprint and high electronegativity offers an opportunity to mimic specific hydroxyl groups on the glycan ($\text{OH} \rightarrow \text{F}$),^{71,72} potentially enhancing the immune response without introducing unacceptable structural alterations. Vaccines based on fluorinated glycan antigens, which are not encountered in biology, offer a tempting opportunity to boost immunogenicity to trigger a stronger specific immune response with the formation of cross-reactive long-lasting antibodies. In this short Perspective, key considerations in vaccine development are presented for a non-specialist audience, and set in a wider discussion of the major milestones in the advent and development of fluorinated glycan vaccines leads. It is hoped that this will foster interest and discussion in this emerging area of targeted immunotherapy.

2. Vaccines and the immune system

From the perspective of function, vaccines are designed to train the adaptive immune system to recognize and eliminate a specific pathogen or cell group swiftly and effectively. The adaptive immune response is mediated by B cells that produce antibodies (humoral immunity) and by T cells (cellular immunity).⁷³ Almost all vaccines in common use are thought to mainly confer protection through the formation of antibodies.⁷⁴

Following administration of a vaccine formulation, antigen-presenting cells (APCs), primarily dendritic cells (DCs), recognise “danger signals” such as those represented by adjuvants

(components of vaccines that trigger a non-specific immune response; please see Section 3.1) (Fig. 2, top).^{75–79} This process triggers a sequence of events beginning with the maturation of DCs. Within the lymph nodes, mature DCs present processed antigen fragments on both major histocompatibility complex (MHC) class I and class II to naïve T cells. Antigen fragments on MHC I can activate CD8^+ T cells that leads to the generation of cytotoxic mediators; these in turn can directly kill infected cells (cellular immunity). Antigen fragments on MHC II activate CD4^+ T helper (Th) cells with subsequent proliferation and secretion of cytokines that drive the immune response toward either a humoral (antibody-driven) or cellular (cytotoxic-driven) profile.

For protective antibody production, B cell activation is crucial.⁷³ Vaccine antigens can trigger a T cell-independent response by binding directly to B cell receptors (BCRs) present on the B cell surface. This leads to the production of less specific IgM-antibodies. To enable a more robust immune response, B cells require T cell help, *i.e.*, a T cell-dependent immune response. This is the basis for glycoconjugate vaccines, where a polysaccharide antigen is chemically linked to a protein carrier (please see Section 3.1),^{80,81} and allows B cells that recognize the polysaccharide to internalize the entire conjugate and process the protein carrier.⁸² Subsequently, they migrate to the border between the B cell and T cell zones in lymphoid tissue and present it on MHC II to previously activated, antigen-specific Th cells. There, if an activated Th cell recognizes the MHC II complex on the B cell, a cognate interaction occurs. This entire coordinated cascade results in B cell proliferation, generation of memory B cells, differentiation into antibody-secreting plasma cells, and antibody class switching: from less specific IgM to high-affinity, specific IgG; that dramatically changes the outcome of vaccination.

IgM is a fast primary responder,^{83,84} which is produced within the first days after the initial exposure. Its large, pentameric structure is ideally suited to bind multiple pathogens at once, agglutinating them for easier clearance by phagocytes.⁸⁵ However it cannot provide a specific reaction. The biological half-life of IgM in the bloodstream is only about 5 to 7 days,^{83,86} which means its concentration drops rapidly after vaccination. For lasting immunological protection that effective vaccines must provide, IgG plays an important role.^{87,88} These antibodies take longer to be produced, but react with high specificity to the selected antigens during subsequent exposures (prophylactic effect) and provide the long-term immunological memory that is sought.^{88,89} It is important to stress, however, that is not sufficient to simply generate IgG antibodies; they must be cross-reactive. Cross-reactivity occurs when an antibody that was generated in response to a specific antigen (*e.g.* from a vaccine) also recognises and binds to a different, but structurally similar, antigen.^{90,91} This phenomenon has been successfully capitalized on through vaccine design, which enables vaccines to be effective against various bacterial or cell subtypes (serogroups/serotypes) and makes structural alterations of antigens possible.⁹² For example, a native pathogen antigen Ag undergoes modifications to give a new vaccine component AgV (Fig. 2, bottom). Antibodies raised after this vaccine injection will be specific for the modified AgV, but they are also required






Fig. 2 Simplified representation of the adaptive immune response to vaccine administration and the concept of cross-reactivity. Top: B cell can be activated without Th cell assistance (T cell-independently) via recognition of vaccine antigen to produce IgM or with the help of previously matured Th cell (T cell-dependently) leading to proliferation and IgG and long-term immunological memory. Bottom: modified antigen AgV induces formation of AgV-specific antibodies, which might recognize the original Ag, thus, be cross-reactive, or be exclusively specific for AgV – not cross-reactive – therefore, providing no protection against the original threatening epitope. DC = dendritic cell; MHC = major histocompatibility complex.

to recognize initial Ag from the target pathogen. Cross-reactivity makes this possible: with proper AgV design, the antibodies raised will recognize the native Ag as well and attack the pathogen, thereby affording protection to the individual.

3. Carbohydrate-based vaccines: the immune response and key components

Carbohydrate-based vaccines have a venerable history that has its roots in the seminal discovery by Avery and Heidelberger in 1923, that the CPS of *Streptococcus pneumoniae* was immunogenic (Fig. 3).^{93,94} This key milestone led to the development of

the first carbohydrate-based vaccine against pneumococcal disease.^{95,96} With the advent of antibiotics, the development of carbohydrate-based vaccines slowed, but the emergence of antibiotic resistance led to renewed interest in this field of research. To date, several carbohydrate-based vaccines have been licensed and are credited with having saved millions of lives. Among them are multivalent vaccines against *Streptococcus pneumoniae*, *Neisseria meningitidis*, and *Haemophilus influenzae* type b.⁵⁶

3.1. Immune response

Soon after the establishment of native CPS based vaccines, it was observed that for certain groups in the population, the






Fig. 3 Timeline showing selected milestones in the field of carbohydrate-based vaccines.^{47,56,93,96,104–106}

elicited immune response was not effective.⁹⁷ This was especially problematic in patients under two years old. Only short-term IgM antibodies were elicited and no long-term protection was achieved. Carbohydrates (pathogen- or tumor-associated) tend to be poorly immunogenic due to the presence of similar or even identical antigens on the human cell surface.^{61,64} Although these glycans usually have lower density or are expressed during early development stages in healthy individuals, the difference is not significant enough to trigger a robust immune response.

3.1.1. Carrier proteins. A major advance in addressing the poor immunogenicity inherent to glycans has been to couple the glycan to a carrier protein to induce more robust immune responses.^{98–104} Carrier proteins increase the immunogenicity of the vaccine as they can initiate a T-cell-dependent immune response when they are processed by APCs and subsequently

presented on MHC II to Th cells.^{107,108} As long as the original APC (B cell) has bound to the vaccine *via* the glycan epitope having been conjugated to the protein, T-cell assistance is associated with this specific sugar. Carrier proteins can be derived from pathogens like diphtheria toxin mutant (CRM₁₉₇),¹⁰⁹ tetanus toxoid (TT),¹⁰⁹ diphtheria toxoid (DT),¹¹⁰ meningococcal outer membrane protein complex (OMPC),¹⁰⁹ *Haemophilus influenzae* protein,¹⁰⁹ and *Pseudomonas aeruginosa* exotoxin A (rEPA).¹¹¹ These proteins can also originate from non-pathogenic (and therefore often more easily accessible) sources like bovine serum albumin (BSA)¹¹² and keyhole limpet hemocyanin (KLH).¹¹³ Glycoconjugate vaccines can therefore be grouped into three different types as summarized in Fig. 4: (1) natural, (2) semi-synthetic, and (3) fully synthetic glycoconjugates.⁹⁸

In the case of natural glycoconjugates, both components are derived from natural sources such as microbial cultures.¹¹⁴ Despite their efficacy against the corresponding pathogens, major issues are associated with this manufacturing method and include complicated and expensive purification procedures still leading to batch-to-batch variability, the presence of cell-derived impurities, and uncontrolled and irreproducible protein conjugation chemistry.⁵⁶ In comparison, synthetically derived carbohydrates are pure, well-defined and can be equipped with a synthetic linker to facilitate controlled coupling to the carrier protein. This type of glycoconjugate is termed semi-synthetic.¹¹⁵ In 2004, a Cuban research team published the first industrial scale synthesis of a semi-synthetic carbohydrate-based vaccine (Quimi-Hib®) that was later included in the Cuban and South American vaccination schedule to address *Haemophilus influenzae* type b (Hib) (Fig. 5).¹⁰⁶

It is important to note that the protein in this approach is essential to ensure a robust immune response, but it suffers from some of the challenges associated with the natural approach mentioned above, such as inconsistent conjugation results.

3.1.2. Adjuvants. In contrast to the approaches mentioned above, fully-synthetic glycoconjugates are free of many of the drawbacks associated with using naturally-sourced components.¹¹⁵ These therapeutics are composed of the antigen glycostructure coupled to a synthetic carrier (such as proteins or nanoparticles¹¹⁶) and combined with adjuvants to ensure




Fig. 4 Schematic representations of the three approaches to glycan-based vaccine synthesis: the natural approach (all components are naturally derived), the semi-synthetic approach (certain components are naturally derived, while others are synthetic), and the fully synthetic approach (all components are synthetic).






Fig. 5 Structure of the semi-synthetic vaccine included in the Cuban and South American vaccination schedule.¹⁰⁶

a robust immune response. Adjuvants are agents that non-specifically augment the immune response to a co-administered antigen, without being the primary target of the adaptive immune response. The major adjuvant groups include: alum-based, bacteria-derived, liposome adjuvants, and adjuvant emulsions.^{117,118} In approved glycoconjugate vaccines, as well as in many vaccine candidates, adjuvants are used in the combination with proteins to achieve the desired immune response.

3.1.3. Other approaches. The field of fully synthetic vaccines is relatively new and there are currently no candidates in clinical trials that the authors are aware of. A conspicuous challenge is imitating the immune response generated by the carrier protein.⁶⁸ Recent findings also indicate that zwitterionic polysaccharides can induce robust antibody production without conjugation to a carrier. Alteration of the 3D structure of the vaccine (*e.g.*, the organisation of glycoclusters on dendrimeric structures) has led to positive outcomes as well.¹¹⁹ The modularity of this approach enables combinations of different carbohydrate antigens to be leveraged in the pursuit of multi-valent vaccines against different serogroups of the same bacterium. A prominent example is a 20-valent pneumococcal vaccine that provides protection against 20 serotypes of *Streptococcus pneumoniae*.¹²⁰ However, it is important to emphasise that a limiting factor in the development of synthetic glycoconjugate vaccines is the chemical synthesis of carbohydrates; this remains challenging and time-consuming. One method to simplify glycan synthesis is automated glycan assembly (AGA),¹²¹ which was formalized by the Seeberger group in 2001¹²² and has undergone rapid growth in recent years.^{123–128} Using pre-functionalized building blocks, AGA allows for a “plug and play” approach to generate the desired oligosaccharides in a rapid and efficient way. Contemporary AGA workflows can be conducted on commercial synthesizers (*e.g.*, the Glyconeer 2.1® instrument)¹²⁹ to produce glycans that are directly amenable to subsequent conjugation.

3.1.4. Structure modification. Strategically introducing non-native antigens is an expansive approach to overcome immune tolerance and induce robust immune responses. However, these antibodies must closely simulate recognition of the original target antigens. In 1986, Jennings and co-workers first demonstrated that the chemical modification of a carbohydrate antigen enhanced antigen immunogenicity, but these early examples were limited by a low cross-reactivity of the elicited antibodies with their native structures.¹³⁰ This discovery

remains a foundational milestone for the field, and its impact is reflected in the chemically modified carbohydrates that have already reached clinical trials. For example, a vaccine targeting small cell lung cancer using *N*-propionyl-modified α -(2 \rightarrow 8)-linked polysialic acid¹³¹ and the melanoma vaccines with GD2- and GD3-lactone derivatives are powerful exemplars.¹³² Given its success in medicinal chemistry, it logically follows that strategic fluorination of antigenic glycans presents a promising direction for native glycan modification. Fluorination of glycostructures is a potentially powerful editing approach that allows a sugar to be subtly modified in a minimally invasive manner whilst offering the possibility to influence physicochemical parameters and immunogenicity. This is largely grounded in the unique properties of the fluorine atom (small size similar to hydrogen and high electronegativity) making it a suitable bioisostere of the H-atom and OH-group.^{71,72} Selective bioisosteric substitution with the F-atom has been successfully integrated into glycoengineering strategies for studying antibody behavior.^{133,134} Fluorinated glycostructures have been shown to be more hydrolytically stable, which have positive consequences for bioavailability: enzymatic degradation is inhibited significantly by destabilisation of the transition state. The effect can be convincingly observed when the C(sp³)-F is incorporated into different positions of the sugar, with the largest stability benefits conferred when fluorine is placed next to the anomeric centre (Fig. 6).^{134–138} The combination of these factors renders fluorinated glycans potentially valuable from the perspective of designing next generation of carbohydrate-based vaccines.

An additional aspect of C2 fluorination that is particularly appealing in synthetic planning is the ability of this small substituent to regulate the selectivity of glycosylation reactions. Not only does this make chemical glycosylation predictable and efficient,^{140–144} the ¹⁹F nucleus facilitates analysis by NMR. The current field of fluorosugar research is multifaceted; it includes a diversity of synthetic methods for fluorine incorporation,^{145–147}




Fig. 6 Simplified representation of a fluorinated glycan in a hydrolase binding pocket. Fluorine destabilizes the transition state and plays an active role in enhancing hydrolytic stability.¹³⁹



the use of fluorine as a directing group in glycosylation,^{140–144} and the practical application of fluorosugars in bio-medicine and chemical biology.^{148–155} In developing lines of research, fluorination can simplify the synthesis of glycan building blocks, rendering them more suitable for automated glycan assembly. In 2021, the Delbianco group presented an automated approach to fluorinated Lewis type 2 antigens.¹⁵⁶ In 2022, the Gilmour group disclosed an automated synthesis of HIV-relevant fluorinated high mannose structures using the Glyco-ner 2.1® synthesizer.¹⁵⁷ Furthermore, in 2026, the same platform was used to generate fluorinated frameshifts of *Klebsiella pneumoniae* antigens to investigate how antigen frameshifts affected binding to the lectin Concanavalin A.¹⁵⁸ These representative examples serve to illustrate the levels of complexity that can be generated in F-glycan mimetics, and the opportunities that this technology affords in vaccine development.

4. Fluorinated carbohydrate vaccines

4.1. Prophylactic vaccines

Prophylactic carbohydrate-based vaccines are administered prior to an initial infection and protect against future infection by the target pathogen. Lacking easily targetable proteins on their surfaces, certain bacteria rely on their unique CPS to infect a host, evade immune responses, and cause disease. This capsule physically shields the bacterium, enabling it to evade initial immune detection by masking underlying antigens. Vaccines targeting this capsule induce highly specific antibodies that can neutralise the pathogen with remarkable precision. Research has consistently shown that antibodies directed at polysaccharides are highly beneficial in preventing bacterial infections.¹⁵⁹ Consequently, vaccines based on these capsular polysaccharides, in both monovalent and multivalent forms, have been approved for use against *Streptococcus pneumoniae*, *Neisseria meningitidis*, *Haemophilus influenzae* type b (Hib), and *Salmonella typhi*.¹⁶⁰ Regrettably, this does not guarantee permanent protection against these infections, and new and modified vaccines are required.

4.1.1. *Neisseria meningitidis*. *Neisseria meningitidis* (meningococcus) is a Gram-negative bacterium known to be a causative agent of meningococcal meningitis and meningococcal septicemia.¹⁶¹ Sialic acids play a significant role in the biology and pathogenicity of *N. meningitidis*, shaping its lipooligosaccharide (LOS) structure. Sialylation of LOS makes the bacterium appear more “host-like”, reducing its recognition by the immune system. *N. meningitidis* serogroup B CPS contains α -(2,8)-linked *N*-acetylneuraminic acid (Neu5Ac),¹⁶² a structure also prevalent in human neuronal glycans.¹⁶³ Due to this molecular mimicry, the epitope is poorly immunogenic and carries risks of inducing autoimmunity. In contrast, the α -(2,9)-linked sialyl epitope is uniquely bacterial and absent from human tissues, making it an attractive target for vaccine design.^{164,165}

This observation inspired the work of the Gilmour and Seeberger groups on a fluorinated sialic acid vaccine lead against meningitis B and C.¹⁶⁶ The shortest immunogenic unit was described as a disaccharide⁴⁴ informing the vaccine design




Fig. 7 Fluorinated sialic acid vaccine lead against meningitis B and C reported by the Gilmour and Seeberger groups and the schedule of mice immunization with candidates 1-CRM₁₉₇ and 1-PorA.¹⁶⁶

which was a fluorinated disialoside (**1**) with a linker for protein coupling at the reducing end (Fig. 7, top). Fluorine was strategically introduced at the C3 position adjacent to the anomeric centre since C3 fluorinated sialic acid derivatives do not undergo elimination during glycosylation and the fluorine also acts as a directing group affording mostly α -product.^{167,168} In addition, the fluorine also functions as a practical and sensitive NMR-active nucleus that expedites structural analysis. Following global deprotection, the glycan was conjugated *via* a *p*-nitrophenyl adipate ester to two different carrier proteins: CRM₁₉₇ and PorA. To investigate the influence of these candidates on the immune response, the study employed either aluminium hydroxide adjuvant (Alum) or Freund's Adjuvant. For immunological studies, four groups of six of C57BL/6 mice were immunized in accordance with the schedule (Fig. 7, bottom). Given the expected increase in immunogenicity from fluorination, a much smaller amount of the conjugate vaccine was used.

All tested glycoconjugates successfully activated the immune system in a T-cell-dependent manner, and this was reflected in the similar long-term IgG responses observed. The highest level of antibodies was achieved on day 70, after which the levels decreased. The booster immunization triggered the rise of IgG levels back to previous levels, revealing production of memory B cells. IgG₁ isotype antigens were predominantly formed, IgG₂, and IgG₃ were indicated in lower amounts. Among those, IgG₁ and IgG₃ can effectively bind and activate complement. All glycoconjugates demonstrated specific binding to *N. meningitidis* C CPS and to whole bacteria. *N. meningitidis* B CPS was






Fig. 8 *L. donovani* trisaccharides with different fluorination patterns of the galactose unit synthesized by Hoffmann-Röder and co-workers.¹⁷⁰

specifically recognized only by antibodies induced by **1-PorA** containing conjugates. Antibodies induced by **1-PorA-Alum** were found to be protective against both *N. meningitidis* B and C and **1-CRM₁₉₇-Alum** displayed the highest level of protection.

4.1.2. Leishmania donovani. *Leishmania donovani* is a parasitic protozoan that causes visceral leishmaniasis, also known as kala-azar or black fever.¹⁶⁹ It is a life-threatening condition because it primarily targets the internal organs, most notably the liver, spleen, and bone marrow.

In 2018, Hoffmann-Röder and co-workers reported the synthesis of fluorinated analogues of the trisaccharide *L. donovani* CPS component 2 (Fig. 8).¹⁷⁰ The authors focussed particular attention on the galactose subunit, which was systematically fluorinated. The synthetic strategy enabled the target disaccharides to be coupled to the terminal mannose subunit to furnish the desired trisaccharides with the NH₂-linker (3–7). In so doing, modified epitopes potentially suitable for the development of diagnostic tools and synthetic anti-leishmanial vaccines were developed. According to the authors, work in the aforementioned direction is in progress.

4.1.3. Streptococcus pneumoniae. *Streptococcus pneumoniae* (pneumococcus) is a Gram-positive bacterium with a dual

nature: it's a common colonizer of the upper respiratory tract, but also a major human pathogen.¹⁷¹ Consequently, *S. pneumoniae* presents a threat to infants, immunocompromised patients, and the elderly. The impact of the bacterium, though reduced, remains significant due to its ability to adapt to antibiotics and serotype replacement.¹⁷¹ *S. pneumoniae* serotype 8 (ST8) is now one of the top-ranking serotypes causing invasive pneumococcal diseases as it is only partially covered by the glycoconjugate vaccines in use (present in PPSV23 and PCV-20 and absent in PCV-7, PCV-13, and PCV-15).^{120,172} This makes the ST8 capsular polysaccharide an important target for vaccine development.

The minimal protective epitope of ST8, which appeared to be a trisaccharide, was identified in 2017 by Schumann and co-workers.¹⁷³ It was found that by conjugating it to a glucose moiety, the relatively low immunogenicity can be enhanced. An important study by Hoffmann-Röder and co-authors validated synthetic routes to fluoro-substituted glycans **8–16** with a linker at the reducing end for subsequent conjugation steps (Fig. 9).¹⁷⁴ The 6-position was chosen as the most exposed site and each of the molecules obtained contained only one fluorinated unit for the sake of subsequent systematic investigation of the heteroatom impact.

In 2026, the same group described a synthetic approach to the minimal protective antigenic structure of CPS of *S. pneumoniae* ST14 (Fig. 10),¹⁷⁵ which has found application as a component of several vaccines in current use.¹⁷⁶ The authors prepared tetrasaccharides **17–22** with allyl groups appended at C1: this is a convenient handle for conjugation to a carrier protein. These materials will certainly be valuable in determining the influence of fluorination on the subsequent antibody production and vaccine design.

4.1.4. Summary. From these selected case studies, it is evident that fluoro-glycans hold great promise in the conception of vaccine leads against bacteria, but that the field is at an early stage. Whilst the sialic acid-based vaccine lead against meningitis B and C demonstrates convincing proof of concept in bolstering immunogenicity, stability, and the induced antibodies' cross-reactivity, there is much to be done. Further advancement of vaccine candidates will require a concerted, interdisciplinary approach that includes preclinical testing, scalable manufacturing, and compelling evidence that they are




Fig. 9 *S. pneumoniae* ST8 tri- and tetrasaccharides with different fluorination patterns at the 6-position of the galactose units synthesized by Hoffmann-Röder and co-workers.¹⁷⁴




Fig. 10 *S. pneumoniae* ST14 tetrasaccharides with different fluorination patterns synthesized by Hoffmann-Röder and co-workers.¹⁷⁵






Fig. 11 Therapeutic carbohydrate-based vaccine leads based on TACAs: mono-epitopic Globo H vaccine by Wong, Danishefsky and co-workers and multi-epitopic pentavalent vaccine by Danishefsky and co-workers; KLH = keyhole limpet hemocyanin.^{177,178}

more effective and safer than the current vaccines. This need to innovate in opening up new areas of glycomimetic chemical space is also evident from how a popular commercial vaccine against pneumococcus – PCV-7 and PCV-13 – having immunized millions of people was futile against ST8 and supported the rise of the ST8 epitope.

4.2. Therapeutic vaccines

In the arena of therapeutic carbohydrate-based vaccines, emphasis has been heavily concentrated on the aberrant glycosylation patterns of malignant cells.^{8,65} Tumor-associated carbohydrate antigens (TACAs) are antigens, which are over- or uniquely expressed on the tumor cell surface.^{45,75,179} TACAs arise from alterations in sugar chain structure, including truncation, sequence changes, or excessive sialylation of glycolipids and glycoproteins. Abnormal glycosylation is shown to be linked to key stages of cancer progression, such as invasion and

metastasis. Since many TACAs are predominantly expressed during cancer development and are largely absent from normal adult tissues, they represent promising targets for therapeutic cancer vaccines.^{180,181} A wide range of TACAs have been characterized, including: the mucin related (*O*-linked) GalNAc (Tn), sialyl Tn (sTn), Thomsen–Friedenreich (TF), sialyl TF; the glycosphingolipids Globo H, stage-specific embryonic antigen-3 (SSEA-3); the gangliosides GM2, GD2, GD3, fucosyl GM1; poly-sialic acid (PSA).^{182,183}

Carbohydrate-based cancer vaccines can be grouped into three different categories: (1) mono-epitopic, (2) mono-epitopic cluster (presents multiple copies of a single epitope), and (3) multi-epitopic vaccines.¹⁸⁴ A mono-epitopic vaccine of Globo-H 23, an unnatural TACA, was developed by Wong, Danishefsky, and co-workers (Fig. 11, top).¹⁷⁸ Overall, five different total syntheses have been reported, of which an enzymatic sequence proved the most cost effective, with a yield of over 80% over only






Fig. 12 Fluorinated Tn analogues with modified *N*-acyl groups by Ye and co-workers and the standard immunization scheme applied by the Ye group.¹⁸⁷

two steps. In Phase I trials, a synthetic Globo-H-KLH conjugate was tested and the study confirmed the immunogenicity and safety of the vaccine candidate, which has now moved to Phase II/III trials. A multi-epitopic, unimolecular, and pentavalent vaccine **24** that targets Globo-H, STn, Tn, TF, and GM2 was developed by Danishefsky and co-workers (Fig. 11, bottom).¹⁷⁷ The vaccine candidate showed induction of IgG and IgM antibodies against all five of the carbohydrate antigens. Unfortunately, these vaccine candidates suffer from non-specific, short-lived immune responses. To overcome this, considerable effort has been invested in increasing immunogenicity *via* sugar structure modification, especially fluorination (for more examples of non-fluorinative modifications please see Section 3.1.4).

4.2.1. Thomsen-nouveau (Tn) antigen. The Thomsen-nouveau (Tn) antigen (**25**) is a truncated *O*-linked glycan bound to mucin proteins *via* serine or threonine. It consists of *N*-acetyl galactosamine.¹⁸⁵ Tn is a known TACA and has been found to be present in up to 90% of cases across a spectrum of cancers.¹⁸⁶

In 2016, Ye and co-workers synthesized and evaluated three fluorinated Tn analogues by modifying the *N*-acyl group (Fig. 12).¹⁸⁷ Analogues with a monofluoromethyl group (**26-CRM₁₉₇**), a difluoromethyl group (**27-CRM₁₉₇**), and a trifluoromethyl group (**28-CRM₁₉₇**) on the *N*-acyl moiety were prepared by reacting a 2-amino galactose derivative with the corresponding methyl ester of the fluorinated motif. The compounds were conjugated to CRM₁₉₇ and combined with the C34 adjuvant for the immunization of mice in accordance with the schedule (Fig. 12, bottom). After 4 vaccinations, the anti-STn IgG antibody titer was greatest for **26-CRM₁₉₇** (2.7 times greater than that of **25-CRM₁₉₇**). However, **27-CRM₁₉₇** and **28-CRM₁₉₇** led to a lower IgG titer than when vaccinating with **25-CRM₁₉₇**. Furthermore, flow cytometry revealed that the immunized (**26-CRM₁₉₇**) mouse sera could recognise Tn in its native




Fig. 13 Fluorinated STn analogues synthesized by modifying the two amide groups of STn by the Ye group.^{47,194}

environment on cancer cells (MCF-7 and Jurkat) after the third vaccination. It was also found that vaccination with **26-CRM₁₉₇** led to a higher proportion of IFN- γ producing splenocytes than for **25-CRM₁₉₇**, indicating a Th1 skewed response and that fluorination patterns offer a potential solution.

4.2.2. Sialyl-Thomsen-nouveau (STn) antigen. The sialyl-Thomsen-nouveau (STn) (**29**, Fig. 13) antigen is a TACA associated with poor prognoses.¹⁸⁸

This truncated *O*-antigen is capped by sialylation (of the Tn antigen) and linked to a protein *via* a serine or threonine residue at the reducing end. There has been interest in exploiting the STn antigen as a potential target for vaccination in oncology. One high-profile example was Theratope®, a conjugate of STn and KLH that was developed as a potential vaccine against metastatic breast cancer.¹⁸⁹ In initial trials this candidate showed great promise with significant anti-STn IgG titer.¹⁹⁰ However, in 2003, a phase III trial with 1028 patients showed that the vaccine didn't significantly prolong survival (23.1 months *vs.* 22.3 months for a control group).¹⁹¹ This unexpectedly poor performance was explained as possibly resulting from low STn expression on the tumors (they were not phenotyped), the trial not being long enough, or an inadequate immune response.^{191,192} This was despite the vaccine being administered after cyclophosphamide treatment to suppress regulatory T cells (Tregs)¹⁹³ and with an adjuvant for the first vaccination to enhance the immune response.

In a bid to develop more immunogenic STn vaccine candidates, in 2010, Ye and co-workers synthesized a panel of STn analogues by modifying the two amide groups of STn (Fig. 13).⁴⁷ Compounds **30–38** have the trifluoro-, difluoro- and monofluoromethyl groups incorporated into the amides. A variety of other amide modifications were investigated but here only those containing fluorine will be discussed. Enzyme-linked immunosorbent assay (ELISA) screening was used to identify promising potential vaccine candidates. The most promising candidates (*i.e.*, strongest binders) were conjugated to keyhole limpet hemocyanin (KLH) for *in vivo* immunological evaluation in mice. Important to note is that no adjuvants were used to enable a more robust comparison between glycan epitopes. After 4 vaccinations (Fig. 12, bottom), pooled sera were analyzed to measure the anti-STn IgG and IgM titers.



This revealed that the CF₃ modifications (**32-KLH**, **33-KLH**, **38-KLH**) were detrimental in all positions. Modification of only the sialic acid amide (**30-KLH**, **31-KLH**, **32-KLH**) also offered no improvement over native STn. However, a monofluoromethyl amide on the galactose (**33-KLH**) or on both sugars (**36-KLH**) led to anti-STn IgG titers 2 times and 2.3 times higher than 1-KLH, respectively. A difluoromethyl amide was beneficial on the galactose moiety (**34-KLH**) resulting in anti-STn IgG titers 2 times higher than 1-KLH. This revealed that **33-KLH**, **34-KLH**, and **36-KLH** have potential as vaccine candidates against STn, and that they may be able to overcome the problems observed when using native STn as a vaccine. The IgG/IgM ratio was found to be higher for the fluorinated compounds compared to the natural **30**. The binding of these antibodies to STn in its natural environment was proven using flow cytometry to confirm binding to human cancer cells expressing STn. Specificity was also demonstrated by the diminished binding to STn deficient human cancer cells.

Candidates **34**, **35**, and **37** underwent further testing in 2017, with a murine tumor model using murine colon cancer cells.¹⁹⁵ During a course of five bi-weekly vaccinations, after the penultimate vaccination, mice were injected with CT-26 cells in a tumor challenge study. Cyclophosphamide was given before the initial vaccination, which used Freund's complete adjuvant. Subsequent vaccinations utilized Freund's incomplete adjuvant. Of the three tested vaccines, **36-KLH** showed a significant reduction in tumor burden in the lungs compared to other vaccines, including **29-KLH**, and significantly extended the survival time.

With the promising activity of fluorinated **36** established, attention turned to investigating other carrier proteins, e.g., CRM₁₉₇.¹⁹⁶ In 2019, Ye and co-workers synthesized **36-CRM₁₉₇** and compared it to **29-CRM₁₉₇** with (Freund's or C34) and without adjuvant.¹⁹⁴ The findings showed that the anti-sera from the fluorinated vaccine were more effective at promoting cancer cell lysis by complement dependent cytotoxicity (CDC) and antibody dependent cellular cytotoxicity (ADCC) pathways. Furthermore, analysis of splenocytes revealed they produced interferon-gamma (IFN-γ) and interleukin 4 (IL-4) upon *in vitro* stimulation by the glycoconjugate, indicating a mixed Th1/Th2 response. This mixed cellular and humoral response was found to be stronger with splenocytes from **36-CRM₁₉₇** vaccinated

mice *versus* those vaccinated with **29-CRM₁₉₇**. Furthermore, the anti-STn IgG titers were higher for **36-CRM₁₉₇** with and without adjuvant, confirming the fluorinated motif **36** yielded a multi-faceted improvement to the immune response.

With the intention of developing another panel of potential STn vaccine candidates with improved immunogenicity, in 2015, Ye and coworkers reported the synthesis and evaluation of fluorinated S-linked STn conjugates (Fig. 14).¹⁹⁷ With the fluorination verified to improve the immune response to STn in previous work,¹⁹⁵ the sulfur linkage was purported to contribute to the "non-self" nature of the glycan and to improve its stability, which has been demonstrated in literature.^{198,199}

The previous conjugates of **29–38** with KLH were prepared by linking the glycan bearing an allyl group to the protein, first by oxidation to the aldehyde and then by reductive amination to couple the glycan to the protein.¹⁹⁵

None of the S-linked STn conjugates resulted in higher anti-STn IgG titers than native STn (**29-KLH**). The lower anti-STn IgG titers for the S-linked conjugates were not a result of poor immunogenicity, but rather of poor cross reactivity of the antibodies with native STn. It was observed that fluorination could increase the anti-native STn IgG titers in mice compared to the non-fluorinated S-linked conjugate **39-KLH** by improving the cross reactivity of the antibodies. The best performing S-linked conjugate was **46-KLH** with two monofluoromethyl amides, though this still resulted in less than 10% of the titer seen for **29-KLH** after 4 vaccinations without adjuvant.

Overall, these studies of modified STn conjugates indicate that fluorinated S-linked STn derivatives (as opposed to S-linked STn) could be promising vaccine candidates owing to their improved immunogenicity compared to **29**. STn derivative **36** with two monofluoromethyl amides showed particularly promising behaviour.

4.2.3. Microvesicles. Microvesicles displaying fluorinated analogues of STn were shown to be potentially useful as cancer vaccine leads by Ye and coworkers in 2023 (Fig. 15).²⁰⁰ The microvesicles bearing modified STn (**50**) were produced by incubating cells with fluorinated carbohydrate precursors and




Fig. 14 Fluorinated S-linked STn analogues synthesized by modifying the two amide groups of STn by the Ye group.¹⁹⁷




Fig. 15 Fluorinated analogues of STn displayed on microvesicles and the immunization schedule.²⁰⁰






Fig. 16 Fluorinated TF analogues synthesized by modifying the amide group of TF by the Ye group.²⁰⁴

isolating the resulting microvesicles. Mice had been immunized with the microvesicles prior to implantation of a tumor (Fig. 15, bottom). Immunization with **50** proved to be more effective at tumor growth inhibition compared to **49** (*i.e.*, CH₃ to CH₂F had a positive impact). Immunization with microvesicles also reduced metastasis compared to wild type microvesicles when mice were challenged with cancer cells. It was reported that one of the reasons for the benefits seen using the fluorinated STn (**50**) was an increased affinity to the major histocompatibility complex I (MHC-I).

4.2.4. Thomsen-Friedenreich (TF) antigen. The Thomsen-Friedenreich (TF) antigen is a disaccharide TACA commonly found in breast and prostate cancers.²⁰¹ TF-KLH was explored as a potential vaccine, with the suite of leads structures including fluorinated derivatives.^{202,203} In 2016, Li, Ye and co-workers synthesized the TF antigen (**51**) as well as modifications of the *N*-acetyl group: monofluoromethyl (**52**), difluoromethyl (**53**) and trifluoromethyl (**54**) (Fig. 16).²⁰⁴ These were conjugated to CRM197 and used to vaccinate mice with C34 as an adjuvant. After 4 vaccinations, **52-CRM**₁₉₇ led to the highest anti-51 IgG titer (2.5 times greater than **51-CRM**₁₉₇). The trifluoromethyl derivative **54-CRM**₁₉₇ led to anti-TF IgG titers 1.6 times larger

than **51-CRM**₁₉₇. However, the difluoromethyl derivative **53-CRM**₁₉₇ performed worse than **51-CRM**₁₉₇. Flow cytometry confirmed that the antibodies could bind tumor cells expressing Tn and that the antibodies from modified Tn vaccines bound more strongly than antibodies from **51-CRM**₁₉₇. Antisera were evaluated on their ability to affect CDC-mediated tumor cell killing. The results showed 74% lysis for pooled sera after **52-CRM**₁₉₇ vaccination compared to 57% after **51-CRM**₁₉₇ vaccination. Collectively, these experiments validated that fluorination, in particular the introduction of the monofluoromethyl amide, could increase the immunogenicity of the TF antigen, producing antibodies with good cross-reactivity for the native Tn antigen. This indicates a potentially productive avenue for anti-cancer vaccine development.

4.2.5. Mucin-associated Thomsen-Friedenreich (TF) antigen. Mucin 1 (MUC1) is a transmembrane glycoprotein, primarily studied in mammals, including humans, mice, rats, and other species.²⁰⁵ It is presented on the surface of a multitude of cells: the respiratory tract, gastrointestinal tract, genitourinary system, *etc.* The primary function of this glycoprotein lies in providing a barrier against pathogens and other environmental factors to protect epithelial surfaces.

MUC1 is well documented to be overexpressed on many types of cancer cells, such as breast, pancreatic, ovarian, or lung.^{179,206} The cancer-associated glycoprotein is distinct from its healthy counterpart not only in the abundance, but also in its structural and functional characteristics (Fig. 17).²⁰⁷ MUC1 covering normal cells is heavily glycosylated: complex and long carbohydrate chains mask the protein core. In an interesting reversal of circumstances, and as a result of altered glycosyltransferase activity, on tumor tissues it undergoes aberrant glycosylation, characterized by shorter, simpler carbohydrate chains (*e.g.*, truncated *O*-glycans like T, Tn, and STn antigens).





Fig. 17 MUC1 covering normal cells (covered with complex and long carbohydrate chains masking the protein core) and tumor-associated MUC1 (aberrant glycosylation characterized by shorter and simpler carbohydrate chains).²⁰⁷



Hoffman-Röder, 2009



Hoffman-Röder, 2010




Fig. 18 First fluorinated glycoprotein possessing a fragment of MUC1 peptide chain synthesized by Hoffmann-Röder and co-workers (top),²⁰⁸ glycoprotein mimetic including full the MUC1 repeat sequence prepared by Hoffmann-Röder and co-workers (bottom).²⁰⁹

These patterns are mainly related to the extracellular region of the glycoprotein consisting of a variable number of tandem repeats, each built of 20 amino acids (the sequence PDTRPAPGSTAPPAHGVTSA). These changes expose peptide epitopes that are normally hidden, making cancer-associated MUC1 immunologically distinct. These key characteristics of MUC1 have shed light on this glycoprotein as a potential target for cancer immunological therapy.

In 2009, Hoffmann-Röder and co-authors described a route to synthesize MUC1 fragments bearing fluorinated analogues of T (55–57) and sialyl-T (58) antigens (Fig. 18, top).²⁰⁸ Di- and trisaccharides were prepared according to standard glycosylation procedures. Galactosamine building blocks were conjugated to threonine to be subsequently incorporated into a peptide chain. Compounds 56 and 57 were then submitted to automated solid-phase peptide synthesis (SPPS) according to Fmoc-strategy to be integrated into a MUC1 tandem repeat-peptide sequence comprising the immunodominant PDTRP epitope (a fragment of the peptide presented on Fig. 18). Ultimately, glycopeptides 59 and 60 were obtained following global deprotection with overall yields of 41% and 18% (based on the loaded resin), respectively.

In 2010, the group expanded their synthesis of glycopeptide analogues to create a mimetic that included the full MUC1 tandem repeat sequence (Fig. 18, bottom).²⁰⁹ Initially, fluorine-modified antigens were prepared according to standard synthetic strategies. The last step of each synthesis was

deprotection of the ester of the Thr moiety for subsequent SPPS conducted according to the standard SPPS method. After the Fmoc-deprotection of the final amino acid, the triethylene glycol spacer was conjugated to the obtained sequence. Following cleavage from the resin led to the formation of 61 and 62 in 29 and 21% yields, correspondingly. Global deprotection of the glycopeptides was conducted to give the mimics in 55 and 60% yields, thereby leading to the synthesis of a convenient platform for ensuing vaccine candidates' preparation.

In 2010, Kunz, Hoffmann-Röder and co-workers prepared a series of fluorinated and non-fluorinated MUC1 glycopeptide analogues conjugated to tetanus toxoid as vaccine candidates (64) to test if enzymatically stable fluorinated mimics could replace their less stable natural counterparts 63 (Fig. 19, top).²⁰² The diethyl ester of squaric acid was used to conjugate the SPPS-synthesized glycopeptides to the carrier proteins (BSA and TT). The immunological properties of the vaccine candidate leads were evaluated *via* immunization of three balb/cJ mice with 63-TT and three mice with 64-TT. The immunization schedule is summarized in Fig. 19 (bottom). Both vaccine candidates induced high antibody levels, although the values for the fluorinated conjugate were lower. Further binding studies revealed that the antibodies formed by 63-TT strongly recognized 63-BSA, mirroring its glycopeptide sequence. Similar affinity was observed for the BSA conjugate containing 2,6-sialyl-T antigen. The antibodies induced by 64-TT showed very similar level of recognition of 63-BSA, 2,6-sialyl-T antigen, sialyl-T_N antigen,






Fig. 19 Conjugation of fluorinated and non-fluorinated MUC1 glycopeptide analogues to the carrier proteins by Kunz, Hoffmann-Röder and co-workers and the immunization schedule.²⁰²

and the conjugate with non-glycosylated MUC1 peptide, whereas shortened to 12 amino acids the glycopeptide sequence was only recognized weakly. An important finding showing cross-reactivity potential is the equally strong binding observed for both the native conjugate **63-BSA** and the fluorinated version **64-BSA**. For both vaccine candidates, minimal IgM, IgA, and IgD antibody production was observed, indicating selective immune responses and initiation of immunological memory. IgG₁ was




Fig. 20 Fluorinated TF antigens by Hoffmann-Röder and co-workers.²¹⁰

predominant for the natural and the fluorinated candidates, whereas IgG_{2a} and IgG_{2b} levels were slightly higher for **64-TT**. Moreover, the authors showed that the antibodies induced by both vaccines strongly bound to MCF-7 (human breast cancer cell line). Noteworthy, the T-antigen-MUC1 glycopeptide, as well as the corresponding difluorinated antigen, neutralized binding of the antibodies from **64-BSA** and **64-TT** when added to the MCF-7 cells. In contrast, MUC4, possessing a different peptide sequence, did not show any signs of neutralization, which suggests selective recognition of tumor-associated antigens.

In 2011, Hoffmann-Röder and co-authors published the syntheses of two new fluorinated TF antigens (Fig. 20).²¹⁰ The synthetic approach enabled α/β mixtures of **65** and **66** to be obtained, which were separated to isolate the β -isomers. Only antigen **65** was used in the subsequent research.

Later, Hoffmann-Röder and co-authors continued their investigation on TF-fluorinated antigen **67–71** development and investigated specific binding to serum antibodies induced by the vaccine candidates **63-TT**, **64-TT**, and **72-TT** presented in Fig. 21.²¹¹ After preparing the fluorinated disaccharides glycosylated to Thr, standard SPPS was applied to obtain the corresponding glycoconjugates **67–71** in 15–44% yields. Neutralization experiments were performed to test if the number and position of the fluorine atoms on the glycan moieties of the glycoconjugates affect binding. For this, antibodies induced by the natural (**63-TT**), mono- (**72-TT**), and difluorinated (**64-TT**) glycopeptide-based vaccine candidates were applied. ELISA plates were coated with the glycopeptide-BSA conjugates reflecting the structure of the vaccines. For **63-TT**, binding of all freshly prepared antibodies was similar and strong as for the native one. Monofluorinated candidate **72-TT**, glycopeptides **67–69** and **71** demonstrated similar affinities, while structure **70** (no fluorine at C2') displayed slightly lower binding. The antibodies induced by **64-TT** showed similar binding with **67**, **68**, and **70**, but **69** and **71** (no fluorine at C6) bound weaker. These recognition patterns suggest that the affinity can be affected by the fluorination pattern difference between the vaccine candidate applied to induce antibodies and the antigens. The next question to be addressed was the effect of an altered peptide sequence on the binding. To investigate this, the MUC4 glycopeptide, possessing a different peptide sequence, was tested with the antibodies induced by vaccine candidate **72-TT**; no recognition was observed. These findings






Fig. 21 Fluorinated MUC1 antigens by Hoffmann-Röder and co-workers (top), synthetic vaccine candidates used for immunization (bottom).²¹¹

indicate that vaccine-induced antibodies primarily bind the whole MUC1 glycopeptide antigen, rather than its substructures like the (fluorinated) carbohydrate moiety.

In 2011, the same group described a promising new vaccine candidate **73-BSA** (Fig. 22).²¹² The disaccharide **73** was synthesized as previously described and subjected to SPPS, giving the product in 50% yield. To conjugate this structure to BSA, a previously established strategy *via* diethyl squarate was employed. Next, the **73-BSA** binding to the antibodies induced by the vaccine candidates **63-TT** and **64-TT** was investigated. The binding of the serum antibodies raised by immunization with **63-TT** was partially neutralized by the natural antigen and fluorinated **73-BSA**. In the case of the vaccine **64-TT**, a notable difference in binding was observed. The non-fluorinated natural antigen showed the lowest affinity, with affinity increasing for the 6,6'-difluorinated antigen and reaching the highest value **73-BSA**. Despite the differences in binding, the observed cross-reactivity underscores the potential of fluorinated glycopeptide conjugate **73-BSA** to become a vaccine candidate.




Fig. 22 Fluorinated MUC1 antigen-BSA conjugate by Hoffmann-Röder and co-workers.²¹²

In 2015, Hoffmann-Röder and co-authors prepared a vaccine candidate, bearing fluorine atom at C4' position.²⁰³ The synthesized fluoro disaccharide **74** and the native antigen **75** were submitted to an *in vitro* stability test against enzymatic degradation (Fig. 23). After incubation with the enzyme in aqueous 2-(*N*-morpholine)ethanesulfonic acid (MES buffer) at pH 4.5 (enzyme activity optimum) in the presence of 2,6-di-*O*-methyl- β -cyclodextrin at 25 °C, the native antigen was half-digested in \sim 3 h, while the fluorinated **74** stayed unreactive over 7 h. This clearly demonstrates the enhanced metabolic stability of the fluorinated glycan.

Subsequently, standard SPPS was used to synthesize the fluorinated glycopeptide **76** (13%), which was conjugated to a carrier protein *via* diethyl squarate (Fig. 24, top). The immunogenicity of the synthesized vaccine candidate was assessed by immunizing three balb/cJ mice with compound **76-TT**, following the schedule detailed in Fig. 24 (bottom). Five days after the final vaccination, blood was collected, and serum antibody levels were measured using ELISA. The ELISA plates were coated with the corresponding conjugate **76-BSA**. The titer level was found to be similar to the results obtained for the difluorinated vaccine **64-TT**, which was used as a positive control in the current study. Carrier protein – tetanus toxoid – was tested in terms of immunogenicity as well and demonstrated a strong immune response. Isotype analysis of the antisera revealed intense formation of IgG₁ and lower levels of IgG_{2a} and IgG_{2b}. On the other hand, IgM, IgA, and IgD were produced in negligible amounts. MCF-7 binding experiments gave positive results, but it is worth to mention that the binding of antibodies from all three mice significantly differed.




Fig. 23 Fluorinated and non-fluorinated glycan stability test against enzymatic degradation.²⁰³






Fig. 24 Fluorinated MUC1-protein conjugates by Hoffmann-Röder and co-workers and the mice immunization schedule.²⁰³

Additionally, the antibody binding levels measured in the sera following the first booster immunization were higher than those observed after the second booster.

4.2.6. GM3. GM3 **77** is one of the most frequently encountered gangliosides in human cells, and it is found on the surface of nearly all vertebrate cells.²¹³ It can be overexpressed and more exposed on tumor cells, making it a prominent target for vaccine development.²¹⁴ In 2015, Ye and co-workers synthesized a panel of GM3 derivatives with various *N*-acyl modifications to the sialic acid moiety, among which were the fluorinated motifs: monofluoromethyl (**78**), difluoromethyl (**79**), and trifluoromethyl (**80**) amides (Fig. 25). Among the non-fluorinated analogues, the ethyl amide **81** performed particularly well in immunological evaluation.²¹⁵ These glycans were conjugated to KLH and used to vaccinate mice, without an adjuvant.

After 4 vaccinations in accordance with the standard scheme (Fig. 12, bottom), the anti-GM3 IgG titers were quantified. The non-fluorinated **81-KLH** led to the highest titer, 3.7 times greater than that of **77-KLH**. The second highest IgG titer resulted from **79-KLH**, 1.6 times greater than for **77-KLH**. These results showed that the difluoromethyl compound **79** could be investigated as a potential vaccine, however the ethyl compound **81** induced a stronger IgG response in this investigation. This did illustrate that certain modifications led to increased immunogenicity compared to the native structure **77** and that




Fig. 25 Fluorinated GM3 derivatives with various *N*-acyl modifications by Ye and co-workers.²¹⁵

some of these structures afforded antibodies with a cross reactivity to **77** that could indicate promising vaccine candidates.

4.2.7. Globo H. Globo H (**82**) is a key TACA with significant implications in cancer diagnosis and immunotherapy. It is overexpressed in a wide range of cancers, including: breast, prostate, ovarian, colorectal, and lung cancer.²¹⁶ Synthetic Globo H based vaccines have progressed to various stages of clinical studies.²¹⁷ Despite the promising start, early-stage clinical results demonstrated that the level of IgG antibodies induced by the vaccine candidates was significantly lower than IgM antibodies.

In 2021, Ye and co-workers replaced the methyl moiety of the native acetamide of **82** with a monofluoromethyl group (**83**), difluoromethyl group (**84**) and a trifluoromethyl group (**85**) (Fig. 26).²¹⁸ Conjugates with CRM₁₉₇ were used as vaccines for mice with C34 as the adjuvant following the standard scheme (Fig. 12, bottom). **84-CRM₁₉₇** and **85-CRM₁₉₇** led to significantly higher anti-Globo-H IgG titers compared to **83-CRM₁₉₇**. Antisera from **84-CRM₁₉₇** and **85-CRM₁₉₇** vaccinated mice led greater cell lysis towards MCF-7 cancer cells by complement-dependent cytotoxicity (CDC).

Wong and co-authors have demonstrated that altering the carrier and the adjuvant used can induce a strong immune response with the antibody class switching from IgM to IgG in case of Globo H and Globo H-related epitopes.²¹⁹ In light of these promising findings, the group decided to investigate the impact of structure modifications with different substituents.




Fig. 26 Fluorinated Globo H-CRM₁₉₇ conjugates with *N*-acyl modifications by Ye and co-workers.²¹⁸






Fig. 27 Globo H fluorinated analogues by Wong and co-workers.²¹⁷

Fluorine was incorporated at the C-6 position of Glc at the reducing end (**86**) or the C-6 position of Fuc at the nonreducing end of Globo H (**87**) (Fig. 27).²¹⁷ Corresponding hexasaccharides bearing an amine-linker at the reducing end were prepared through chemoenzymatic synthesis strategies. Unfortunately, an attempt to obtain difluoro-substituted molecule **88** failed giving only trace amounts of the desired product. Molecules **86** and **87** were conjugated to the DT (carrier protein) *via p*-nitrophenyl adipate diester and later combined C34 adjuvant. For immunological investigation 10 groups of 5 female balb/c mice were immunized intramuscularly with Globo H analogues **86-DT** and **87-DT** according to the Fig. 12 (bottom). The collected sera were tested on the glycan microarray consisting of 94 different TACAs.

Antibodies generated by the tested candidates were recognized by Globo H, Globo H analogues, and Globo H fragments, but not by other TACAs and functional linkers. As Globo H, Gb5, and SSEA4 were found to be overexpressed on breast cancer stem cells and possess rather similar core structures, these TACAs were selected as standard antigens for the DT-conjugates. In the binding tests of antibodies induced against the chosen antigens high IgG antibodies titer, indicating a T-cell-dependent immune response, was obtained. No significant IgM antibody formation was indicated in case of both candidates. IgG tests revealed regioselectivity of the

immunogenicity modulation by the fluorine substituents. The F moiety at the C-6 position of Glc at the reducing end of Globo H (**86-DT**) demonstrated a comparable titer to the natural Globo H, whereas **87-DT**, bearing F group at the C-6 position of Fuc at the nonreducing end of Globo H, showed significantly weaker interaction with Globo H. Rather similar results were obtained during the studies with Gb5 and SSEA4. The ratio of IgG/IgM for **86-DT** and **87-DT** was found to be 78.07 and 11.28, respectively. IgG subclass test showed that the antibodies had a significant amount of IgG₁, IgG_{2b}, IgG_{2c}, and IgG₃ and low level of IgG_{2a}. It is noteworthy that the glycans bearing the azide-group instead of the fluorine-atom performed better.

4.2.8. KH-1. In 2023, Ye and co-workers synthesized CRM₁₉₇ conjugated analogues of KH-1 where the methyl moiety of the native acetamide of **89** was replaced by a difluoromethyl group (**90**) and a trifluoromethyl group (**91**) (Fig. 28).²²⁰ Mice were vaccinated 4 times with glycoconjugates (Fig. 12, bottom) and Freund's adjuvant then anti-KH-1 IgG titers were determined. The titer for **91-CRM197** was 26 times greater than that of **89-CRM197**. **90-CRM197** led to 2.2 times higher anti-KH-1 IgG titers than the native structure. Analysis of the IgG subtypes revealed a mixed Th1/Th2 response by the presence of IgG₁, IgG_{2a}, IgG_{2b}, and IgG₃. The IgG antibodies were capable of binding native KH-1 on MCF-7 cancer cells, with the antibodies from the fluorinated analogues displaying a higher binding affinity than those from **89-CRM197**.

4.2.9. Pentamannose. The Man₅ motif is found on the gp120 protein of HIV and it therefore logically serves as an appealing target for vaccine lead validation.²²¹ Previous carbohydrate vaccine investigations focused on Man₉ without success.²²² In 2022, Ye and co-workers synthesized the natural Man₅ (**92**), an S-linked analogue (**93**), a monofluorinated analogue (**94**), and a trifluorinated analogue (**95**) (Fig. 29).²²³ These modifications were with the aim of increasing immunogenicity whilst enabling cross reactivity of the antibodies with native Man₅. The glycans were conjugated to CRM₁₉₇ and used to vaccinate mice with ODN1826 as an adjuvant (Fig. 12, bottom). **92-CRM197** did not lead to production of Man₅-specific antibodies. The results of **93-CRM197**, **94-CRM197** and **95-CRM197** were identical in that just 1 of the 6 mice produced significant amounts of anti-Man₅ antibodies. The antisera




Fig. 28 CRM₁₉₇ conjugated fluorinated analogues of KH-1 by Ye and co-workers.²²⁰




Fig. 29 Natural Man₅, an S-linked analogue and fluorinated analogues by Ye and co-workers.²²³



bound 92-CRM₁₉₇ strongly, indicating that the antibodies were directed mainly towards CRM₁₉₇.

4.2.10. Summary. Fluorinated carbohydrate-based therapeutic cancer vaccines are part of an emerging field in immunological studies. A wide range of TACAs have been synthetically modified: Tn, STn, TF, sialyl-TF, MUC1, GM3, Globo H, KH-1, and a pentamannose epitope. The amide moiety of these molecules has been systematically modified with fluoroacetyl groups with different fluorination patterns by Ye and co-workers and this has now matured into a valuable design strategy. Studies have revealed that the effect of modification is highly target-dependable. The concept has been demonstrated in the MUC1 glycopeptide by Hoffmann-Röder and colleagues. Reproducible synthetic approaches to modified MUC1 have been validated and this lowers the barrier to immunological investigations. Glycopeptides with varying sites of fluorination have been tested in the context of both antibody formation and recognition. Comparatively, fluorine incorporation into carbohydrate antigens affected immunogenicity much more than in the case of glycopeptide (MUC1). This may be a consequence of the less pronounced changes caused by fluorine atoms in the interior of the glycopeptide. Wong and co-workers' work on Globo H analogues serves to demonstrate the importance of the modification site choice in the initial design of the vaccine. Overall, strategic incorporation of fluorine atoms into antigen structures continues to enrich the goal of immunogenicity modulation and induction of potent cross-reactive antibodies in certain cases.

5. Conclusions and future directions

Carbohydrate-based vaccines continue to play a foundational role in public health strategies across the globe. Pathogens that are familiar to us all include *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Neisseria meningitidis*, which are omnipresent and continually trying to evade detection by our immune system. There is a risk of taking vaccines for granted and it is hoped that this short Perspective serves to highlight the challenges and opportunities that exist in this important area of contemporary research. Carbohydrates are a crucially important structural sub-set of vaccine design modules, and their clinical success is grounded in the diversity and complexity that they afford. The natural role of glycans in modulating molecular and cellular recognition events renders these biomolecules ideally suited to the task of triggering the immune system, and moving to unnatural analogues offers the opportunity to further enhance immunogenicity and fine-tune their physicochemical profiles. Broadening the spectrum of available antigens may reveal new correlations and causations, thereby expediting rational design in the future. Although the validation of a safe and fully functional vaccine is incredibly complex, looking at epitope design through the lens of fluorination offers a new and exciting perspective. What is clear is that the introduction of fluorine into carbohydrate-based vaccine leads can lead to discernible patterns that enable structure–activity relationships to be extrapolated.

Author contributions

The manuscript was written with contributions from all authors.

Conflicts of interest

There are no conflicts to declare.

List of abbreviations

CPS	Capsular polysaccharide
TACA	Tumor-associated carbohydrate antigen
APC	Antigen-presenting cell
DC	Dendritic cell
MHC	Major histocompatibility complex
Th	T helper
BCR	B cell receptor
CRM ₁₉₇	Diphtheria toxin mutant
TT	Tetanus toxoid
DT	Diphtheria toxoid
OMPC	Meningococcal outer membrane protein complex
rEPA	<i>Pseudomonas aeruginosa</i> exotoxin A
BSA	Bovine serum albumin
KLH	Keyhole limpet hemocyanin
Hib	<i>Haemophilus influenzae</i> type b
AGA	Automated glycan assembly
LOS	Lipooligosaccharide
PorA	Porin A
Alum	Aluminium hydroxide adjuvant
ST	Serotype
ELISA	Enzyme-linked immunosorbent assay
CDC	Complement dependent cytotoxicity
ADCC	Antibody dependent cellular cytotoxicity
MUC1	Mucin 1
CFA	Complete Freund's Adjuvant
IFA	Incomplete Freund's Adjuvant
ODN	Oligonucleotide
Tn	Thomsen-nouveau
STn	Sialyl-Thomsen-nouveau
TF	Thomsen–Friedenreich
sialyl-TF	Sialyl-Thomsen–Friedenreich
PSA	Polysialic acid

Data availability

The data underlying this study are available in the published article.

Acknowledgements

We acknowledge the University of Münster for generous financial support.



Notes and references

- 1 J. E. Salk, P. L. Bazeley, B. L. Bennett, U. Krech, L. J. Lewis, E. N. Ward and J. S. Youngner, *Am. J. Public Health Nation's Health*, 1954, **44**, 994–1009.
- 2 J. E. Salk, *Am. J. Public Health Nation's Health*, 1955, **45**, 285–297.
- 3 J. E. Salk, *J. Am. Med. Assoc.*, 1955, **158**, 1239–1248.
- 4 D. A. Montero, R. M. Vidal, J. Velasco, L. J. Carreño, J. P. Torres, M. A. Benachi O., Y.-Y. Tovar-Rosero, A. A. Oñate and M. O'Ryan, *Front. Public Health*, 2024, **11**, 1326154.
- 5 P. Bonanni, *Vaccine*, 1999, **17**, S120–S125.
- 6 A. J. Shattock, H. C. Johnson, S. Y. Sim, A. Carter, P. Lambach, R. C. W. Hutubessy, K. M. Thompson, K. Badizadegan, B. Lambert, M. J. Ferrari, M. Jit, H. Fu, S. P. Silal, R. A. Hounsell, R. G. White, J. F. Mosser, K. A. M. Gaythorpe, C. L. Trotter, A. Lindstrand, K. L. O'Brien and N. Bar-Zeev, *Lancet*, 2024, **403**, 2307–2316.
- 7 H. Zhang, B. Patenaude, H. Zhang, M. Jit and H. Fang, *Bull. W. H. O.*, 2024, 276–287.
- 8 I. N. Weerarathna, E. S. Doelakeh, L. Kiwanuka, P. Kumar and S. Arora, *Mol. Biomed.*, 2024, **5**, 57.
- 9 S. A. Plotkin, *Nat. Med.*, 2005, **11**, S5–S11.
- 10 A. Hinman, *Annu. Rev. Public Health*, 1999, **20**, 211–229.
- 11 R. Rappuoli, S. Black and P. H. Lambert, *Lancet*, 2011, **378**, 360–368.
- 12 D. A. Kennedy and A. F. Read, *Proc. Natl. Acad. Sci. U. S. A.*, 2018, **115**, 12878–12886.
- 13 K. U. Jansen and A. S. Anderson, *Hum. Vaccines Immunother.*, 2018, **14**, 2142–2149.
- 14 R. Laxminarayan, I. Impalli, R. Rangarajan, J. Cohn, K. Ramjeet, B. W. Trainor, S. Strathdee, N. Sumpradit, D. Berman, H. Wertheim, K. Outterson, P. Srikantiah and U. Theuretzbacher, *Lancet*, 2024, **403**, 2534–2550.
- 15 P. Van Damme, *J. Infect. Dis.*, 2016, **214**, 1–3.
- 16 R. Brosch, S. V Gordon, T. Garnier, K. Eiglmeier, W. Frigui, P. Valenti, S. Dos Santos, S. Duthoy, C. Lacroix, C. Garcia-Pelayo, J. K. Inwald, P. Golby, J. N. Garcia, R. G. Hewinson, M. A. Behr, M. A. Quail, C. Churcher, B. G. Barrell, J. Parkhill and S. T. Cole, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 5596–5601.
- 17 R. W. Sutter, O. M. Kew, S. L. Cochi and R. B. Aylward, in *Plotkin's Vaccines (Seventh Edition)*, ed. S. A. Plotkin, W. A. Orenstein, P. A. Offit and K. M. Edwards, Elsevier, 7th edn, 2018, pp. 866–917.
- 18 R. Rappuoli and E. Malito, in *Corynebacterium diphtheriae and Related Toxigenic Species: Genomics, Pathogenicity and Applications*, ed. A. Burkovski, Springer Netherlands, Dordrecht, 2014, pp. 225–238.
- 19 S. A. Plotkin, *Clin. Infect. Dis.*, 2006, **43**, S164–S168.
- 20 A. H. Ellebedy and R. J. Webby, *Vaccine*, 2009, **27**, D65–D68.
- 21 S. A. Rella, Y. A. Kulikova, A. R. Minnegaliev and F. A. Kondrashov, *Evolution*, 2024, **78**, 1722–1738.
- 22 S. L. Takala and C. V Plowe, *Parasite Immunol.*, 2009, **31**, 560–573.
- 23 D. J. Smith, A. S. Lapedes, J. C. de Jong, T. M. Bestebroer, G. F. Rimmelzwaan, A. D. M. E. Osterhaus and R. A. M. Fouchier, *Science*, 2004, **305**, 371–376.
- 24 S. Cele, L. Jackson, D. S. Khoury, K. Khan, T. Moyo-Gwete, H. Tegally, J. E. San, D. Cromer, C. Scheepers, D. G. Amoako, F. Karim, M. Bernstein, G. Lustig, D. Archary, M. Smith, Y. Ganga, Z. Jule, K. Reedoy, S.-H. Hwa, J. Giandhari, J. M. Blackburn, B. I. Gosnell, S. S. Abdool Karim, W. Hanekom, M.-A. Davies, M. Hsiao, D. Martin, K. Mlisana, C. K. Wibmer, C. Williamson, D. York, R. Harrichandparsad, K. Herbst, P. Jeena, T. Khoza, H. Kløverpris, A. Leslie, R. Madansein, N. Magula, N. Manickchund, M. Marakalala, M. Mazibuko, M. Moshabela, N. Mthabela, K. Naidoo, Z. Ndhlovu, T. Ndung'u, N. Ngcobo, K. Nyamande, V. Patel, T. Smit, A. Steyn, E. Wong, A. von Gottberg, J. N. Bhiman, R. J. Lessells, M.-Y. S. Moosa, M. P. Davenport, T. de Oliveira, P. L. Moore, A. Sigal, NGS-SA and C.-K. Team, *Nature*, 2022, **602**, 654–656.
- 25 L. A. Hicks, L. H. Harrison, B. Flannery, J. L. Hadler, W. Schaffner, A. S. Craig, D. Jackson, A. Thomas, B. Beall, R. Lynfield, A. Reingold, M. M. Farley and C. G. Whitney, *J. Infect. Dis.*, 2007, **196**, 1346–1354.
- 26 F. Ganaie, S. Jamil, L. McGee, J. van T. Andries, S. Bentley, S. Lo, R. Gladstone, P. Turner, J. Keenan, R. Breiman and M. Nahm, *mBio*, 2020, **11**, e00937–20.
- 27 S. Gandon, M. J. Mackinnon, S. Nee and A. F. Read, *Nature*, 2001, **414**, 751–756.
- 28 M. G. Milgroom, in *Biology of Infectious Disease: From Molecules to Ecosystems*, ed. M. G. Milgroom, Springer International Publishing, Cham, 2023, pp. 175–192.
- 29 R. Strugnelli, F. Zepp, A. Cunningham and T. Tantawichien, *Perspect. Vaccinol.*, 2011, **1**, 61–88.
- 30 F. Zepp, *Vaccine*, 2010, **28**, C14–C24.
- 31 J. A. Berzofsky, S. B. Howe and P. B. Olkhanud, in *Encyclopedia of Infection and Immunity*, ed. N. Rezaei, Elsevier, Oxford, 2022, pp. 76–89.
- 32 A. Sharma, S. Gautam and N. Bandyopadhyay, in *Encyclopedia of Food Microbiology (Second Edition)*, ed. C. A. Batt and M. Lou Tortorello, Academic Press, Oxford, 2014, pp. 680–687.
- 33 A. Varki, R. D. Cummings, J. D. Esko, P. Stanley, G. W. Hart, M. Aebi, A. G. Darvill, T. Kinoshita, N. H. Packer, J. H. Prestegard and others, *Essentials of Glycobiology*, Cold spring harbor laboratory Press, 4th edition, 2015.
- 34 A. Varki, *Glycobiology*, 2017, **27**, 3–49.
- 35 K. J. Ryan, G. C. Ray and J. C. Sherris, *Sherris Medical Microbiology: An Introduction to Infectious Diseases*, McGraw-Hill Education, 4th edn, 2004.
- 36 J. M. Daubenspeck, D. S. Jordan and K. Dybvig, in *Mollicutes: Molecular Biology and Pathogenesis*, ed. G. F. Browning and C. Citti, Caister Academic Press, 1st edn, 2014, pp. 131–147.
- 37 W. Vollmer, D. Blanot and M. A. De Pedro, *FEMS Microbiol. Rev.*, 2008, **32**, 149–167.
- 38 B. Zhang, Y. Gao, L. Zhang and Y. Zhou, *J. Integr. Plant Biol.*, 2021, **63**, 251–272.



- 39 J. Chen, J. Gao, M. Zhang, M. Cai, H. Xu, J. Jiang, Z. Tian and H. Wang, *Sci. Rep.*, 2016, **6**, 30247.
- 40 F. Di Lorenzo, K. A. Duda, R. Lanzetta, A. Silipo, C. De Castro and A. Molinaro, *Chem. Rev.*, 2022, **122**, 15767–15821.
- 41 M. Christodoulides and J. Heckels, *Pathog. Dis.*, 2017, **75**, ftx033.
- 42 J. A. Jaurigue and P. H. Seeberger, *Front. Cell. Infect. Microbiol.*, 2017, **7**, 248.
- 43 L. Del Bino, K. E. Østerlid, D.-Y. Wu, F. Nonne, M. R. Romano, J. Codée and R. Adamo, *Chem. Rev.*, 2022, **122**, 15672–15716.
- 44 C. Anish, B. Schumann, C. L. Pereira and P. H. Seeberger, *Chem. Biol.*, 2014, **21**, 38–50.
- 45 S. S. Pinho and C. A. Reis, *Nat. Rev. Cancer*, 2015, **15**, 540–555.
- 46 S. Mereiter, M. Balmaña, D. Campos, J. Gomes and C. A. Reis, *Cancer Cell*, 2019, **36**, 6–16.
- 47 F. Yang, X.-J. Zheng, C.-X. Huo, Y. Wang, Y. Zhang and X.-S. Ye, *ACS Chem. Biol.*, 2011, **6**, 252–259.
- 48 S. J. Danishefsky and J. R. Allen, *Angew. Chem., Int. Ed.*, 2000, **39**, 836–863.
- 49 J. Zhu, J. D. Warren and S. J. Danishefsky, *Expert Rev. Vaccines*, 2009, **8**, 1399–1413.
- 50 M.-M. Wei, Y.-S. Wang and X.-S. Ye, *Med. Res. Rev.*, 2018, **38**, 1003–1026.
- 51 K. Furukawa, Y. Ohmi, O. Tajima, Y. Ohkawa, Y. Kondo, J. Shuting, N. Hashimoto and K. Furukawa, in *Progress in Molecular Biology and Translational Science*, ed. R. L. Schnaar and P. H. H. Lopez, Academic Press, 2018, vol. 156, pp. 265–287.
- 52 K. Kappler and T. Hennet, *Genes Immun.*, 2020, **21**, 224–239.
- 53 R. Misasi, S. Dionisi, L. Farilla, B. Carabba, L. Lenti, U. Di Mario and F. Dotta, *Diabetes/Metab. Rev.*, 1997, **13**, 163–179.
- 54 C.-H. Kim, in *Glycobiology of Innate Immunology*, ed. C.-H. Kim, Springer Singapore, Singapore, 2022, pp. 115–198.
- 55 J. A. Goodfellow and H. J. Willison, in *Progress in Molecular Biology and Translational Science*, ed. R. L. Schnaar and P. H. H. Lopez, Academic Press, 2018, vol. 156, pp. 355–382.
- 56 R. D. Astronomo and D. R. Burton, *Nat. Rev. Drug Discovery*, 2010, **9**, 308–324.
- 57 L. Sun, D. R. Middleton, P. L. Wantuch, A. Ozdilek and F. Y. Avci, *Glycobiology*, 2016, **26**, 1029–1040.
- 58 Y. Narimatsu, H. J. Joshi, R. Nason, J. Van Coillie, R. Karlsson, L. Sun, Z. Ye, Y.-H. Chen, K. T. Schjoldager, C. Steentoft, S. Furukawa, B. A. Bensing, P. M. Sullam, A. J. Thompson, J. C. Paulson, C. Büll, G. J. Adema, U. Mandel, L. Hansen, E. P. Bennett, A. Varki, S. Y. Vakhrushev, Z. Yang and H. Clausen, *Mol. Cell*, 2019, **75**, 394–407.
- 59 N. Banahene, H. W. Kavunja and B. M. Swarts, *Chem. Rev.*, 2022, **122**, 3336–3413.
- 60 H.-K. Guttormsen, L. C. Paoletti, K. G. Mansfield, W. Jachymek, H. J. Jennings and D. L. Kasper, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 5903–5908.
- 61 K. Kappler and T. Hennet, *Genes Immun.*, 2020, **21**, 224–239.
- 62 G. Ragupathi, D. M. Coltart, L. J. Williams, F. Koide, E. Kagan, J. Allen, C. Harris, P. W. Glunz, P. O. Livingston and S. J. Danishefsky, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 13699–13704.
- 63 R. Roy, *Drug Discovery Today:Technol.*, 2004, **1**, 327–336.
- 64 J. Hütter and B. Lepenies, in *Carbohydrate-Based Vaccines: Methods and Protocols*, ed. B. Lepenies, Springer New York, New York, NY, 2015, pp. 1–10.
- 65 M. Kaczmarek, J. Poznańska, F. Fechner, N. Michalska, S. Paszkowska, A. Napierała and A. Mackiewicz, *Cells*, 2023, **12**, DOI: [10.3390/cells12172159](https://doi.org/10.3390/cells12172159).
- 66 G. A. Lemieux and C. R. Bertozzi, *Chem. Biol.*, 2001, **8**, 265–275.
- 67 D. H. Dube and C. R. Bertozzi, *Curr. Opin. Chem. Biol.*, 2003, **7**, 616–625.
- 68 R. Mettu, C.-Y. Chen and C.-Y. Wu, *J. Biomed. Sci.*, 2020, **27**, 9.
- 69 A. H. Lucas, M. A. Apicella and C. E. Taylor, *Clin. Infect. Dis.*, 2005, **41**, 705–712.
- 70 B. Linclau, A. Ardá, N.-C. Reichardt, M. Sollogoub, L. Unione, S. P. Vincent and J. Jiménez-Barbero, *Chem. Soc. Rev.*, 2020, **49**, 3863–3888.
- 71 S. Leusmann, P. Ménová, E. Shanin, A. Titz and C. Rademacher, *Chem. Soc. Rev.*, 2023, **52**, 3663–3740.
- 72 D. O'Hagan, *Chem. Soc. Rev.*, 2008, **37**, 308–319.
- 73 I. J. Amanna and M. K. Slifka, *Virology*, 2011, **411**, 206–215.
- 74 S. A. Plotkin, *Clin. Vaccine Immunol.*, 2010, **17**, 1055–1065.
- 75 C.-C. Liu and X.-S. Ye, *Glycoconjugate J.*, 2012, **29**, 259–271.
- 76 D. J. Irvine, M. A. Swartz and G. L. Szeto, *Nat. Mater.*, 2013, **12**, 978–990.
- 77 T. J. Moyer, A. C. Zmolek and D. J. Irvine, *J. Clin. Invest.*, 2016, **126**, 799–808.
- 78 P. Mistry, F. Barmania, J. Mellet, K. Peta, A. Strydom, I. M. Viljoen, W. James, S. Gordon and M. S. Pepper, *Front. Immunol.*, 2022, **12**, 809244.
- 79 G. Wang, Y. Wang and F. Ma, *Cell Biosci.*, 2024, **14**, 24.
- 80 P. Costantino, R. Rappuoli and F. Berti, *Expert Opin. Drug Discovery*, 2011, **6**, 1045–1066.
- 81 R. Rappuoli, *Sci. Transl. Med.*, 2018, **10**, eaat4615.
- 82 F. Y. Avci, X. Li, M. Tsuji and D. L. Kasper, *Nat. Med.*, 2011, **17**, 1602–1609.
- 83 B. A. Keyt, R. Baliga, A. M. Sinclair, S. F. Carroll and M. S. Peterson, *Antibodies*, 2020, **9**, DOI: [10.3390/antib9040053](https://doi.org/10.3390/antib9040053).
- 84 S. Gong and R. M. Ruprecht, *Front. Immunol.*, 2020, **11**, 1943.
- 85 J. K. Actor, in *Introductory Immunology*, ed. J. K. Actor, Academic Press, Amsterdam, 2014, pp. 28–41.
- 86 D. K. Flaherty, in *Immunology for Pharmacy*, ed. D. K. Flaherty, Mosby, Saint Louis, 2012, pp. 70–78.
- 87 G. Vidarsson, G. Dekkers and T. Rispens, *Front. Immunol.*, 2014, **5**, 520.
- 88 J. K. Actor, in *Introductory Immunology*, ed. J. K. Actor, Academic Press, 2nd edn, 2019, pp. 111–121.
- 89 I. L. Baine and C. A. Tormey, in *Immunologic Concepts in Transfusion Medicine*, ed. R. W. Maitta, Elsevier, 2020, pp. 1–14.



- 90 I. Vojtek, P. Buchy, T. M. Doherty and B. Hoet, *Vaccine*, 2019, **37**, 539–549.
- 91 J. K. Actor, in *Elsevier's Integrated Review Immunology and Microbiology (Second Edition)*, ed. J. K. Actor, W.B. Saunders, Philadelphia, 2nd edn., 2012, pp. 17–24.
- 92 B. Schumann, C. Anish, C. L. Pereira and P. H. Seeberger, in *Biotherapeutics: Recent Developments using Chemical and Molecular Biology*, eds. L. H. Jones and A. J. McKnight, The Royal Society of Chemistry, 2013, pp. 68–104.
- 93 M. Heidelberger and O. T. Avery, *J. Exp. Med.*, 1923, **38**, 73–79.
- 94 M. Heidelberger, M. M. Dilapi, M. Siegel and A. W. Walter, *J. Immunol.*, 1950, **65**, 535–541.
- 95 C. M. MacLeod, R. G. Hodges, M. Heidelberger and W. G. Bernhard, *J. Exp. Med.*, 1945, **82**, 445–465.
- 96 W. S. Tillett and T. Francis Jr, *J. Exp. Med.*, 1929, **50**, 687–701.
- 97 J. F. G. Vliegthart, *FEBS Lett.*, 2006, **580**, 2945–2950.
- 98 F. Berti and F. Micoli, *Curr. Opin. Immunol.*, 2020, **65**, 42–49.
- 99 M. Baker, H. M. Reynolds, B. Lumicisi and C. J. Bryson, *Self. Nonself*, 2010, **1**, 314–322.
- 100 R. Adamo, A. Nilo, B. Castagner, O. Boutureira, F. Berti and G. J. L. Bernardes, *Chem. Sci.*, 2013, **4**, 2995–3008.
- 101 L. P. Deimel, X. Xue and Q. J. Sattentau, *Trends Microbiol.*, 2022, **30**, 866–881.
- 102 C. E. Martina, J. E. Crowe and J. Meiler, *Front. Immunol.*, 2023, **14**, 1126034.
- 103 D. Goldblatt, *Clin. Exp. Immunol.*, 2000, **119**, 1–3.
- 104 O. T. Avery and W. F. Goebel, *J. Exp. Med.*, 1931, **54**, 437–447.
- 105 J. D. Grabenstein and K. P. Klugman, *Clin. Microbiol. Infect.*, 2012, **18**, 15–24.
- 106 V. Verez-Bencomo, V. Fernández-Santana, E. Hardy, M. E. Toledo, M. C. Rodríguez, L. Heynngnezz, A. Rodríguez, A. Baly, L. Herrera, M. Izquierdo, A. Villar, Y. Valdés, K. Cosme, M. L. Deler, M. Montane, E. Garcia, A. Ramos, A. Aguilar, E. Medina, G. Toraño, I. Sosa, I. Hernandez, R. Martínez, A. Muzachio, A. Carmenates, L. Costa, F. Cardoso, C. Campa, M. Diaz and R. Roy, *Science*, 2004, **305**, 522–525.
- 107 S. Nishat and P. R. Andreana, *Vaccines*, 2016, **4**, 19.
- 108 P. J. Carter and V. Quarmby, *Nat. Rev. Drug Discovery*, 2024, **23**, 898–913.
- 109 A. Finn, *Br. Med. Bull.*, 2004, **70**, 1–14.
- 110 E. Juhani, K. Helena, T. A. K, P. Heikki, R. Pirjo-Riitta, K. Eija, P. Eeva, M. P. H and M. P. Helena, *N. Engl. J. Med.*, 1990, **323**, 1381–1387.
- 111 H. Clayton, B. Robert, O. Walter, K. Eun-Jeong, G. H. E, O. M. T, H. Jon, L. Joy, D. M. J and K. Nicholas, *Clin. Vaccine Immunol.*, 2010, **17**, 1868–1874.
- 112 I. Jacques, V. Olivier-Bernardin and G. Dubray, *Vaccine*, 1991, **9**, 896–900.
- 113 N. K. Ibrahim and J. L. Murray, *Clin. Breast Cancer*, 2003, **3**, S139–S143.
- 114 P. H. Seeberger, *Chem. Rev.*, 2021, **121**, 3598–3626.
- 115 M. E. Zaslona, A. M. Downey, P. H. Seeberger and O. Moscovitz, *Biochem. Soc. Trans.*, 2021, **49**, 2411–2429.
- 116 Y. Chen and K. Cheng, *Nano Res.*, 2020, **13**, 2617–2624.
- 117 D. T. O'Hagan, R. N. Lodaya and G. Lofano, *Semin. Immunol.*, 2020, **50**, 101426.
- 118 N. Petrovsky and J. C. Aguilar, *Immunol. Cell Biol.*, 2004, **82**, 488–496.
- 119 F. Peri, *Chem. Soc. Rev.*, 2013, **42**, 4543–4556.
- 120 M. Shirley, *Drugs*, 2022, **82**, 989–999.
- 121 S. Djalali, N. Yadav and M. Delbianco, *Nat. Rev. Mater.*, 2024, **9**, 190–201.
- 122 O. J. Plante, E. R. Palmacci and P. H. Seeberger, *Science*, 2001, **291**, 1523–1527.
- 123 J. Suri and R. Gilmour, *Angew. Chem., Int. Ed.*, 2025, **64**, e202422766.
- 124 M. Guberman and P. H. Seeberger, *J. Am. Chem. Soc.*, 2019, **141**, 5581–5592.
- 125 C. J. Crawford and P. H. Seeberger, *Chem. Soc. Rev.*, 2023, **52**, 7773–7801.
- 126 M. Panza, S. G. Pistorio, K. J. Stine and A. V Demchenko, *Chem. Rev.*, 2018, **118**, 8105–8150.
- 127 A. Pardo-Vargas, M. Delbianco and P. H. Seeberger, *Curr. Opin. Chem. Biol.*, 2018, **46**, 48–55.
- 128 C.-H. Hsu, S.-C. Hung, C.-Y. Wu and C.-H. Wong, *Angew. Chem., Int. Ed.*, 2011, **50**, 11872–11923.
- 129 H. S. Hahm, M. K. Schlegel, M. Hurevich, S. Eller, F. Schuhmacher, J. Hofmann, K. Pagel and P. H. Seeberger, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, E3385–E3389.
- 130 H. J. Jennings, R. Roy and A. Gamian, *J. Immunol.*, 1986, **137**, 1708–1713.
- 131 L. M. Krug, G. Ragupathi, C. Hood, C. George, F. Hong, R. Shen, L. Abrey, H. J. Jennings, M. G. Kris and P. O. Livingston, *Cancer Immunol., Immunother.*, 2012, **61**, 9–18.
- 132 G. Ragupathi, P. O. Livingston, C. Hood, J. Gathuru, S. E. Krown, P. B. Chapman, J. D. Wolchok, L. J. Williams, R. C. Oldfield and W.-J. Hwu, *Clin. Cancer Res.*, 2003, **9**, 5214–5220.
- 133 X. Wang, M. Gomozkova, S. Li, B. Banerjee, G. Zong, D. Missiakas and L.-X. Wang, *Proc. Natl. Acad. Sci. U. S. A.*, 2026, **123**, e2530653123.
- 134 H.-J. Lo, L. Krasnova, S. Dey, T. Cheng, H. Liu, T.-I. Tsai, K. B. Wu, C.-Y. Wu and C.-H. Wong, *J. Am. Chem. Soc.*, 2019, **141**, 6484–6488.
- 135 A. Axer, R. P. Jumde, S. Adam, A. Faust, M. Schäfers, M. Fobker, J. Koehnke, A. K. H. Hirsch and R. Gilmour, *Chem. Sci.*, 2021, **12**, 1286–1294.
- 136 J. A. Delbrouck, L. P. Chêne and S. P. Vincent, in *Fluorine in Life Sciences: Pharmaceuticals, Medicinal Diagnostics, and Agrochemicals*, ed. G. Haufe and F. R. Leroux, Academic Press, 2019, pp. 241–279.
- 137 J. M. Keil, G. R. Rafn, I. M. Turan, M. A. Aljohani, R. Sahebjam-Atabaki and X.-L. Sun, *J. Med. Chem.*, 2022, **65**, 13574–13593.
- 138 C.-H. Wong and H.-J. Lo, WO2020/205034A1, 2020.
- 139 S. G. Withers, I. P. Street, P. Bird and D. H. Dolphin, *J. Am. Chem. Soc.*, 1987, **109**, 7530–7531.



- 140 K. Siebold, E. Chikunova, N. Lorz, C. Jordan, A. D. Gossert and R. Gilmour, *Angew. Chem., Int. Ed.*, 2025, **64**, e202423782.
- 141 S.-A. Zhai, Z. Ai, Q. Zhang, Q. Jiao, S. Wang and C. Xu, *Org. Lett.*, 2026, **28**, 995–1001.
- 142 C. Bucher and R. Gilmour, *Angew. Chem., Int. Ed.*, 2010, **49**, 8724–8728.
- 143 C. Jordan, T. Hayashi, A. Löbber, J. Fan, C. S. Teschers, K. Siebold, M. Auferio, F. Pape, E. Campbell, A. Axer, K. Bussmann, K. Bergander, J. Köhnke, A. D. Gossert and R. Gilmour, *ACS Cent. Sci.*, 2024, **10**, 1481–1489.
- 144 N. Aiguabella, M. C. Holland and R. Gilmour, *Org. Biomol. Chem.*, 2016, **14**, 5534–5538.
- 145 M. L. Uhrig, B. Lantaño and A. Postigo, *Org. Biomol. Chem.*, 2019, **17**, 5173–5189.
- 146 J. P. Dolan, S. T. Evans, C. M. M. Benckendorff, S. Sari, A. Ní Cheallaigh and G. J. Miller, *JACS Au*, 2025, **5**, 3994–4001.
- 147 K. Dax, M. Albert, J. Ortner and B. J. Paul, *Carbohydr. Res.*, 2000, **327**, 47–86.
- 148 B. Linclau, A. Ardá, N.-C. Reichardt, M. Sollogoub, L. Unione, S. P. Vincent and J. Jiménez-Barbero, *Chem. Soc. Rev.*, 2020, **49**, 3863–3888.
- 149 C. E. Council, K. J. Kilpin, J. S. Gusthart, S. A. Allman, B. Linclau and S. S. Lee, *Org. Biomol. Chem.*, 2020, **18**, 3423–3451.
- 150 X. Wei, P. Wang, F. Liu, X. Ye and D. Xiong, *Molecules*, 2023, **28**, 6641.
- 151 R. Hevey, *Chem.–Eur. J.*, 2021, **27**, 2240–2253.
- 152 J. P. Dolan, C. M. M. Benckendorff, R. A. Field and G. J. Miller, *Future Med. Chem.*, 2023, **15**, 1111–1114.
- 153 C.-S. Tsai, H.-Y. Yen, M.-I. Lin, T.-I. Tsai, S.-Y. Wang, W.-I. Huang, T.-L. Hsu, Y.-S. E. Cheng, J.-M. Fang and C.-H. Wong, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 2466–2471.
- 154 S. J. Moons, E. Rossing, J. J. A. Heming, M. A. C. H. Janssen, M. van Scherpenzeel, D. J. Lefeber, M. I. de Jonge, J. D. Langereis and T. J. Boltje, *Bioconjugate Chem.*, 2021, **32**, 1047–1051.
- 155 E. Campbell, C. Jordan and R. Gilmour, *Chem. Soc. Rev.*, 2023, **52**, 3599–3626.
- 156 G. Fittolani, E. Shanina, M. Guberman, P. H. Seeberger, C. Rademacher and M. Delbianco, *Angew. Chem., Int. Ed.*, 2021, **60**, 13302–13309.
- 157 C. S. Teschers and R. Gilmour, *Angew. Chem., Int. Ed.*, 2023, **62**, e202213304.
- 158 J. Suri, C. Jordan, C. S. Teschers, K. Schlangen, S. H. Rüdiger, A. D. Gossert and R. Gilmour, *Angew. Chem., Int. Ed.*, 2026, **65**, e8014647.
- 159 P. H. Seeberger, *Chem. Rev.*, 2021, **121**, 3598–3626.
- 160 G. Stefanetti, F. Borriello, B. Richichi, I. Zanoni and L. Lay, *Front. Cell. Infect. Microbiol.*, 2022, **11**, 808005.
- 161 D. S. Stephens, B. Greenwood and P. Brandtzaeg, *Lancet*, 2007, **369**, 2196–2210.
- 162 O. Harrison, H. Claus, Y. Jiang, J. Bennett, H. Bratcher, K. Jolley, C. Corton, R. Care, J. Poolman, W. Zollinger, C. Frasc, D. Stephens, I. Feavers, M. Frosch, J. Parkhill, U. Vogel, M. Quail, S. Bentley and M. C. J. Maiden, *Emerg. Infect. Dis.*, 2013, **19**, 566.
- 163 R. D. McCoy, E. R. Vimr and F. A. Troy, *J. Biol. Chem.*, 1985, **260**, 12695–12699.
- 164 G. Liao, Z. Zhou and Z. Guo, *Chem. Commun.*, 2015, **51**, 9647–9650.
- 165 G. Liao, Z. Zhou, S. Suryawanshi, M. A. Mondal and Z. Guo, *ACS Cent. Sci.*, 2016, **2**, 210–218.
- 166 C. Jordan, K. Siebold, P. Priegue, P. H. Seeberger and R. Gilmour, *J. Am. Chem. Soc.*, 2024, **146**, 15366–15375.
- 167 T. Hayashi, G. Kehr, K. Bergander and R. Gilmour, *Angew. Chem., Int. Ed.*, 2019, **58**, 3814–3818.
- 168 T. Hayashi, A. Axer, G. Kehr, K. Bergander and R. Gilmour, *Chem. Sci.*, 2020, **11**, 6527–6531.
- 169 F. Chappuis, S. Sundar, A. Hailu, H. Ghalib, S. Rijal, R. W. Peeling, J. Alvar and M. Boelaert, *Nat. Rev. Microbiol.*, 2007, **5**, 873–882.
- 170 A. Baumann, S. Marchner, M. Daum and A. Hoffmann-Röder, *Eur. J. Org. Chem.*, 2018, **2018**, 3803–3815.
- 171 F. A. Ganaie, B. W. Beall, J. Yu, M. van der Linden, L. McGee, C. Satzke, S. Manna, S. W. Lo, S. D. Bentley, N. Ravenscroft and M. H. Nahm, *Clin. Microbiol. Rev.*, 2025, **38**, e00175–24.
- 172 C. Pérez-García, A. González-Díaz, M. Domenech, M. Llamós, A. Úbeda, J. C. Sanz, E. García, C. Ardanuy, J. Sempere and J. Yuste, *Emerging Microbes Infect.*, 2025, **14**, 2521845.
- 173 B. Schumann, H. S. Hahm, S. G. Parameswarappa, K. Reppe, A. Wahlbrink, S. Govindan, P. Kaplonek, L. Pirofski, M. Witzernath, C. Anish, C. L. Pereira and P. H. Seeberger, *Sci. Transl. Med.*, 2017, **9**, eaaf5347.
- 174 D. Gast, S. Neidig, M. Reindl and A. Hoffmann-Röder, *Int. J. Mol. Sci.*, 2025, **26**, 1535.
- 175 M. Reindl, A. Baumann and A. Hoffmann-Röder, *ACS Org. Inorg. Au*, 2026, **6**, 161–168.
- 176 P. Kaplonek, N. Khan, K. Reppe, B. Schumann, M. Emmadi, M. P. Lisboa, F.-F. Xu, A. D. J. Calow, S. G. Parameswarappa, M. Witzernath, C. L. Pereira and P. H. Seeberger, *Proc. Natl. Acad. Sci. U. S. A.*, 2018, **115**, 13353–13358.
- 177 J. Zhu, Q. Wan, D. Lee, G. Yang, M. K. Spassova, O. Ouerfelli, G. Ragupathi, P. Damani, P. O. Livingston and S. J. Danishefsky, *J. Am. Chem. Soc.*, 2009, **131**, 9298–9303.
- 178 S. J. Danishefsky, Y.-K. Shue, M. N. Chang and C.-H. Wong, *Acc. Chem. Res.*, 2015, **48**, 643–652.
- 179 D. W. Kufe, *Nat. Rev. Cancer*, 2009, **9**, 874–885.
- 180 S. Ingale, M. A. Wolfert, J. Gaekwad, T. Buskas and G.-J. Boons, *Nat. Chem. Biol.*, 2007, **3**, 663–667.
- 181 R. Abrantes, C. Forcados, D. J. Warren, L. Santos-Ferreira, K. G. Fleten, E. Senra, A. F. Costa, K. Krpina, R. Henrique, A. M. Liberg, P. Rawat, P. Gelebart, E. McCormack, L. Bjørge, B. Davidson, V. Greiff, D. E. Costea, F. Pinto, K. Flatmark, C. Gomes, E. M. Inderberg, C. A. Reis and S. Wälchli, *Cell Rep. Med.*, 2025, **6**, 102350, DOI: [10.1016/j.xcrm.2025.102350](https://doi.org/10.1016/j.xcrm.2025.102350).



- 182 J. Heimbürg-Molinario, M. Lum, G. Vijay, M. Jain, A. Almogren and K. Rittenhouse-Olson, *Vaccine*, 2011, **29**, 8802–8826.
- 183 C. Fu, H. Zhao, Y. Wang, H. Cai, Y. Xiao, Y. Zeng and H. Chen, *HLA*, 2016, **88**, 275–286.
- 184 Z. Guo and Q. Wang, *Curr. Opin. Chem. Biol.*, 2009, **13**, 608–617.
- 185 T. Ju, V. I. Otto and R. D. Cummings, *Angew. Chem., Int. Ed.*, 2011, **50**, 1770–1791.
- 186 G. F. Springer, *J. Mol. Med.*, 1997, **75**, 594–602.
- 187 C. Song, S. Sun, C.-X. Huo, Q. Li, X.-J. Zheng, G. Tai, Y. Zhou and X.-S. Ye, *Bioorg. Med. Chem.*, 2016, **24**, 915–920.
- 188 J. Munkley, *Int. J. Mol. Sci.*, 2016, **17**, 275.
- 189 L. A. Holmberg and B. M. Sandmaier, *Expert Opin. Biol. Ther.*, 2001, **1**, 881–891.
- 190 G. D. MacLean, M. A. Reddish, R. R. Koganty and B. M. Longenecker, *J. Immunother.*, 1996, **19**, 59–68.
- 191 D. Miles, H. Roché, M. Martin, T. J. Perren, D. A. Cameron, J. Glaspy, D. Dodwell, J. Parker, J. Mayordomo, A. Tres, J. L. Murray, N. K. Ibrahim and the T. S. Group, *Oncologist*, 2011, **16**, 1092–1100.
- 192 S. B. Zeichner, *J. Am. Osteopathic Assoc.*, 2012, **112**, 482–483.
- 193 D. T. Le and E. M. Jaffee, *Cancer Res.*, 2012, **72**, 3439–3444.
- 194 C. Song, X.-J. Zheng, H. Guo, Y. Cao, F. Zhang, Q. Li, X.-S. Ye and Y. Zhou, *Glycoconjugate J.*, 2019, **36**, 399–408.
- 195 C. Song, X.-J. Zheng, C.-C. Liu, Y. Zhou and X.-S. Ye, *Oncotarget*, 2017, **8**, 47330–47343.
- 196 E. Malito, B. Bursulaya, C. Chen, P. Lo Surdo, M. Picchianti, E. Balducci, M. Biancucci, A. Brock, F. Berti, M. J. Bottomley, M. Nissum, P. Costantino, R. Rappuoli and G. Spraggon, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 5229–5234.
- 197 C.-X. Huo, X.-J. Zheng, A. Xiao, C.-C. Liu, S. Sun, Z. Lv and X.-S. Ye, *Org. Biomol. Chem.*, 2015, **13**, 3677–3690.
- 198 C. A. De Leon, P. M. Levine, T. W. Craven and M. R. Pratt, *Biochemistry*, 2017, **56**, 3507–3517.
- 199 D. R. Bundle, J. R. Rich, S. Jacques, H. N. Yu, M. Nitz and C.-C. Ling, *Angew. Chem., Int. Ed.*, 2005, **44**, 7725–7729.
- 200 J. Mo, Y. Zou, B.-H. Li, G. Li, X.-J. Zheng, Y. Liu and X.-S. Ye, *ACS Appl. Mater. Interfaces*, 2023, **15**, 40201–40212.
- 201 V. V. Glinsky, G. V. Glinsky, K. Rittenhouse-Olson, M. E. Huflejt, O. V. Glinskii, S. L. Deutscher and T. P. Quinn, *Cancer Res.*, 2001, **61**, 4851–4857.
- 202 A. Hoffmann-Röder, A. Kaiser, S. Wagner, N. Gaidzik, D. Kowalczyk, U. Westerlind, B. Gerlitzki, E. Schmitt and H. Kunz, *Angew. Chem., Int. Ed.*, 2010, **49**, 8498–8503.
- 203 M. Johannes, M. Reindl, B. Gerlitzki, E. Schmitt and A. Hoffmann-Röder, *Beilstein J. Org. Chem.*, 2015, **11**, 155–161.
- 204 S. Sun, X.-J. Zheng, C.-X. Huo, C. Song, Q. Li and X.-S. Ye, *ChemMedChem*, 2016, **11**, 1090–1096.
- 205 U. S. Grewal, R. Kurzrock and N. P. J. Preci, *Oncol.*, 2025, **9**, 218.
- 206 M. A. Hollingsworth and B. J. Swanson, *Nat. Rev. Cancer*, 2004, **4**, 45–60.
- 207 I. Radziejewska, *Biomed. Pharmacother.*, 2024, **174**, 116619.
- 208 C. W. Mersch Sarah and H.-R. Anja, *Synlett*, 2009, **2009**, 2167–2171.
- 209 S. Wagner, C. Mersch and A. Hoffmann-Röder, *Chem.–Eur. J.*, 2010, **16**, 7319–7330.
- 210 M. Johannes, T. Oberbillig and A. Hoffmann-Röder, *Org. Biomol. Chem.*, 2011, **9**, 5541–5546.
- 211 T. Oberbillig, C. Mersch, S. Wagner and A. Hoffmann-Röder, *Chem. Commun.*, 2012, **48**, 1487–1489.
- 212 A. Hoffmann-Röder and M. Johannes, *Chem. Commun.*, 2011, **47**, 9903–9905.
- 213 S. Sipione, J. Monyor, D. Galleguillos, N. Steinberg and V. Kadam, *Front. Neurosci.*, 2020, **14**, 572965, DOI: [10.3389/fnins.2020.572965](https://doi.org/10.3389/fnins.2020.572965).
- 214 D. Dorvignit, K. F. Boligan, E. Relova-Hernández, M. Clavell, A. López, M. Labrada, H.-U. Simon, A. López-Requena, C. Mesa and S. von Gunten, *Sci. Rep.*, 2019, **9**, 9921.
- 215 X.-J. Zheng, F. Yang, M. Zheng, C.-X. Huo, Y. Zhang and X.-S. Ye, *Org. Biomol. Chem.*, 2015, **13**, 6399–6406.
- 216 D. S. Sigal, D. J. Hermel, P. Hsu and T. Pearce, *Future Oncol.*, 2022, **18**, 117–134.
- 217 H.-Y. Lee, C.-Y. Chen, T.-I. Tsai, S.-T. Li, K.-H. Lin, Y.-Y. Cheng, C.-T. Ren, T.-J. R. Cheng, C.-Y. Wu and C.-H. Wong, *J. Am. Chem. Soc.*, 2014, **136**, 16844–16853.
- 218 C. Zhai, X.-J. Zheng, C. Song and X.-S. Ye, *RSC Med. Chem.*, 2021, **12**, 1239–1243.
- 219 Y.-L. Huang, J.-T. Hung, S. K. C. Cheung, H.-Y. Lee, K.-C. Chu, S.-T. Li, Y.-C. Lin, C.-T. Ren, T.-J. R. Cheng, T.-L. Hsu, A. L. Yu, C.-Y. Wu and C.-H. Wong, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 2517–2522.
- 220 Y. Liu, B. Li, X. Zheng, D. Xiong and X. Ye, *Molecules*, 2023, **28**, 1934, DOI: [10.3390/molecules28041934](https://doi.org/10.3390/molecules28041934).
- 221 C. A. Bewley, *Structure*, 2001, **9**, 931–940.
- 222 J. Ni, H. Song, Y. Wang, N. M. Stamatou and L.-X. Wang, *Bioconjugate Chem.*, 2006, **17**, 493–500.
- 223 C.-C. Liu, C.-X. Huo, C. Zhai, X.-J. Zheng, D.-C. Xiong and X.-S. Ye, *Bioconjugate Chem.*, 2022, **33**, 807–820.

