

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: N. M. Alcock, M. Huzan, T. Burrow, M. O.J.Y Hunault, C. Tamain, M. Autillo, T. Dumas and M. L. Baker, *Chem. Sci.*, 2026, DOI: 10.1039/D6SC02413K.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Journal Name

ARTICLE TYPE

Cite this: DOI: 00.0000/xxxxxxxxxx

Resolving the inner *versus* outer component of the 5*f* radial wavefunction in the early actinidesNathan M. Alcock,^{a,b,c} Myron S. Huzan,^{a,b,c} Timothy G. Burrow,^d Myrtille O.J.Y. Hunault,^d Christelle Tamain,^e Matthieu Autillo,^e Thomas Dumas,^e Michael L. Baker.^{a,b,c}Received Date
Accepted Date

DOI: 00.0000/xxxxxxxxxx

Actinide chemistry is largely defined by 5*f* orbital behaviour, but experimental measurements of their role in bonding are scarce, despite their importance for understanding chemical reactivity. Here, we demonstrate that M₄-edge 3*d*4*f* resonant inelastic X-ray scattering can discern the averaged 5*f* radial wavefunction, revealing how 5*f* electron density varies with distance from the nucleus. This approach enables differential quantification of how the inner and outer components of the 5*f* radial wavefunction respond to ligand bonding. Applied to a series of actinide hexachloride complexes, [AnCl₆]²⁻ (An = U, Np, Pu), it is shown that bonding-induced expansion of the 5*f* radial wavefunction increases along the series. However, this expansion is non-uniform; the outer region expands more when going from U to Pu than the inner region, which remains constrained by increasing nuclear charge. This identified sensitivity is used to rationalise 5*f* covalency trends across the early actinides. The results provide needed experimental insight into the mechanisms by which early actinide 5*f* orbitals engage in covalent bonding.

in actinide ligand bonds underpins the interpretation of physicochemical properties. The proposed presence of 5*f* metal-ligand covalency dates back to work by G. T. Seaborg and co-workers who rationalized ion-exchange behavior in early actinides as evidence of covalent mixing between actinide 5*f* and Cl 3*p* orbitals.²¹ This notion was supported further by Streitwieser and colleagues following the seminal discovery of the actinocenes.²² In more recent years, actinide computational chemistry studies have exposed deeper questions concerning 5*f* covalency, since different bonding analyses provide contrasting perspectives on both the degree of covalency present and the mechanism by which its presence occurs.^{23–26}

There is, therefore, a pressing need for high-quality experimental data to evaluate 5*f* bonding behavior. Experimental measures of 5*f* orbital engagement in bonding are required to correlate with physicochemical properties, validate computational predictions, and develop improved models. There is a paucity of spectroscopic studies that have evaluated early actinide bonding trends. Electron paramagnetic resonance (EPR)²⁷ and nuclear magnetic resonance (NMR)^{28–30} have been applied in cases where ligand ions are nuclear spin active, however studies into covalent contributions to bonding have thus far been limited to U and Th complexes. Chlorine ligand K-edge X-ray absorption near edge structure (XANES) has provided important insight into trends in actinide 5*f* and 6*d* contributions to bonding.^{31–35} This includes a study of [AnCl₆]²⁻ complexes (An = Th, U, Np, Pu) by Batista, Clark, Kozimor, Yang and co-workers that provides

Introduction

The physical and chemical properties of the actinides are influenced by the extent of 5*f* orbital involvement in chemical bonding.^{1–3} Actinide coordination chemistry research is rapidly expanding, revealing rich and varied chemistry including an increasing diversity of transuranium studies,^{4–12} multiple bonds,^{13–15} wide-ranging redox behaviour,^{16–18} and catalysis.^{19,20} In much of this work, the degree of covalency present

^a Department of Chemistry, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK

^b Centre for Radiochemistry Research, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK

^c The University of Manchester at Harwell, Diamond Light Source, Harwell Campus, OX11 0DE, UK

^d Synchrotron SOLEIL, L'Orme des Merisiers, Départementale 128, 91190, Saint-Aubin, France

^e CEA, DES, ISEC, DMRC, Univ. Montpellier, Marcoule 30207, France

† Electronic Supplementary Information (ESI) available: Includes further comments on RIXS experimental details, including instrument scattering geometry and radioactive sample confinement. Further details on ligand field DFT calculations, including AOC calculated active spaces and all calculated LFDFT and multiplet simulation parameters, and calculated radial wavefunctions for each complex. Ligand field multiplet simulations are included showing RIXS planes for [NpCl₆]²⁻ and [PuCl₆]²⁻ simulated with unscaled LFDFT parameters, a partial excitations analysis, as well as simulations showing the spectral dependence on 5*f* ligand field splitting, 5*f* spin-orbit coupling, and Slater integrals. A summary of the X-ray crystallographic analysis of the newly reported compounds is also included. See DOI: 00.0000/00000000.



unique evidence of a $5f$ orbital covalency contribution to Cl bonding.³⁴ Additionally, ligand K-edge XANES has also been proven as a probe of actinide carbon and actinide oxygen ligand bonding.^{36–41}

Resonant inelastic X-ray scattering at actinide $M_{4,5}$ edges ($3d4f$ RIXS) is being increasingly applied as a probe of $5f$ electronic structure and bonding.^{42–52}

Dipole selection rules enable resonant excitation into states of unoccupied $5f$ character, making $3d4f$ RIXS a promising route to exclusively evaluate $5f$ contributions to actinide bonding. There are clear advantages to $3d4f$ RIXS since it can be applied to actinide compounds with any ligand atom identity, nuclear spin, actinide oxidation state, or coordination symmetry. However, $3d4f$ RIXS is considerably less mature than established spectroscopies, with many remaining questions concerning the method's sensitivity to $5f$ bonding.

In the present study, we report M_4 -edge $3d4f$ RIXS on $[AnCl_6]^{2-}$ complexes ($An = U, Np, Pu$) and show how the method carries a previously unidentified sensitivity to the distribution of the averaged $5f$ radial wavefunction, and hence, provides previously lacking insight into the influence of $5f$ orbitals in bonding. It is demonstrated that $3d4f$ RIXS can be analyzed to determine how bonding differentially influences the outer *versus* inner components of the $5f$ radial wavefunction. RIXS fine structure associated with $4f$ - $5f$ spin-exchange is a unique probe of the inner component of the $5f$ radial wavefunction, while fine structure associated with $5f$ inter-electron repulsion is sensitive to the outer component of the $5f$ radial wavefunction. It is shown that this differential sensitivity to the distribution of the $5f$ radial wavefunction provides new insights into early actinide bonding trends. M_4 -edge $3d4f$ RIXS analysis shows that $5f$ radial expansion in going from an An^{4+} free ion to a $[AnCl_6]^{2-}$ complex increases along the actinide series from U to Np and to Pu. However, this expansion becomes increasingly non-uniform. The outer component of the $5f$ radial wavefunction undergoes covalent expansion, due to proximity with the ligand orbitals, while the inner component expands less, due to increasing nuclear charge (Z_{eff}) on traversing the series.

Results and discussion

Experimental RIXS data

The $3d4f$ RIXS measurements were conducted at the M_4 -edge ($3d_{3/2} \rightarrow 5f$) and the $M\beta$ ($4f_{5/2} \rightarrow 3d_{3/2}$) emission line (see Supporting Information for additional details on $3d4f$ RIXS spectroscopy). The M_4 absorption profile is obtained by taking a constant emission energy cut through the two-dimensional RIXS plane, known as high-energy resolution fluorescence detection (HERFD). The $M\beta$ emission profile can be resolved from a vertical cut through the RIXS plane at the resonant excitation energy, known as resonant X-ray emission spectroscopy (RXES). The M_4 HERFD and $M\beta$ RXES spectra for $[UCl_6]^{2-}$, $[NpCl_6]^{2-}$, and $[PuCl_6]^{2-}$ are shown in Fig. 1. The HERFD spectra (Fig. 1a) are dominated by the resonant M_4 absorption feature I_a , around 3725 eV (U), 3846 eV (Np), and 3967 eV (Pu), and exhibit additional fine structure at higher incident energies, labelled I_b . The

two-peak HERFD fine structure for $[PuCl_6]^{2-}$ is similar to that reported for PuO_2 .^{47,53} For $[UCl_6]^{2-}$ only, an additional higher energy satellite feature is detected, labelled I_c . We previously assigned features I_b and I_c to originate from $5f$ inter-electron repulsion interactions.^{51,54}

The RXES spectra (Fig. 1b) are dominated by the actinide $M\beta$ emission line ($4f_{5/2} \rightarrow 3d_{3/2}$) centered around 388 eV (U), 412 eV (Np), and 431 eV (Pu) energy transfer. Low energy transfer satellite features, labelled II_a , are detected for $[UCl_6]^{2-}$, $[NpCl_6]^{2-}$, and $[PuCl_6]^{2-}$ at around 382.5 eV, 405 eV, and 424 eV energy transfer, respectively. An additional satellite feature is shown at higher transfer energies, labelled II_b . We previously assigned feature II_a for $[UCl_6]^{2-}$ to originate from $4f$ - $5f$ spin-exchange interactions.⁵¹ The energy position of satellite feature II_a relative to the maximum of the RXES increases from 6 eV in $[UCl_6]^{2-}$, to 6.3 eV in $[NpCl_6]^{2-}$, and 7.0 eV in $[PuCl_6]^{2-}$, along with a corresponding increase in broadness and intensity relative to the maximum of the RXES.

Computational analysis of RIXS

Uranium(IV) $3d4f$ RIXS fine structure carries a distinct sensitivity to $5f$ inter-electron repulsion and $4f$ - $5f$ inter-shell spin-exchange.⁵¹ In a previous $3d4f$ RIXS study on $[UX_6]^{2-}$ complexes ($X = F, Cl, Br$) we showed that $5f$ inter-electron repulsion and $4f$ - $5f$ spin-exchange both decrease from $[UF_6]^{2-}$ to $[UCl_6]^{2-}$ and to $[UBr_6]^{2-}$. The trend follows the nephelauxetic series,⁵⁵ reflecting reduced inter-electron repulsion and spin-exchange due to expansion of the $5f$ radial wavefunction as covalency increases in going from a hard to soft donor ligand. The $3d4f$ RIXS fine structure of uranium hexahalides is dominated by $5f$ inter-electron repulsion and $4f$ - $5f$ inter-shell spin-exchange, which can be quantified *via* Slater integrals F_{5f5f}^k and G_{4f5f}^k , respectively.⁵¹ The extent by which bonding within a complex reduces F_{5f5f}^k and G_{4f5f}^k for an actinide ion can be expressed in terms of the nephelauxetic reduction parameter (β):⁵⁵ e.g.

$$\beta(F_{5f5f}^k) = \frac{F_{5f5f}^k(\text{complex})}{F_{5f5f}^k(\text{free ion})} \quad \text{and} \quad \beta(G_{4f5f}^k) = \frac{G_{4f5f}^k(\text{complex})}{G_{4f5f}^k(\text{free ion})}, \quad (1)$$

hence a decrease in β is observed as metal-ligand bond covalency increases.

To predict the influence of bonding across the early actinides in $[AnCl_6]^{2-}$ complexes ($An = U, Np, Pu$), we perform ligand field density functional theory (LFDFT) calculations implemented in the Amsterdam Density Functional (ADF) code.^{56,57} LFDFT utilizes an average of configuration (AOC) DFT method to define an active space *via* the selective occupation of frozen Kohn-Sham molecular orbitals. Slater integrals and ligand field parameters are calculated from first principles, including the Coulomb (F^k) and exchange (G^k) Slater integrals, relativistic spin-orbit coupling constants (ζ_{nl}), and the $5f$ ligand field splittings.⁵⁸ All calculated ligand field parameters are listed in Supporting Information Tables S3 to S6. The octahedral ligand field transforms the $5f$ orbitals into three irreducible representations, t_{1u} , t_{2u} and a_{2u} . The total calculated ligand field energy splitting does not ex-






Fig. 1 Experimental (solid colored) and simulated (dashed black) M_4 -edge HERFD (a) and $M\beta$ RXES (b) spectra for $[\text{UCl}_6]^{2-}$ (left), $[\text{NpCl}_6]^{2-}$ (middle), and $[\text{PuCl}_6]^{2-}$ (right), with inplots highlighting satellite features I_c and II_a . Simulated M_4 -edge $3d4f$ RIXS planes (c) are shown for $[\text{UCl}_6]^{2-}$ (left), $[\text{NpCl}_6]^{2-}$ (middle), and $[\text{PuCl}_6]^{2-}$ (right), with the corresponding simulated HERFD and RXES cuts shown as diagonal and vertical red lines, respectively.

ceed 0.365 eV, and there are only minor differences in the ligand field splitting for $[\text{UCl}_6]^{2-}$, $[\text{NpCl}_6]^{2-}$, and $[\text{PuCl}_6]^{2-}$. Calculated ligand field parameters describing $5f$ inter-electron repulsion (F_{5f5f}^k) for the $[\text{AnCl}_6]^{2-}$ complexes are shown in Table 1, expressed in the terms of the nephelauxetic reduction factor β (Eq. 1). The calculated nephelauxetic $\beta(F_{5f5f}^k)$ values ($k = 2, 4, 6$) in Table 1 show a sequential decrease in $5f$ inter-electron repulsion in going from $[\text{UCl}_6]^{2-}$ to $[\text{NpCl}_6]^{2-}$ and to $[\text{PuCl}_6]^{2-}$ relative to their free ion values, indicating that $5f$ radial wavefunction expansion from free ion to complex increases across the series. The calculated $5f$ spin-orbit coupling constants are also reduced relative to their free-ion values due to the relativistic nephelauxetic effect.⁵⁹ Relativistic $\beta(\zeta_{5f})$ decreases slightly, consistent with the trends identified in $\beta(F_{5f5f}^k)$, from $[\text{UCl}_6]^{2-}$ to $[\text{NpCl}_6]^{2-}$ and $[\text{PuCl}_6]^{2-}$ with values of 0.84, 0.83 and 0.81, respectively (See Supporting Information Table S3).

$\beta(F_{5f5f}^k)$	$[\text{UCl}_6]^{2-}$	$[\text{NpCl}_6]^{2-}$	$[\text{PuCl}_6]^{2-}$
$k = 2$	0.80	0.77	0.72
$k = 4$	0.79	0.76	0.71
$k = 6$	0.79	0.75	0.71

Table 1 Nephelauxetic β factors for the F_{5f5f}^k ($k = 2, 4, 6$) Slater integrals describing $5f$ inter-electron repulsion calculated by LFDFT.

To model the $3d4f$ RIXS, further LFDFT calculations are conducted for both RIXS intermediate ($3d^9 4f^{14} 5f^{n+1}$) and final ($3d^{10} 4f^{13} 5f^{n+1}$) state electron configurations and the resulting ligand field parameters are implemented in RIXS multiplet simulations.⁵¹ The calculated $5f$ ligand field splitting energies in the intermediate and final states are similar to the initial states for all three compounds and do not exceed 0.38 eV. The $5f$ ligand field splitting is therefore rather small relative to the measured RIXS energy resolution. LFDFT predicted parameters accurately reproduce the HERFD and RXES spectra, except the G_{4f5f}^0 values which, to obtain agreement with experiment, require a reduction to 62% of the LFDFT calculated value for $[\text{NpCl}_6]^{2-}$ and






Fig. 2 Experimental (a) and simulated (b) $M\beta$ RXES satellite feature II_a for $[UCl_6]^{2-}$, $[NpCl_6]^{2-}$, and $[PuCl_6]^{2-}$, plotted relative to the energy and intensity of the maximum of the RXES.

69% of the LFDFT calculated value for $[PuCl_6]^{2-}$, compared to a reduction to 54% of the LFDFT calculated value for $[UCl_6]^{2-}$ (see Fig. S6 for LFDFT RIXS simulations excluding reduction of G_{4f5f}^0).^{51,54} No other changes to the LFDFT parameters are required for good agreement with experiment. The simulated RIXS planes are shown in Fig. 1(c), and the corresponding HERFD and RXES simulations are compared with experiment in Fig. 1(a) and (b), respectively. Good reproduction of the general features is observed. The simulated HERFD captures the general shape of the spectra, including features I_a and I_b which result in an asymmetry in the HERFD shape for $[NpCl_6]^{2-}$ and the double peak structure for $[PuCl_6]^{2-}$. To gain insight into the sensitivity of the RIXS spectra to the obtained LFDFT parameters, a series of systematic calculations were conducted, identifying significant sensitivity to $5f$ inter-electron repulsion, inter-shell spin-exchange, and spin-orbit coupling, and a very limited influence from $5f$ ligand field splitting (see Supporting Information Fig. S8 to S12). It was previously shown that RIXS fine-structure for $[UCl_6]^{2-}$ has minimal sensitivity to $5f$ spin-orbit coupling.⁵¹ In contrast, the simulated RIXS fine structure for $[NpCl_6]^{2-}$ and $[PuCl_6]^{2-}$ shows a more significant sensitivity to $5f$ spin orbit coupling (see Supporting Information Fig. S12). The relative energy and intensity of spec-

tral features I_a and I_b for $[NpCl_6]^{2-}$ and $[PuCl_6]^{2-}$ are identified to depend on the magnitude of the F_{5f5f}^k Slater integrals and $5f$ spin-orbit coupling constant, ζ_{5f} . Varying these parameters away from those predicted by LFDFT worsens the agreement with experiment (See Supporting Information Fig. S9 and S12).

The experimental spectra, therefore, provide an independent confirmation for the LFDFT predicted values of F_{5f5f}^k and ζ_{5f} , confirming the predicted increase in covalency in moving across the series from U to Np and Pu. The simulated RXES, including satellite features II_a and II_b , are reproduced for all three complexes. The RXES satellites II_a and II_b are narrower than observed in experiment. Due to the different nature of the RXES satellite excitations relative to the main emission line, different line shape broadening can be expected. However, a constant line shape was used for the calculations of the RIXS planes, in order to describe electronic fine structure in the most transparent way. The simulated RXES show increasing intensity and energy separation of satellite II_a relative to the maximum of the RXES in going from $[UCl_6]^{2-}$ to $[NpCl_6]^{2-}$ to $[PuCl_6]^{2-}$, reproducing the experimental spectral trend (Fig. 2). The calculations confirm that satellite features II_a and II_b originate from $4f$ - $5f$ spin-exchange interactions, and most strongly depend on G_{4f5f}^0 (see Supporting Information Fig. S11). The relative energy and intensity between the maximum of the RXES and II_a is therefore a direct probe of G_{4f5f}^0 .^{51,52}

The simulations reveal that the spectral features in the $3d4f$ RIXS of $[AnCl_6]^{2-}$ are insensitive to the relatively small $5f$ ligand field splitting generated by the coordinated chloride ions (see further analysis in Supporting Information Section S3.2). The simulations reveal that $5f$ a_{2u} , t_{2u} , and t_{1u} orbital character is strongly mixed over the full energy range of the absorption edge (see transition dipole tensor analysis, Fig. S7). Therefore, while symmetry-restricted mixing of metal and ligand orbitals will contribute to the reduction of inter-electron repulsion, $5f$ spin-orbit coupling and spin-exchange, the spectra are insensitive to the relative amounts of σ versus π covalency⁶⁰ and $[AnCl_6]^{2-}$ ligand field splitting.

Decomposition of the $5f$ radial wavefunction

Since the RIXS of all three compounds are found to carry a strong sensitivity to F_{5f5f}^k and G_{4f5f}^0 , we focus on these Slater integrals for the $[AnCl_6]^{2-}$ complexes in Table 2. For a given actinide, the nephelauxetic reduction parameters (β) express the amount the $5f$ radial wavefunction expands in going from free ion to $[AnCl_6]^{2-}$ complex.

The trend in $\beta(F_{5f5f}^k)$ when moving from An = U to Np to Pu indicates an increasing $5f$ radial expansion in response to bonding across the $[AnCl_6]^{2-}$ series and, therefore, an increase in covalency. Conclusions from earlier Cl ligand K-edge XANES, DFT and relativistic multiconfigurational wavefunction theory studies on the same complexes identified that An $5f$ - Cl $3p$ covalency either increases slightly or does not significantly change across the series.^{25,34} A more recent DFT analysis of covalent bond formation in $[UCl_6]^{2-}$, $[NpCl_6]^{2-}$ and $[PuCl_6]^{2-}$, applying a prebonding assessment of atomic orbital overlap versus atomic orbital energy



	[UCl ₆] ²⁻	[NpCl ₆] ²⁻	[PuCl ₆] ²⁻
$\beta(F_{5f5f}^2)$	0.77	0.71	0.64
$\beta(F_{5f5f}^4)$	0.76	0.70	0.64
$\beta(F_{5f5f}^6)$	0.76	0.70	0.63
$\beta(G_{4f5f}^0)$	0.45	0.50	0.53

Table 2 RIXS final state configuration ($4f^{13}5f^{n+1}$) F_{5f5f}^k and G_{4f5f}^0 Slater integrals consistent with experimental $3d4f$ RIXS data given in terms of the nephelauxetic reduction factor β .

matching, identify a small but non negligible An($5f$)-Cl($3p$) orbital overlap throughout the series²⁶. The computational analyses in these studies relate to symmetry-adapted mixing of Cl $3p$ and An $5f$ character (symmetry-restricted covalency). In contrast, $3d4f$ RIXS and LDFDT based determination of nephelauxetic β incorporates the effect of both average $5f$ radial wavefunction expansion (central-field covalency) and symmetry-restricted covalency. Since Cl K-edge XANES and theoretical calculations^{25,26,34} infer that symmetry-restricted covalency does not vary significantly along the early actinide series, we assign an increase in central-field covalency from U to Pu to explain the identified reductions in $\beta(F_{5f5f}^k)$. The $\beta(G_{4f5f}^0)$ values show an opposing trend relative to $\beta(F_{5f5f}^k)$ in moving from An = U to Np to Pu. This difference may seem counter intuitive, however the sensitivities of F_{5f5f}^k and G_{4f5f}^0 to bonding inherently differ since G_{4f5f}^0 is linked to the $4f$ core sub-shell.⁵¹ The $4f$ core sub-shell has a contracted radial wavefunction and is unaffected by bonding (Fig. S13). Consequently, G_{4f5f}^0 is linked to the behaviour of the inner part of the $5f$ radial wavefunction. In contrast, F_{5f5f}^k carries greater sensitivity to the behavior of the outer part of the $5f$ radial wavefunction. For example, the distinct radial sensitivities of $F_{5f5f}^2(r)$ and $G_{4f5f}^0(r)$ can be demonstrated by comparing them with their limiting values $F_{5f5f}^2(\infty)$ and $G_{4f5f}^0(\infty)$ alongside the associated $5f$ or $4f$ radial wavefunctions (Fig. 3). The F_{5f5f}^2 integral gains magnitude at radial distances greater than $r \sim 0.3$ Å and reaches its terminal value ($F_{5f5f}^2(r)/F_{5f5f}^2(\infty) \rightarrow 1$) at large radial distances, showing the sensitivity of F_{5f5f}^2 to the outer component of the $5f$ radial wavefunction. In contrast, the G_{4f5f}^0 Slater integral gains magnitude at low radial distances and reaches its terminal value ($G_{4f5f}^0(r)/G_{4f5f}^0(\infty) \rightarrow 1$) by $r \sim 0.4$ Å.

A subtle sensitivity of different Slater integrals to inner *versus* outer component of radial wavefunctions has been identified in transition metal UV-Vis spectra^{61,62} and, more recently, high-resolution $2p3d$ valence-to-core RIXS.^{63,64} For transition metals, F_{3d3d}^2 is slightly more sensitive to the outer part of the $3d$ radial wavefunction than F_{3d3d}^4 .⁶⁵ Consequently, a greater reduction from the free ion value of F_{3d3d}^2 is observed relative to F_{3d3d}^4 in covalent complexes.⁶¹ By contrast, a very distinct sensitivity of G_{4f5f}^0 to the inner components of the $5f$ radial wavefunction and F_{5f5f}^2 to the outer components enables deconvolution of the influence of bonding inside *versus* outside the $5f$ radial node.




Fig. 3 (a) The averaged $5f$ radial wavefunction for $[\text{PuCl}_6]^{2-}$ plotted with the normalized $F_{5f5f}^2(r)/F_{5f5f}^2(\infty)$ rolling integral, showing its strong sensitivity to the outer component of the $5f$ radial wavefunction. The rolling integral behaviour of $F_{5f5f}^{k=4,6}$ are very similar to $F_{5f5f}^{k=2}$, see Supporting Information Fig. S14. (b) $4f$ and $5f$ radial wavefunctions for $[\text{PuCl}_6]^{2-}$ plotted with the normalized $G_{4f5f}^0(r)/G_{4f5f}^0(\infty)$ rolling integral, showing its strong sensitivity to the inner component of the $5f$ radial wavefunction. Schematics showing cross-sections of a $5f_{x(x^2-3y^2)}$ atomic orbital highlight the sensitivity the radial integrals have to the outer (a) and inner (b) components.

On traversing the actinide series, radial contraction of the $5f$ orbitals is understood to occur due to imperfect screening of the increasing nuclear charge by successive addition of electrons in $5f$ orbitals.² The consequence of imperfect screening of nuclear charge is the so-called actinide contraction, the sequential radial contraction of the atomic $5f$ radial wavefunction. The opposing trends in β values demonstrate that distortion of the $5f$ radial wavefunction increases from U to Pu, providing a partial counteraction of the actinide contraction, by maintaining a radially contracted electron density inside the $5f$ node, thereby permitting the outer component of the $5f$ radial wavefunction to expand for



participation in covalent metal-ligand bonding.

The specific sensitivity of G_{4f5f}^0 to orbital behaviour at low radial distances provides an explanation for the required reduction of the LFDFT predicted G_{4f5f}^0 value to match experiment. The exclusion of semi-core correlation effects in the *ab initio* determination of ligand field parameters may lead to an underestimation of the nephelauxetic effect,⁶⁶ and hence, an overestimation of Slater integral values. Since G_{4f5f}^0 depends on both the core $4f$ and the inner "semi-core" component of the $5f$ radial wavefunction, it follows that it will be more sensitive to these dynamic correlation effects than the outer "valence" component of the $5f$ radial wavefunction. Specific inclusion of semi-core correlation effects in the LFDFT model may, therefore, correct overestimation of G_{4f5f}^0 in the spectroscopic final state.

Conclusion

Our analysis reveals that $3d4f$ RIXS possesses exceptional sensitivity to the influence of bonding on the inner *versus* outer component of the actinide $5f$ radial wavefunction. This sensitivity is so distinct that it enables deconvolution of the response to bonding inside *versus* outside the $5f$ radial node. For $[\text{AnCl}_6]^{2-}$ (An = U, Np, Pu), it is found that An-Cl bonding increasingly distorts the $5f$ radial wavefunction in going from U to Np and to Pu. Bonding induced expansion of the outer component of the $5f$ radial wavefunction increases from U to Pu whereas the inner component decreases, due to the increase in nuclear charge. This opposing trend in the behavior of the inner *versus* outer components of the $5f$ radial wavefunction contribute to partly counteract actinide contraction, aiding the availability of the $5f$ orbitals to participate in covalent metal-ligand bonding.

There is a limited number of experimental methods available for quantifying trends in actinide electronic structure and bonding. The approach presented here is particularly versatile as it is applicable to actinide systems irrespective of ligand element, nuclear spin, oxidation state, and coordination symmetry. The identified $3d4f$ RIXS sensitivity to covalency and the $5f$ radial wavefunction significantly enhances our ability to establish structure–function relationships and physicochemical trends in actinide chemistry, and provides valuable experimental data to evaluate the predictive power of theoretical models.

Methods

Experimental

Caution! ^{237}Np and ^{239}Pu isotopes are high specific activity α -particle emitting radionuclides, and their use presents extreme hazards to human health. ^{238}U isotopes are low specific-activity α -particle emitting radionuclides, and their use presents hazards to human health. This research was conducted in a radiological facility with appropriate analyses of these hazards and implementation of controls for the safe handling and manipulation of these toxic and radioactive materials.

Synthesis of $[(\text{C}_2\text{H}_5)_4\text{N}]_2\text{PuCl}_6$ (1) and $[(\text{C}_2\text{H}_5)_4\text{N}]_2\text{NpCl}_6$ (2)

All manipulations involving radioactive materials were conducted at the ATALANTE facility (CEA-ISEC, France) under strictly controlled conditions. Due to the high radiotoxicity of neptunium

and plutonium, all experiments were carried out in a negative-pressure glovebox under ambient air, with stringent operational protocols in place. All chemicals were used as received: nitric acid (HNO_3 , 65%) and hydrochloric acid (HCl , 37%) were obtained from Sigma-Aldrich, along with tetraethylammonium chloride (TEAC) and boron nitride (BN). Sodium hydroxide (NaOH) pellets were purchased from VWR and hydroxylamine hydrochloride ($\text{NH}_2\text{OH}\cdot\text{HCl}$) from Merck.

The syntheses of **1** and **2** were adapted from procedures previously reported for analogous actinide complexes of the type M_2AnCl_6 , typically prepared *via* slow evaporation from aqueous solutions.^{67,68} For compound **1**, the plutonium precursor was Pu(IV) in 1.5 M HNO_3 . 20 mg ($V = 133 \mu\text{L}$) was precipitated by the addition of concentrated NaOH . The resulting solid was washed three times with deionized water and subsequently dissolved in 4 M HCl (200 μL). After complete dissolution, two equivalents of TEAC (27.7 mg), pre-dissolved in 4 M HCl (100 μL) were added. Crystals of **1** suitable for single crystal X-ray diffraction (SC-XRD) were obtained by slow evaporation of the solution. For compound **2**, the neptunium precursor was Np(V) in 3 M nitric acid. A 20 mg aliquot ($V = 369 \mu\text{L}$) was used. The reduction of Np(V) to Np(IV) was achieved by adding 100 equivalents of $\text{NH}_2\text{OH}\cdot\text{HCl}$ followed by heating at 70°C for several hours. The reduced Np(IV) species was then precipitated by addition of concentrated NaOH , was washed three times with water, and dissolved in 4 M HCl (200 μL) containing 4 equivalents of $\text{NH}_2\text{OH}\cdot\text{HCl}$. After complete dissolution, 2 equivalents of TEAC (27.7 mg) dissolved in 4 M HCl (100 μL) were added. Crystals of **2** suitable for SC-XRD were obtained by slow evaporation of the solution.

Single crystal X-ray diffraction

Single crystal X-ray diffraction (SC-XRD) measurements were performed using a Bruker D8 Quest diffractometer equipped with a PHOTON II charge-coupled device (CCD) detector and a Mo-target $\text{I}\mu\text{S}$ microfocus radiation source ($\lambda = 0.71073 \text{ \AA}$). Data were collected at 200 K using an Oxford Cryosystems 800 series cryostream cooler. Individual crystals were mounted on a MiTeGen MicroMount and enclosed in MicroRT capillaries, which were sealed to the goniometer base using adhesive to prevent the release of radioactive material and ensure safe handling. Data were collected using ϕ and Ω scan strategies. Diffraction intensities were integrated using the SAINTplus software package,⁶⁹ and multi-scan absorption corrections were performed.⁷⁰

Unit cell parameters were refined from the complete dataset. Structural solution and refinement were carried out using SHELX-2017.⁷¹ Heavy atoms were located by direct methods, and the remaining atoms were identified through successive Fourier map analyses. All non-hydrogen atoms were refined anisotropically. Due to disorder in one of the TEA cations, hydrogen atoms on the ethyl chains could not be located from Fourier map analyses and were not included in the refinement. Hydrogen atoms bonded to the ordered TEA cation were placed in idealized positions using a riding model and assigned fixed isotropic displacement parameters.

Crystallographic data and refinement details for both struc-



tures (Fig. S15–S16) are summarized in Table S11. The structures have been deposited on the Cambridge Crystallographic Data Centre (CCDC) structural database with deposition numbers 2466992 (1) and 2466993 (2). These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

Resonant Inelastic X-ray Scattering

For RIXS measurements, samples were prepared as pellets by finely grinding *ca.* 6 mg of single crystals with *ca.* 15 mg of BN. The pellets were mounted in special stainless steel sample holders enabling double confinement of the samples following the requirements of the French Nuclear Safety authorization at the MARS beamline. Further description of the sample holders can be found in the Supporting Information.

Neptunium and plutonium M_4 -edge RIXS measurements were carried out at ambient temperature using the CX3 end-station of the MARS beamline⁷² at Synchrotron SOLEIL (Saint-Aubin, France). The storage ring operated in top-up mode at 2.5 GeV with an electron current of 500 mA. The incident photon energy was selected using a double-crystal monochromator (DCM) equipped with two Si(111) crystals. Vertical focusing and harmonic rejection were achieved using Si mirror strips positioned before and after the DCM at an incidence angle of 4 mrad. Energy calibration was performed using the L_3 absorption edge of iodine in a KI pellet (4557 eV). The incident X-ray flux at 3700 eV was 1.9×10^9 photons s^{-1} , with a beam size of $350 \times 140 \mu\text{m}$ (horizontal \times vertical, FWHM) at the sample position. The effects of photon-induced radiolysis for each sample were quantified before being mitigated by minimizing the scanning time per sample position on the pellets. The $M\beta$ emission lines of Np (3435 eV) and Pu (3534 eV) were detected using a spherically bent Si(220) crystal analyzer (1 m bending radius) and a KETEK silicon drift detector arranged in a vertical Rowland circle geometry, with Bragg angles of 70.0° and 66.0° , respectively. Further details concerning the precise experimental scattering geometry for each measurement are found in the Supporting Information Section S1.1. Emission intensity was normalized to the incident flux. Further details on the experimental geometry can be found in Supporting Information, S1.1. A He-filled chamber was used to minimize air scattering between the sample, analyzer, and detector. HERFD spectra were normalized to 1 in the post-edge region (3905–3940 eV for Np, 4015–4030 eV for Pu) using a linear fitting function. Background-subtracted RXES spectra were scaled so that their resonant intensity matched the maximum intensity of the HERFD spectra.

Uranium M_4 -edge RIXS measurements were performed at the European Synchrotron Research Facility (ESRF, Grenoble, France) as previously reported.⁵¹ HERFD and RXES spectra were normalized as above, using a HERFD post-edge region of 3820–3840 eV.

Computational

Electronic structure calculations were performed on $[\text{AnCl}_6]^{2-}$ fragments (An = U, Np, Pu) based on single-crystal X-ray diffraction (XRD) crystallographic data.

Ligand field density functional theory calculations

Average of configuration (AOC) DFT calculations were performed using the Amsterdam Density Functional (ADF) code implemented in the Amsterdam Modeling Suite (AMS, version 2023.104).⁵⁶ The B3LYP functional was used for all calculations. Equivalent calculations were also conducted with the PBE0 function. There are negligible differences in LFDFT parameters between PBE0 and B3LYP; see Supporting Information Tables S7–8 for β values using PBE0 *versus* B3LYP. Scalar relativistic effects were treated with the zeroth order relativistic approximation (ZORA). Given known incompleteness errors originating in inner shells using ZORA⁷³, we also conducted calculations using X2C. LFDFT parameters obtained using X2C are very similar, and related RIXS simulations are nearly identical. Supporting Information Tables S9–S10 show that there are negligible differences in nephelauxetic β values using ZORA *versus* X2C.^{74,75} All calculations employed Slater type orbital (STO) triple- ζ -plus double polarisation all-electron basis sets from the ZORA/TZ2P database.⁷⁶ The active spaces in the AOC calculations were defined as follows for An(IV) with ground state electron occupations of $5f^n$ for An = U ($n = 2$), Np ($n = 3$), Pu ($n = 4$). For the ground state, each of the seven $5f$ MOs were fractionally populated with $n/7$ electrons, whilst for the intermediate and final states, the $5f$ MOs were fractionally populated with $(n + 1)/7$ electrons. The core holes were defined by fractionally occupying each actinide core $3d$ orbital with $9/5$ electrons for the intermediate state and each core $4f$ orbital with $13/7$ electrons in the final state. The AOC DFT results were treated by the ligand field density functional theory^{57,58} (LFDFT) implemented in ADF to obtain ligand field parameters for each of the initial ($3d^{10}4f^{14}5f^n$), intermediate ($3d^94f^{14}5f^{n+1}$), and final ($3d^{10}4f^{13}5f^{n+1}$) RIXS electronic configurations. Isosurface plots of the associated Kohn-Sham MOs are provided in Supporting Information Fig. S5 (isosurface level = 0.03). The plotted $4f$ and $5f$ radial wavefunctions are represented by radial distribution functions (RDFs) extracted from the AOC Kohn-Sham orbitals centered on the actinide ion and printed by inclusion of the *debug* keyword in the LFDFT input file.

Ligand field multiplet simulations

Multiplet simulations of $3d4f$ RIXS planes were carried out using Quanty Version 0.6.⁷⁷ The simulated RIXS planes include linewidth broadening to incorporate the effect of energy broadening due to core-hole lifetimes combined with spectrometer resolution. Lorentzian full-width half-maximum broadenings of 4.4 eV and 0.6 eV are applied to the incident energy ($3d^9$) dimension and the energy transfer dimension ($4f^{13}$), respectively. Additionally, a Gaussian full-width half-maximum broadening of 0.8 eV is applied to both incident and transfer energy dimensions.

The selected broadenings were determined empirically. The Lorentzian broadenings applied are slightly larger than available core-hole lifetime broadening data,⁷⁸ since they also incor-



porate instrumental broadening contributions to the linewidth. Improved fits can be obtained by including slightly different linewidths for each element, however, for simplicity and transparency, only one set of linewidth parameters is used for all compounds.

Hamiltonians for the initial, intermediate, and final electronic configurations of the $3d4f$ RIXS process were constructed using the results of LFDFT calculations, including $5f$ ligand field splitting, electron-electron interactions, and spin-orbit coupling. The influence of each term in the Hamiltonian was systematically investigated via an analysis of simulated RIXS fine structure to determine the relationship between the observed spectral features and the differences in electronic structure and bonding in going from $[\text{UCl}_6]^{2-}$ to $[\text{NpCl}_6]^{2-}$ and $[\text{PuCl}_6]^{2-}$ (see Supporting Information S3.2 for further details).^{79,80} The G_{4f5f}^0 Slater integral required adjustment to obtain RIXS simulations that best fit the experimental results. For consistency, the other $4f-5f$ Slater integrals, including F_{4f5f}^k ($k=2,4,6$) and G_{4f5f}^k ($k=2,4$) are treated with the same scaling, although RIXS spectral simulations are less sensitive to these Slater integrals (Supporting Information Fig. S10 and S11), as previously established.^{51,52} Other Slater integrals determined from LFDFT analysis (including F_{5f5f}^k , F_{3d5f}^k and G_{3d5f}^k) were found to be consistent with experiment. The 2D RIXS planes were calculated via a Green's theorem approach to the Kramers-Heisenberg equation, incorporating the experimental geometries⁸¹ (see Supporting Information S1.1 for further details).

Author contributions

MLB supervised and conceptualized the study. Np and Pu RIXS measurements were conducted by NMA, MSH, TGB, MLB, TD and MOJYH. NMA performed the data analysis and calculations under the supervision of MLB. Ligand field multiplet simulations were performed by NMA and MSH under the supervision of MLB. MA and CT synthesized compounds and determined their structures by SC-XRD under supervision of TD. The manuscript was written by NMA and MLB with review from all co-authors.

Conflicts of interest

There are no conflicts to declare.

Data availability

Data for this paper, including RIXS data and calculations are available at Figshare at <https://doi.org/DOI>

Acknowledgements

We thank Synchrotron SOLEIL (20190807 and 20230706) and the European Synchrotron Radiation Facility (CH-6518) for the experimental time. This research was supported by the EPSRC (grant number EP/V029347/1 and EP/W029057/1). MLB was supported by the Analytical Chemistry Trust Fund and a Community for Analytical Measurement Science (CAMS) Fellowship. This work was supported by computing resources provided by the STFC Scientific Computing Department SCARF cluster and the University of Manchester Computational Shared Facility, and associated support services. We extend thanks to William Breton

at MARS for his technical assistance, as well as to Bruno Forgeat (SOLEIL) of the Radioprotection Services for their help.

Notes and references

- S. T. Liddle, *Angewandte Chemie International Edition*, 2015, **54**, 8604–8641.
- M. L. Neidig, D. L. Clark and R. L. Martin, *Coordination Chemistry Reviews*, 2013, **257**, 394–406.
- S. T. Liddle, *Inorganic Chemistry*, 2024, **63**, 9366–9384.
- J. Murillo, J. A. Seed, A. J. Wooles, M. S. Oakley, C. A. Goodwin, M. Gregson, D. Dan, N. F. Chilton, A. J. Gaunt, S. A. Kozimor *et al.*, *Journal of the American Chemical Society*, 2024, **146**, 4098–4111.
- C. A. Goodwin, M. T. Janicke, B. L. Scott and A. J. Gaunt, *Journal of the American Chemical Society*, 2021, **143**, 20680–20696.
- M. S. Dutkiewicz, C. A. Goodwin, M. Perfetti, A. J. Gaunt, J.-C. Griveau, E. Colineau, A. Kovács, A. J. Wooles, R. Caciuffo, O. Walter *et al.*, *Nature Chemistry*, 2022, **14**, 342–349.
- T. N. Poe, H. Ramanantoanina, J. M. Sperling, H. B. Wineinger, B. M. Rotermund, J. Brannon, Z. Bai, B. Scheibe, N. Beck, B. N. Long, S. Justiniano, T. E. Albrecht-Schönart and C.-B. Cristian, *Nature Chemistry*, 2023, **15**, 722–728.
- J. E. Niklas, K. S. Otte, C. M. Studvick, S. Roy Chowdhury, B. Vlaisavljevich, J. Bacsá, F. Kleemiss, I. A. Popov and H. S. La Pierre, *Nature Chemistry*, 2024, **16**, 1490–1495.
- D. R. Russo, A. N. Gaiser, A. N. Price, D.-C. Sergentu, J. N. Wacker, N. Katzer, A. A. Peterson, J. A. Branson, X. Yu, S. N. Kelly, E. T. Ouellette, J. Arnold, J. R. Long, W. W. Lukens Jr, S. J. Teat, R. J. Abergel, P. L. Arnold, J. Autschbach and S. G. Minasian, *Science*, 2025, **387**, 974–978.
- K. P. Carter, K. M. Shield, K. F. Smith, Z. R. Jones, J. N. Wacker, L. Arnedo-Sanchez, T. M. Mattox, L. M. Moreau, K. E. Knope, S. A. Kozimor, C. H. Booth and R. J. Abergel, *Nature*, 2021, **590**, 85–88.
- N. B. Beck, C. Celis-Barros, M. C. Martelles, J. M. Sperling, Z. Bai, J. P. Brannon, D. G. Martinez, Z. K. Huffman, B. N. Long, K. N. Mendoza, R. W. Merinsky, B. M. Rotermund, T. N. Poe and T. E. Albrecht, *Nature Communications*, 2025, **16**, 7759.
- H. Zhang, A. Li, K. Li, Z. Wang, X. Xu, Y. Wang, M. V. Sheridan, H.-S. Hu, C. Xu, E. V. Alekseev, Z. Zhang, P. Yan, K. Cao, Z. Chai, T. E. Albrecht-Schönart and S. Wang, *Nature*, 2023, **616**, 482–487.
- D. M. King, F. Tuna, E. J. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Science*, 2012, **337**, 717–720.
- H.-S. Hu, Y.-H. Qiu, X.-G. Xiong, W. E. Schwarz and J. Li, *Chemical Science*, 2012, **3**, 2786–2796.
- A. R. Fox and C. C. Cummins, *Journal of the American Chemical Society*, 2009, **131**, 5716–5717.
- C. L. Tracy, M. Lang, J. M. Pray, F. Zhang, D. Popov, C. Park, C. Trautmann, M. Bender, D. Severin, V. A. Skuratov and R. C. Ewing, *Nature Communications*, 2015, **6**, 6133.
- C. Deng, J. Liang, R. Sun, Y. Wang, P.-X. Fu, B.-W. Wang,



- S. Gao and W. Huang, *Nature Communications*, 2023, **14**, 4657.
- 18 L. S. Natrajan, A. N. Swinburne, M. B. Andrews, S. Randall and S. L. Heath, *Coordination Chemistry Reviews*, 2014, **266–267**, 171–193.
- 19 P. L. Arnold, T. Ochiai, F. Y. Lam, R. P. Kelly, M. L. Seymour and L. Maron, *Nature Chemistry*, 2020, **12**, 654–659.
- 20 X. Liu, Y. Xie, M. Hao, Y. Li, Z. Chen, H. Yang, G. I. Waterhouse, X. Wang and S. Ma, *Nature Communications*, 2024, **15**, 7736.
- 21 R. Diamond, K. Street Jr and G. T. Seaborg, *Journal of the American Chemical Society*, 1954, **76**, 1461–1469.
- 22 A. Streitwieser Jr and U. Müller-Westerhoff, *Journal of the American Chemical Society*, 1968, **90**, 7364–7364.
- 23 N. Kaltsoyannis and A. Kerridge, *Nature Reviews Chemistry*, 2024, **8**, 701–712.
- 24 N. Kaltsoyannis, *Inorganic Chemistry*, 2013, **52**, 3407–3413.
- 25 D. C. Sergentu and J. Autschbach, *Chemical Science*, 2022, **13**, 3194–3207.
- 26 A. Sen, A. Obeng, K. Stanistreet-Welsh and J. Autschbach, *Inorganic Chemistry*, 2025, **64**, 23787–23803.
- 27 A. Formanuiik, A.-M. Ariciu, F. Ortu, R. Beekmeyer, A. Kerridge, F. Tuna, E. J. McInnes and D. P. Mills, *Nature Chemistry*, 2017, **9**, 578–583.
- 28 J. Du, J. A. Seed, V. E. Berryman, N. Kaltsoyannis, R. W. Adams, D. Lee and S. T. Liddle, *Nature Communications*, 2021, **12**, 5649.
- 29 J. Du, J. Hurd, J. A. Seed, G. Balázs, M. Scheer, R. W. Adams, D. Lee and S. T. Liddle, *Journal of the American Chemical Society*, 2023, **145**, 21766–21784.
- 30 T. W. Hayton and J. Autschbach, *Accounts of Chemical Research*, 2025, 3259–3263.
- 31 S. A. Kozimor, P. Yang, E. R. Batista, K. S. Boland, C. J. Burns, D. L. Clark, S. D. Conradson, R. L. Martin, M. P. Wilkerson and L. E. Wolfsberg, *Journal of the American Chemical Society*, 2009, **131**, 12125–12136.
- 32 S. G. Minasian, J. M. Keith, E. R. Batista, K. S. Boland, D. L. Clark, S. D. Conradson, S. A. Kozimor, R. L. Martin, D. E. Schwarz, D. K. Shuh, G. L. Wagner, M. P. Wilkerson, L. E. Wolfsberg and P. Yang, *Journal of the American Chemical Society*, 2012, **134**, 5586–5597.
- 33 L. P. Spencer, P. Yang, S. G. Minasian, R. E. Jilek, E. R. Batista, K. S. Boland, J. M. Boncella, S. D. Conradson, D. L. Clark, T. W. Hayton, S. A. Kozimor, R. L. Martin, M. M. MacInnes, A. C. Olson, B. L. Scott, D. K. Shuh and M. P. Wilkerson, *Journal of the American Chemical Society*, 2013, **135**, 2279–2290.
- 34 J. Su, E. R. Batista, K. S. Boland, S. E. Bone, J. A. Bradley, S. K. Cary, D. L. Clark, S. D. Conradson, A. S. Ditter, N. Kaltsoyannis, J. M. Keith, A. Kerridge, S. A. Kozimor, M. W. Löble, R. L. Martin, S. G. Minasian, V. Mocko, H. S. La Pierre, G. T. Seidler, D. K. Shuh, M. P. Wilkerson, L. E. Wolfsberg and P. Yang, *Journal of the American Chemical Society*, 2018, **140**, 17977–17984.
- 35 J. N. Cross, J. Su, E. R. Batista, S. K. Cary, W. J. Evans, S. A. Kozimor, V. Mocko, B. L. Scott, B. W. Stein, C. J. Windorff and P. Yang, *Journal of the American Chemical Society*, 2017, **139**, 8667–8677.
- 36 S. G. Minasian, J. M. Keith, E. R. Batista, K. S. Boland, D. L. Clark, S. A. Kozimor, R. L. Martin, D. K. Shuh and T. Tylliszczak, *Chemical Science*, 2014, **5**, 351–359.
- 37 J. A. Branson, P. W. Smith, J. Arnold and S. G. Minasian, *Inorganic Chemistry*, 2024, **63**, 15557–15562.
- 38 Y. Zhang, W. Duan, Y. Yang, T. Jian, Y. Qiao, G. Ren, N. Zhang, L. Zheng, W. Yan, J. Wang *et al.*, *Inorganic Chemistry*, 2021, **61**, 92–104.
- 39 R. G. Denning, J. C. Green, T. E. Hutchings, C. Dallera, A. Tagliaferri, K. Giarda, N. B. Brookes and L. Braicovich, *The Journal of Chemical Physics*, 2002, **117**, 8008–8020.
- 40 Y. Qiao, G. Ganguly, C. H. Booth, J. A. Branson, A. S. Ditter, D. J. Lussier, L. M. Moreau, D. R. Russo, D.-C. Sergentu, D. K. Shuh *et al.*, *Chemical Communications*, 2021, **57**, 9562–9565.
- 41 D. E. Smiles, E. R. Batista, C. H. Booth, D. L. Clark, J. M. Keith, S. A. Kozimor, R. L. Martin, S. G. Minasian, D. K. Shuh, S. Chantal, E. Stieber and T. Tylliszczak, *Chemical Science*, 2020, **11**, 2796–2809.
- 42 K. O. Kvashnina and S. M. Butorin, *Chemical Communications*, 2022, **58**, 327–342.
- 43 R. Caciuffo, G. H. Lander and G. van der Laan, *Rev. Mod. Phys.*, 2023, **95**, 015001.
- 44 K. O. Kvashnina, S. M. Butorin, P. Martin and P. Glatzel, *Physical Review Letters*, 2013, **111**, 253002.
- 45 T. Vitova, I. Pidchenko, D. Fellhauer, P. S. Bagus, Y. Joly, T. Pruessmann, S. Bahl, E. Gonzalez-Robles, J. Rothe, M. Altmaier, M. A. Denecke and H. Geckeis, *Nature Communications*, 2017, **8**, 16053.
- 46 D.-C. Sergentu, T. J. Duignan and J. Autschbach, *The journal of physical chemistry letters*, 2018, **9**, 5583–5591.
- 47 P. S. Bagus, B. Schacherl and T. Vitova, *Inorganic Chemistry*, 2021, **60**, 16090–16102.
- 48 R. Polly, B. Schacherl, J. Rothe and T. Vitova, *Inorganic chemistry*, 2021, **60**, 18764–18776.
- 49 J. Tobin, S. Nowak, S. Yu, R. Alonso-Mori, T. Kroll, D. Nordlund, T. Weng and D. Sokaras, *Journal of Physics: Condensed Matter*, 2022, **34**, 505601.
- 50 C. Silva, L. Amidani, M. Retegan, S. Weiss, E. Bazarkina, T. Graubner, F. Kraus and K. Kvashnina, *Nature Communications*, 2024, **15**, 6861.
- 51 T. G. Burrow, N. M. Alcock, M. S. Huzan, M. A. Dunstan, J. A. Seed, B. Detlefs, P. Glatzel, M. O. J. Y. Hunault, J. Bendix, K. S. Pedersen and M. L. Baker, *Journal of the American Chemical Society*, 2024, **146**, 22570–22582.
- 52 B. Schacherl, M. Tagliavini, H. Kaufmann-Heimeshoff, J. Göttlicher, M. Mazzanti, K. Popa, O. Walter, T. Pruessmann, C. Vollmer, A. Beck, R. S. K. Ekanayake, J. A. Branson, T. Neill, D. Fellhauer, C. Reitz, D. Schild, D. Brager, C. Cahill, C. Windorff, T. Sittel, H. Ramanantoanina, M. W. Haverkort and T. Vitova, *Nature Communications*, 2025, **16**, 1221.
- 53 K. O. Kvashnina, A. Y. Romanchuk, I. Pidchenko, L. Amidani, E. Gerber, A. Trigub, A. Rossberg, S. Weiss, K. Popa, O. Walter,



- R. Caciuffo, A. C. Scheinost, S. M. Butorin and S. N. Kalmykov, *Angewandte Chemie*, 2019, **131**, 17722–17726.
- 54 T. G. Burrow, N. M. Alcock, M. S. Huzan, M. A. Dunstan, J. A. Seed, B. Detlefs, P. Glatzel, M. O. J. Y. Hunault, J. Bendix, K. S. Pedersen *et al.*, *Journal of the American Chemical Society*, 2025, **147**, 37876–37878.
- 55 C. K. Jørgensen, *Orbitals in Atoms and Molecules*, Academic Press, London, United Kingdom, 1962, p. 59.
- 56 G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, *J. Comput. Chem.*, 2001, **22**, 931–967.
- 57 H. Ramanantoanina, W. Urland, F. Cimpoesu and C. Daul, *Physical Chemistry Chemical Physics*, 2013, **15**, 13902–13910.
- 58 M. Atanasov, C. Daul and C. Rauzy, *Optical Spectra and Chemical Bonding in Inorganic Compounds: Special Volume dedicated to Professor Jørgensen I*, 2004, 97–125.
- 59 J. Jung, M. Atanasov and F. Neese, *Inorganic Chemistry*, 2017, **56**, 8802–8816.
- 60 E. C. Wasinger, F. M. De Groot, B. Hedman, K. O. Hodgson and E. I. Solomon, *Journal of the American Chemical Society*, 2003, **125**, 12894–12906.
- 61 J. Griffith and L. Orgel, *Journal of The American Chemical Society*, 1956, 4981–4983.
- 62 J. Ferguson and D. Wood, *Australian Journal of Chemistry*, 1970, **23**, 861–871.
- 63 M. O. J. Y. Hunault, Y. Harada, J. Miyawaki, J. Wang, A. Meijerink, F. M. De Groot and M. M. Van Schooneveld, *The Journal of Physical Chemistry A*, 2018, **122**, 4399–4413.
- 64 A. W. Hahn, B. E. Van Kuiken, M. Al Samarai, M. Atanasov, T. Weyhermüller, Y.-T. Cui, J. Miyawaki, Y. Harada, A. Nicolaou and S. DeBeer, *Inorganic Chemistry*, 2017, **56**, 8203–8211.
- 65 M. M. Gerloch and R. C. Slade, *Ligand-field parameters*, Cambridge University Press, London, 1973, pp. 200–228.
- 66 S. V. Rao, D. Maganas, K. Sivalingam, M. Atanasov and F. Neese, *Inorganic Chemistry*, 2024, **63**, 24672–24684.
- 67 S. G. Minasian, K. S. Boland, R. K. Feller, A. J. Gaunt, S. A. Kozimor, I. May, S. D. Reilly, B. L. Scott and D. K. Shuh, *Inorganic Chemistry*, 2012, **51**, 5728–5736.
- 68 M. Autillo and R. E. Wilson, *European Journal of Inorganic Chemistry*, 2017, **2017**, 4834–4839.
- 69 Bruker Analytical X-ray Systems, *SAINTPlus*, version 6.22, Madison, WI, 2001.
- 70 Bruker Analytical X-ray Systems, *SADABS*, version 2.03, Madison, WI, 2001.
- 71 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 2008, **64**, 112–122.
- 72 P. L. Solari, S. Schlutig, H. Hermange and B. Sitaud, *Journal of Physics: Conference Series*, 2009, **190**, 12042.
- 73 D. A. Pantazis and F. Neese, *Journal of Chemical Theory and Computation*, 2011, **7**, 677–684.
- 74 E. van Lenthe, E. J. Baerends and J. G. Snijders, *The Journal of Chemical Physics*, 1994, **101**, 9783–9792.
- 75 E. van Lenthe, A. Ehlers and E.-J. Baerends, *The Journal of Chemical Physics*, 1999, **110**, 8943–8953.
- 76 E. van Lenthe and E. J. Baerends, *Journal of Computational Chemistry*, 2003, **24**, 1142–1156.
- 77 M. W. Haverkort, M. Zwierzycki and O. K. Andersen, *Physical Review B*, 2012, **85**, 165113.
- 78 J. Campbell and T. Papp, *Atomic Data and Nuclear Data Tables*, 2001, **77**, 1–56.
- 79 F. De Groot and A. Kotani, *Core level spectroscopy of solids*, CRC press, 2008.
- 80 S. M. Butorin, *Inorganic Chemistry*, 2020, **59**, 16251–16264.
- 81 T. G. Burrow, M. O. J. Y. Hunault, F. Besnard, A. Juhin and C. Brouder, *Physical Review B*, 2026, **113**, 115131.



The data supporting this article have been included as part of the Supplementary Information. Numerical data for this paper, including experimental data and calculation files, will be made available on Figshare before publication.

[View Article Online](#)

DOI: 10.1039/D6SC02413K

