

# Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Y. Wei, K. Kasama, A. Igartua, D. B. Yildiz, A. Ceuninck, T. D. Santos, C. Büttner, D. Thevenet, M. Bossart, V. Derdau, M. Mendez-Perez, P. Jubault, T. Poisson, B. Roure and D. Leonori, *Chem. Sci.*, 2026, DOI: 10.1039/D6SC03470E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## ARTICLE

## A photochemical strategy for pyrazole to imidazole conversion

Youhao Wei,<sup>a#</sup> Kengo Kasama,<sup>a,b#</sup> Antòn Igartua,<sup>a</sup> Dilara Berna Yildiz,<sup>a,c</sup> Aurore Ceuninck,<sup>d</sup> Thiago dos Santos,<sup>a</sup> Cornelia Büttner,<sup>a</sup> Damien Thevenet,<sup>e</sup> Martin Bossart,<sup>f</sup> Volker Derdau,<sup>f</sup> María Méndez,<sup>f</sup> Philippe Jubault,<sup>d</sup> Thomas Poisson,<sup>d\*</sup> Baptiste Roue,<sup>a\*</sup> Daniele Leonori<sup>a\*</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Heteroaromatic scaffolds are central to modern medicinal chemistry. Methods that can reconfigure the core heterocycle of a molecule while preserving its substitution pattern would greatly streamline analogue synthesis and bioisosteric replacement. Yet, direct heterocycle-to-heterocycle interconversions remain rare. Here we report a photochemical strategy that directly converts pyrazoles into imidazoles with broad functional-group tolerance and full retention of peripheral substitution. The reaction is effective across densely substituted and bicyclic systems and extends to pyrazolo[1,5-a]azines, a class of high-value heteroaromatics that have never previously been reconfigured. This photochemical strategy is readily translated to continuous flow, confirming its potential for scalable applications. Our mechanistic studies support an N–N bond homolysis pathway in which solvent-dependent conformational changes govern the reactivity of the ensuing bi-nitrogen-radical intermediates. Overall, this work establishes a practical platform for direct core reconfiguration, providing modular access to imidazole analogues of pyrazoles that are otherwise difficult to prepare or very expensive.

## Introduction

Heteroaromatic scaffolds are ubiquitous in bioactive molecules, where the identity and spatial arrangement of their heteroatoms and ring-substituents play a decisive role in governing binding interactions, physicochemical properties, and metabolic stability.<sup>1</sup> As a result, medicinal chemists routinely explore closely related heterocyclic cores to fine-tune biological activity and selectivity.<sup>2</sup> However, modifying a heteroaromatic framework while preserving its substitution pattern remains a major synthetic challenge. In practice, such transformations typically require complete *de novo* synthesis, slowing structure–activity relationship (SAR) campaigns and limiting access to promising regions of chemical space.<sup>3</sup> Direct heterocycle-to-heterocycle interconversions would provide a powerful solution, provided they can operate with broad functional group tolerance and high fidelity in retaining substitution patterns.<sup>4</sup>

Nitrogen-containing heteroaromatics are among the most prevalent motifs in drug discovery. In particular, pyrazoles and imidazoles rank among the most frequently encountered five-

membered heterocycles in approved pharmaceuticals (Scheme 1A).<sup>5</sup> Although they share a common ring size and contain two nitrogen atoms, their electronic structures and physicochemical properties differ substantially. Imidazoles possess a pyridine-like nitrogen that confers higher basicity and stronger hydrogen-bond acceptor ability, whereas the adjacent nitrogen arrangement in pyrazoles results in lower basicity and pronounced tautomerism.<sup>6</sup> These differences translate into distinct binding modes, pharmacokinetic profiles, and biological activities.<sup>7</sup> Consequently, strategic pyrazole→imidazole replacement is a common medicinal chemistry tactic to fine-tune biological properties (Scheme 1B). Prominent recent examples include the development of aplithianines, chimeric J-PKAc kinase inhibitors with potential applications in carcinoma treatment,<sup>8</sup> and RP-1664, the first-in-class orally available PLK4 inhibitor.<sup>9</sup> Importantly, such scaffold modifications are not limited to isolated pyrazole/imidazole rings but also extend to annulated systems, such as pyrazolo[1,5-a]pyridine→imidazo[1,2-a]pyridine transformations,<sup>10</sup> which, for example, was crucial to the development of the anti-tuberculosis drug telacebec.<sup>11</sup>

Despite their close structural relationship, no general and practical method exists for the direct interconversion of pyrazoles into imidazoles. Early studies by Barltrop, Pavlik, and co-workers established that UV irradiation of simple pyrazoles can induce rearrangement processes.<sup>12</sup> These seminal investigations provided important mechanistic insight into the photochemical behaviour of azoles, but the reactions typically resulted in complex mixtures, low conversions, and significant decomposition, with product formation often inferred indirectly from mass balance rather than isolation. Consequently, these transformations have not been adopted as synthetic methods, and current approaches to access

<sup>a</sup> Institute of Organic Chemistry, RWTH Aachen University, Landoltweg 1, 52054 Aachen, Germany.

<sup>b</sup> Faculty of Pharmaceutical Sciences, University of Toyama, Sugitani, Toyama 930-0194, Japan.

<sup>c</sup> Department of Chemistry, Faculty of Science, Gazi University, Teknikokullar, Türkiye.

<sup>d</sup> INSA Rouen Normandie, Univ. Rouen Normandie, Univ. Caen Normandie, ENSICAEN, CNRS, Institut CARMEN UMR 6064, F-76000 Rouen, France

<sup>e</sup> Minakem Recherche, 145 Chemin des Lilas, 59310 Beuvry-La-Forêt, France

<sup>f</sup> Integrated Drug Discovery, R&D, Sanofi Germany, Industriepark Hoechst, 65926 Frankfurt am Main, Germany


# These authors contributed equally to this work.



imidazoles from pyrazole precursors rely instead on stepwise construction from acyclic building blocks.

Our group has recently started to work on establishing the concept of photochemical permutation as a potential strategy

to streamline heteroatomic synthesis and bioactive molecule diversification.<sup>13</sup> In this approach, selective excitation of a



**Scheme 1.** A. Pyrazole and imidazole are two relevant N-heteroaromatics, but their N-atom dispositions provide key chemical difference. B. Examples of pyrazole $\rightarrow$ imidazole conversions leading to improved bioactive molecules. C. Previous permutation work on other azoles features mechanistically distinct pathways and intermediates. D. This work demonstrates a direct pyrazole $\rightarrow$ imidazole conversion that can be applied to pyrazolo[1,5a]azines.

heteroaromatic framework enables direct reconfiguration into a different heteroaromatic while preserving its substitution pattern. Importantly, these transformations do not operate through a unified mechanism but instead proceed through distinct excited-state manifolds that depend on the heterocycle type. For example, thiazoles & isothiazoles undergo reversible rearrangement via 4 $\pi$ -electrocyclization to Dewar intermediates followed by "sulfur-shift".<sup>13a</sup> In contrast, indazoles require excited-state tautomerization prior to 4 $\pi$ -electrocyclization, imposing strict constraints on substitution patterns as only NH-derivatives with C3 alkyl groups can be engaged.<sup>13c, 14</sup> Isoxazoles proceed through N–O bond cleavage leading to vinylnitrene intermediates that display strong dependence on further chromophore-controlled excitation for


oxazole formation whereby a C5 aromatic group is necessary in order to observe high reactivity.<sup>15</sup> These findings highlight that photochemical heterocycle permutation is governed by multiple, mechanistically distinct regimes rather than a single general pathway. Hence, no general framework can be used to generalize reactivity, scope and mechanism but rather a case-by-case study is required to inform on all these aspects. In this context, we questioned whether pyrazoles would follow any of the previously identified paradigms or instead access a distinct reactivity mode. In this full paper, we report a general and operationally simple photochemical strategy for the direct conversion of pyrazoles into imidazoles (Scheme 1D). The transformation proceeds under mild, additive-free conditions and accommodates a wide range of substitution patterns,



including mono-, di-, and trisubstituted pyrazoles as well as diverse annulated systems. Notably, the method enables the first direct reconfiguration of pyrazolo[1,5-a]azines, despite the presence of competing chromophores that would be expected to dominate photoexcitation. Mechanistically, this pyrazole→imidazole conversion requires a fine interplay between excited and high-energy intermediates. Our studies revealed the process proceeds via N–N bond homolysis to an iminyl/aminyl diradical species followed by conformationally controlled radical cyclization, with solvent polarity and H-bonding playing decisive roles in product selectivity. Such mechanism is intrinsically irreversible and, in contrast to the indazole→benzimidazole case, does not require tautomerization, thus allowing broader tolerance of N-substitution patterns and expanded substrate scope. The reaction can be translated to continuous-flow conditions, underscoring its practical potential for scalable applications. Collectively, these findings establish a distinct mechanistic regime for heterocycle permutation and provide a direct entry to imidazole analogues that are otherwise difficult to access.

## Results

**Reaction Development.** We initiated our studies by examining the photochemical conversion of *N*-Ph-pyrazole **1a** into imidazole **1b**, which can be visualized as a direct “positional swap” between the N atom and the C2 unit (Scheme 2). All reactions were carried out at room temperature under dilute conditions (0.025 M) using  $\lambda = 254$  nm as the irradiation source. Notably, the reaction displayed a pronounced solvent dependence, providing early evidence for the involvement of high-energy intermediates whose fate is strongly influenced by the reaction medium. Specifically, the reaction proceeded in polar solvents such as CH<sub>3</sub>CN (entry 1) and 1,4-dioxane (entry 2), affording **1b** in moderate yield but along the ring-opened by-product **1c**. When DMSO (entry 3) or *i*-PrOH (entry 4) were employed, complete material decomposition was observed. A notable shift in product distribution was achieved using the more Brønsted acidic solvent TFE (entry 5), which favored the formation of **1b** over **1c**, albeit in moderate efficiency. This effect was further amplified with HFIP (entry 6), that provided **1b** in 70% yield with no detectable formation of **1c**. This solvent effect can be potentially rationalized with stabilization of reactive intermediates through H-bonding interactions. The reaction efficiency improved significantly upon reducing the irradiation time (entries 7 and 8), with 1 h being optimum. In contrast, irradiation at  $\lambda = 310$  nm (entry 9) resulted in no product formation, likely due to insufficient absorption at this wavelength. A comprehensive summary of the solvent screening and additional optimization studies is provided in the Supplementary Information.



**Scheme 2.** Optimization of the pyrazole-to-imidazole conversion using **1a**.

**Reaction Scope.** With optimized conditions in hand, we examined the influence of substitution on the pyrazole scaffold to delineate the structural features governing reactivity (Scheme 3). For each substrate, we screened five different conditions that varied in solvent composition (A–D) and irradiation wavelength (E). UV/vis absorption spectroscopy studies revealed that nearly all substrates absorb in the UV-B region, so we performed all reactions at room temperature under  $\lambda = 254$  and 310 nm irradiation. In general, HFIP (conditions A and E) was identified as the optimum medium for various of these examples, albeit specific substrates performed better in other solvents.

*N*-Aryl pyrazoles (**1a–5a**) showed fast and efficient reactivity under these conditions. The conversion of **2a** into **2b** converts a cheap material (**2a**: 61 \$/g) in a rather expensive building block (**2b**: 426 \$/g). Interestingly, the *para*-Ph derivative (**3a**) gave significantly lower yield, which we attribute to competitive excitation of the extended aromatic system. Given the frequent presence of azine-*N*-substituted azoles in bioactive molecules, we next examined substrates bearing C4, C3, and C2 pyridine substituents (**6a–8a**). These were successfully engaged in the permutation reaction giving **6b–8b**, albeit with moderate to low efficiency.


We also identified several *N*-groups incompatible with the permutation chemistry. In general, we believe reactivity to be governed by the ability of the pyrazole core to compete for photoexcitation with other chromophores present. Indeed, the strongly absorbing *N*-1-naphthyl unit (**9a**) led to full starting material recovery due to its dominant light absorption, which outcompetes the pyrazole core. *N*-alkyl (e.g., Me, **10a**), *N*-Bz (**11a**), and *N*-Ts (**12a**) substituents resulted in no reactivity and significant *N*-deprotection in the case of **12a**. As we will discuss below, *N*-alkyl pyrazoles can be transformed into *N*-alkyl imidazoles upon simple substitution on the azole nucleus (see Scheme 4). We therefore attribute the lack of reactivity of **10a** primarily to its insufficient absorption in the irradiation range employed.



We next investigated the influence of different substituents on the *N*-Ph pyrazole scaffold. Introduction of Me (**13a–15a**) and ester (**16a–18a**) groups at the C3, C4, and C5 positions afforded the corresponding imidazoles (**13b–17b**) in good yields, except for **18a**, which gave **18b** with low efficiency. Electron-donating OMe (**19–21a**) and electron-withdrawing CF<sub>3</sub> (**22–24a**) groups were also examined. Imidazole formation was successfully achieved for the C3 (**19a** and **22a**) and C4 (**20a** and **23a**) isomers, providing the desired products (**19b**, **20b**, **22b** and **23b**) in good to moderate yields. In contrast, C5-substituted derivatives (**21a** and **24a**) proved unreactive.

Introduction of a Ph group on the pyrazole core induced a bathochromic shift in the absorption profile, and these reactions were therefore conducted using  $\lambda = 310$  nm irradiation. Under these conditions, a marked positional effect

was observed. Specifically, C3-Ph substitution (**25a**) resulted in full mass recovery with no product formation, whereas C4-Ph (**26a**) and C5-Ph (**27a**) derivatives afforded the corresponding imidazoles (**26b** and **27b**) in excellent and moderate yields, respectively. These observations underscore the strong sensitivity of the reaction to structural changes in the heteroaromatic that have strong impact in excited-state dynamics. This reinforces the need for experimental guidelines as *a priori* generalization of reaction scope can be challenging. Interestingly, the presence of the Ph group on the pyrazole core enabled reaction on some *N*-Me substituted systems. Specifically, while **28a** did not lead to productive rearrangement (which is in line with **25a**), **29a** and **30a** gave **29b** and **30b** in high and low yields respectively.



**Scheme 3.** Substrate scope for the pyrazole-to-imidazole conversion using mono-substituted derivatives. E = CO<sub>2</sub>Me. The yield for substrate **12** corresponds to the detosylated product.

Hence, using the C4-Ph substitution pattern as a benchmark, we re-evaluated the compatibility of various *N*-substituents in the permutation reaction (Scheme 4). Both *para*-substituted electron-rich (**31a**) and electron-poor (**32a**) aryl groups, as well as *ortho*-substituted variants (**33a**), were well tolerated,

delivering products **31–33b** in high yields. Notably, *N*-vinyl (**34a**) and *N*-alkynyl (**35a**) substrates also participated, affording **34b** and **35b** in moderate and good yields, respectively. *N*-Bz pyrazole (**36a**) led to complete mass recovery, but both *N*-Bn and NH derivatives (**37a** and **38a**) were successfully engaged in




the reaction, giving the corresponding imidazoles (**37b** and **38b**).

Building on these examples, we next examined the effect the ester substituents on the reactivity of *N*-Me pyrazole core. In this case, a different trend was observed as both the C3 (**39a**) and C5 (**41a**) derivatives underwent efficient transformation, providing the **39b** and **41b** in high yields, while the C4 derivative (**40a**) was unreactive.

Finally, we assessed the impact of other types of substituents using *N*-Ph substituted pyrazoles. In this case, we successfully demonstrated compatibility with a series of polar functionalities spanning C4-amide (**42a**), -nitrile (**43a**), hydroxymethyl (**44a**) and -fluorine (**45a**). Interestingly, substrate **46a** features two

pyrazoles unit and lead exclusively to the rearrangement of the *N*-Ph substituted fragment providing **46b** in high yield. We believe this isomerization selectivity to be influenced by the *N*-Ph substituents that likely modulates the photochemical behavior of the attached pyrazole core (see below for comparison the high reactivity of **51a**).

Importantly, most pyrazoles included in this study are commercially available and inexpensive, whereas the corresponding imidazoles typically require multi-step synthesis and are often unavailable from vendors (see Supplementary material). The transformation therefore represents a direct and practical entry to valuable heterocycles that would otherwise demand significant synthetic and/or economic investment.



**Scheme 4.** Substrate scope for the pyrazole-to-imidazole conversion using mono-substituted derivatives. E = CO<sub>2</sub>Me.

We next examined the scope of the transformation on disubstituted pyrazoles to assess its compatibility with increased substitution density (Scheme 5). Using *N*-Ph substitution pattern, we investigated a series of derivatives bearing either two Ph groups at C3,C4 (**47a**) and C4,C5 (**48a**), mixed Ph and Me substituents (**49a** and **50a**), two Me groups (**51a**), or Me and ester functionalities (**52a** and **53a**). In all cases, the desired disubstituted imidazoles (**47b–53b**) were obtained in good to excellent yields. Notably, the method was also applicable to a substrate push-pull system based on a C4 nitrile and a C5 free amine functionality (**54a**), affording **54b** in moderate yield.

We then turned our attention to NH-pyrazoles to probe the influence of substituents in the absence of *N*-protection or substitution. Here, substrate availability was more restricted because certain isomers cannot be isolated due to rapid


tautomerization.<sup>5a, 7a</sup> Among the accessible derivatives (**55a–60a**), both Me- and Ph-substituted pyrazoles underwent efficient conversion, delivering the corresponding imidazoles **55b–57b** in good yields. In contrast, among the ester-containing substrates, only the C3-ester/C4-methyl pyrazole **58a** afforded the desired product **58b**, while the remaining analogues **59a** and **60a** were unreactive. Finally, we demonstrated that the method tolerates further substitution as trisubstituted *N*-Me pyrazole **61a** underwent smooth rearrangement to give **61b** in good yield.

Having established the generality of the process on mono-, di-, and trisubstituted systems, we next explored its applicability to annulated pyrazoles, which are easily accessible through condensation chemistry.<sup>16</sup> These substrates are synthetically valuable, as the corresponding imidazoles are typically not commercial and more challenging to prepare. Hence, their



direct conversion would offer a strategic shortcut to privileged building blocks suitable for library synthesis and high-throughput screening. Pleasingly, the methodology proved compatible with a variety of bicyclic frameworks annulated at C4 and C5. Successful examples included both carbocyclic systems (**62a** and **63a**) and heterocycles based on *N*-Boc-

piperidine (**64a**), *N*-Boc-pyrrolidine (**65a**), tetrahydropyran (**66a**), and tetrahydrofuran (**67a**) that gave **62b–67b** in high yields. The method was also effective for the synthesis of bicyclic trisubstituted imidazoles, as exemplified by the conversion of **68a** into **68b** in good yield.



**Scheme 5.** Substrate scope for the pyrazole-imidazole conversion using di- and tri-substituted derivatives. <sup>A–D</sup> see Scheme 3 for the legend with the specific conditions used.

Finally, we assessed the applicability of the transformation to complex, bioactive molecules. Stanozolol (**69a**), a drug used clinically for the treatment of hereditary angioedema, contains a pyrazole core annulated across C3 and C4 onto a steroidal framework. Under our optimized conditions, it was directly and efficiently converted into **69b** in 40% yield. This compound had

previously required a four-step sequence during the drug development campaign,<sup>17</sup> an effort now bypassed in a single step by our methodology. Furthermore, withasomnine (**70a**) and iso-withasomnine (**71a**), two natural alkaloids isolated from the roots of the ashwagandha plant,<sup>18</sup> feature a bicyclic linkage between C5 and the pyrazole nitrogen. Both substrates were



smoothly converted to the corresponding imidazoles (**70b** and **71b**) in good and moderate yields, respectively, thus enabling the fast preparation of natural product-like isomers.

Also in this case, the methodology enabled direct access to building blocks that are either extremely expensive (e.g. **54a**: 10 \$/g while **54b**: 2370 \$/g) or currently not commercially available and require individual synthesis (see supplementary material for more information).


The successful conversion of pyrazoles into imidazoles across bicyclic systems prompted us to evaluate whether this reactivity could be extended to more complex heterocyclic frameworks. In particular, we targeted pyrazolo[1,5-a]azines, a class of motifs frequently embedded in bioactive molecules and blockbuster drugs.<sup>10b</sup> However, these systems present a different photochemical scenario compared to simple pyrazoles. The presence of an annulated azine introduces a competing chromophore, which typically displays stronger absorption and would be expected to dominate photoexcitation. Moreover, such substrates have not been previously investigated in the context of photochemical permutation, raising the possibility that they may follow entirely different excited-state pathways or fail to undergo productive rearrangement altogether. We therefore questioned whether selective excitation of the pyrazole unit could still be achieved in these systems, or whether the intrinsic photochemical behaviour of the azine would suppress or divert reactivity.

We initiated our investigation with a pyridine derivative **72a**, which underwent smooth conversion under standard conditions B to the corresponding **72b** in good yield (Scheme 6). Notably, the reaction tolerated various substitution patterns, including a C2-methyl group (**73a**) and C2- and C3-ester (**74a** and **75a**), affording **72b–75b** in high to excellent yields.

The rearrangement was next applied to pyrimidine derivatives **76a** and **77a**, bearing C2- and C3-ester as well as nitrile

substituents (**78a**), which gave **76b–78b** in moderate yields. Pyrazolo[1,5-a]pyrimidines are common scaffolds in bioactive molecules. Notably, **79a** is used in the manufacture of the anticancer agents dinaciclib via  $S_NAr$  chemistry at the activated aryl chloride functionalities.<sup>19</sup> Pleasingly, these groups were fully tolerated under our conditions, delivering **79b** in useful yield, which can be integrated into existing synthetic routes for the preparation of drug analogues. We further examined the scope of the reaction with pyrazine and pyridazine derivatives **80a** and **81a**, which afforded **80b** and **81b** in high and moderate yields, respectively. It is important to note that also cross these azine classes, permutation provided direct access to imidazo[1,5-a]azine scaffolds that are either prohibitively expensive or entirely unavailable commercially (e.g., **76a**: 78 \$/g, while **76b**: 1962 \$/g). This underscores the practical value of the method that converts widely available building blocks into elusive analogues in a single step.

Given the prevalence of pyrazolo[1,5-a]azine motifs in bioactive molecules, we next evaluated the potential of this method for late-stage ring reconfiguration of complex substrates. Zaleplon (**82a**), a hypnotic agent used for the treatment of severe insomnia, features a C3-nitrile-substituted pyrazolo[1,5-a]pyrimidine core with a pendant amide-substituted benzene ring. Under our photochemical conditions, we successfully obtained the corresponding imidazole derivative **82b** in 23% yield. Furthermore, the structurally complex anticancer drug selpercatinib (**83a**) underwent selective isomerization of its pyrazolo[1,5-a]pyridine core, delivering the rearranged product **83b** in 12% yield. Although the conversions in these two cases are currently moderate, they demonstrate the ability of this methodology to reconfigure core heterocyclic frameworks in highly functionalized molecules in a single step, thus bypassing the need for lengthy and resource-intensive total syntheses.



**Scheme 6.** Substrate scope for the pyrazole to imidazole conversion using pyrazolo[1,5-a]azine derivatives. <sup>A–D</sup> see Scheme 3 for the legend with the specific conditions used.



## ARTICLE


**Reaction scalability.** To demonstrate that the rearrangement is not limited to small-scale batch photolysis, we translated the reaction into a continuous-flow photochemical process (Scheme 7). Using **1a** as the benchmark substrate, initial settings closely mirrored the batch setup (HFIP,  $\lambda = 254$  nm, 0.0125 M). Under these conditions, a 10 mL UV reactor afforded **1b** in 81% yield with 19% recovered **1a** at 20 °C and a 20 min residence time (entry 1). Optimization revealed that reaction temperature was the key determinant of throughput. Increasing the reactor temperature to 70 °C while reducing the residence time to 11 min delivered complete conversion (entry 2). This temperature effect was fully consistent with the reaction's photochemical nature and not attributable to thermal activation. To reduce solvent cost while maintaining efficiency, we replaced HFIP with a 1:1 HFIP/CH<sub>2</sub>Cl<sub>2</sub> mixture, which delivered the same conversion and identical quantum yields ( $\Phi = 0.064$  vs 0.063),<sup>20</sup> and afforded **1b** in 79% yield and a productivity of 0.078 g/h (entry 3). Reactivity was maintained upon doubling the substrate concentration (96% yield, 0.094 g/h productivity), albeit at a slightly longer residence time (22 min), demonstrating that the transformation is not inherently limited by light penetration under these conditions (entry 4).

A second representative substrate, **26a**, also reached full conversion to the imidazole product **26b** under similar conditions, with a 30 min residence time (productivity 0.055 g/h). The robustness of the method across substrates confirms that the rearrangement is flow-compatible, scalable, and operationally simple, suitable for both library synthesis and gram-scale analogue production.

**Mechanistic proposal.** In principle, two mechanistic pathways, both originating from the S<sub>1</sub>-excited pyrazole (S<sub>1</sub>-**a**), can account for the observed transformation (Scheme 8a). By analogy with our previous work on thiazole/isothiazole and indazole/benzimidazole interconversions,<sup>13a, 13c</sup> an excited-state 4 $\pi$  electrocyclicization could form a Dewar intermediate **d**, which after an N-shift would furnish intermediate **e** and ultimately the imidazole **b'**. Alternatively, direct N–N bond homolysis<sup>21</sup> could generate a di-N-radical **f**. This species would afford, upon cyclization, the imine-azirine intermediate **g** and, subsequently, the methaniminium species **h**, capable of delivering **b** by 6 $\pi$  electrocyclic ring closure.

Although both pathways converge to the same imidazole framework, they involve fundamentally different bond rearrangements with clear structural consequences. The radical pathway leading to **b** can be viewed as an “N–C3 swap,” preserving the original C4–C5 connectivity. In contrast, formation of a Dewar intermediate **d** would entail complete positional reorganization of the carbon framework in **b'**. Our

experimental results show no evidence of positional scrambling, providing compelling support that the reaction proceeds through N–N bond cleavage rather than Dewar-type electrocyclicization. To further support the proposed homolytic mechanism experimentally, C5- and C3-deuterated **1a** were prepared and subjected to the photochemical conditions (Figure 7b). In both cases, **1b** was obtained in 70% yield. However, complete deuterium retention was observed for C5-labelled **1a**, whereas C3-labelled **1a** furnished non-deuterated **1b**. This latter outcome is consistent with the intermediacy of methaniminium species **h**, from which C3-deuterium labeling is lost.



**Scheme 7.** Reaction scale-up using photo-flow equipment. <sup>a</sup> Reaction conducted at 0.025 M instead of 0.0125 M. nd: not determined.

To obtain further evidence on the reaction mechanism we conducted computational studies using *N*-Ph pyrazole **1a** as the model (Scheme 8b). Upon photoexcitation this species populates the S<sub>1</sub> state ( $E = 122.4$  kcal mol<sup>-1</sup>), which has  $\pi, \pi^*$  character. From this excited state, a conical intersection (**CX1**) was identified at  $E = 88.6$  kcal mol<sup>-1</sup>. This structure features a complete breakage of the N–N bond and therefore funnels the



system into N–N bond homolysis pathway, generating the diradical intermediate **A1**. In HFIP solvent, this intermediate can cyclize to **A2** ( $\Delta G^\circ = -20.0 \text{ kcal mol}^{-1}$ ) via a shallow transition state ( $\Delta G^\ddagger = 2.8 \text{ kcal mol}^{-1}$ ). We initially hypothesized that **A2** might thermally interconvert into **A3**.<sup>40</sup> However, our calculations revealed that this pathway is associated with a prohibitively high kinetic barrier ( $\Delta G^\ddagger = 46.4 \text{ kcal mol}^{-1}$ ), rendering it inaccessible under the experimental conditions. Instead, time-dependent DFT studies demonstrated that **A2** absorbs light in the same wavelength range as our irradiation source. Hence, we propose that a second photochemical event, involving **A2** could populate the excited state  $S_1$ -**A2** ( $E = 104.9 \text{ kcal mol}^{-1}$ ), which subsequently evolves through a conical intersection (**CX2**,  $E = 68.6 \text{ kcal mol}^{-1}$ , see Supplementary material) to form **A3**. From this intermediate, a  $6\pi$  electrocyclization would deliver **1b** in a kinetically facile and exothermic manner ( $\Delta G^\ddagger = 4.9 \text{ kcal mol}^{-1}$ ;  $\Delta G^\circ = -62.9 \text{ kcal mol}^{-1}$ ). Thus, in contrast to our previous work on thiazoles and indazoles, which proceed via pericyclic excited-state pathways, or isoxazoles, which proceed through the formation of a nitrene intermediate, the present transformation operates through N–N bond homolysis via di-N-radical species, establishing a distinct mechanistic regime for this heterocycle.

We then became interested in understanding the crucial role of HFIP in selectively providing access to the desired imidazole product. Indeed, as discussed above, when switching from H-bond-donating HFIP to 1,4-dioxane, the nitrile by-product **1c** was formed in a 1:1 ratio with **1b** (see Scheme 2). We propose this solvent dependence can be rationalized by conformational effects on **A1**. In HFIP, **A1** should be stabilized by H-bonding interactions in an *s-cis* conformation that can readily cyclize to **A2**. In contrast, in 1,4-dioxane, a C–C bond rotation leads to a nearly isoenergetic *s-trans* conformer **A1'**. Crucially, this isomerization has the same kinetic barrier of the cyclization to **A2**, so we propose can compete with it. Species **A1'** is unable to cyclize and instead can undergo 1,4-HAT between the iminyl-type radical and the acyl-type  $C(sp^2)$ –H bond to give by-product **1c**. Although this HAT step exhibits a high calculated barrier, it is characterized by a large imaginary frequency, indicative of significant quantum tunnelling,<sup>22</sup> which we propose can enable the process to compete under the reaction conditions. Overall, it is noteworthy that the striking solvent effect exerted by HFIP in both the imidazole→benzimidazole and pyrazole→imidazole permutations arises from fundamentally different mechanistic origins. In the case of NH-indazoles, the acidity of HFIP is required to promote excited-state tautomerization,<sup>13c</sup> whereas for pyrazoles it plays a key role in locking the conformation of the resulting di-N radical species **A1**.

The formation of by-product **1c** warrants further discussion. According to our mechanistic proposal, this species arises from a parasitic pathway that diminishes the efficiency of the desired pyrazole→imidazole conversion when  $\text{CH}_3\text{CN}$  solvent is used. Consistent with this, irradiation of an authentic sample of **1c** did not yield **1b**, but instead led to partial  $Z \rightarrow E$  olefin isomerization

(Figure 1D). However, the substitution pattern of this species can significantly influence its photochemical behaviour. This was demonstrated by the preparation of **53c** via irradiation of **53a** in  $\text{CH}_3\text{CN}$ , which afforded **53b** (22%) and **53c** (44%). Further irradiation of **53c** produced **53b** in 23% yield, accompanied by olefin isomerization. Therefore, we cannot exclude that, in certain cases, the pathway leading to the enamine–nitrile by-products may also positively contribute to the overall productivity of the process. These observations reinforce the notion that photochemical permutation mechanisms are highly sensitive to heteroaromatic substitution patterns. Such behavior is difficult to rationalize based on current structure–reactivity paradigms and highlights a gap in our understanding of the excited-state behavior of these heteroaromatic systems.

## Conclusions

We have developed a general photochemical strategy that transforms pyrazoles into imidazoles in a single step and fully preserve the substitution pattern of the parent heterocycle. The method tolerates a wide spectrum of electronic and steric environments, operates reliably across densely substituted and annulated systems, and, critically, enables the first direct reconfiguration of pyrazolo[1,5-*a*]azines. The successful translation of the reaction to continuous flow underscores its practical potential for scalable analogue generation. Our mechanistic studies suggest the heterocycle rearrangement is initiated by N–N bond homolysis and controlled by H-bonding interactions with the solvent. Taken together, these findings establish a blueprint for direct core reconfiguration as a versatile tool for accessing otherwise inaccessible heterocycles and accelerating bioisosteric exploration in medicinal chemistry.

## Author contributions

B.R. and D.L. designed the project. Y.W., K.K., A.I., T.d.S. and C.B. performed all synthetic experiments. D.B.Y. performed all computational studies. T.P. designed the flow experiments and quantum yield determinations. A.C. performed all flow experiments. All authors analysed the results and wrote the manuscript.

## Conflicts of interest


M. B., V. D. and M. M. are employees of Sanofi and might have shares and/or stock options in the company. The other authors do not declare competing interests.

## Data availability

All data is available in the Supporting Information.



## ARTICLE



**Scheme 8. A.** Proposed mechanistic pathway for the pyrazole-imidazole conversion and computational studies. **B.** Experiments on deuterated **1a**. **C.** Computational profile for the conversion of **1a** into **1b** and **1c**. All energy values are in kcal mol<sup>-1</sup>. **D.** Photochemical experiments on the reactivity of enamine nitriles by-products.



## ARTICLE

## Acknowledgements

T.d.S. thanks the Alexander von Humboldt – CAPES for a fellowship (88881.699295/2022-01). C.S.B. thanks the Marie Curie Actions for a Fellowship (B-STRAIN 101102819). D.B.Y. acknowledges the financial support of the Study Abroad Postgraduate Education Scholarship (YLSY) awarded by the Republic of Türkiye Ministry of National Education. Computations were performed with computing resources granted by RWTH Aachen University under projects RWTH1268 and P0021519, and with the High Performance and Grid Computing Center (TRUBA resources) at TUBITAK ULAKBIM. A. C. thanks INSA Rouen Normandie for funding through the COMP. T. P. acknowledge the PHOTODERACS program (Normandy Region, no. 0015464) for funding. A.C., P.J. and T.P. acknowledge the API program and NFC (NormandyFlowChem platform) supported by the European Union through the operational program (Normandy Region, ERDF/FSE 2014–2020, no. 20E04976), and FNADT-DRACCARE (program 2020–2023 (N)15179) for funding. Normandie Univ. (NU), Région Normandie, CNRS, Univ. Rouen Normandie, INSA Rouen Normandie, Labex SynOrg (ANR-11-LABX0029), the graduate school for research XL-Chem (ANR-18- EURE-0020 XL CHEM) and Innovation Chimie Carnot (I2C) are acknowledged for support.

## Notes and references

# These authors contributed equally to this work.

- (a) S. D. Roughley and A. M. Jordan, *J. Med. Chem.*, 2011, **54**, 3451-3479; (b) J. Shearer, J. L. Castro, A. D. G. Lawson, M. MacCoss and R. D. Taylor, *J. Med. Chem.*, 2022, **65**, 8699-8712.
- (a) N. A. Meanwell, *J. Med. Chem.*, 2011, **54**, 2529-2591; (b) M. A. M. Subbaiah and N. A. Meanwell, *J. Med. Chem.*, 2021, **64**, 14046-14128; (c) P. Ertl, E. Altmann, S. Racine and R. Lewis, *European Journal of Medicinal Chemistry*, 2022, **238**, 114483.
- (a) D. C. Blakemore, L. Castro, I. Churcher, D. C. Rees, A. W. Thomas, D. M. Wilson and A. Wood, *Nat. Chem.*, 2018, **10**, 383-394; (b) K. R. Campos, P. J. Coleman, J. C. Alvarez, S. D. Dreher, R. M. Garbaccio, N. K. Terrett, R. D. Tillyer, M. D. Truppo and E. R. Parmee, *Science*, 2019, **363**, 244.
- (a) B. W. Joynson and L. T. Ball, *Helv. Chim. Acta*, 2023, **106**, e202200182; (b) J. Jurczyk, J. Woo, S. F. Kim, B. D. Dherange, R. Sarpong and M. D. Levin, *Nature Synthesis*, 2022, **1**, 352-364; (c) G. L. Bartholomew, L. J. Karas, R. M. Eason, C. S. Yeung, M. S. Sigman and R. Sarpong, *J. Med. Chem.*, 2025, **68**, 6027-6040.
- (a) G. Li, Y. Cheng, C. Han, C. Song, N. Huang and Y. Du, *RSC Medicinal Chemistry*, 2022, **13**, 1300-1321; (b) E. Vitaku, D. T. Smith and J. T. Njardarson, *J. Med. Chem.*, 2014, **57**, 10257-10274; (c) A. Siwach and P. K. Verma, *BMC Chemistry*, 2021, **15**, 12; (d) C. M. Marshall, J. G. Federice, C. N. Bell, P. B. Cox and J. T. Njardarson, *J. Med. Chem.*, 2024, **67**, 11622-11655.
- I. I. Gerus, R. X. Mironetz, I. S. Kondratov, A. V. Bezdudny, Y. V. Dmytriv, O. V. Shishkin, V. S. Starova, O. A. Zaporozhets, A. A. Tolmachev and P. K. Mykhailiuk, *The Journal of Organic Chemistry*, 2012, **77**, 47-56.
- (a) J. V. Faria, P. F. Vegi, A. G. C. Migueta, M. S. dos Santos, N. Boechat and A. M. R. Bernardino, *Biorg. Med. Chem.*, 2017, **25**, 5891-5903; (b) L. Zhang, X.-M. Peng, G. L. V. Damu, R.-X. Geng and C.-H. Zhou, *Medicinal Research Reviews*, 2014, **34**, 340-437.
- L. Du, B. A. P. Wilson, W. J. Moore, M. Dalilian, S. R. Shenoy, N. Li, J. A. Martinez Fiesco, J.-Y. Hwang, E. A. Smith, A. Wamiru, E. I. Goncharova, A. Alvarez de la Cruz, K. Pagadala, H. K. Piswa, V. Pattet, V. P. Jampana, P. Manepalli, R. Nimmala, N. R. Marri, M. Gunuguntla, J. J. Reddy, J. S. Schneekloth, Jr., P. Zhang and B. R. O'Keefe, *J. Med. Chem.*, 2025, **68**, 12756-12785.
- F. Vallée, M. Casás-Selves, M. Bubenik, M. Duplessis, B. Sow, C. Suarez, B. Sangiorgi, L. Li, M. Hyer, R. Papp, M.-E. Leclaire, A. L. Perryman, B. Liu, S. Surprenant, P. Mochirian, V. Pau, Z. Maderova, P. Mader, S. Y. Yin, E. Goodfellow, A. Roulston, R. Stocco, C. Godbout, P. Baruah, A. Bonneu-Fortin, J. D. Schonhoft, P. Nejad, D. Norman, V. L. Truong, S. Crane, M. A. Attia, D. Mao, F. Sicheri, C. G. Marshall, M. Zimmermann, D. Bendahan, M. Gallant and W. C. Black, *J. Med. Chem.*, 2025, **68**, 10631-10647.
- (a) D. Devi Priya, M. Nandhakumar and S. Mohana Roopan, *Synth. Commun.*, 2020, **50**, 3535-3562; (b) E.-G. Cecile and G. Alain, *Mini-Reviews in Medicinal Chemistry*, 2007, **7**, 888-899.
- O. Kovar, M. Kufa, V. Finger, O. Soukup, M. Kratky, C. Torruellas, J. Roh and J. Korabecny, *J. Med. Chem.*, 2025, **68**, 20903-20929.
- (a) J. W. Pavlik and N. Kebede, *The Journal of Organic Chemistry*, 1997, **62**, 8325-8334; (b) J. W. Pavlik, R. E. Connors, D. S. Burns and E. M. Kurzweil, *Journal of the American Chemical Society*, 1993, **115**, 7645-7652; (c) J. W. Pavlik, P. Tongcharoensirikul, N. P. Bird, A. C. Day and J. A. Barltrop, *Journal of the American Chemical Society*, 1994, **116**, 2292-2300; (d) J. W. Pavlik, N. Kebede, N. P. Bird, A. C. Day and J. A. Barltrop, *The Journal of Organic Chemistry*, 1995, **60**, 8138-8139; (e) J. W. Pavlik and P. Tongcharoensirikul, *The Journal of Organic Chemistry*, 2000, **65**, 3626-3632; (f) J. A. Barltrop, A. C. Day, A. G. Mack, A. Shahrissa and S. Wakamatsu, *J. Chem. Soc., Chem. Commun.*, 1981, DOI: 10.1039/C39810000604, 604-606; (g) H. Tiefenthaler, W. Dörscheln, H. Göth and H. Schmid, *Helv. Chim. Acta*, 1967, **50**, 2244-2258; (h) P. Beak, J. L.



- Miesel and W. R. Messer, *Tetrahedron Lett.*, 1967, **8**, 5315-5318.
13. (a) B. Roure, M. Alonso, G. Lonardi, D. B. Yildiz, C. S. Buettner, T. dos Santos, Y. Xu, M. Bossart, V. Derdau, M. Méndez, J. Llaveria, A. Ruffoni and D. Leonori, *Nature*, 2025, **637**, 860-867; (b) M. Alonso, G. Lonardi, E. M. Arpa, B. Roure, A. Ruffoni and D. Leonori, *Nature Communications*, 2025, **16**, 7502; (c) T. dos Santos, C. S. Buettner, D. B. Yildiz, M. Mamone, A. Ruffoni and D. Leonori, *Angew. Chem. Int. Ed.*, 2025, **64**, e202423804.
14. G. L. Bartholomew, S. F. Kim, Y. Oyamada, F. Sbordone, J. A. Carroll, J. E. Jurczyk, C. S. Yeung, C. Barner-Kowollik and R. Sarpong, *Angew. Chem. Int. Ed.*, 2025, **64**, e202423803.
15. Y. Xu, L. Poletti, E. M. Arpa, B. Roure, A. Ruffoni and D. Leonori, *Nature Communications*, 2026, **17**, 2141.
16. M. Ye, A. J. F. Edmunds, J. A. Morris, D. Sale, Y. Zhang and J.-Q. Yu, *Chemical Science*, 2013, **4**, 2374-2379.
17. G. Ohta, K. Koshi and K. Obata, *CHEMICAL & PHARMACEUTICAL BULLETIN*, 1968, **16**, 1487-1497.
18. H. B. Schröter, D. Neumann, A. R. Katritzky and F. J. Swinbourne, *Tetrahedron*, 1966, **22**, 2895-2897.
19. K. Paruch, M. P. Dwyer, C. Alvarez, C. Brown, T.-Y. Chan, R. J. Doll, K. Keertikar, C. Knutson, B. McKittrick, J. Rivera, R. Rossman, G. Tucker, T. Fischmann, A. Hruza, V. Madison, A. A. Nomeir, Y. Wang, P. Kirschmeier, E. Lees, D. Parry, N. Sgambellone, W. Seghezzi, L. Schultz, F. Shanahan, D. Wiswell, X. Xu, Q. Zhou, R. A. James, V. M. Paradkar, H. Park, L. R. Rokosz, T. M. Stauffer and T. J. Guzi, *ACS Medicinal Chemistry Letters*, 2010, **1**, 204-208.
20. (a) N. El Achi, Y. Bakkour, L. Chausset-Boissarie, M. Penhoat and C. Rolando, *RSC Advances*, 2017, **7**, 29815-29820; (b) M. Roseau, L. Chausset-Boissarie, S. Gremetz, P. M. C. Roth and M. Penhoat, *Photochemical & Photobiological Sciences*, 2022, **21**, 421-432.
21. (a) D. J. Hughes, W. B. Ng, R. T. Chapman, G. Healing, M. A. Parkes, J. Rigden, O. J. Smith, E. Springate, J. O. F. Thompson, T. Walmsley, J. L. Woodhouse and R. S. Minns, *Physical Chemistry Chemical Physics*, 2025, **27**, 19762-19771; (b) R. E. Connors, J. W. Pavlik, D. S. Burns and E. M. Kurzweil, *The Journal of Organic Chemistry*, 1991, **56**, 6321-6326.
22. P. R. Schreiner, *Journal of the American Chemical Society*, 2017, **139**, 15276-15283.

View Article Online  
DOI: 10.1039/D6SC03470E



All data is available in the Supporting Information.

