

# RSC Sustainability

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Y. Mahamat Ahmat, M. El Kaddouri, L. Fradette and S. Kaliaguine, *RSC Sustainability*, 2026, DOI: 10.1039/D6SU00222F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## Sustainability Spotlight Statement

This work analyzes advances in the selective epoxidation of bio-olefins, highlighting the transition from classical stoichiometric oxidants to catalytic systems that employ greener oxidants and an improved green process. It discusses mechanistic insights into active oxygen species formation, and strategies that enhance activity, selectivity, and process efficiency. All reagents used in the process, including catalysts, are fully recyclable, very inexpensive, much less hazardous, and driven under ambient conditions while no byproducts or waste are produced. This process is essential to reduce fossil olefins dependency and environmental impact in the polymer industry and supporting circular and green chemical manufacturing.



## Sustainable Green Epoxidation of Terpenes and other Olefins by Dioxirane Generated from H<sub>2</sub>O<sub>2</sub>

View Article Online  
DOI: 10.1039/D6SU00222F

Yacoub Mahamat Ahmat<sup>1</sup>, Malak El Kaddouri<sup>1,2</sup>, Louis Fradette<sup>3</sup>, Serge Kaliaguine<sup>1</sup>

<sup>1</sup>Department of Chemical Engineering, Laval University, Quebec City, Quebec Canada

<sup>2</sup>Euromed University of Fes, UEMF, Morocco

<sup>3</sup>Polytechnique Montreal, Montreal, Canada

### Abstract

This study examines the catalytic synergy between a nitrile and a ketone for the formation of dioxirane from hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) for the purpose of epoxidizing a wide variety of terpenes and other olefins. The influence of factors such as the amount of nitrile, ketone, reaction time and the pH of the reaction medium on the conversion rate and epoxide yield were systematically evaluated to determine optimal conditions. Experimental findings reveal that under ambient conditions and at pH values above 11, nearly complete conversion and epoxide yields (~100%) are achieved across diverse terpene and other olefin substrates. The reaction process is simple and efficient, proceeding without any pre-synthesized solid catalyst, whereas readily available acetonitrile and acetone act as co-catalysts and can be recycled indefinitely, facilitating large-scale implementation. The combination of complete substrate conversion and quantitative epoxide yield, together with the use of low-cost raw materials that generate no waste, ambient reaction conditions and the perpetual recyclability of the reagents, renders this process more pertinent than other epoxidation techniques and easier to implement on an industrial scale in response to the current industrial demand for epoxides.

**Keywords:** epoxidation, terpenes, H<sub>2</sub>O<sub>2</sub>, nitrile, ketone, dioxirane



## Introduction

The epoxidation of olefins is an important primary step in the production of a wide variety of high-value products in fine chemicals, polymers and pharmaceuticals [1-5]. Literature discusses a variety of epoxidation pathways, some of which being currently under development. The latter include homogeneous and heterogeneous catalytic ones using molecular oxygen, hydrogen peroxide or other oxidizing agents [5-10]. From environmental considerations, molecular oxygen and hydrogen peroxide are considered the most desired oxidants as they generate no waste or only water. However, most of the catalytic pathways developed to date fail to meet all the essential criteria for an ideal epoxidation process, including high turnover numbers, excellent product selectivity, compatibility with environmentally benign solvents and catalyst stability with effective recovery and reusability [11-13]. For example, cyclic compounds bearing trisubstituted double bonds, such as limonene,  $\alpha$ -pinene, terpinene and their epoxides are chemically unstable in typical aerobic epoxidation media, promoting the formation of undesirable secondary oxidation products [6]. Additionally, some studies have observed a trade-off between selectivity and conversion, with higher conversion generally corresponding to lower selectivity [14, 15].


In response to these challenges, our recent research was focused on developing an efficient and sustainable method for terpene epoxidation. Our findings demonstrated that the dioxirane pathway represents one of the most promising and industrially viable route for achieving highly selective terpene epoxidation [16-18]. A dioxirane is an oxidant usually formed from a ketone and oxone as the primary oxidant in an aqueous medium. This reaction medium was proven to generate the least secondary reactions with substrates and their corresponding epoxides, resulting in high selectivity toward the desired epoxide products. Beyond its outstanding selectivity, this pathway embodies the principles of green chemistry and offers significant potential for large scale implementation. This process high selectivity to the target epoxide eliminates the need for post-epoxidation separation, while employing low-cost and recyclable reagents [16]. In our previous investigations, we further demonstrated that the reaction can be carried out entirely in an aqueous medium without the use of organic solvents. This represents a major advancement in process sustainability and scalability. The only remaining drawback is the formation of  $3\text{KHSO}_4 \cdot \text{K}_2\text{SO}_4$  sulfate waste from oxone by the reaction [16].

To overcome this drawback and further enhance the environmental performance of the process, a more attractive option for generating dioxirane would be the in situ formation of this type of oxidant from hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) in the presence of a ketone. Oxone itself is synthesized



using  $\text{H}_2\text{O}_2$  as oxidant (Scheme 1), potassium hydroxide and sulfuric acid [19, 20]. Therefore, the direct generation of a dioxirane from  $\text{H}_2\text{O}_2$  represents a clean and promising alternative to the conventional oxone route. Since no persulfate would be generated in the reaction, the new route would produce no waste, and the liquid reagents involved in the reaction could be recyclable, further enhancing the process sustainability.

Building on this concept, Shi et al. developed a system for the in situ formation of a dioxirane from  $\text{H}_2\text{O}_2$  for the asymmetric epoxidation of a series of olefins [21, 22]. In their approach, both a ketone and a nitrile are required. A peroxyamine intermediate, which is formed through the reaction between the nitrile and  $\text{H}_2\text{O}_2$ , acts as an active oxidant, playing a role analogue to that of the peroxymonosulfate anion ( $\text{HSO}_5^-$ ) in the conventional synthesis pathway. This intermediate subsequently reacts with the ketone to produce the oxidizing dioxirane species, which finally epoxidizes the olefin. Scheme 1 provides a comparative description of these two pathways for the generation of dioxirane (exemplified here as dimethyldioxirane) to produce the epoxide.



**Scheme 1.** Comparison between the generation of a dioxirane either from oxone or directly from  $\text{H}_2\text{O}_2$

This study aims at investigating in greater detail the synergistic effect between a nitrile and a ketone in the epoxidation of a series of olefins, mainly terpenes, using hydrogen peroxide as an oxidant, with the ultimate goal of developing an efficient process for synthesis of renewable and sustainable epoxides. Particular attention is given to the aspects of green chemistry, emphasizing the use of low-cost materials and operational simplicity, seeking even compliance with the twelve principles of green chemistry.



## Materials and methods

### Materials

All substrates subjected to epoxidation (R-(+)-limonene, (-)- $\alpha$ -pinene, (-)- $\beta$ -pinene, 3-carene,  $\alpha$ -terpinene,  $\gamma$ -terpinene,  $\beta$ -myrcene, carvone, carveol and styrene) were purchased from Sigma Aldrich with a purity of nearly 100%. The other products were purchased from Alpha Ceaser. The oxidant used for the reaction is 30 % hydrogen peroxide in aqueous volume. Potassium carbonate ( $K_2CO_3$ ) was used to adjust the pH in order to conduct the reaction in a basic medium. Diethyl ether is used for liquid-liquid extraction due to its high affinity for epoxides and its low boiling point (34°C). All products were used as received without any preliminary purification.

### Methods

For a typical test, the epoxidation reaction was carried out under ambient conditions in a 250 mL round-bottom flask. The reaction mixture consisted of 15 mL of acetonitrile, 10 mL of acetone and 12 mL of an aqueous potassium carbonate ( $K_2CO_3$ ) solution (1M), which were mixed thoroughly. Subsequently, 1 mL (6.2 mmol) of the substrate to be epoxidized ( $\alpha$ -pinene) was added to the reaction mixture, followed by the addition of 1.8 mL (i.e., 18 mmol) of 30 % hydrogen peroxide. The mixture was stirred using a magnetic stirrer at 500 rpm for 2 hours.

At the end of the test, the reaction mixture was transferred to a separating funnel for liquid-liquid extraction. Approximately 50 mL of diethyl ether was added to the separating funnel containing the reaction mixture, resulting in the formation of two distinct phases. The lower phase, which is the aqueous phase, was discarded, and the upper phase is the diethyl ether phase containing the epoxidized olefin. The organic phase was then recovered and dried using magnesium sulfate ( $MgSO_4$ ) and transferred to a rotary evaporator to isolate the oxidized olefin. The final product is recovered for analysis to quantify the conversion of  $\alpha$ -pinene and the corresponding epoxide yield.

### Characterization

The reaction products were first identified by gas chromatography–mass spectrometry (GC–MS) to confirm the presence of terpene epoxides. The GC–MS comprised a Hewlett-Packard GC System HP 5890 series coupled to a Hewlett-Packard MSD Model 5970. The instrument was equipped with a Zebron ZB- 5MS capillary column (30 m  $\times$  0.25 mm  $\times$  0.25 mm).



For quantitative analysis, measurements were then conducted using a CP-3800 gas chromatograph (Varian Inc.) equipped with a flame ionization detector (FID) and coupled to a 5 m long Stabilwax column (30 m × 0.53 mm × 1 μm). The final products were quantified based on the calibration curves of the compounds identified by their retention time. Methyl benzoate was used as an internal standard. The conversion of terpenes, yield and selectivity of terpene epoxides were calculated on the basis of chromatographic results using Eqs 1, 2 and 3 respectively. The oxygen yield is calculated as the ratio of the number of oxygen atoms in the epoxidized olefin (N) to the number of moles of H<sub>2</sub>O<sub>2</sub> used (Eq. 4).

$$\text{Conversion}_{\text{terpene}} (\%) = \frac{n_{i(\text{olefin})} - n_{f(\text{olefin})}}{n_{i(\text{olefin})}} * 100 \quad (1)$$

$$\text{Yield}_{\text{terpene epoxide}} (\%) = \frac{n_{(\text{olefin epoxide})}}{n_{i(\text{olefin})}} * 100 \quad (2)$$

$$\text{Selectivity}_{\text{epoxide}} (\%) = \frac{n_{\text{epoxide}}}{n_{i(\text{olefin})} - n_{f(\text{olefin})}} * 100 \quad (3)$$

$$\text{Yield}_{\text{oxygen}} (\%) = \frac{N_{\text{Oxygen in epoxide}}}{n_{\text{H}_2\text{O}_2}} * 100 \quad (4)$$

The <sup>1</sup>H-NMR spectra were recorded on a Varian Inova at 400 MHz spectrometer using 32 scans and a relaxation time of 2 s. Approximately, 10 mg of each sample was dissolved in approximately 1 g of deuterated chloroform (CDCl<sub>3</sub>).

## Results and discussion




Traditionally, in order to form a dioxirane, the source used as an oxidant is the stable commercial form of potassium persulfate, oxone (2KHSO<sub>5</sub>·K<sub>2</sub>SO<sub>4</sub>·KHSO<sub>4</sub>) [20, 23]. The latter provides the active oxygen of the peroxy monosulfate anion (HSO<sub>5</sub><sup>-</sup>) capable of reacting directly with a ketone to generate dioxirane. This method yields an effective and clean oxidizing intermediate. Several epoxidation studies have been carried out using this method of dioxirane in situ generation [16].

It would, however, be particularly interesting if this dioxirane could be generated directly from H<sub>2</sub>O<sub>2</sub> while maintaining the same reaction efficiency in terms of operational simplicity, conversion and epoxide yield.



According to preliminary tests,  $H_2O_2$  alone was shown not to react directly with acetone to produce dimethyldioxirane. Based on the results summarized in Table 1, it was found necessary to use a nitrile that acts synergistically with the ketone to epoxidize the double bond. To demonstrate and confirm this catalytic synergy between a nitrile and a ketone in performing epoxidation through dioxirane generation in the presence of  $H_2O_2$ , a series of experiments was performed, and results are summarized in Table 1.

**Table 1.** Synergistic effect of a nitrile and a ketone in epoxidation of  $\alpha$ -pinene with  $H_2O_2$ <sup>a</sup>.

					
Entry	Catalyst #1	Catalyst #2	Conv (%) <sup>b</sup>	Yield (%) <sup>c</sup>	Selectivity (%) <sup>d</sup>
1	CH <sub>3</sub> CN	CH <sub>3</sub> COCH <sub>3</sub>	100	96	96
2	CH <sub>3</sub> CH <sub>2</sub> CN	CH <sub>3</sub> COCH <sub>3</sub>	100	95	96
3	CH <sub>3</sub> CN	-	2	-	-
4	CH <sub>3</sub> CN	CH <sub>3</sub> OH	12	8	66
5	CH <sub>3</sub> CN	Ethyl acetate	5	-	-
6	-	CH <sub>3</sub> COCH <sub>3</sub>	3	-	-
7	Triethylamine 	CH <sub>3</sub> COCH <sub>3</sub>	4	-	-
8	 Pyridine	CH <sub>3</sub> COCH <sub>3</sub>	3	-	-
9	-	-	0	-	-


<sup>a</sup>Catalyst #1 (15 ml), catalyst #2 (10 ml),  $\alpha$ -pinene (1 ml, 6.2 mmol), reaction time (2h),  $H_2O_2$  (30 %; 1.8 ml, 18 mmol),  $K_2CO_3$  (12 ml of 1M solution), room temperature. Calculated using <sup>b</sup>Equation (1) <sup>c</sup>Equation (2) <sup>d</sup>Equation (3).

According to the results summarized in Table 1, high epoxide conversions and yields are achieved only when both a nitrile and a ketone are used simultaneously (Entries 1 and 2). The selectivity for alpha-pinene epoxide is always very high, due to the absence or suppression of acid-catalyzed epoxide decomposition which poses a challenge in the case of heterogeneous epoxidation using  $O_2$  or  $H_2O_2$ . When only a nitrile is used, the conversion of  $\alpha$ -pinene drops drastically to





2% (Entry 3). To determine whether acetone plays a catalytic or solubilizing role in the reaction, another solvent with properties similar to acetone (methanol) was tested (Entry 4). In this case, the conversion of  $\alpha$ -pinene was only 12 %. Similarly, when ethyl acetate was used, the conversion decreased to 5 % (CH<sub>3</sub>CN/Ethyl acetate, Entry 5).

Furthermore, when the reaction was carried out in the absence of nitrile but in the presence of acetone, the conversion was limited to 3 % (Entry 6). To verify whether the presence of nitrile itself or simply a nitrogen-containing compound was required, additional tests were conducted using triethylamine and pyridine instead of acetonitrile (Entries 7 and 8). In both cases, the conversion of  $\alpha$ -pinene remained very low, confirming that the nitrile functionality plays a specific and essential role in the reaction mechanism. These observations strongly support the existence of a catalytic synergy between the nitrile and the ketone during the epoxidation process, which can be rationalized by the successive reactions illustrated in Scheme 2.




**Scheme 2.** Proposed successive reactions during epoxidation with H<sub>2</sub>O<sub>2</sub> in the presence of the nitrile/ketone pair.

A test conducted in absence of  $\alpha$ -pinene allowed identify acetonitrile as the product of reaction (2) instead of imidic acid. Reactions (1-3) are accompanied by decomposition reactions possibly of the three oxidizing agents (4-6) as indicated by gaseous oxygen evolution from the reaction medium.



To demonstrate that the reaction involves the formation of a dioxirane that reacts with the olefin to form the epoxide and to support the proposed mechanism (Scheme 2), we conducted new tests using a chiral ketone that we had previously studied [17] to determine whether a chiral yield could be achieved. To do this, we used a chiral molecule, R-limonene (Scheme 3), while replacing acetone (which is not a chiral ketone) with the chiral ketone described in Table 2. The results are summarized in Table 2.



**Scheme 3.** Epoxidation of R-limonene by a dioxirane to produce its chiral epoxides

Indeed, according to the results in Table 2, when the chiral ketone is used instead of acetone (which is non-chiral), the yield of trans-limonene dioxide reached up to 92%. Only 5% of cis-limonene dioxide was formed. Whereas if acetone is used, a mixture of cis and trans is obtained with a trans yield of only 39%. In the absence of any ketone, the conversion is only 3%. This is sufficient evidence that the ketone group is indeed involved in the reaction through the formation of a dioxirane, which in turn acts to epoxidize the double bond.

**Table 2.** Effect of the chiral ketone on the chiral yield of R-limonene dioxide.

Ketone Catalyst		Conv (%)	Yield of Trans (%)	Yield of cis (%)
Name	structure			
<b>Catalyst Free</b>	-	3	-	-
<b>Acetone</b>		100	39	60
<b>Chiral ketone</b>		100	92	5


<sup>a</sup>Acetonitrile (15 ml), acetone (10 ml) or chiral ketone (2 g), R-limonene (0.5 ml, 3.1 mmol), reaction time (2h), H<sub>2</sub>O<sub>2</sub> (30 %; 3.6 ml, 36 mmol), K<sub>2</sub>CO<sub>3</sub> (12 ml of 1M solution), room temperature



## Optimization of acetonitrile and ketone concentrations

View Article Online  
DOI: 10.1039/D6SU00222F

Since it was confirmed that acetonitrile and acetone act synergistically in the formation of epoxides using  $\text{H}_2\text{O}_2$  as the oxidant, a study was performed in which the amounts of acetonitrile and acetone were independently varied to assess their influence on the epoxidation reaction. The results for  $\alpha$ -pinene conversion and epoxide yield are presented in Figure 1. Optimal conversion (98%) and epoxide partial conversion (96%) were obtained when the volume of acetonitrile ranged between 15 and 20 mL. In contrast, both conversion and yield decreased gradually when the acetonitrile volume exceeded 25 mL (Figure 1). This behavior suggests that an excess of acetonitrile in the reaction medium promotes the catalytic decomposition of  $\text{H}_2\text{O}_2$  into oxygen rather than its utilization for dioxirane formation. Consequently, the reduced availability of the active oxidizing species accounts for the observed decline in conversion and epoxide yield (Figure 1).




**Figure 1.** Variation in  $\alpha$ -pinene conversion and its epoxide yield as a function of the amount of acetonitrile<sup>a</sup>

<sup>a</sup>Acetone (10 ml, 172 mmol),  $\alpha$ -pinene (1 ml, 6.2 mmol), reaction time (2h),  $\text{H}_2\text{O}_2$  (1.8 ml, 18 mmol),  $\text{K}_2\text{CO}_3$  (12 ml of 1M solution), room temperature.

To evaluate the effect of acetone, the amount of acetonitrile was kept constant at 15 mL to observe the variation in  $\alpha$ -pinene conversion and epoxide yield was examined as a function of the amount of acetone. The results are shown in Figure 2. Both conversion and yield increased progressively with increasing the amount of acetone, reaching nearly 100 % at an acetone volume of 10 mL. Beyond this value, no decrease in conversion or yield was observed, in contrast to behavior noted of acetonitrile, where a decrease was observed starting at 25 mL.



View Article Online  
DOI: 10.1039/D6SU00222F

**Figure 2.** Variation in  $\alpha$ -pinene conversion and its epoxide yield as a function of the amount of acetone<sup>a</sup>

<sup>a</sup>Acetonitrile (15 ml, 314 mmol),  $\alpha$ -pinene (1 ml, 6.2 mmol), reaction time (2h),  $H_2O_2$  (1.8 ml, 18 mmol),  $K_2CO_3$  (12 ml of 1M solution), room temperature.

To minimize the overall reaction volume of the reagents involved in the epoxidation reaction, a series of experiments was conducted in which the volumes of acetonitrile, acetone and water were systematically varied. The corresponding results are summarized in Table 3. When the total reaction volume that previously yielded nearly complete  $\alpha$ -pinene conversion and epoxide formation (entry 1) was reduced by half (entry 2), both conversion and yield decreased by approximately 10 %. Further reduction of the reaction volume to one-third led to a reduction of approximately 25 % in both conversion and yield, resulting in values of 72 % and 69 % respectively (entry 3, Table 3). Therefore, the reagent quantities used in entry 1 are considered at optimal values for achieving maximum conversion and epoxide yield.

**Table 3.** Effect of total reaction volume on  $\alpha$ -pinene conversion and epoxide yield<sup>a</sup>.

Entry	$V_{\text{acetonitrile}}$ (mL)	$V_{\text{acetone}}$ (mL)	$K_2CO_3$ (mL) <sup>b</sup>	Conv (%) <sup>c</sup>	Yield (%) <sup>c</sup>	Selectivity (%) <sup>c</sup>
1	15	10	12	98	96	98
2	7.5	5	6	87	86	98
3	5	3.3	3	72	69	95

<sup>a</sup> $\alpha$ -pinene (1 ml, 6.2 mmol), reaction time (2h),  $H_2O_2$  (1.8 ml, 18 mmol), <sup>b</sup> $K_2CO_3$  (1M of aqueous solution), room temperature. <sup>c</sup>See characterization section.


In terms of mechanism, effectiveness, and scalability, reducing the reaction volume from 15/10 mL (acetonitrile/acetone) to 5/3.3 mL led to a decrease in  $\alpha$ -pinene conversion from 98% to



72%, while selectivity remained high (95–98%). It is our experience that the epoxidation reaction should be considered as in competition with oxidizing agents decomposition. A lower epoxide yield at full H<sub>2</sub>O<sub>2</sub> conversion indicates higher rate of this decomposition. It is likely that reducing the total volume as indicated in Table 3 decreases the relative rate of epoxidation, for example by decreasing the liquid-liquid interphase area

### Optimization of reaction time

After optimizing the amounts of acetonitrile and acetone, the effect of reaction time on  $\alpha$ -pinene conversion and yield was evaluated. The results are presented in Figure 3. Both conversion and yield gradually increase with reaction time and stabilized after approximately 90 minutes. The  $\alpha$ -pinene epoxide obtained was stable in the reaction medium, no decomposition was observed even when the reaction was extended to 4 hours. This shows that the reaction medium is chemically stable with respect to the epoxide formed. As shown in Figure 3, the conversion of  $\alpha$ -pinene to epoxide was very rapid, reaching nearly 80% within the first 30 minutes of reaction.



**Figure 3.** Effect of reaction time on  $\alpha$ -pinene conversion and epoxide yield <sup>a</sup>

<sup>a</sup>Acetonitrile (15 ml), acetone (10 ml),  $\alpha$ -pinene (1 ml, 6.2 mmol), reaction time (2h), H<sub>2</sub>O<sub>2</sub> (1.8 ml, 18 mmol), K<sub>2</sub>CO<sub>3</sub> (12 ml of 1M solution), room temperature.





### H<sub>2</sub>O<sub>2</sub> concentration effect

Table 4 shows the effect of hydrogen peroxide concentration on the conversion and yield to epoxides. According to the results listed, it is observed that for the trisubstituted epoxide ( $\alpha$ -pinene), a molar ratio of H<sub>2</sub>O<sub>2</sub> to  $\alpha$ -pinene of 3 was sufficient to achieve total conversion and yield, whereas for styrene, a ratio of 4 was required. This higher ratio is due to the greater




nucleophilicity of the trisubstituted double bond in  $\alpha$ -pinene compared to styrene [24]. Higher excess concentrations of  $\text{H}_2\text{O}_2$  (3 for  $\alpha$ -pinene and 6 for styrene) had no effect on the epoxides formed. The epoxides were obtained in a highly pure form, as shown in Figures 4 and 5.

**Table 4.** Effect of  $\text{H}_2\text{O}_2$  concentration on olefin conversion and yield<sup>a</sup>


Olefin	Epoxide	$n_{\text{H}_2\text{O}_2}/n_{\text{olefin}}$	Conv (%)	Yield (%)	Selectivity (%)	Entry
		2	71	70	99	1
		3	100	97	97	2
		4	100	97	97	3
		3	91	88	97	4
		4	100	97	98	5
		5	100	97	98	6

<sup>a</sup>Acetonitrile (15 ml), acetone (10 ml), olefin (1 ml, 6.2 mmol), reaction time (2h),  $\text{K}_2\text{CO}_3$  (12 ml of 1M solution), room temperature



**Figure 4.** Proton NMR spectrum of  $\alpha$ -pinene and its corresponding epoxide.





**Figure 5.** Proton NMR spectrum of styrene and its styrene epoxide obtained.

### Effect of pH on the epoxidation reaction

The influence of pH on the epoxidation reaction was also evaluated based on the conversion to  $\alpha$ -pinene and its epoxide yield. Potassium carbonate ( $K_2CO_3$ ) was chosen to adjust the pH of the reaction medium. For this purpose, a series of 12 mL of aqueous  $K_2CO_3$  solutions with different concentrations was prepared (column 2, Table 5). Each solution was used in a separate reaction under identical conditions and the corresponding results are summarized in Table 5. The overall pH of the reaction mixture ( $pH_{\text{whole}}$ ) was measured at the end of each experiment.

**Table 5.** Influence of reaction pH on  $\alpha$ -pinene conversion and epoxide yield<sup>a</sup>

Entry	$pH_{\text{whole}}$	$K_2CO_3$ (M) <sup>b</sup>	Conv (%) <sup>c</sup>	Yield (%) <sup>c</sup>	Selectivity (%) <sup>c</sup>	Yield <sub>oxygen</sub> (%) <sup>c</sup>
1	11,4	1	96	94	98	32
2	9,8	0,6	90	87	97	30
3	9	0,4	55	50	91	17
4	8	0,1	42	38	90	13
5	7,3	0,05	30	26	86	9
6	6	0	0	0	0	0

<sup>a</sup>Acetonitrile (15 ml), acetone (10 ml),  $\alpha$ -pinene (1 ml, 6.2 mmol), reaction time (2h),  $H_2O_2$  (1.8 ml, 18 mmol), <sup>b</sup> $K_2CO_3$  (12 ml of aqueous solution), room temperature. <sup>c</sup>See characterisation section



It can be observed that when the reaction is carried out without adding potassium carbonate, no epoxide is formed, indicating that  $\alpha$ -pinene has not been converted (Entry 6). A sharp increase in both conversion and epoxide yield was observed as the aqueous concentration of  $K_2CO_3$  increased, nearly 100% conversion and yield are achieved under strongly basic conditions, corresponding to a pH of 11.4 with a  $K_2CO_3$  concentration of 1M (Entry 1).

The following explanation is based on literature [25-28]. The positive effect of pH on  $\alpha$ -pinene conversion and epoxide selectivity can be explained by the mechanism of peroxyimidic acid formation in the acetonitrile/ $H_2O_2$  system. Under basic conditions, hydrogen peroxide is partially converted into the hydroperoxide anion ( $HOO^-$ ), a more nucleophilic species that reacts readily with acetonitrile to form peroxyimidic acid, the active oxidant in the Payne epoxidation. As the pH increases, the concentration of this oxidizing intermediate rises, leading to a higher epoxidation rate and thus greater  $\alpha$ -pinene conversion via the dioxirane pathway. In addition, alkaline conditions suppress acid-catalyzed ring opening and rearrangement of  $\alpha$ -pinene oxide, thereby improving epoxide selectivity. Consequently, both conversion and selectivity increase markedly as the pH is raised from 6.0 to 11.4.

### Comparison of different epoxidation techniques

To demonstrate the efficiency of the present epoxidation method employing dioxirane generated in situ from  $H_2O_2$ , with acetonitrile and acetone as co-catalysts, a comparative analysis was conducted against various reported epoxidation techniques, as summarized in Table 6. The comparison considers several key parameters including the type of catalyst used, reaction temperature, reaction time, oxidant type and quantity, as well as the resulting conversion and epoxides selectivity.

**Table 6.** Comparison of different limonene epoxidation methods

Entry	Catalyst	T (°C)	t <sub>r</sub> (h)	$\frac{n_{H_2O_2}}{n_{limonene}}$	Conv (%)	Sel (%) <sup>a</sup>	Ref
1	CH <sub>3</sub> CN-acetone	25	2	3	100	98	This work
2	Al <sub>2</sub> O <sub>3</sub>	80	10	5	92,2	80,4	[5]
3	Mo/Hf-Mof	70	24	TBHP	85	91	[29]
4	Mo-TUD-1	70	24	THBP	95	60	[30]
5	Ti-SBA-16	75	24	TBHP	80	79	[31]



6	BW <sub>12</sub> /Mo <sub>72</sub> Fe <sub>30</sub>	75	6	3	89	58	[32] <small>View Article Online DOI: 10.1039/D6SJ00222F</small>
7	W/SiO <sub>2</sub> -Liq	90	6	2	55	56,4	[33]
8	SiMo <sub>12</sub> / Mo <sub>72</sub> Fe <sub>30</sub>	75	6	3	92	46	[32]
9	W/SiO <sub>2</sub> -Imp	90	6	2	68	63,2	[33]

<sup>a</sup>Monoepoxide and diepoxide of limonene

Based on the conditions employed for the epoxidation of limonene and the corresponding results for conversion and epoxide selectivity, it can be observed that in most cases, achieving satisfactory performance often requires catalyst development, high temperatures and prolonged reaction times. In some cases, it is even necessary to use an oxidant such as *tert*-butyl hydroperoxide (TBHP), which is a relatively expensive (Entries 3, 4 and 5). Under these conditions, both conversion and yield tend to be lower, and in some cases, catalyst deactivation results in limited recyclability.

In contrast, the epoxidation technique presented in this work achieved complete substrate conversion and close to maximum epoxide yield under ambient reaction conditions. The catalyst is indefinitely recyclable, the reaction time is three to twelve times shorter than the results reported in the literature and no waste is generated during the reaction. These advantages clearly demonstrate that this innovative method is more efficient and sustainable than previously reported ones.

### Epoxidation of terpenes and other olefins

After systematically optimizing the reaction parameters for the epoxidation reaction by evaluating various factors such as the amounts of acetonitrile, acetone, the concentration of potassium carbonate solution and the reaction time, a broad series of olefins, including terpenes, was epoxidized under these optimized conditions. The corresponding results are summarized in Table 7.




**Table 7.** Epoxidation of terpenes and other olefins under optimized conditions<sup>a</sup>View Article Online  
DOI: 10.1039/D6SU00222F<sup>a</sup>Acetonitrile (15 ml), acetone (10 ml), olefin (1 ml, 6.2 mmol), reaction time (2h), <sup>b</sup>H<sub>2</sub>O<sub>2</sub> (1.8

Entry	Olefin	Epoxide	$\frac{n_{\text{H}_2\text{O}_2}}{n_{\text{olefin}}}$ <sup>b</sup>	Conv (%) <sup>b</sup>	Yield (%) <sup>c</sup>	Selectivity (%) <sup>c</sup>	Yield <sub>oxygen</sub> (%) <sup>c</sup>
1			3	100	97	97	33
2			3	98	91	93	32
3			3	100	98	98	33
4			6	100	68/38 <sup>d</sup>	98	
5			6	100	83/16 <sup>e</sup>	83	28
6			6	100	85	85	29
7			3	100	97	97	33
8			6	100	96	96	33
9			6	100	100	100	33
10A			4	99	97	98	25
10B			3	91	88	97	33

mL, 18 mmol for one double bond or 3.6 mL, 36 mmol for two or three double bounds), K<sub>2</sub>CO<sub>3</sub> (12 ml of 1M solution), room temperature. <sup>c</sup>See characterization section, <sup>d</sup>diepoxide yield, p-cymene yield. <sup>1</sup>H NMR analysis are available in the supporting information.



It can be observed that for all epoxidized olefins containing a single double bond, an  $\text{H}_2\text{O}_2$ /olefin molar ratio of 3 is more than sufficient to achieve nearly complete conversion and yield (entries 1, 2, 3, and 10). For olefins possessing two double bonds, doubling this molar ratio is necessary to reach full conversion and obtain higher yields (entries 4, 5, 6, 8, and 9). For the epoxidation of limonene, both monoepoxide and diepoxide products are formed, with yields of 60% and 38%, respectively (entry 4). In the case of  $\beta$ -myrcene, a terpene containing three double bonds (entry 9), only double epoxidation occurs, due to the conjugation of two of its double bonds. This observation is consistent with previously reported findings [16, 34]. Some of the previously reported catalysts were able to epoxidize only one double bond of  $\beta$ -myrcene [11, 12, 35]. The epoxides produced at the end of the epoxidation reaction of all these olefins are almost pure, in contrast to those typically formed using most heterogeneous catalysts. In the case of the epoxidation of corvone (entry 7), only the outer double bond was epoxidized. This is due to the presence of the ketone group, which forms a conjugated double bond with the endocyclic double bond, thereby preventing its epoxidation. A phenomenon observed in a previous study [36, 37]. For other olefins with two double bonds (entries 4, 5, 6, 8 and 9), double epoxidation is achieved by the epoxidation of the endocyclic monoepoxide (Scheme 4), which is the first product formed during the reaction; due to the strong nucleophilicity of the double bond compared to the external one [24]. However, diepoxides cannot be formed without sufficient use of the oxidizing agent, which is hydrogen peroxide.




**Scheme 4.** Successive epoxidation of limonene to mono- and diepoxide

The exception is for  $\gamma$ -terpinene (entry 5), which undergoes partial aromatization of its ring, resulting in the formation of p-cymene with a yield of 15 % (Scheme 5). This side reaction has also been observed when performing epoxidation with dioxirane generated from oxone and following other epoxidation pathways [16, 38]. Some studies have not confirmed this, but still show a relatively low yield of  $\gamma$ -terpinene epoxide compared to other epoxides; this could be related to the aromatization of the ring (yielding p-cymene) [7]. Olefins other than terpenes



such as styrene can also be efficiently epoxidized (entry 10), but a higher molar ratio was required, primarily due to the terminal and monosubstituted double bond.



**Scheme 5.** Products obtained by epoxidation of  $\gamma$ -terpinene (entry 5 of Table 7)

### Proposed reaction mechanism

As previously illustrated in Scheme 2, the epoxidation reaction proceeds through the formation of a reactive oxygen transfer species generated in situ from  $H_2O_2$  in the presence of acetonitrile and acetone. Initially, acetonitrile activated  $H_2O_2$  to form a peroxyimide intermediate, which subsequently oxidizes acetone to generate dioxirane. This dioxirane species is capable of transferring oxygen atom to the olefinic substrate to yield the corresponding epoxide. During this process acetone and acetonitrile are both regenerated, allowing the reaction to proceed catalytically without generating by-products. This synergistic sequence between the nitrile and ketone accounts for the high selectivity, complete conversion and environmentally benign nature of the epoxidation method.

### Conclusion

This study reports the efficient epoxidation of terpenes and other olefins using dioxirane generated directly from  $H_2O_2$  as the oxidant, highlighting the synergistic catalytic role of acetonitrile and acetone. The results demonstrate that the simultaneous presence of a nitrile and a ketone under strongly basic conditions ( $pH > 10$ ) is essential for the reaction to proceed effectively. Acetone and acetonitrile recovery tests from the two phases obtained (both aqueous and organic) upon solvent extraction indicated that acetonitrile/acetone indeed act as homogeneous catalyst and are fully recovered after reaction. After optimizing the reaction conditions, a broad range of olefins was successfully epoxidized with complete conversion and high yield under ambient conditions. All the reagents used in the reaction are very low cost, readily available and can in principle be recycled indefinitely, while the process generates no waste [7].



A comparative analysis with other reported epoxidation methods confirms that the proposed approach appears to be most relevant and effective in meeting the current industrial demand for epoxides. The sustainability and green aspect of this methodology are evidenced by the absence of toxic solvents, the use of recyclable acetone and acetonitrile as a pair of catalysts and the employment of hydrogen peroxide, an inexpensive, stable and environmentally benign oxidant widely used in industry. Moreover, the resulting epoxides are obtained in high purity and can be isolated simply by liquid-liquid extraction using diethyl ether, without the need for further purification.

Considering these advantages, the developed methodology represents a clean, economical and scalable process that aligns strongly with green chemical engineering principles and exhibits strong potential for industrial implementation.

### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### **Acknowledgments**

The authors would like to acknowledge the Natural Sciences and Engineering Council of Canada (NSERC) and Consortium de Recherche et Innovation en Biotechnologie Industrielle du Québec (CRIBIQ) for financial support. We would also like to thank our industrial sponsor, Cycle Carbone inc.



## Reference

- [1] F. Orabona, S. Napolitano, V.D. Badazhkova, W. Perez-Sena, K. Eränen, R. Tesser, M. Di Serio, V. Russo, T. Salmi, CO<sub>2</sub> cycloaddition to limonene diepoxide: Kinetic and mass transfer analysis, *Chemical Engineering Journal*, (2025) 168292.
- [2] A. Mouren, L. Avérous, Sustainable cycloaliphatic polyurethanes: from synthesis to applications, *Chemical Society Reviews*, 52 (2023) 277-317.
- [3] M. Bähr, A. Bitto, R. Mülhaupt, Cyclic limonene dicarbonate as a new monomer for non-isocyanate oligo-and polyurethanes (NIPU) based upon terpenes, *Green Chem.*, 14 (2012) 1447-1454.
- [4] E. Louisy, V. Khodyrieva, S. Olivero, V. Michelet, A. Mija, Use of limonene epoxides and derivatives as promising monomers for biobased polymers, *ChemPlusChem*, 87 (2022) e202200190.
- [5] A.J. Bonon, J.O. Bahu, B.C. Klein, D. Mandelli, R. Maciel Filho, Green production of limonene diepoxide for potential biomedical applications, *Catal. Today*, 388 (2022) 288-300.
- [6] L.A. Gallego-Villada, E.A. Alarcón, A.d.L. Villa, Versatile heterogeneous catalytic system for the selective synthesis of limonene epoxide and diepoxide, *Industrial & Engineering Chemistry Research*, 62 (2023) 20152-20169.
- [7] C.R. Ramaswamy, I. Kopparapu, A. Raykar, A. Kulkarani, V.S. AV, S. Ayyanar, A. Nagarjuna, R.K. Cheedarala, A continuous protocol for the epoxidation of olefins, monocyclic terpenes, and Alpha Beta Unsaturated Carbonyl Synthons using eco-friendly Flow Reactor Conditions, *Results in Engineering*, 16 (2022) 100652.
- [8] J. Liu, H. Tang, P. Jian, B. Liu, Oxygen-vacancy defect engineering to boost the aerobic oxidation of limonene over Co<sub>3</sub>O<sub>4</sub> nanocubes, *Appl. Catal., B*, 334 (2023) 122828.
- [9] J. Liu, X. Ji, C. Wang, L. Wang, P. Jian, Beneficial Synergistic Intermetallic Effect in ZnCo<sub>2</sub>O<sub>4</sub> for Enhancing the Limonene Oxidation Catalysis, *Inorganic Chemistry*, 62 (2023) 18750-18757.
- [10] S. Salaria, B.N. Ruggiero, A. Deberghes, L.C. Seitz, J.M. Notestein, Biphasic Selective Oxidation of Cyclohexene with Dilute Aqueous Hydrogen Peroxide Using Phase Transfer Catalysts, *Industrial & Engineering Chemistry Research*, 63 (2024) 14074-14082.
- [11] W.B. Cunningham, J.D. Tibbetts, M. Hutchby, K.A. Maltby, M.G. Davidson, U. Hintermair, P. Plucinski, S.D. Bull, Sustainable catalytic protocols for the solvent free epoxidation and anti-dihydroxylation of the alkene bonds of biorenewable terpene feedstocks using H<sub>2</sub> O<sub>2</sub> as oxidant, *Green Chem.*, 22 (2020) 513-524.
- [12] J.D. Tibbetts, W.B. Cunningham, M. Vezzoli, P. Plucinski, S.D. Bull, Sustainable catalytic epoxidation of biorenewable terpene feedstocks using H<sub>2</sub> O<sub>2</sub> as an oxidant in flow microreactors, *Green Chem.*, 23 (2021) 5449-5455.
- [13] W. Cunningham, *Catalytic Conversion of Terpene Feedstocks into Value-Added Chemicals and Commodity Chemicals*, University of Bath, 2018.
- [14] J. Van den Reijen, S. Kanungo, T. Welling, M. Versluijs-Helder, T. Nijhuis, K. De Jong, P. De Jongh, Preparation and particle size effects of Ag/ $\alpha$ -Al<sub>2</sub>O<sub>3</sub> catalysts for ethylene epoxidation, *Journal of catalysis*, 356 (2017) 65-74.
- [15] Y. Meng, F. Taddeo, A.F. Aguilera, X. Cai, V. Russo, P. Tolvanen, S. Leveneur, The Lord of the chemical rings: Catalytic synthesis of important industrial epoxide compounds, *Catalysts*, 11 (2021) 765.



- [16] Y. Mahamat Ahmat, H.M. Kamkui, S. Kaliaguine, Scaled up epoxidation of terpenes in microemulsion, *Catal. Today*, 449 (2025) 115202. View Article Online  
DOI: 10.1039/D6SU00222F
- [17] Y. Mahamat Ahmat, S. Kaliaguine, Oxone in microemulsion for diastereoselective epoxidation of R-limonene to trans-limonene dioxide, *Chemical Engineering Journal*, 483 (2024) 149178.
- [18] Y. Mahamat Ahmat, S. Kaliaguine, Epoxidation of limonene and pinenes by dimethyldioxirane in microemulsions, *Catal. Today*, 407 (2023) 146-155.
- [19] B.R. Travis, B.P. Ciaramitaro, B. Borhan, Preparation of purified  $\text{KHSO}_5 \cdot \text{H}_2\text{O}$  and  $\text{nBu}_4\text{NHSO}_5$  from oxone by simple and efficient methods, *Eur. J. Org. Chem.*, 2002 (2002) 3429-3434.
- [20] H. Hussain, I.R. Green, I. Ahmed, Journey describing applications of oxone in synthetic chemistry, *Chem. Rev.*, 113 (2013) 3329-3371.
- [21] L. Shu, Y. Shi, Asymmetric epoxidation using hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) as primary oxidant, *tetrahedron Lett.*, 40 (1999) 8721-8724.
- [22] M. Frohn, Y. Shi, Chiral ketone-catalyzed asymmetric epoxidation of olefins, *Synthesis*, 2000 (2000) 1979-2000.
- [23] R.W. Murray, Chemistry of dioxiranes. 12. Dioxiranes, *Chem. Rev.*, 89 (1989) 1187-1201.
- [24] G. Strukul, Transition metal catalysis in the Baeyer–Villiger oxidation of ketones, *Angewandte Chemie International Edition*, 37 (1998) 1198-1209.
- [25] Y. Sawaki, Y. Ogata, Mechanism of the reaction of nitriles with alkaline hydrogen peroxide. Reactivity of peroxy-carboximidic acid and application to superoxide ion reaction, *Bulletin of the Chemical Society of Japan*, 54 (1981) 793-799.
- [26] G. Grigoropoulou, J. Clark, J. Elings, Recent developments on the epoxidation of alkenes using hydrogen peroxide as an oxidant, *Green Chem.*, 5 (2003) 1-7.
- [27] G.B. Payne, P.H. Deming, P.H. Williams, Reactions of hydrogen peroxide. VII. Alkali-catalyzed epoxidation and oxidation using a nitrile as co-reactant, *J. Org. Chem.*, 26 (1961) 659-663.
- [28] G. Laus, Kinetics of acetonitrile-assisted oxidation of tertiary amines by hydrogen peroxide, *Journal of the Chemical Society, Perkin Transactions 2*, (2001) 864-868.
- [29] D.P. Gomes, A.F. Silva, A.C. Gomes, P. Neves, A.A. Valente, I.S. Gonçalves, M. Pillinger, Epoxidation catalysts prepared by encapsulation of molybdenum hexacarbonyl in UiO-66 (Zr/Hf)-type metal-organic frameworks, *Microporous and Mesoporous Materials*, 330 (2022) 111603.
- [30] D.M. Gomes, P. Neves, M.M. Antunes, A.J. Fernandes, M. Pillinger, A.A. Valente, Post-Synthesis Strategies to Prepare Mesostructured and Hierarchical Silicates for Liquid Phase Catalytic Epoxidation, *Catalysts*, 12 (2022) 1513.
- [31] L. Charbonneau, X. Foster, D. Zhao, S. Kaliaguine, Catalyst-free epoxidation of limonene to limonene dioxide, *ACS Sustainable Chem. Eng.*, 6 (2018) 5115-5121.
- [32] H. Taghiyar, B. Yadollahi, New perspective to catalytic epoxidation of olefins by Keplerate containing Keggin polyoxometalates, *Polyhedron*, 156 (2018) 98-104.
- [33] C. Bisio, A. Gallo, R. Psaro, C. Tiozzo, M. Guidotti, F. Carniato, Tungstenocene-grafted silica catalysts for the selective epoxidation of alkenes, *Appl. Catal., A*, 581 (2019) 133-142.
- [34] A. Brandolese, D.H. Lamparelli, M.A. Pericàs, A.W. Kleij, Synthesis of Biorenewable Terpene Monomers Using Enzymatic Epoxidation under Heterogeneous Batch and Continuous Flow Conditions, *ACS Sustainable Chem. Eng.*, 11 (2023) 4885-4893.



[35] A.A.H. Haidar, P. Guillo, D. Agustin, Polyoxometalates Surrounded by Organic Cations or Immobilized on Functionalized Merrifield Resin as Catalysts for Oxidation of  $\beta$ -Myrcene and  $\beta$ -Caryophyllene, *Applied Sciences*, 15 (2025) 7981. View Article Online  
DOI: 10.1039/D5SU00222F

[36] L. Charbormeau, X. Foster, S. Kaliaguine, Ultrasonic and Catalyst-Free Epoxidation of Limonene and Other Terpenes Using Dimethyl Dioxirane in Semibatch Conditions, *ACS Sustainable Chem. Eng.*, 6 (2018) 12224-12231.

[37] A. Zeroual, M. Ríos-Gutiérrez, O. Amiri, M. El Idrissi, L.R. Domingo, A molecular electron density theory study of the mechanism, chemo- and stereoselectivity of the epoxidation reaction of R-carvone with peracetic acid, *RSC advances*, 9 (2019) 28500-28509.

[38] T. Szuppa, A. Stolle, B. Ondruschka, W. Hopfe, Solvent-free dehydrogenation of  $\gamma$ -terpinene in a ball mill: investigation of reaction parameters, *Green Chem.*, 12 (2010) 1288-1294.



## Data Availability Statement

We declare that all data presented in the journal article are available upon request.

