

Environmental Science Advances

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Q. Ullah, W. Haider, M. Qasim, M. Waqar, T. Khomphet, M. Farid and Z. U. Zaman Asam, *Environ. Sci.: Adv.*, 2026, DOI: 10.1039/D5VA00274E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Department of Environmental Sciences

University of Gujrat, Pakistan



Nano-charged resilience: Harnessing chitosan-based nanomaterials for enhanced vegetable crop adaptation in sustainable agriculture

Environmental Significance Statement

Chitosan-based nanomaterials offer a biodegradable, non-toxic alternative to synthetic agrochemicals, addressing major environmental challenges in agriculture. By enhancing nutrient uptake, photosynthesis, and stress resilience in vegetable crops, they reduce reliance on chemical fertilizers and pesticides while mitigating climate-induced stresses such as drought, salinity, and heavy metal toxicity. Their rapid biodegradation minimizes soil and water contamination, aligning with sustainable agriculture and global food security goals. This review highlights their potential as green nanotechnologies to increase crop yields, safeguard ecosystems, and support the United Nations Sustainable Development Goals.



1 Nano-charged resilience: Harnessing chitosan-based nanomaterials for 2 enhanced vegetable crop adaptation in sustainable agriculture

3 Qudrat Ullah^a, Waqas Haider^a, Muhammad Qasim^a, Muhammad Waqar^a, Thanet Khomphet^a,
4 Mujahid Farid^{b*}, Zaki ul Zaman Asam²

5 *^aFood Technology and Innovation Research Center of Excellence, School of Agricultural
6 Technology and Food Industry, Walailak University, Tha Sala, Nakhon Si Thammarat 80160,
7 Thailand*

8 *^bDepartment of Environmental Sciences, University of Gujrat, Hafiz Hayat Campus, Gujrat,
9 50700, Pakistan*

10 Corresponding Author: mujahid.farid@uog.edu.pk

11 Abstract

12 Vegetable crops are increasingly exposed to new environmental conditions, including elevated
13 temperatures, erratic rainfall patterns, and declining soil fertility, which threaten global food
14 security. Traditional synthetic fertilizers and pesticides exacerbate environmental degradation.
15 Chitosan, a biodegradable and non-toxic biopolymer derived from chitin, has been developed
16 into nanomaterials such as nanoparticles and nanofibers. These chitosan-based nanomaterials,
17 typically less than 100 nm in size, exhibit high biocompatibility and bioactivity, enhancing
18 chlorophyll content, nutrient uptake, and disease resistance in crops. Nonetheless, differences
19 in synthetic processes and composition may cause unstable efficacy, and field-level increase in
20 yield is between 5-20% in comparison with 15-25% in controlled settings. This review explores
21 current advances in chitosan nanomaterials for vegetable crop improvement under biotic and
22 abiotic stress, focusing on crops like tomatoes, potatoes, and lettuce. It critically evaluates
23 benefits and limitations while emphasizing nanotechnology's role in achieving higher yields
24 and environmental sustainability.

25 **Keywords:** Crop; Chitosan; Agriculture; Nanotechnology; Fertilizers

26 1. Introduction

27 Nanotechnology has become a disruptive technology in contemporary agriculture that allows
28 the use of nanoparticles with high accuracy in delivering nutrients, pesticides, or biostimulants
29 ¹. Nanotechnology ensures a higher bioavailability, decreases chemicals, and increases stress-
30 resistance against crops by designing materials at 1-100 nm ². Chitosan-based nanoparticles

31 (ChNPs) and nanofibers are unique nanomaterials since they are biocompatible, biodegradable,
32 and applicable in multifunctional applications in the promotion of plant growth and in
33 controlling pathogens ³. The vegetable crops that are very sensitive to the stresses caused by
34 climatic conditions are a major area of application of nano-chitosan technologies ⁴.

35 Vegetable crop production faces significant challenges due to climate change, including global
36 warming, temperature increases, erratic rainfall, and soil degradation ⁵. Due to
37 biodegradability, non-cytotoxicity, and natural chitin origin, chitosan has been developed to
38 become more sophisticated nanomaterials, such as nanoparticles and nanofibers, appropriate
39 for use in vegetable crops ⁶. Based on the accumulated data, one can assume that opportunities
40 for enhancing the resistance of vegetable crops to various forms of stress exist in the case of
41 using chitosan-based nanomaterials ⁷. ChNPs are capable of raising tomatoes and lettuce
42 chlorophyll by 15-25 percent under salinity stress when used at 50-100 mg/L during 7-14 days
43 ^{8,9}. Since they are cationic, they can bind to the negatively charged plant cell wall, but nutrient
44 uptake efficiency is 25-30% higher compared to the chemical fertilizers ¹⁰. For example,
45 chitosan nanofibers elicitor, as mentioned earlier, has the capability of activating systemic
46 resistance and reducing the rate of diseases such as Phytophthora that affects potatoes to a range
47 of 50-60 percent through enhancing the defense genes. Chitosan's deacetylation degree,
48 ranging from 70-95%, leads to variability in the deacetylation process outcomes ¹¹. As a result,
49 field formulations may only achieve yield increases of approximately 10-15%.

50 This review analyzes how chitosan-based nanomaterials enhance vegetable crop resilience
51 under various biotic and abiotic stressors, using evidence from peer-reviewed studies.

52 **2. Synthesis and Properties of Chitosan-Based Nanomaterials**

53 The development of chitosan-based nanomaterials, such as chitosan nanoparticles (ChNPs), is
54 vital for improving vegetable crop resistance ¹². Bottom-up methods build nanomaterials from
55 smaller units, like assembling Lego blocks. The most common is ionotropic gelation, where
56 positively charged chitosan binds with negatively charged molecules (e.g., TPP) to form tiny
57 spherical nanoparticles (50–100 nm) in a simple, water-based process ^{13, 14}. This is a relatively
58 easy method that is approximately 80% efficient on the bioactive compounds ¹⁵. On the other
59 hand, the bottom-up techniques such as milling and ultrasonication mainly involve the
60 mechanical breakdown of chitosan particles into particles of size in the range of 200-300 nM
61 but do not alter the morphological homogeneity of chitosan and there is only about 20-30%



62 variation in particle size distribution ¹⁶. This is why the bottom-up strategies can offer a high
63 level of accuracy; however, these methods cannot be implemented in crop farming at a large
64 scale since they require some heavy mechanical energy.

65 Chemical synthesis approaches expand the fabrication of chitosan to a higher degree following
66 the chemical process where chitin is liquefied by the usage of strong acids and bases to convert
67 to chitosan and then in nanoscale form ¹⁷. Another conventional method of thermo-chemical
68 hydrolysis to get soluble and biologically active products is capable of producing an extent of
69 deacetylation between 70-95% the result is a non-uniform molecular size of polymer of
70 between 50-1000 kDa ¹⁵. Although this kind of variability is industrially feasible, it reduces the
71 effect in vegetable crops relative to fresh weight by up to 15-20% due to variability in
72 deacetylation ¹⁸. The enzymatic methods for chitin deacetylation include the use of purified
73 chitin deacetylase, which is found in microorganisms such as *Bacillus* spp., and at moderate
74 conditions, the degree of deacetylation aimed at 85% was achieved. However, it is reported to
75 be expensive and estimated to be 38–73% cheaper than chemical methods when using agro-
76 industrial waste such as shrimp shells or crab waste ¹⁹.

77 The preparation of chitosan nanomaterials through microbial fermentation and
78 biotechnological application of various enzymes also involves an environmental factor in the
79 process of microbial fermentation ²⁰. Proteolytic microorganisms and organic acids enhance
80 deproteinization and demineralization, and chitosan has a low molecular weight of 100-300
81 KDa and a low particle size of below 150 nm ²¹. By synthesizing ChNPs biologically, there is
82 a great potential to control plant fungal pathogens, with the example of growth inhibition of
83 solani's mycelium to 70-80% by the ChNPs, which is higher than the chemically synthesized
84 ChNPs by 10-15 ²². However, there are also some limitations of this type of production, such
85 as the yields, which are in the range of 20%-30% lower than those of chemical synthesis, which
86 is due to slow fermentation ²³.

87 According to the literature, in this method of thermo-chemical hydrolysis, the degree of
88 deacetylation ranges from 70-95% to increase the solubility and bioactivity of chitosan; the
89 polymers' non-uniformity in molecular weight of 50 to 1000 kDa. This is still industrially
90 possible but decreases the performance of vegetable crops since bioactivity is reduced by 15-
91 20% due to irregular deacetylation ²⁴. Alternatively, an enzymatic process that employs chitin
92 deacetylase from microbial sources, particularly *Bacillus* spp, achieves at most 85%
93 deacetylation under gentle conditions. The high cost of the process was estimated to be 38-



94 73% less expensive than that of the chemical method in utilizing agro-industrial waste
95 Therefore, there is a need to balance the improvement of real-life productive, efficient, and
96 sustainable bioprocesses that cater to the agriculture sector. However, their high surface area
97 can cause agglomeration, reduce germination by up to 80%, and impact droplet size by 10-20%
98 under field conditions unless stabilized with surfactants.

99 Further, enhanced controlled release is beneficial in the use of ChNPs as nutrient release
100 sustains for about 30-60 days, unlike the 5-10 days observed for regular fertilizers. For instance,
101 chitosan NPs encapsulated with indole-3-acetic acid enhance the hydroponic lettuce's growth
102 rate by 20%-25% because of the duration of IAA release²⁶. However, the release kinetics are
103 influenced by the particle size and pH of the environment, which shows that 50 nm is
104 marginally greater than 200 nm, which might be hypothetical and challenging to standardize.
105 This implies that there exists a large variation whereby accuracy when determined under
106 laboratory conditions differs from that of field cropping; this deserves a boost for vegetable
107 crops^{27, 28}.

108 This one can be considered as a conjugation with metals, particularly copper, as a technique of
109 "nanoengineering" to enhance the function of chitosan nanomaterials²⁹. This biocontrol
110 system relates the chitosan's biocompatibility with copper to control the growth of *Fusarium*
111 *oxysporum*, decreasing the growth by 85-90% in tomato crops. While Chitosan nanoparticles
112 organically reduce the crop growth by 70-75%. This entry on copper loading of nano-fertilizer
113 of between 5-10% w/w enhances the enzyme activation of pathogenicity and raises the defense
114 of plants by 25-30 %³⁰. However, toxicity is observed at a higher concentration of copper of
115 more than 15% w/w because it accumulates at the soil level, at which microbial population
116 may be reported to have been reduced by 5-10%. Often, such a trade-off is made to achieve
117 perfect coordination with the necessary potency and non-carcinogenic effect on the natural
118 environment to some extent.

119 Nano-fibering, an additional nano-engineering technique, enhances vegetable crop resilience
120 by improving structural and functional properties. Chitin nanofiber with a size ranging from
121 10-20 nm in diameter exhibits eliciting activities that enhance the defense gene expression of
122 cabbage and its resistance to *Alternaria brassicicola* by 60-70 %³¹. This has enhanced their
123 mechanical properties; their tensile strengths are 2-3 times those of ChNPs, hence enhancing
124 bioactivity with durability. However, the costs of fabricating covalent CNTs-TiO₂
125 nanocomposites are still higher by 20-30% than the costs of nanoparticles, and this is a problem



126 of marketability. This could be a new method of employing nanoparticles and nanofibers to
127 enhance the durability of vegetable crops irrespective of the kind in question³².

128 Consequently, they have the capability of achieving outstanding impacts, for example, an
129 increased yield by 20-40 % on the foliar applied systems because of their surface area and
130 controlled release²⁶. The second type of improvement is nano-tailoring, which is also used
131 where certain areas need changes. There also needs to be strict controls in terms of quality and
132 the work achieved. In vegetable production, which, among many other agricultural
133 productions, is often affected massively by climate stressors, these nanomaterials are in a
134 privileged position to transform sustainable agricultural output if only the synthesis of these
135 nanomaterials can meet the conditions in the field. The preparation of chitosan-based
136 nanomaterials is fundamental to enhancing the resistance of vegetable crops since the various
137 synthesis leads to different properties of the nanomaterials³³. This variability is seen in **Table**
138 **1**; Ionotropic gelation offers 70-85% of Fusarium control in tomato at an optimal size of 50-
139 100 nm ChNPs, while enzyme hydrolysis gives 30-40% Ni removal in lettuce at a 1.4 cost
140 factor more. **Table 1** presents a detailed comparison of synthesis techniques for chitosan-based
141 nanomaterials applied to vegetable crops, encompassing methods such as ionotropic gelation,
142 enzymatic hydrolysis, and chemical deacetylation. It includes columns for nanomaterial type,
143 vegetable crop examples, application methods, particle size, deacetylation percentage, yield
144 increase, pathogen control, stress mitigation, scalability score, and cost factors, offering a
145 comprehensive dataset derived from key studies. The purpose is to link specific synthesis
146 approaches to their practical outcomes in enhancing vegetable resilience, highlighting both
147 efficacy and scalability challenges.

148 **3. Mechanisms of Resilience Enhancement in Vegetable Crops**

149 Nanomaterials from chitosan, such as ChNPs, are more effective against biotic stresses, which
150 result in enhanced resistance of vegetable crops³⁴. Current research demonstrates that chitosan
151 nanoparticles (ChNPs) inhibit 70-85% of the mycelial growth of *Fusarium oxysporum* Schldl.
152 In potato and tomato systems, outperforming bulk chitosan by 20-25% due to their nanoscale
153 size, which enhances penetration of fungal cell walls³⁵. This efficiency is attributed to the
154 cationic nature of chitosan, and it interferes with the pathogen membranes as well as being a
155 germination inhibitor of the spore in *Phytophthora infestans* by 90%³⁶. However, in the case
156 when the size of particles varies between 50 and 200 nm, the actions are unstable; this is due
157 to the reasons that 10-15% action of smaller particles and at the same time the process of



158 synthesis must be very accurate ³⁷. This brings ChNPs as the green solution to synthetic
159 fungicides; however, the stability of these ChNPs at the field level remains a great challenge.
160 As concluded from **Table 2**, a test of pathogen control efficiency for lettuce with *Botrytis*
161 *cinerea* was between 60-70% while for okra it was 85-90% against *Fusarium oxysporum* when
162 using Ch-CuNPs. The result confirmed the role of chitosan nanomaterials in biological stress
163 management. **Table 2** details the biotic stress resistance mediated by chitosan nanomaterials
164 across various vegetable crops, synthesizing data on pathogens, nanomaterials, application
165 methods, and effects such as growth inhibition and enzyme induction. Columns include
166 pathogen reduction percentages, enzyme activity increases, yield impacts, and field variability,
167 offering a comprehensive view of efficacy and challenges. The purpose is to highlight specific
168 pathogen control outcomes, facilitating comparisons across crops and nanomaterials, with
169 scalability scores reflecting practical deployment potential.

170 In addition to repelling invaders at the physical level, chitosan nanomaterials trigger the
171 biochemical defense mechanisms that would improve the ability of vegetable crops to resist
172 biotic stresses. General findings: By applying the ChNPs on the foliage of discomfort, the
173 defense enzymes and activities have been enhanced, where chitinase and peroxidase of
174 tomatoes have increased by 30- 40% in 48 h ²⁶. This induction is in concordance with the
175 increase of the endochitinase genes to decrease *Ralstonia solanacearum* by 50-60%. They also
176 realized that it increased phenolic compounds by 25-35% which boosted its systemic resistance
177 ³⁸. There is, however, the variance of nanomaterial concentration that ranges from 50 to 75 mg
178 / L because beyond this range the efficiency drops by 10-15% due to phytotoxicity ³⁹. This
179 ability of ChNPs to be both antimicrobial and an elicitor is proving the versatility of the
180 compound; however, the need arises for an implementation of a proper amount of the ChNPs
181 to elicit the required response ⁴⁰.

182 As far as abiotic stress is concerned, the use of chitosan nanomaterials has the potential to
183 enhance the water relations of vegetable crops under drought stress. In basil, when applied as
184 a foliar spray, it was found that ChNPs reduced transpiration rates by 20-30% and, on the other
185 hand, enhanced the water use efficiency by 15-25% under water deficit conditions ⁴¹. This
186 could be attributed to the hydrophilic characteristic of chitosan to form a layer on the surfaces
187 of the leaves. This is in concordance with the findings of studies on potatoes, whereby the
188 writers observed that the amount of chitosan must be dried to 50 mg/L and increased the root
189 biomass of plants by 20-30% ⁴². **Table 3** spells out the impact of chitosan nanomaterials on



190 stress factors that affect vegetable crops, including the types of stress, the nanomaterial used
191 application method, and results, including water use efficiency and nutrient absorption. More
192 longitude columns represent stress reduction percentages, yield increase, nutrient uptake, and
193 field variability, so there is a good check on the efficiency and the problems encountered. The
194 objective is to present the effects of nanomaterials on stressors in crop plants and provide a
195 comparison basis among them and the approaches, with scaling scores feasibility to be utilized.

196 The second one, which is highly associated with the salty stress test, is only comparable to
197 salinity tolerance and mitigated by chitosan nanomaterials. ChNPs also have a defensive role
198 in lettuce to decrease the adverse impact of sodium toxicity resulting from a 25-35 % reduction
199 in ion leakage concentration at 100mM NaCl ⁴³. This became a result of the enhancement of
200 the chlorophyll component by 15 to 20% through photosynthesis when the plant undergoes salt
201 stress ⁴⁴. However, if the concentration of water is above 150 mM NaCl, these benefits are
202 reduced up to 10-15%, as OS prevails over the positive impact of the nanomaterials ⁴⁵. Based
203 on the comparative assessment, it has been found that the nano-chitosan is comparatively 20-
204 25% more saline than the bulk forms, with restriction to the upper limit in saline areas ⁴⁶.

205 Apart from improving the nutrient intake through its ChNP-based nano-carriers, it also confers
206 an additional advantage of stress tolerance on the vegetables. As stated by ³⁰ ChNPs used in
207 onion systems to apply NPK fertilizers enhance the nutrient uptake by 30-40 % and bulb yield
208 increases by between 20-25%. This is because of slow release, for it is processed gradually in
209 a period of about one month to two months, compared to 5-10 days for normal fertilizers ⁴⁷.
210 Similarly, nitrogen use efficiency has been improved by 25 to 30 % in Wheat Trials, yet the
211 data on vegetable production are also variable. It shows a 10 to 15% times increase, particularly
212 in nutrient-deficient soils ⁴⁸. Such differences particularly confirm the parameter of soil type as
213 a constraint, the formulation of which demands the development of an effective nano-carrier
214 to be used in the field.

215 These and other biotic things of resilience supplement the abiotic aspects to demonstrate a
216 diverse utilization of chitosan nanomaterials, which, however, has a notable lack of research in
217 past literature ⁴⁹. Antimicrobial effectiveness range is 70 to 90 percent; all the microorganisms
218 are killed, but they can only work with a certain number of particles and in a certain fraction.
219 Field results of the experiments are 10 to 20 percent less than the laboratory experiments ^{50, 51}.
220 Concomitant to abiotic gains of between 15-25%, the effectiveness of water also depends on
221 the climate, the lower being where temperatures are quite high ⁵². Nutrient delivery works well



222 in controlled conditions and badly in dynamic conditions, and that is why one has to work with
223 adaptive strategies³⁰. This variation implies that the ChNPs cannot be implemented as a generic
224 concept, and thus, more attention will be paid to the usage of ChNPs in vegetable crops.

225 In summing up, the theoretical construction of the “nano-mediated stress shield” may be
226 described as an attempt to expand the understanding of a means by which chitosan
227 nanomaterials enhance the firmness of this material. This model defines ChNPs to encompass
228 structures that help discourage pathogen invasions, for instance, the inability of Fusarium to
229 penetrate by 70 – 85% and signaling molecules that trigger defense response, which are
230 enzymes in the range of 30 – 40%²⁶. For abiotic stress, the size of the shield addresses water
231 loss (biotic and abiotic) in a range of 20-30% and nutrition lock, an improvement to nutrition
232 availability by a range of 30-40%. It occupies plant tissue to form a shield³⁸. However, it is
233 moderate most of the time – highest at moderate stress levels and decreases by 10-20% at
234 higher stress levels, which confirms the conditionality of the shield as provided above⁴⁵.

235 While biotic resistance helps in controlling pathogens, the problem with the method is that it
236 highly depends on the synthesis consistency, and it offers only 15-20% less effect if the
237 formulation is not standardized^{53, 54}. Monogenic abiotic stress yield loss avoidance is
238 especially profitable in low-stress zones, which can be interpreted as low-stress yield
239 improvement even such as between 20-25% in onions, in contrast to the stress zone assays in
240 which it is unprofitable⁴⁷. The nutrient enhancements would aid in sustainable yields;
241 nevertheless, owing to the inconsistency in the texture of the soil, there is a 10-15 % lesser
242 yield augmentation that needs calibration for the site⁴⁸. This powerful lens establishes the
243 model as the basis for moving forward, thus assisting in guaranteeing that vegetable resilience
244 enhances actual-life aspects of nanomaterial applications⁵⁵.

245 **4. Application Methods and Delivery Systems**

246 Therefore, the concept of applying Chitosan-based nanomaterials (ChNPs) to seeds as seed
247 coats could be deemed a common practice in enhancing the tolerance of vegetable crops,
248 mainly in the seedling stage⁵⁶. The process of applying ChNP suspension is done at a
249 concentration level of 50-100 mg/L through dip or vacuum infiltration tools, depending on the
250 homogeneity in seed adhesion^{30,57}. There is an increase in germination by 20-30% in chickpeas
251 and cucumbers, reducing factors such as water diffusivity and activation of enzymes, including
252 amylases, by 25-35%⁴³. It acts early It also inhibits the defensive mechanism, reducing the



253 infection of *Fusarium* spp by 60-70 through antimicrobial membrane break, as several authors
254 noted⁵⁸. The problem emerges with the least coated amounts because the germination
255 decreases to 10-15 % for non-optimized batches; the work has to use professional instruments
256 such as rotary coaters⁵⁹.

257 Spraying the chlorides of Ni, Co, and Ni-Co mixed NPs on the foliage of the plant is a novel
258 technique for delivering nutrients to the plant and protecting vegetables like okra and tomatoes
259 from diseases and pests. The technique employed in this study is the sprayers that are used to
260 spray ChNP solutions, most of which are in the concentration of 50-75 mg/L singly or in
261 combination with nutrients such as NPK on the leaves³⁸. This mechanism involves the stomata
262 uptake and slow releasing ability, which enhances the nutrient uptake by 25-40% in 30-60 days,
263 in contrast to 5-10 days in foliar sprays⁴⁷. They are expected to increase crop yield by 20-25%
264 at optimal timing and reduce *Phytophthora infestans* infection by 70-85% through the
265 activation of antioxidant enzymes, such as peroxidase, which exhibit up to a 30% increase in
266 activity²⁶. Nanomaterials of chitosan, used either by such means as foliar spraying, offer a
267 broad spectrum of advantages to vegetable crops (tomatoes, potatoes, and lettuce) improving
268 their growth and resistance, as shown in **Fig. 1**. As an example, foliar-applied ChNPs suppress
269 *Fusarium* wilt by 70-85% by being used to enhance the activity of enzymes; late blight
270 (*Phytophthora infestans*) by a factor of 50-60 by being used to systemic resistance; and nutrient
271 uptake in lettuce by 15-20% by being used to increase chlorophyll content. This value point
272 places emphasis on these crop-specific vigor increases, such as biomass increase and stress
273 resistance. **Table 4** depicts the application methods of the chitosan nanocomposites and their
274 effects on vegetable crop resilience, with information about crops, the kind of nanomaterials,
275 and some benefits that include pathogen control, tolerance to dryness, and enhanced nutrient
276 uptake. Other headings are on yield increases, pathogenic control efficiency, stress effects on
277 yield, variability, and scalability, which allow evaluation of a delivery system's effectiveness
278 and potential difficulties. This is to ensure that each method can be traced to the desired
279 outcome around resilience and enable crop and nanomaterial comparison, with cost analysis
280 also considering the aspect of feasibility in terms of cost.

281 Foliar application's rationale for rapid response centers on delivering nutrients and defenses
282 under stress. The spraying tools of ChNPs involved the use of indole-3-acetic acid growth
283 hormones, which enhanced okra shoot growth by 20-30%. The resistance mechanisms
284 embrace leaf invasion as well as systemic defensive mechanisms, which diminish *Fusarium*

View Article Online
DOI: 10.1039/DSEA00274E



oxysporum levels down to 70-80% ³⁶. The effects are likely to be moderate during conditions of relative humidity between 50 – 70 percent. Evaporation of solution nutrients costs between 10 and 15% of the entire nutrient application, as nutrients from the solution could soil through evaporation. Plants become phytotoxic when the solution exceeds 100 mg/L concentration. Researchers aim to achieve two outcomes by adjusting the solution pH between 5 and 6 and developing solution 2 for better leaf retention and stability.

Soil incorporation's purpose long-term support targets nutrient efficiency and soil health in vegetable systems. Drip irrigation tools apply ChNPs, and the controlled release mechanisms maintain nutrient accessibility within the property to result in 20-25% heavier tomato fruits than the standard control ⁴⁷. ChNPs exhibited equal importance in remediation by adsorbing cadmium from lettuce, making it 25-35% less available to uptake ⁶⁰. The degradation time spans from 60-90 days, together with minimal functionality under acidic conditions, having a pH level below 5, creates a 10-20% reduction of benefits, thus hindering broader ChNP adoption ⁵⁰. ChNPs require a combination with organic amendments for successful implementation since these organic materials maintain optimal soil pH levels and enhance the release mechanism properties. **Fig. 2** shows a flow diagram of the step-by-step processes of chitosan nanoparticles (ChNPs) in increasing nutrient accessibility and uptake in vegetable crops, including application techniques (foliar, soil, seed) and interaction processes (binding, controlled release, chelation) to resulting benefits (improved yield and stress resistance) ⁶¹. In particular: (1) Foliar spray application can be used to rapidly deliver the product to the stomata; (2) Incorporation into the soil can be used to release the product slowly over 30-60 days through pH-triggered chelation; (3) Coating of the seed can be used to increase the rate at which the product attaches to the root; resulting in (4) Depending on the crop, either 20-25% higher nutrient efficiency in tomatoes or 15-20% lower transpiration in drought-stressed potatoes. This brings out the aspect of nanotechnology in accurate and sustained delivery.

Nanomaterials of chitosan are currently being applied as edible coatings to harvested vegetables in order to increase shelf life and minimize post-harvest losses ⁶². Applied as a dipping or spray solution, chitosan films (1-2% w/v) create a semi-permeable barrier that inhibits respiration, ethylene production, and microbial growth ⁶³. As an example, tomatoes covered with nano-chitosan coatings accelerated the loss of weight by 30-40 percent and postponed the process of ripening by 7-10 days at 20 °C ⁶⁴. Chitosan-nanoZnO films reduced



316 the *E. coli* and *L. monocytogenes* by 85-95 percent in 14 days in lettuce⁶⁵. These applications
317 are in line with the zero-waste and limit the usage of artificial preservatives.

318 The versatile nature of chitosan nanocarriers becomes a drawback because they demonstrate
319 antimicrobial action and enzyme induction alongside nutrient delivery⁴⁰. The physical defense
320 layer created through seed coating ranges between 50 to 100 nm in thickness, yet becomes
321 ineffective because of inconsistent application methods⁶⁶. The enzyme enhancement activities
322 of ChNPs applied to plant leaves reach 30% intensification, and their dose-dependent effects
323 open concerns about excessive plant stimulation. Soil chitosan NPs effectively remediate and
324 fertilize the ground while facing challenges with gradual chemical release during conditions of
325 high stress⁶⁰. The implementation of remote sensors intends to improve the performance of
326 techniques by adjusting application amounts through a system of measurements.

327 A new intelligent nano-delivery system combines sensors with the vegetable crops' stress
328 response, ChNPs for application. The smart nano-delivery system combines ChNPs with
329 embedded sensors (pH, moisture, or conductivity) that can be released only under the
330 conditions of sensing stress (e.g., soil moisture below 50%)⁶⁷. As an illustration, a lowering of
331 the pH to less than 5.5 may trigger nutrient release within 48 hours and enhance efficiency by
332 15-25%, and lessen waste^{38, 68}. Research indicates that pathogenic levels would improve by 20
333 to 30 % while nutrient utilization would increase by at least 15 to 25 % without sensing
334 difficulties⁴⁷. Biodegradable polymers remain usable for designing prototype smart clothing
335 systems that monitor elderly health status according to⁵⁵.

336 The delivery systems containing ChNP ensure different mechanisms for seeds and soils, and
337 plants show significant variations, thus achieving numerous changes, including yield increases
338 between 20-25 % with pathogen control reaching between 60-85%³⁶. Smart nano-delivery
339 provides a solution due to its ability to address application inconsistency issues, which cause
340 degradation rates between 10- 20% while also resolving environmental dependency problems
341 as mentioned in³⁰. The present research explores how to optimize the positioning of ChNPs
342 for vegetable production harmony as a resilience strategy.

343 5. Crop-Specific Case Studies

344 Tomato (*Solanum lycopersicum* L.) provides an optimal example of how chitosan nanoparticles
345 (ChNPs) affect growth in controlled environment agriculture. The use of 50 mg/L substance
346 concentration leads to a 20-25% increase in shoot biomass due to enhanced photosynthesis



347 rates, which correspond to elevated chlorophyll content levels of 15-20% ⁴³. Plant nutrient
348 uptake is enhanced via stomatal absorption, with efficiency improving by 25–30% compared
349 to conventional fertilizers ⁶⁹. The variable elevation of 10-15% detected in randomized field
350 trials reflects the light sensitivity of this factor, which reduces its effectiveness ³⁸. The
351 phenomenal prospects of ChNPs demonstrated earlier require precise usage when employing
352 this pest management technique, particularly within tomato horticultural applications.
353 ChNPs demonstrate excellent effectiveness as a suppressant against the bacterial wilt pathogen
354 *Ralstonia solanacearum* present in tomatoes. The peroxidase enzyme activity improves by 30-
355 40% after ChNPs reach a 75 mg/L concentration in the soil, according to ²⁶Through their SAR
356 response mechanism. British scientists found that ChNPs destroy Gram-positive bacteria by
357 crumbling their cell walls, which leads to 70-80% bacteria death in 48 hours ⁷⁰. The stability
358 of ChNP experiences a decrease of 10-15% in acidic soiling conditions with pH levels below
359 5 units ⁵⁰. The test results demonstrate the defensive role of ChNPs as the soil composition
360 remains an unmanageable factor.

361 The incorporation of chitin nanofibers into tomato plants boosts their natural resistance against
362 Fusarium wilt better than using ChNPs independently. The *F. oxysporum* infection rate remains
363 between 70-85% when the nanofiber treatment reaches 100 mg/L, while gene expression of
364 chitinase reaches between 40-50% levels ⁷¹. The fibrous 10–20 nm structure of ChNPs
365 produces superior eliciting results than ChNPs according to laboratory tests by about 15–20%
366 ⁷². Their high production costs, which amount to 20-30% more than ChNPs, combined with
367 50-60% reduced effectiveness in field conditions, according to ⁷³. Represent the main
368 limitations for their practical use. The situation emphasizes both nanofiber resistance to
369 biological agents and cost-effectiveness, together with scalability in real-world operations.

370 The soil application of nano-chitosan shows high performance in areas with water scarcity
371 problems in drought conditions on potatoes (*Solanum tuberosum* L.). A 50 mg/L guttation
372 spraying results in a 20-30% decrease in transpiration rate and enhances water use efficiency
373 by 15-25% during conditions of 50% dry soil ⁷⁴. It has been proven that this micro-coating
374 method increases leaf water retention, so plants produce 20-25% additional tubers ⁴¹. The
375 osmotic stress experienced during excessive dry soil conditions results in decreased benefits of
376 5-10% ⁴⁵. This system provides favorable drought resistance, although it functions in a specific,
377 limited time frame and needs irrigation systems to function.



378 Nano-chitosan provides effective biological management of Phytophthora infestans, which
379 threatens potatoes as a major disease agent. The fertilization of soil at 75 mg/L ChNPs results
380 in minimal mycelial expansion of 70-80 percent and time-limited spore germination of 85-90
381 percent due to membrane breakdown processes ³⁶. The microbiological activity of peroxidase
382 defense enzymes rises by 25-35% while late blight disease incidence decreases by 50-60% ³⁸.
383 The current field application rate of between ten and fifteen percent reduction in efficacy stems
384 from the recent variation of particle size from 50 – 200 nm, which requires further improvement
385 in synthesis techniques ⁷⁵. Compatibility between the delivery mode and dual functionality is
386 established for both nanoparticles, but their performance depends heavily on developing more
387 accurate delivery systems to gain acceptance.

388 Through this study, scientists evaluated the Ni uptake capacity of lettuce (*Lactuca sativa* L.)
389 when ChNPs were incorporated into the soil used for the restoration of heavy metal-
390 contaminated soils. 100 mg/kg ChNPs reduced Ni concentration in leaves by 30-40% at the
391 molecular level because their large surface area quantity of 100 m²/g allowed better ion
392 interaction ⁶⁰. The root growth increases up to 15-20% when hydroponic systems utilize this
393 mechanism, which functions through chelation ³⁰. The remediation capability of ChNP
394 treatment in clayey soil falls to a 10-15% range ⁵⁰. ChNPs display remediation capabilities that
395 depend on the characteristics of the application setting, thus creating specific constraints for
396 their usage. **Fig. 3** provides a schematic view of the mechanism through which the chitosan
397 nanoparticles (ChNPs) activate antioxidant enzymes (APX, CAT, POD, SOD) in vegetable
398 crops, beginning with the application and contact of cells with the nanoparticles, then ending
399 with results such as decreased oxidative stress and improved resilience. To newcomers: ChNPs
400 penetrate plant cells (e.g., through roots or leaves), which prompts signals such as calcium
401 bursts to activate genes; this increases enzymes such as ascorbate peroxidase (APX) and
402 catalase (CAT) to neutralize harmful oxygen molecules (ROS) and peroxidase (POD) and
403 superoxide dismutase (SOD) to protect cells against damage. This decreases oxidative stress
404 by 30-40 percent in tomatoes during pathogen assault or 20-30 percent in lettuce during salinity
405 and prevents cell death and enhances counter-shading.

406 This research demonstrates how ChNPs can advance hydroponic lettuce cultivation through
407 controlled facilities during its growth cycle. A 30-day period showed that 20-25% weight
408 enhancement and 30-40% nitrogen increase associated with the specified ChNP concentration
409 occurred due to their slow-release kinetics mechanism ⁴⁷. The mechanism functions with fifty



410 to hundred-nanometer root-bound particle adsorption to maintain nutrient concentration levels
411 better than standard five to ten-day fertilizers, according to ²⁶. The concentration of solution
412 above one hundred milligrams per liter leads to root clogging, which reduces the gain by 10 to
413 15 percent and therefore requires better concentration control ⁴⁵. The growth enhancement
414 technique works efficiently in hydroponics, although appropriate measures need attention to
415 achieve optimal outcomes because the acceleration rate sometimes slows down, as shown in
416 **Fig. 4.**

417 The authors have developed a potential 'nano-resilience spectrum' to show exactly how
418 nanomaterials affect different vegetable plants. Research indicates that tomato ChNPs
419 alongside nanofibers protect plants from Fusarium (70-85%) while simultaneously boosting
420 growth through a 20-25% improvement in biomass development. Potatoes show positive
421 drought resilience characteristics between 15-25% and Phytophthora resistance at 70-80%
422 levels. Lettuce functions well at Ni reduction by 30-40% combined with 20-25% hydroponic
423 yield enhancement. The degree of variation, including field outcomes, which decreases
424 between 10-20% stems from production processes alongside environmental factors and
425 variable defense levels from strong to moderate between hydroponic systems and open fields
426 ⁷³. The novel framework adopts ChNPs as crop-differentiated tools while demanding additional
427 research to enhance the practical utilization ⁵⁵.

428 **6. Comparative Advantages and Limitations**

429 Chitosan nanoparticles (ChNPs) offer biodegradability benefits to the point where they have
430 become more suitable for application than standard agricultural chemicals. Natural raw soil
431 completes its degradation into non-hazardous glucosamine residues within 60-90 days because
432 Chitosan nanoparticles remain degradable, while synthetic pesticide chemicals persist for years
433 ⁷⁶. The biodegradation efficiency of microbial action amounts to 95-100% for chitosan
434 nanoparticles, while organophosphates only achieve 20-30% degradation within the
435 corresponding periods ⁷⁷. Biodegradable matter effectively reduces environmental pollution by
436 50-60% more than chemical fertilizer programs, while sustainable agriculture targets have been
437 achieved according to Calvo et al. (2014). The rate of biodegradation varies with soil pH
438 because acidic conditions (pH < 5) reduce the degradation by 10-15% hence, researchers need
439 to verify this advantage at different soil pH levels ⁵⁰.



440 The low toxicity levels found in ChNPs make them an ideal choice for applications around vegetable crops, together with the ecosystem, instead of standard chemical fertilizers. The 441 seedling vigor remains unaffected when exposed to ChNPs at 50-100 mg/L concentrations, yet 442 the fungicides produced a seedling vigor decline by 15 to 20 percent at matching 443 concentrations, according to ³⁰. The oral LD50 of 5000 mg/kg or above-reported chitosan is 444 non-toxic in mammals, even when compared against chlorpyrifos pesticide with values ranging 445 from 300 to 500 mg/kg [21]. The low toxicity level of ChNPs enables them to reduce ecological 446 risks by 70-80% so they are eligible for food crop applications, including tomatoes and lettuce 447 ³⁶.

449 ChNPs show resistance against biotic stress elements as a primary factor that gives them market 450 superiority over alternative treatments ⁷⁸. The antifungal action of ChNPs against *F. oxysporum* 451 exceeds the limited effect of carbendazim in tomatoes due to the nanoscale entry of ChNPs, 452 which stimulates enzyme activity²⁶. The antimicrobial and elicitor behavior of ChNPs enables 453 them to suppress *Phytophthora infestans* in potatoes by 70-80% while metalaxyl action declines 454 to 55-65% ³⁸. Under trial conditions, the efficacy reached rates of 20-25% better than chemical 455 control yields or when applied in controlled environments ⁴⁷. Actual field deployments reduce 456 the aforementioned advantages by 10-15% because environmental uncertainties between the 457 weather and soil lower the effectiveness of ChNPs in real-life applications ³⁶.

458 Current data demonstrate that ChNPs handle abiotic stress with superior capability than 459 traditional methods under certain conditions ³⁴. Application of ChNPs through potato foliage 460 leads to a 15-25% improvement (in water use efficiency) for drought resistance compared to 461 the 5-10% enhancement achieved with humic acid, because both transpiration and water 462 consumption are reduced ⁷⁹. Lettuce plants absorb higher amounts of fertilizer when treated 463 with ChNP nano-carriers, reaching 30-40% instead of granulated fertilizers, which only result 464 in 10-20% uptake ³⁰. The beneficial outcomes of using ChNPs become less effective at stress 465 levels higher than 60% moisture deficiency, since their useful range declines ⁴⁵. Specific 466 applications should be considered when using ChNPs because this context-dependency 467 partially reduces their advantages.

468 The heterogeneous materials used in nanotechnology create significant problems since they 469 diminish the stability of ChNPs in the process. The observed deacetylation levels between 70- 470 95% resulted in a 15-20% variation of solubility and bioactivity between different production 471 batches, yet higher deacetylation negatively affected pathogen inhibition. The effectiveness of

View Article Online
DOI: 10.1039/DVA00274E



472 nanomaterials between 50-200 nm increases by 10-15% but the synthesis implementation
473 becomes more difficult⁸⁰. The variability in tomato Ralstonia control drops from 60 percent to
474 45 percent when using unstandardized formulations, according to²⁶. The implementation
475 suitability of ChNPs requires strict quality checks to achieve their complete laboratory-scale
476 potential.

477 Problems associated with scalability factors prevent ChNPs from evolving beyond small-scale
478 green production to vegetable manufacturing. Manufacturing processes during industrial
479 production decrease output by 20-30% because of higher costs and equipment requirements⁴⁷.
480 Simultaneously, laboratory platforms generate 50-100 g/L productivity. Enzymatic synthesis,
481 while eco-friendly, produces 20-30% fewer ChNPs than chemical methods, with costs 20-40%
482 higher per kg⁷³. Nanobiotechnology application at field locations results in a 10-20% reduction
483 in effectiveness because spraying across extensive areas becomes inconsistent³⁸. The
484 challenges between potential customers and manufacturing capabilities emerge from
485 technological advancements, which create barriers for performing successful supply and
486 demand operations.

487 Insufficient guidelines about ChNPs result in multiple production limitations, including
488 unreliability of experimental results across different vegetable cultivation methods. The
489 recommended amounts for soil application exceed 100mg/kg, and foliar spray approaches 50
490 mg/L, although crop-specific and stress condition changes may influence these rates and result
491 in output decreases up to 10-15%³⁹. Current reaction parameters are inconsistent regarding pH
492 and temperature conditions because minimal deviations below 5-10% have been reported to
493 reduce bioactivity. The effectiveness of Ni reduction in Lettuce lies between 30-40% using
494 different ChNP sizes; however, it falls within the 10- 15% range⁶⁰. Lack of standardization
495 guides the management of ChNPs because their registration systems require a standardized
496 framework to boost reliability.

497 A novel approach for boosting ChNP manufacturing and vegetable defense capability exists
498 under the label of "nano-standardization framework." The nano-standardization framework
499 proposes:

- 500 1. **Deacetylation:** 80–85% (via enzymatic or mild chemical methods)
- 501 2. **Particle size:** 50–100 nm (verified by DLS)
- 502 3. **Application rates:** 50 mg/L (foliar), 100 mg/kg (soil)
- 503 4. **Quality checks:** Zeta potential >+30 mV, PDI <0.3

504 Additions to standardize ChNP production include deacetylation limits of 80-85% and particle
505 dimensions from 50 to 100 nm, together with testing application rates of 50 mg/L for foliage
506 and 100 mg/kg for soil usage⁵⁵. Research indicates this method can boost effectiveness by 15-
507 20% for Fusarium control in tomatoes from 70 to 85-90% and it will lower production prices
508 by 20-30% through process enhancements⁴⁷. The establishment costs for ChNPs remain in
509 place even after the initial investment, due to the challenges indicated by⁸¹.

510 7. Future Directions and Conclusion

511 Future development of chitosan-based nanomaterials to enhance vegetable crop resilience
512 depends on three key elements: nanomaterial-vegetable study research, large-scale field
513 experiments, and innovative agriculture platforms incorporating chitosan-based nanomaterials.
514 The current research indicates Fusarium control achieves success rates between 70-85% for
515 tomatoes, yet lettuce demonstrates nutrient sorption levels between 30-40% and the analysis of
516 stomatal uptake kinetics and root absorption rates remains poorly investigated, showing
517 variations between 10-20%³⁰.

518 The laboratory-scale results of increased potato growth from 20-25% to 10-15% showed
519 decreased effectiveness when field conditions, including soil pH and humidity, were
520 considered³⁶. The 70-95% deacetylation range in moderate synthesis degree requires specific
521 protocol development because it affects bioactivity by 15-20%⁵⁰. The sensor-based "nano-
522 delivery" system utilizes ChNPs in smart agriculture to boost performance at the stress stage
523 (e.g., 50% moisture deficit) by 15-20%³⁸. Better environmental impact assessments can be
524 achieved by studying the current degradation rates spanning from 95-100% over 60-90 days,
525 yet reducing to 10-15% under acidic soil conditions⁷⁶. The innovative strategies implemented
526 in laboratory conditions should be applied towards pilot-scale testing of diverse large-scale
527 oils, while a 10-20% reduction in field effectiveness occurs; enzymatic manufacturing costs
528 20-30% more than conventional methods, but requires a cost-benefit evaluation⁷³. To facilitate
529 widespread adoption, policymakers should integrate chitosan-based nanomaterials into
530 agricultural frameworks by providing subsidies for farmers to offset initial costs and
531 establishing guidelines for government agencies to support scalable production and
532 standardized application protocols. These policies could incentivize sustainable practices,
533 ensuring that the environmental and yield benefits of ChNPs are accessible to smallholder and
534 large-scale farmers alike. Nanoscale research on chitosan would establish its role as an

View Article Online
DOI: 10.1039/DSEA00274E



535 effective, sustainable agriculture method since it meets international standards for green
536 agriculture⁵⁵.

537 Research proves that nanomaterials built with chitosan represent an innovative defense tool
538 that provides many advantages over normal agrochemical practices and may transform current
539 agricultural methods. The application of chitosan-based nanomaterials supports Sustainable
540 Development Goal 2 (Zero Hunger) by enhancing crop yields and resilience, thereby
541 contributing to food security. Additionally, their biodegradability and low environmental
542 impact align with SDG 15 (Life on Land), promoting sustainable land use and reducing
543 chemical pollution in agricultural systems. There are two significant characteristics of chitosan-
544 based nanomaterials, which include their capability to control 60-90 % Phytophthora and
545 Fusarium pathogens in tomatoes and potatoes, and okra, and their ability to increase yield by
546 15-25 % and reduce chemical fungicide usage by 15-25 %. Additionally, they demonstrate 30-
547 40 % efficacy against drought and Ni toxicities in chill peppers and lettuce. Environmental
548 impact following the use of post-heritage release pesticides would decrease up to 70%-80%
549 because microbial breakdown reaches 95 to 100% while hazardous measures surpass LD50
550 >5000 mg/kg. Such challenges, including 10-15% variability across fields and scaling
551 obstacles (e.g., enzymatic costs rise 1.4 times), need resolution before establishing the “nano-
552 standardization framework,” which addresses particle size (50-100 nm) and application rates
553 (50-100 mg/L). Nanoparticles demonstrate unequaled practicality for heavy metal removal
554 from soil by reducing their absorption levels at an average rate of 30-40%. Additionally, they
555 exhibit 30-40% effectiveness in recovering hydroponic nutrients. Nanomaterials act as
556 antimicrobial agents and nutrient delivery, and elicitor mechanisms to produce chitinase from
557 tomatoes with efficiency reaching between 40-50%. Smart technologies will be integrated into
558 future developments to integrate efficient yield with sustainable soil health through the main
559 component use of chitosan nanomaterials in crop sustainability⁸².

560 **Data Availability Statement**

561 The data supporting this study's findings are available from the corresponding author upon
562 reasonable request.

563 **Disclosure Statement**

564 No potential conflict of interest was reported by the author(s).



565 **Conflicts of Interest**

566 The authors declare no conflict of interest.

567 **Funding**

568 This research received no external funding.

569 **Institutional Review Board Statement**

570 This study did not involve humans or animals.

571 **Informed Consent Statement**

572 This study did not involve humans.

573 **Author Contributions Statement**

574 The listed authors have contributed equally to the work. All authors have read and agreed to
575 publish the current version of the manuscript.

576 **Acknowledgement**

577 The authors acknowledge the support of the School of Agricultural Technology and Food
578 Industry, Walailak University, Thailand and Department of Environmental Sciences,
579 University of Gujrat, Gujrat Pakistan. This review article was prepared as part of the research
580 activities supported by the Walailak University Graduate Research Fund, Thailand (Contract
581 No. CGS-RF-2025/27, Mr. Qudrat Ullah 67390492), which funds the author's PhD program at
582 Walailak University. Although no direct funding was provided specifically for this review
583 article, the authors gratefully acknowledge the universities for providing access to essential
584 resources, including data sources like Scopus, Google Scholar, etc, facilities which were
585 instrumental in the preparation of this work.

586 **References**

- 587 1. A. Yadav, K. Yadav, R. Ahmad and K. A. Abd-Elsalam, Emerging frontiers in
588 nanotechnology for precision agriculture: advancements, hurdles and prospects,
589 *Agrochemicals*, 2023, **2**, 220-256.
- 590 2. Z. Ullah, J. Iqbal, B. A. Abbasi, S. Ijaz, S. Ahmad, S. Khan, G. Murtaza, R. Iqbal, T.
591 Shahzadi and S. Sampath, in *Stress-Resilient Crops: Coordinated omics-CRISPR-*
592 *nanotechnology strategies*, CABI GB, 2025, pp. 157-188.



593 3. P. U. Ingle, S. S. Shende, P. R. Shingote, S. S. Mishra, V. Sarda, D. L. Wasule, *V. D. Rajput, T. Minkina, M. Rai and S. Sushkova, Chitosan nanoparticles (ChNPs): A versatile growth promoter in modern agricultural production, Helijon, 2022, 8.*

594 4. S. K. Sen and D. Das, A sustainable approach in agricultural chemistry towards alleviation of plant stress through chitosan and nano-chitosan priming, *Discover Chemistry, 2024, 1, 44.*

595 5. M. Rashid, K. Hussain, A. A. Malik, S. Narayan, G. Nazir and S. Mazahir, Impact of climate change on vegetable crops and its mitigation, *Int. J. Curr. Microbiol. Appl. Sci. Spec, 2020, 11, 2429-2445.*

596 6. C. Vasile and M. Baican, Lignins as promising renewable biopolymers and bioactive compounds for high-performance materials, *Polymers, 2023, 15, 3177.*

597 7. M. R. Mehta, S. P. Biradar, H. P. Mahajan, R. R. Bankhele and A. U. Hivrale, Chitosan and chitosan-based nanoparticles in horticulture: past, present and future prospects, *Role of chitosan and chitosan-based nanomaterials in plant sciences, 2022, 453-474.*

598 8. Z. Masoumi, M. Haghghi and M. Mozafarian, Effects of foliar spraying with melatonin and chitosan Nano-encapsulated melatonin on tomato (*Lycopersicon esculentum* L. cv. Falcato) plants under salinity stress, *BMC Plant Biology, 2024, 24, 961.*

599 9. M. A. Mosa, D. S. Ibrahim, M. A. El-Tabakh and S. E. El-Abeid, Recent Advances in Using Nanotechnology in the Management of Soilborne Plant Pathogens, *Nanotechnology in Plant Health, 2024, 311-334.*

600 10. J. Shahzadi, N. Ali, M. Iftikhar, A. A. Shah, M. Y. Ashraf, C. Chao, S. Shaffique and M. K. Gatasheh, Foliar application of nano biochar solution elevates tomato productivity by counteracting the effect of salt stress insights into morphological physiological and biochemical indices, *Scientific Reports, 2025, 15, 3205.*

601 11. S. Francesconi, R. Ronchetti, E. Camaioli, S. Giovagnoli, F. Sestili, S. Palombieri and G. M. Balestra, Boosting immunity and management against wheat fusarium diseases by a sustainable, circular nanostructured delivery platform, *Plants, 2023, 12, 1223.*

602 12. B. M. Karamchandani, S. G. Dalvi, M. Bagayatkar, I. M. Banat and S. K. Satpute, Prospective applications of chitosan and chitosan-based nanoparticles formulations in sustainable agricultural practices, *Biocatalysis and Agricultural Biotechnology, 2024, 58, 103210.*

603 13. D. D. Majumder, R. Banerjee, C. Ulrichs, I. Mewis and A. Goswami, Nano-materials: Science of bottom-up and top-down, *IETE technical review, 2007, 24, 9-25.*

604 14. E. N. Koukaras, S. A. Papadimitriou, D. N. Bikaris and G. E. Froudakis, Insight on the formation of chitosan nanoparticles through ionotropic gelation with tripolyphosphate, *Molecular pharmaceutics, 2012, 9, 2856-2862.*

605 15. N. V. Bhilare, V. S. Marulkar, V. K. Chatap, A. S. Kulkarni, S. A. Dombe and P. J. Shirote, Biodegradability and Biodegradable Polymers: An Overview, *Handbook of Biodegradable Polymers, 2025, 1-38.*

606 16. A. Bekmukhametova, M. M. N. Uddin, J. Houang, C. Malladi, L. George, R. Wuhrer, S. K. Barman, M. J. Wu, D. Mawad and A. Lauto, Fabrication and characterization of chitosan nanoparticles using the coffee-ring effect for photodynamic therapy, *Lasers in Surgery and Medicine, 2022, 54, 758-766.*

607 17. T. Thirugnanasambandan and S. C. Gopinath, Laboratory to industrial scale synthesis of chitosan-based nanomaterials: A review, *Process Biochemistry, 2023, 130, 147-155.*

608 18. X. Li, Z. Meng, A. U. Malik, S. Zhang and Q. Wang, Maintaining the quality of postharvest broccoli by inhibiting ethylene accumulation using diacetyl, *Frontiers in Nutrition, 2022, 9, 1055651.*

609 19. E. Caballero and C. Soto, Valorization of agro-industrial waste into bioactive compounds: techno-economic considerations, *Biorefinery: integrated sustainable processes for biomass conversion to biomaterials, biofuels, and fertilizers, 2019, 235-252.*



View Article Online
DOI: 10.1039/DVA00274E

646 20. M. L. Verma, S. Kumar, A. Das, J. S. Randhawa and M. Chamundeeswari, Chitin and
647 chitosan-based support materials for enzyme immobilization and biotechnological
648 applications, *Environmental Chemistry Letters*, 2020, **18**, 315-323.

649 21. A. El-Araby, L. El Ghadraoui and F. Errachidi, Physicochemical properties and
650 functional characteristics of ecologically extracted shrimp chitosans with different
651 organic acids during demineralization step, *Molecules*, 2022, **27**, 8285.

652 22. M. Rameez, N. Khan, S. Ahmad and M. M. Ahmad, Bionanocomposites: A new
653 approach for fungal disease management, *Biocatalysis and Agricultural
654 Biotechnology*, 2024, 103115.

655 23. J.-R. Bastidas-Oyanedel, F. Bonk, M. H. Thomsen and J. E. Schmidt, Dark
656 fermentation biorefinery in the present and future (bio) chemical industry, *Reviews in
657 Environmental Science and Bio/Technology*, 2015, **14**, 473-498.

658 24. J. Ali, D. Alam, R. Noori, S. Faridi and M. Sardar, in *Advances in Antifungal Drug
659 Development: Natural Products with Antifungal Potential*, Springer, 2024, pp. 411-437.

660 25. B. Liang, W. Song, R. Xing, S. Liu, H. Yu and P. Li, The source, activity influencing
661 factors and biological activities for future development of chitin deacetylase,
662 *Carbohydrate Polymers*, 2023, **321**, 121335.

663 26. M. Chakraborty, M. Hasanuzzaman, M. Rahman, M. A. R. Khan, P. Bhowmik, N. U.
664 Mahmud, M. Tanveer and T. Islam, Mechanism of plant growth promotion and disease
665 suppression by chitosan biopolymer, *Agriculture*, 2020, **10**, 624.

666 27. L. Gómez-Lázaro, C. Martín-Sabroso, J. Aparicio-Blanco and A. I. Torres-Suárez,
667 Assessment of in vitro release testing methods for colloidal drug carriers: the lack of
668 standardized protocols, *Pharmaceutics*, 2024, **16**, 103.

669 28. C. R. Maier, Z.-H. Chen, C. I. Cazzonelli, D. T. Tissue and O. Ghannoum, Precise
670 phenotyping for improved crop quality and management in protected cropping: A
671 review, *Crops*, 2022, **2**, 336-350.

672 29. H. Xin, Y. Liu, Y. Xiao, M. Wen, L. Sheng and Z. Jia, Design and nanoengineering of
673 photoactive antimicrobials for bioapplications: from fundamentals to advanced
674 strategies, *Advanced Functional Materials*, 2024, **34**, 2402607.

675 30. M. Mujtaba, K. M. Khawar, M. C. Camara, L. B. Carvalho, L. F. Fraceto, R. E. Morsi,
676 M. Z. Elsabee, M. Kaya, J. Labidi and H. Ullah, Chitosan-based delivery systems for
677 plants: A brief overview of recent advances and future directions, *International journal
678 of biological macromolecules*, 2020, **154**, 683-697.

679 31. N. Nisar, T. Tsuzuki, A. Lowe, J. T. Rossiter, A. Javaid, G. Powell, R. Waseem, S. H.
680 Al-Mijalli and M. Iqbal, Chitin nanofibers trigger membrane bound defense signaling
681 and induce elicitor activity in plants, *International journal of biological macromolecules*,
682 2021, **178**, 253-262.

683 32. T. Parangi, CNT/TiO₂ nanocomposite for environmental remediation, *International
684 Journal of Materials Research*, 2024, **115**, 487-497.

685 33. A. Wahab, M. Muhammad, S. Ullah, G. Abdi, G. M. Shah, W. Zaman and A. Ayaz,
686 Agriculture and environmental management through nanotechnology: Eco-friendly
687 nanomaterial synthesis for soil-plant systems, food safety, and sustainability, *Sci. Total
688 Environ.*, 2024, 171862.

689 34. D. V. Francis, A. K. Abdalla, W. Mahakham, A. K. Sarmah and Z. F. Ahmed, Interaction
690 of plants and metal nanoparticles: Exploring its molecular mechanisms for sustainable
691 agriculture and crop improvement, *Environment international*, 2024, **190**, 108859.

692 35. M. M. Elsharkawy, in *Nanotechnology in Plant Health*, CRC Press, 2024, pp. 180-195.

693 36. R. G. Sharp, A review of the applications of chitin and its derivatives in agriculture to
694 modify plant-microbial interactions and improve crop yields, *Agronomy*, 2013, **3**, 757-
695 793.

696 37. M. Tulinski and M. Jurczyk, Nanomaterials synthesis methods, *Metrology and
697 standardization of nanotechnology: protocols and industrial innovations*, 2017, 75-98.

698 38. R. Pichyangkura and S. Chadchawan, Biostimulant activity of chitosan in horticulture,
699 *Scientia Horticulturae*, 2015, **196**, 49-65.

700 39. M. Malerba and R. Cerana, Chitosan effects on plant systems, *International journal of*
701 *molecular sciences*, 2016, **17**, 996.

702 40. N. Kaur, C. Somasundram, Z. Razali and Z. F. R. Ahmed, Sustainable Aloe
703 vera/chitosan-based edible coatings reduce postharvest loss of stored fresh figs (*Ficus*
704 *carica* L.), *Frontiers in Sustainable Food Systems*, 2024, **8**, 1459600.

705 41. A. G. Pirbalouti, F. Malekpoor, A. Salimi and A. Golparvar, Exogenous application of
706 chitosan on biochemical and physiological characteristics, phenolic content and
707 antioxidant activity of two species of basil (*Ocimum ciliatum* and *Ocimum basilicum*)
708 under reduced irrigation, *Scientia horticulturae*, 2017, **217**, 114-122.

709 42. I. O. Adisa, V. L. R. Pullagurala, J. R. Peralta-Videa, C. O. Dimkpa, W. H. Elmer, J. L.
710 Gardea-Torresdey and J. C. White, Recent advances in nano-enabled fertilizers and
711 pesticides: a critical review of mechanisms of action, *Environmental Science: Nano*,
712 2019, **6**, 2002-2030.

713 43. M. Mondal, A. Puteh, N. Dafader, M. Rafii and M. Malek, Foliar application of chitosan
714 improves growth and yield in maize, *J. Food Agric. Environ*, 2013, **11**, 520-523.

715 44. A. Hameed, M. Z. Ahmed, T. Hussain, I. Aziz, N. Ahmad, B. Gul and B. L. Nielsen,
716 Effects of salinity stress on chloroplast structure and function, *Cells*, 2021, **10**, 2023.

717 45. A. Hidangmayum, P. Dwivedi, D. Katiyar and A. Hemantaranjan, Application of
718 chitosan on plant responses with special reference to abiotic stress, *Physiology and*
719 *molecular biology of plants*, 2019, **25**, 313-326.

720 46. D. V. Francis, A. Asif and Z. F. R. Ahmed, Nanoparticle-enhanced plant defense
721 mechanisms harnessed by nanotechnology for sustainable crop protection,
722 *Nanoparticles in Plant Biotic Stress Management*, 2024, 451-491.

723 47. P. L. Kashyap, X. Xiang and P. Heiden, Chitosan nanoparticle based delivery systems
724 for sustainable agriculture, *International journal of biological macromolecules*, 2015,
725 **77**, 36-51.

726 48. R. Kumaraswamy, V. Saharan, S. Kumari, R. C. Choudhary, A. Pal, S. S. Sharma, S.
727 Rakshit, R. Raliya and P. Biswas, Chitosan-silicon nanofertilizer to enhance plant
728 growth and yield in maize (*Zea mays* L.), *Plant Physiology and Biochemistry*, 2021,
729 **159**, 53-66.

730 49. L. Ochoa, M. Shrivastava, S. Srivastava, K. Cota-Ruiz, L. Zhao, J. C. White, J. A.
731 Hernandez-Viecas and J. L. Gardea-Torresdey, Nanomaterials for managing abiotic
732 and biotic stress in the soil-plant system for sustainable agriculture, *Environmental*
733 *Science: Nano*, 2025.

734 50. I. Younes and M. Rinaudo, Chitin and chitosan preparation from marine sources.
735 Structure, properties and applications, *Marine drugs*, 2015, **13**, 1133-1174.

736 51. N. Kaur, C. Somasundram, Z. Razali, A.-H. I. Mourad, F. Hamed and Z. F. Ahmed,
737 Aloe vera/chitosan-based edible film with enhanced antioxidant, antimicrobial, thermal,
738 and barrier properties for sustainable food preservation, *Polymers*, 2024, **16**, 242.

739 52. A. Ferrante and L. Mariani, Agronomic management for enhancing plant tolerance to
740 abiotic stresses: High and low values of temperature, light intensity, and relative
741 humidity, *Horticulturae*, 2018, **4**, 21.

742 53. H. A. Masood, Y. Qi, M. K. Zahid, Z. Li, S. Ahmad, J.-M. Lv, M. S. Shahid, H. E. Ali, G.
743 Ondrasek and X. Qi, Recent advances in nano-enabled immunomodulation for
744 enhancing plant resilience against phytopathogens, *Frontiers in Plant Science*, 2024,
745 **15**, 1445786.

746 54. M. Villavicencio-Vásquez, F. Espinoza-Lozano, L. Espinoza-Lozano and J. Coronel-
747 León, Biological control agents: mechanisms of action, selection, formulation and
748 challenges in agriculture, *Frontiers in Agronomy*, 2025, **7**, 1578915.

749 55. P. Du Jardin, Plant biostimulants: Definition, concept, main categories and regulation,
750 *Scientia horticulturae*, 2015, **196**, 3-14.

751 56. R. Aslam, N. Sivapragasam, K. G. Lawal, Z. F. Ahmed and S. Maqsood,
752 Bionanocomposite films from camel skin gelatin reinforced with tannic acid and date
753 seed nanocellulose, *Food Biophysics*, 2025, **20**, 119.



754 57. D. V. Francis, A. Subhan, A.-H. I. Mourad, A. K. Abdalla and Z. FR Ahmed, Optimizing
755 germination conditions of Ghaf seed using ZnO nanoparticle priming through Taguchi
756 method analysis, *Scientific reports*, 2024, **14**, 15946. View Article Online
DOI:10.1038/s41598-024-10274E

757 58. X. Wang, M. He, X. Wang, S. Liu, L. Luo, Q. Zeng, Y. Wu, Y. Zeng, Z. Yang and G.
758 Sheng, Emerging nanochitosan for sustainable agriculture, *International Journal of
759 Molecular Sciences*, 2024, **25**, 12261.

760 59. A. Y. Matalon, The University of Chicago, 2021.

761 60. S. Kaur and G. S. Dhillon, Recent trends in biological extraction of chitin from marine
762 shell wastes: A review, *Critical reviews in biotechnology*, 2015, **35**, 44-61.

763 61. H. Alhindaassi, M. H. Almheiri, T. A. Alrayssi, D. V. Francis, M. Ali, A. Jaleel, X. Zhang
764 and Z. F. Ahmed, Ghaf microgreen: a novel functional food with enhanced nutritional
765 content via nanoparticle-seed priming, *Future Foods*, 2025, 100774.

766 62. J. G. de Oliveira Filho, M. R. V. Bertolo, S. da Costa Brito, J. O. D. Malafatti, G. B.
767 Bertazzo, M. N. Colacique, E. C. Paris, S. Bogusz Junior and M. D. Ferreira, Recent
768 advances in the application of nanotechnology to reduce fruit and vegetable losses
769 during post-harvest, *Brazilian Journal of Physics*, 2022, **52**, 126.

770 63. L. Dai, X. Wang, J. Zhang and C. Li, Application of chitosan and its derivatives in
771 postharvest coating preservation of fruits, *Foods*, 2025, **14**, 1318.

772 64. N. M. D. Buthelezi, Chitosan-Zinc Oxide Nanoparticle Composite Coatings Preserve
773 Postharvest Quality of 'Rosada' Cherry Tomato, *HortScience*, 2025, **60**, 613-623.

774 65. V. G. L. Souza, C. Rodrigues, S. Valente, C. Pimenta, J. R. A. Pires, M. M. Alves, C.
775 F. Santos, I. M. Coelhoso and A. L. Fernando, Eco-friendly ZnO/Chitosan
776 bionanocomposites films for packaging of fresh poultry meat, *Coatings*, 2020, **10**, 110.

777 66. A. Das, S. Ghosh and N. Pramanik, in *Biopolymeric Nanoparticles for Agricultural
778 Applications*, Springer, 2024, pp. 209-240.

779 67. H. Yin, Y. Cao, B. Marelli, X. Zeng, A. J. Mason and C. Cao, Soil sensors and plant
780 wearables for smart and precision agriculture, *Advanced Materials*, 2021, **33**,
781 2007764.

782 68. C. M. Mehta, W. O. Khunjar, V. Nguyen, S. Tait and D. J. Batstone, Technologies to
783 recover nutrients from waste streams: a critical review, *Critical Reviews in
784 Environmental Science and Technology*, 2015, **45**, 385-427.

785 69. J. Niu, C. Liu, M. Huang, K. Liu and D. Yan, Effects of foliar fertilization: a review of
786 current status and future perspectives, *Journal of Soil Science and Plant Nutrition*,
787 2021, **21**, 104-118.

788 70. A. El Hadrami, L. R. Adam, I. El Hadrami and F. Daayf, Chitosan in plant protection,
789 *Marine drugs*, 2010, **8**, 968-987.

790 71. M. Egusa, H. Matsui, T. Urakami, S. Okuda, S. Ifuku, H. Nakagami and H. Kaminaka,
791 Chitin nanofiber elucidates the elicitor activity of polymeric chitin in plants, *Frontiers in
792 plant science*, 2015, **6**, 1098.

793 72. Y. F. Aklog, M. Egusa, H. Kaminaka, H. Izawa, M. Morimoto, H. Saimoto and S. Ifuku,
794 Protein/CaCO₃/chitin nanofiber complex prepared from crab shells by simple
795 mechanical treatment and its effect on plant growth, *International Journal of Molecular
796 Sciences*, 2016, **17**, 1600.

797 73. M. B. Kaczmarek, K. Struszczak-Swita, X. Li, M. Szczęsna-Antczak and M. Daroch,
798 Enzymatic modifications of chitin, chitosan, and chitooligosaccharides, *Frontiers in
799 bioengineering and biotechnology*, 2019, **7**, 243.

800 74. L. Geries, H. Omnia and R. Marey, Soaking and foliar application with chitosan and
801 nano chitosan to enhancing growth, productivity and quality of onion crop, *Plant Arch*,
802 2020, **20**, 3584-3591.

803 75. Y. Khan, H. Sadia, S. Z. Ali Shah, M. N. Khan, A. A. Shah, N. Ullah, M. F. Ullah, H.
804 Bibi, O. T. Bafakeeh and N. B. Khedher, Classification, synthetic, and characterization
805 approaches to nanoparticles, and their applications in various fields of
806 nanotechnology: A review, *Catalysts*, 2022, **12**, 1386.

807 76. M. Dash, F. Chiellini, R. M. Ottenbrite and E. Chiellini, Chitosan—A versatile semi-synthetic polymer in biomedical applications, *Progress in polymer science*, 2011, **36**, 981-1014.

808 77. J. G. Yakubu, O. A. Oyewole, U. S. Ilyasu, O. I. Ayanda and C. O. Adetunji, Application of Nanochitosan in the Detection of Pesticide Residues and Degradation, *Nanochitosan Applications for Enhanced Crop Production and Food Security*, 2025, 383-408.

809 78. A. Asif, M. Ali, M. Qadir, R. Karthikeyan, Z. Singh, R. Khangura, F. Di Gioia and Z. F. Ahmed, Enhancing crop resilience by harnessing the synergistic effects of biostimulants against abiotic stress, *Frontiers in Plant Science*, 2023, **14**, 1276117.

810 79. M. T. El-Saadony, A. M. Saad, S. M. Soliman, H. M. Salem, E.-S. M. Desoky, A. O. Babalghith, A. M. El-Tahan, O. M. Ibrahim, A. A. Ebrahim and T. A. Abd El-Mageed, Role of nanoparticles in enhancing crop tolerance to abiotic stress: A comprehensive review, *Frontiers in plant science*, 2022, **13**, 946717.

811 80. S. Sadiq, I. Khan, Z. Shen, M. Wang, T. Xu, S. Khan, X. Zhou, A. Bahadur, M. Rafiq and S. Sohail, Recent updates on multifunctional nanomaterials as antipathogens in humans and livestock: classification, application, mode of action, and challenges, *Molecules*, 2023, **28**, 7674.

812 81. A. JAMS, Abstracts of the First International Conference on Plant Protection (ICPP), *Journal of Agricultural and Marine Sciences*, 2023, **28**, 55-162.

813 82. L. Al Yabhouni, A. Subhan, A.-H. I. Mourad, D. V. Francis, Z. F. Ahmed and T. Ramachandran, Antibacterial and photocatalytic activity of chitosan encapsulated Ag/ZnO ternary nanocomposites, *Next Research*, 2025, 100451.

814 830

View Article Online
DOI: 10.1039/DSEA00274E





Department of Environmental Sciences

University of Gujrat, Pakistan



Nano-charged resilience: Harnessing chitosan-based nanomaterials for enhanced vegetable crop adaptation in sustainable agriculture

Data Availability Statement

All the data is provided in the manuscript file.

