

# Environmental Science Advances

Accepted Manuscript

View Article Online  
View Journal

This article can be cited before page numbers have been issued, to do this please use: R. Lougkovois, K. Parinos, G. Gkotsis, I. Hatzianestis, M. Nika, N. S. Thomaidis and A. Pavlidou, *Environ. Sci.: Adv.*, 2026, DOI: 10.1039/D5VA00287G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## Environmental significance statement

This is the first study investigating the chemical imprint of human-related activities in Saronikos Gulf, Greece, setting the baseline of chemical contamination of the study area, through novel high resolution mass spectrometric workflows. Detected substances were prioritized based on their persistent, bioaccumulative and toxic properties, as well as EU legislation values (where applicable). A total of 20 compounds presented high-risk for studied seawater, while 12 compounds presented high-risk for sediments. Among the latter, perfluorooctanesulfonic acid (PFOS) and di-(2-ethylhexyl)-phthalate (DEHP) exhibited the highest extent of predicted no-effect concentration (PNEC) exceedance in sediments. This is the first step towards unravelling the chemical imprint of major anthropogenic activities in the Eastern Mediterranean marine environment.



1 **Occurrence, distribution and risk assessment of organic micropollutants in**  
2 **the Saronikos Gulf, Greece, utilizing LC-TIMS-HRMS**

3  
4  
5  
6 Rallis Lougkovois<sup>1,2</sup>, Constantine Parinos<sup>1,\*</sup>, Georgios Gkotsis<sup>2</sup>, Maria-Christina Nika<sup>2</sup>,  
7 Ioannis Hatzianestis<sup>1</sup>, Alexandra Pavlidou<sup>1</sup>, Nikolaos Thomaidis<sup>2,\*\*</sup>  
8  
9  
10

11 <sup>1</sup> Hellenic Centre for Marine Research, Institute of Oceanography, 46.7 Km Athens-Sounio  
12 av., Mavro Lithari, 19013, Anavyssos, Attiki, Greece

13 <sup>2</sup> National and Kapodistrian University of Athens, Department of Chemistry, Laboratory of  
14 Analytical Chemistry, University Campus, Zografou, 15771, Athens, Greece  
15  
16

17  
18 **Corresponding authors:**

19 \* Hellenic Centre for Marine Research, Institute of Oceanography, 46.7 km Athens-Sounion  
20 Ave., Anavyssos, 19013, Athens, Greece - Constantine Parinos, email: [ksparinos@hcmr.gr](mailto:ksparinos@hcmr.gr), tel:  
21 +30 2291076372

22 \*\* National and Kapodistrian University of Athens, School of Science, Department of  
23 Chemistry, Laboratory of Analytical Chemistry, Panepistimiopolis Zografou  
24 157 71 Athens, Greece - Nikolaos Thomaidis, email: [ntho@chem.uoa.gr](mailto:ntho@chem.uoa.gr), tel: +30 2107274317



## Abstract

This study investigates the occurrence, distribution and ecological risk of emerging contaminants (ECs) and priority pollutants (PPs) in seawater and sediments of the Saronikos Gulf, and Elefsis Bay, Greece, an area continuously impacted by wastewater treatment plant (WWTP) effluents, industrial discharges and maritime traffic. Utilizing novel liquid chromatography tandem ion-mobility spectrometry and high-resolution mass spectrometry the occurrence of more than 4,000 LC-amenable organic micropollutants was investigated through wide-scope target and suspect screening. A total of 171 analytes were detected in marine samples, with pharmaceuticals identified as the most prevalent class (36% in seawater, 41% in sediments) followed by plant protection products (18% in seawater, 27% in sediments). Per- and polyfluoroalkyl substances (PFAS) were also detected in both matrices. Semi-polar ECs with higher molar mass were determined exclusively in sediments near WWTPs, possibly due to their high log P values, reflecting their affinity for particulate matter. Additionally, seawater circulation pattern was found to play a significant role in controlling the spatial distribution of ECs. Comparison with earlier studies in the area suggests a clear shift in pharmaceutical usage by the local population. Risk assessment, based on risk quotient calculations and environmental quality standards (EQS) set by EU legislation, revealed that PFAS exceeded annual average environmental quality standard values in 92% of seawater samples, whereas 20 ECs in seawater and 12 in sediments exceeded predicted no-effect concentrations (PNECs), indicating potential adverse effects on marine biota.

**Keywords:** ion mobility spectrometry, wastewater, seawater, sediments, wide-scope target screening, suspect screening



## 50 Highlights

View Article Online  
DOI: 10.1039/D5VA00287G

- 51 • Baseline organic pollution status in the Saronikos Gulf and Elefsis Bay
- 52 • WWTPs were identified as the main sources of pollution in the study area
- 53 • Suspect screening of endocrine-disrupting surfactants was conducted
- 54 •  $\Sigma$ PFAS concentrations exceeded AA-EQS values in 92% of tested seawater samples
- 55 • Circulation and precipitation patterns are discussed

56



## 1. Introduction

Over the past years, emerging contaminants (ECs) have gained the wide interest of the scientific community, as well as national and international environmental agencies (US-EPA; EU-EEA), because of the possible risks they pose towards both human health and environmental ecological balance<sup>1,2</sup>. Among others, ECs include pharmaceutically active compounds (PhACs), personal care products, artificial sweeteners, industrial chemicals, per- and polyfluorinated alkyl substances (PFAS), plant protection products and surfactants, along with their transformation products (TPs)<sup>1,3,4</sup>.

Potential sources of ECs in the marine environment include a wide variety of human-related activities, such as landfill leachates, surface run-offs, atmospheric deposition, application of biosolids, metallurgy, distilleries, textile/paper/dye/pharmaceutical industries, manure of agricultural land and shipping activities. Moreover, small- and large-scale wastewater treatment plants (WWTPs) have been highlighted as major sources of ECs in the marine environment, since the complete removal of ECs even with tertiary treatment technologies is yet to be achieved. It is reported that in WWTPs, ECs removal efficiency varies in the range of 20-50%, 30-70%, and, in some cases, >90% during the primary, secondary, and tertiary treatment steps, respectively, depending on their physicochemical properties and utilized treatment technologies<sup>5,6</sup>. The main mechanisms for removal of ECs occurring during the secondary treatment at WWTPs are biological and/or chemical transformation and sorption. The most common employed processes in WWTPs are activated sludge (CAS) and membrane biological reactors (MBRs). The efficiency of a CAS system depends on the physicochemical characteristics of the substances and on the nature of the microbial community. MBRs emerged as an alternative to CAS, integrating aerobic biodegradation and membrane separation, modestly more efficient than CAS in the extent of removal of several CECs<sup>5</sup>.



Following the release of ECs in the marine environment, biotic and abiotic processes take place, leading to their degradation in seawater, sediments and biofilms, while their distribution depends on their varying physicochemical properties, such as water solubility, solid-liquid distribution, polarity, vapor pressure and environmental conditions like pH values, organic matter content and water temperature, leading to the formation of metabolites and TPs<sup>7</sup>. These compounds can potentially pose even greater adverse effects than parent substances towards human health and marine environmental quality status<sup>8,9</sup>.

Sediments constitute a significant long-term repository for moderate to high lipophilicity ECs in the marine environment considering their high logP values<sup>10</sup>. Their accumulation in sediments may affect benthic communities, due to bioaccumulation and bio-transfer along the food chain<sup>11,12</sup>. Sediment and benthic communities serve as comprehensive indicators of anthropogenic impacts in ecological and environmental status assessments, as outlined by EU directives (Marine Strategy Framework Directive - MSFD; Water Framework Directive - WFD)<sup>13</sup>. To that end, the environmental protection agency (EPA) of the United States has been recently collecting environmental monitoring data to develop exposure computation mechanisms, aiming to facilitate quicker strategy development and decision-making by regulatory authorities worldwide<sup>4</sup>.

Since 2015, the European Commission has established watch-lists of substances for Union-wide monitoring in the field of water policy regarding the occurrence of ECs in aquatic environments<sup>14–16</sup>. These watch-lists are updated at least every two years, and recently, the EU Commission adopted a new watchlist of substances in surface waters (Decision EU 2025/439), suspected of posing a risk to the environment and human health<sup>16,17</sup>. As of now, a total of 67 different chemicals have been included in watch-lists since 2015 (five different EU decisions, namely: 2015/495, 2018/840, 2020/1161, 2022/1307 and the latest, 2025/439). The latest update included 29 substances to be investigated in the field of water policy. Member States



should monitor listed substances to provide data on their concentrations and environmental presence, thus helping regulators to determine whether these substances pose a widespread threat. For substances posing such a risk, maximum allowable threshold values should be set, and measures should be taken to reduce or phase out their emissions. Thus, it is highly important for member states to monitor the occurrence of ECs in the marine environment. However, only a few related studies have been published to date, focusing on a limited number of compounds from specific chemical groups, like industrial chemicals and pharmaceuticals<sup>18–21</sup>.

Following recent advances in high resolution mass spectrometry (HRMS), the interest of many scientific groups has shifted towards simultaneous determination of numerous ECs with various physicochemical properties, belonging to different chemical groups<sup>22–24</sup>. Besides broadening the target group of ECs, HRMS provides the ability to perform suspect and non-target screening, further multiplying the number of ECs and their TPs that can be traced<sup>25,26</sup>. To the best of our knowledge, published data on the occurrence of ECs in the Greek Seas are scarce, focusing either on a limited number of pharmaceuticals, along with their human metabolites and TPs in seawaters, via low resolution mass spectrometry (LRMS)<sup>27,28</sup>, and on the utilization of HRMS techniques on both seawaters and marine sediments<sup>29,30</sup>.

## 2. Study aim

Herein, a wide-scope target screening of more than 2,500 semi-polar to polar LC-amenable ECs from different chemical classes is conducted in the water column and sediments of the Saronikos Gulf and Elefsis Bay, Greece, by applying a novel analytical methodology utilizing liquid chromatography trapped ion mobility spectrometry tandem high-resolution mass spectrometry (LC-TIMS-HRMS). Additionally, a suspect screening was performed, covering a wide variety of surfactants, cleansers and antiseptic substances, some of which





exhibit endocrine disrupting properties<sup>31</sup>, further leveraging the capabilities of HRMS. These compounds cover a wide chemical domain with different physicochemical properties, such as molecular weight and polarity. The targeted ECs' occurrence and distribution patterns in the water column and sediments are studied in relation to anthropogenic activities, their physicochemical properties and the characteristics of the studied marine environment, facilitating the investigation of possible point sources and accumulation mechanisms. An ecological risk assessment was also conducted, aiming to prioritize detected chemicals risk-wise.

This study establishes a baseline assessment of the environmental quality of the Saronikos Gulf and Elefsis Bay regarding ECs, thus setting the baseline for future scientific research in the area. Additionally, it explores the connection between specific substances and wastewater treatment plant (WWTP) effluents. To support the discussion on EC sources, influent and effluent wastewater samples from the two major WWTPs operating in the region - Thriassio and Psyttalia - were also analyzed, simultaneously evaluating both the removal efficiencies and chemical impacts of these WWTPs on the studied marine environment.

### 3. Study area and sampling design

The Saronikos Gulf is located in the central Aegean Sea, northeastern Mediterranean Sea [Figure 1]. With a coastline of ~270 km, a sea surface area of ~2,866 km<sup>2</sup> and an average water depth of ~100 m, the gulf constitutes the coastal receptor area of the metropolitan city of Athens, capital city of Greece, with more than 5,000,000 inhabitants. The Saronikos Gulf is subjected to intense anthropogenic pressure from several point and non-point pollution sources. In detail, these include intense maritime traffic, industrial activities, oil refineries, marinas, touristic facilities, the industrial zone of Elefsis Bay and Piraeus port (the largest port in Greece and one of the biggest in Europe) affect the marine environment of the area via runoff and/or



submarine groundwater discharges, as well as atmospheric deposition of ECs from adjacent urban areas<sup>32–34</sup> **[Figure 1]**.

Most importantly, the inner Saronikos Gulf, at the northern part of its eastern basin, receives effluent wastewater from the second largest WWTP (Psyttalia island) in Europe and one of the biggest worldwide (capacity of 5,600,000 p.e and flow of approximately 800,000 m<sup>3</sup> d<sup>-1</sup>; provision of 1,000,000 m<sup>3</sup> d<sup>-1</sup>)<sup>35,36</sup>. The Psyttalia WWTP carries out pre-treatment, primary and secondary treatment with advanced biological nitrogen removal, sludge treatment and cogeneration of electrical and thermal energy. Until 1994, sewage from Athens was discharged as untreated wastewater in the Saronikos Gulf, north of Psyttalia, directly affecting Elefsis Bay. During the period 1994 - 2004 the sewage was primarily treated, while from 2004 onwards, an advanced secondary sewage treatment plant is also operating. Psyttalia WWTP uses advanced secondary biological treatment based on the activated sludge process. The treated sewage effluent is dispersed into the Saronikos Gulf at 63m depth, following the prevailing circulation, and is traced to the south, southeast and southwest of the sewage outfall<sup>37</sup>, below the pycnocline (extending between depths of ~40 and ~70 m) during the stratified period (May - November), or in the surface layer during the cold period (December - April). Lower dissolved oxygen (~3.00 mL L<sup>-1</sup>) values were recorded near the bottom of the sampling stations located mainly southwest of Psyttalia. This decrease had been systematically observed in the past and is probably due to the oxidation of organic matter and has been associated with the treated wastewater field and its dispersion in the inner Saronikos Gulf depending on the prevailing circulation. The dissolved oxygen concentrations usually recorded in the study area during winter-early march period are ~6.00 mL L<sup>-1</sup> <sup>13,38</sup>.

Elefsis Bay is an almost enclosed small and shallow embayment, with an area of 71.5 km<sup>2</sup> and an average water column depth of 33 m, which is connected to the Saronikos Gulf by two narrow and shallow channels **[Figure 1]**. Elefsis Bay is differentiated from the Saronikos



180 Gulf not only in its morphology, but also in the extent of the chemical pollution it receives. A  
181 significant number of industries (oil refineries, shipyards, chemical plants, food, metal cement  
182 industries, etc.) are located along its northern coastline. Elefsis Bay is considered a heavily  
183 impacted marine area, whereas the marine environment of the Saronikos Gulf is characterized  
184 as heavily (near the Psytalia sewage outfall) to moderately polluted<sup>13</sup>. The treated sewage of  
185 the Thriassio WWTP affects the eastern part of Elefsis Bay. The Thriassio WWTP has been  
186 operating since 2012 and treats wastewater of the surrounding municipalities (Aspropyrgos,  
187 Elefsis, Mandra and Magoula), as well as the surrounding industrial and craft areas (capacity  
188 of 117,000 p.e and flow  $\sim 21,000 \text{ m}^3 \text{ d}^{-1}$ ). Thriassio WWTP uses secondary biological treatment  
189 based on the activated sludge process.

190 In March 2023, the Hellenic Centre for Marine Research (HCMR) conducted a sampling  
191 campaign using the research vessel (R/V) “Aegaeo” in the wider Saronikos Gulf and Elefsis  
192 Bay. The campaign involved the collection of 13 seawater samples from various depths  
193 (surface, 50 m and near bottom) and 4 surface sediment samples. Additionally, influent (n=2)  
194 and corresponding effluent (n=2) wastewater samples were obtained from both wastewater  
195 treatment plants (WWTPs) operating in the region (Thriassio and Psytalia). The sampling  
196 followed the treatment stage timeframes at each facility, ensuring a 24-hour interval between  
197 influent and effluent collection for both WWTPs. Seawater samples were collected with 12 L  
198 Niskin bottles mounted on a rosette sampling system. In each case, 2.5 L were transferred to  
199 amber-glass bottles with Teflon-lined caps and stored in the fridge at 4°C. Marine sediments  
200 were collected using a stainless-steel Box Corer with a surface area of 40×40 cm, wrapped in  
201 pre-combusted aluminum foil and stored in the freezer at -20°C. Wastewater samples were  
202 received in 1 L HDPE bottles and immediately stored in the freezer at -20°C. The sampling grid  
203 was designed primarily to assess the impact of wastewater treatment plants (WWTPs) on the  
204 marine environment. Additionally, other influencing factors, such as maritime traffic, industrial



activities, and land-derived pressures, were also considered. The sampling stations were named in accordance with the National Monitoring Project for the implementation of the WFD in Greece. Sampling stations S7, S8 and S11 are positioned in the Saronikos Gulf [Figure 1], S7 is located very close to the mouth of the pipe discharging from the Psytalia WWTP, while stations S8 and S11 are located 4.0 nmi southwest and southeast of the WWTP. These two stations are also influenced by various anthropogenic pressures [Figure 1]. Finally, station S1 is located at the north-eastern part of the Elefsis Bay and is affected by the Thriassio WWTP, as it is located ~1 nm from the sewage pipe. Stations S1-biol and S7-biol are located at the sewage pipes, receiving direct discharges from the Thriassio and Psytalia WWTPs respectively [Figure 1]. Details regarding sampling stations and information for their physicochemical characteristics are presented in the Supplement (Sect. 1: Sampling stations) [Tables S1 & S2].

## 4. Materials and methods

### 4.1. Sample preparation

Upon reaching the lab, samples were immediately treated through generic sample preparation workflows, based on previously reported methods<sup>28,29,39–42</sup>. These methods consist in solid phase extraction (SPE) for seawater and wastewater samples, and an in-house developed ultrasonic assisted extraction (UAE) method for sediments with minor modifications. The utilization of untargeted protocols aimed to extract as many LC-amenable, semi-polar to polar, thermally unstable, organic ECs from a variety of chemical groups, along with their respective human metabolites and TPs. Studied chemical classes include coffee and tobacco related compounds, artificial sweeteners, illicit drugs, industrial chemicals, PFAS, personal care products, pharmaceuticals, plant protection products and surfactants.

In brief, pH of 1L seawater and 100 mL wastewater samples was initially adjusted to  $6.5 \pm 0.2$  and then spiked with a mixture of deuterated internal standards (I.S.), recorded in the



Supplement [Table S4]. Samples were then loaded onto a mixed-mode cartridge SPE system. Elution was performed with 6 mL of a basic solution, followed by 4 mL of an acidic solution. The eluents were evaporated to dryness under a gentle N<sub>2</sub> stream and reconstituted to a final volume of 200 µL with a mixture of H<sub>2</sub>O/MeOH 50/50 v/v, thus attaining a 5,000-fold pre-concentration factor for seawater samples and a 500-fold pre-concentration factor for wastewater samples.

Regarding marine sediment samples, 1 g of each lyophilized sample was initially transferred to a 15 mL polyethylene centrifuge tube and spiked with the same I.S. mixture as seawater samples. Afterwards, 3 mL of an extraction mixture, consisting of H<sub>2</sub>O/MeOH 50/50 v/v, 0.1% EDTA, and 0.5% formic acid, were added, and each tube was shaken by hand, then via vortex, then inserted in a 50°C ultrasonic bath for 15 min. Samples were centrifuged for 10 min at 4,000 rpm, and supernatant liquids were collected in glass tubes. The whole process was repeated two more times until a final extract of 9 mL was collected for each sample. Extracts were again evaporated to dryness and reconstituted to a final volume of 200 µL with a mixture of H<sub>2</sub>O/MeOH 50/50 v/v, thus attaining a 5-fold pre-concentration factor. In this calculation, we assumed that 1 g of sediment is equivalent to 1 mL of sample, considering the bulk density of sediments, which is close to 1 g mL<sup>-1</sup>.

Further information regarding sample preparation procedures is presented in the Supplement (Sect. 2: Sample preparation).

#### 4.2. Instrumentation

The separation of the analytes was carried out using an ultra-high performance liquid chromatograph (UHPLC) (Elute LC, Bruker, Bremen, Germany) with a 1.8 µm C18-2 100×2.1 column (Intensity Solo, Bruker, Bremen, Germany). A hybrid trapped ion mobility spectrometer (TIMS) coupled to a hybrid quadrupole time-of-flight mass spectrometer (QTOF-MS) was employed (timsTOF Pro 2, Bruker, Bremen, Germany). A novel ionization source,



vacuum insulated probe heated electrospray ionization (VIP-HESI) was employed, aiming to enhance ionization efficiency without in-source fragmentation of thermally unstable analytes.

The HRMS system operated in both positive and negative ionization modes, producing data from data independent acquisition (DIA), in which extraction of both full scan MS and MS/MS, without pre-selection of ions, was conducted. This mode, also referred to as broad band Collision Induced Dissociation (bbCID), was utilized for quantification purposes. Auto sampler injection volume was set at 1  $\mu$ L for positive and at 3  $\mu$ L for negative ionization mode, ensuring acceptable peak intensity both for identified analytes' parent ions and their respective fragments. Samples were analyzed in TIMS-ON mode, making use of the instrumentation's added value.

#### 4.3. QA/QC procedures

To ensure the instrumentation's proper function, a system suitability test (SST) was analyzed at the beginning of the sequence. This aliquot contains a number of compounds representative of the database, which are eluted throughout the whole chromatogram, while its analysis provides data regarding elution status, retention time (RT) shift and mass error. A reference standard solution (STD) was analyzed throughout the sequence, followed by pure solvent aliquots. Analyses of said solutions facilitate the evaluation of the instrument's sensitivity and the assessment of possible carry-over effects, respectively. Compounds detected in the pure solvent mixture were not reported if their intensity surpassed 5% of the previous injection. Detailed information is available in the Supplement (Sect. 4.2. Instrumentation performance evaluation).

A field blank sample was generated by leaving a 2.5 L amber-glass bottle filled with milli-Q water with the cap off during seawater sampling. For wastewater analysis, two 1L HDPE bottles were filled with milli-Q water as field blanks and placed close to the sewage pipes with the caps off during influent and effluent wastewater sampling, respectively. Three



procedural blank samples were generated, one for each environmental matrix analyzed. As a seawater procedural blank, 1 L of milli-Q water was utilized, 9 mL of pure extraction solvents mixture were used as a procedural blank for sediment analysis, while 100 mL of milli-Q water were used as a procedural blank for wastewater analysis. Field and procedural blank samples were treated and analyzed with the same procedures and parameters as real samples. For quantification purposes, analytes with relative areas not exceeding 150% of either blank sample's relative area were considered contaminations occurring during sampling and/or sample preparation, thus not reported in this study.

Calibration of mass and mobility accuracy was performed before analysis. Sodium formate ion cluster masses produced in the ion source were measured and compared to theoretical values. Acceptable mass accuracy was considered only if experimental mass readings matched theoretical ones by more than 99%. After that, mass readings were corrected to match the theoretical ones. Regarding mobility calibration, CCS values of compound ions included in an IMS tuning mix were evaluated and accepted if experimental ion mobility readings matched theoretical ones by more than 99%. Again, ion mobility values, expressed as  $k_0^{-1}$  were corrected to match theoretical ones.

Method accuracy was evaluated by calculating the recovery range for each tested matrix. To that end, matrix spiked samples were analyzed for 3 concentration levels, along with matrix-matched samples. All aliquots were also spiked with the I.S. mix to correct signal suppression or enhancement. For the seawater, sediment and wastewater matrices included herein, calculated recovery ranges were equal to 62 - 109%, 60 - 122% and 71 - 118%, respectively.”)

#### 4.4. Data treatment parameters

All analytes determined through wide-scope target screening are included in the target screening database developed by the Laboratory of Analytical Chemistry of the NKUA<sup>43</sup>. Analytes from the surfactants' chemical group, which were determined through suspect





screening, were included in a previously developed suspect screening dataset<sup>44–46</sup> were treated using the TASQ software (Bruker Daltonics). Strict identification criteria were applied to HRMS data, ensuring accuracy of analyte determination. These criteria concern mass-to-charge ratio ( $m/z$ ) accuracy, retention time shift, isotope profile matching, qualifier ions detection and CCS value matching. Detailed information is included in the Supplement (Sect. 5.1 Screening parameters).

#### 4.5. Analyte quantification

For target screening quantification purposes, a 6-level spike curve was produced, and relative areas of analytes were utilized for concentration calculation, as explained in the Supplement (Sect. 5.3 Quantification procedure). Quantification of surfactant compounds included in the suspect list was realized by semi-quantification procedure, based on available reference standards, and is further explained in the Supplement (Sect. 5.3 Quantification procedure). Consequently, signal-to-noise ratios (S/N) generated by the software were used to calculate the limits of detection (LOD) and quantification (LOQ), which correspond to S/N values equal to 3 and 10, respectively. Analyte concentrations not exceeding LOD values were not considered in quantification, while those with concentration levels between LOD and LOQ values were classified as “below quantification level (BQL)”. Statistical compromises were applied, in which BQL concentrations were replaced with LOQ/2 values, in the interest of calculating cumulative concentrations of chemical classes in each sample.

#### 4.6 Risk assessment

To conduct a risk assessment of the study area, predicted no-effect concentrations (PNEC) in seawater and sediment samples were collected from the NORMAN database<sup>47</sup> [Table S5]. Risk quotient factors (RQF) were calculated based on the following equation:

$$\text{RQF} = \text{MEC} / \text{PNEC}$$

MEC = measured concentration.





As MEC values, the average measured concentration was used to highlight the general risk factors, while maximum determined concentrations reflected the worst-case scenario. Compounds were consequently categorized based on calculated RQFs, as follows:

1. **RQF < 0.1**: Low risk, exposure is unlikely to promote adverse effects
2. **0.1 < RQF < 1**: Moderate risk, signifying the need for further investigation
3. **RQF > 1**: High risk, necessitating appropriate risk reduction measures

## 5. Results & Discussion

### 5.1. Chemical classification of the determined compounds

A total of 171 analytes were detected in marine samples of the Saronikos Gulf and Elefsis Bay areas, 128 of which were determined in seawater samples, while 70 were determined in sediment samples. Out of these substances, 101 were uniquely detected in seawater samples, 43 were uniquely detected in sediment samples, and 27 were distributed in both environmental compartments. Their occurrence, based on their chemical classification and sub-classification, is depicted in pie charts [Figures 2A & 2B]. Regarding wastewater analysis, 189 compounds were determined in influents and 162 in effluents. Among them, 134 analytes were commonly detected both in influent and effluent wastewater, while 52 were uniquely determined in influents, and 25 were solely detected in effluents, possibly due to transformation mechanisms of influent compounds during the cleanup process<sup>48</sup>. The occurrence of chemicals in wastewater samples is also presented in pie charts [Figures 2C & 2D]. Detailed analysis results, regarding concentration levels of ECs, are presented in the Supplement [Tables S6 & S7].

In general, more polar ECs were present in aqueous matrices, in contrast to the sediment layer, which is generally comparted by semi-polar lipophilic ECs with higher log P values [Tables S5, S6 & S7]. A similar trend is observed, regarding the molar mass of detected



compounds, with low molar mass compounds being generally present in seawater and wastewater samples, while heavier ECs tend to precipitate to the sediment layer. Literature suggests that semi- to non-polar chemicals form aggregations with microplastics through sorption effects<sup>49</sup>. These aggregations contribute to the formation of new sediment layer over a span of many years, further degrading sea bottom quality. A critical threat they pose is the recontamination of near bottom seawater, as ECs are continuously freed from microplastics due to the latter's (bio)degradation and return to the aquatic or sediment compartments.

Regarding seawater samples, the largest proportion of detected compounds were pharmaceuticals (36%, n=46) and plant protection products (18%, n=23). It is noted that a total of 12 out of the 55 PFAS included in the wide-scope target screening database were determined (9% of total compounds). A total of 19 industrial chemicals (15%), 17 surfactants (13%), 6 coffee and tobacco related compounds (5%), 3 personal care products (2%) and 2 artificial sweeteners (2%) were also present in the study area during the sampling period [Figure 2A; Table S6]. Comparing detected chemicals in different tested matrices, 58 compounds were simultaneously determined in wastewater effluents and seawater samples of the study area, reflecting on the continuous encumbrance that local WWTPs impose on the marine environment [Figure 3].

Regarding sediments, pharmaceuticals (41%, n=29) and plant protection products (27%, n=19) were the dominant chemical classes of determined analytes. A total of 9 PFAS (13%) were determined in the sampling area, along with 6 additional industrial chemicals (9%), 4 surfactants (6%), 2 coffee and tobacco related compounds (3%), and 1 personal care product (1%) [Figure 2B; Table S7]. A total of 14 compounds were simultaneously determined in wastewater effluents, seawater and sediment samples of the study area, reflecting the WWTPs' chronic chemical impact on the Saronikos Gulf and Elefsis Bay area. However, 28 substances



were commonly determined in seawater and sediment samples, which could be attributed to different point sources of contamination, besides WWTP discharges [Figure 3].

Wastewater analysis indicates similar trends, regarding determined compound classification between influents and effluents. Specifically, proportions of detected substances in influent and effluent wastewater were equal to 52% (n=98) and 53% (n=86) for pharmaceuticals, 13% (n=25) and 12% (n=20) for surfactants, 12% (n=22) and 12% (n=20) for plant protection products, 7% (n=14) and 6% (n=10) for industrial chemicals, 5% (n=9) and 9% (n=14) for PFAS, 4% (n=7) and 3% (n=5) for coffee and tobacco related compounds, 4% (n=7) and 1% (n=1) for illicit drugs, 2% (n=4) and 2% (n=3) for personal care products, as well as 1% (n=3) and 2% (n=3) for artificial sweeteners, respectively [Figures 2C & 2D; Table S8].

## 5.2. Occurrence and spatial distribution of ECs and PPs in seawater samples

Holistic results regarding seawater sample analysis can be found in the Supplement [Table S6]. Determined ECs and PPs per chemical class and sampling depth is visualized as a heatmap [Figure 4]. The most chemically burdened seawater samples, in terms of cumulative concentration of compounds, were seawater collected at the sewage discharge points of the Thriassio (S1-biol) and Psyttalia (S7-biol) WWTPs. Specifically, 45 compounds amounted to a cumulative concentration of 13.8  $\mu\text{g L}^{-1}$  in sampling station S1-biol, whereas 64 compounds amounted to a cumulative concentration of 12.0  $\mu\text{g L}^{-1}$  in sampling station S7-biol. Subsequently, 47 analytes were detected in seawater at the surface (2m) of station S1, with a cumulative concentration of 2.17  $\mu\text{g L}^{-1}$ . In seawater close to the surface (2m) of station S7, a total of 42 compounds were detected in cumulative concentration equaling 970  $\text{ng L}^{-1}$ , while close to the bottom, 29 detected substances cumulated a concentration of 846  $\text{ng L}^{-1}$ . The sewage from the Psyttalia WWTP (station S7) outflows at 63 m depth (max depth of station S7 is ~70 m) affecting the near bottom layer and probably the sediment<sup>50</sup>. During March, prevailing



hydrological conditions result in a homogenized water column, facilitating the dispersion of sewage from the Psyttalia and Thriassio WWTPs into the surface layer [48]. Although the prevailing circulation in the inner Saronikos Gulf typically follows a southwestward direction from Psyttalia<sup>35,36</sup>, surface seawater currents during the sampling days were directed southeastward (current speed: 2 - 12 cm sec<sup>-1</sup>). This suggests that ECs discharged from the Psyttalia WWTP would be more abundant at the surface layer of station S11, located southeast of the outfall, rather than at the surface layer of station S8, situated to the southwest. The current measurements (direction and speed) are further supported by the lower salinity and reduced % light transmission values recorded at station S11, confirming the dispersion of ECs southeast of Psyttalia **[Figure S1]**. According to the results, the majority of the studied ECs were indeed dispersed toward the southeastern region of Psyttalia (Figure 5) confirming the important role of hydrological conditions in controlling the spatial distribution of ECs in the study area. Moreover, SFCs were found at station S11, in higher concentrations than at station S7, probably indicating other sources than the WWTP sewage. Specifically, shipping and anthropogenic activities along the coastline, appear to contribute to their presence in the study area. Some ECs depending on their characteristics (e.g log P), may be adsorbed onto particles and start sinking in deeper layers, following the prevailing circulation patterns in the near bottom layer. ECs were distributed towards southeast and southwest of Psyttalia at the deep water layers, as it was found that in this layer, the sewage plume was dispersed towards both southeast and southwest of Psyttalia **[Figure S1]**. The majority of ECs, especially those with relatively low logP values, appear at lower concentrations at the near-bottom layer compared to the surface. **[Figure 6]**.

The highest concentration levels regarding coffee and tobacco related compounds were determined in seawater close to the surface (2m) of stations S1 and S7. Stimulants caffeine (85% FoD) and cotinine (77% FoD) were determined at concentrations ranging between 16.8 and 580 ng L<sup>-1</sup>. Caffeine and its metabolites theobromine and theophylline were determined in

influent wastewater from the Thriassio WWTP at 49.8, 29.5 and 8.14  $\mu\text{g L}^{-1}$ , respectively. Consequently, caffeine was not detected in S1-biol seawater, while its concentration levels in S1 seawater were determined at 206  $\text{ng L}^{-1}$  (2m) and 16.8  $\text{ng L}^{-1}$  (18m). Theophylline was also detected in concentrations equal to 27.8  $\text{ng L}^{-1}$  (2m) and 3.26  $\text{ng L}^{-1}$  (18m). This could be an indication of water column contamination by external factors besides WWTP effluents, such as industrial area operations in the Elefsis Bay area. The presence of compounds linked to human habits like coffee consumption and smoking is likely indicative of the vast urbanization in the Attica region and have also been reported in the past in various aquifer datasets<sup>51–54</sup>.

Corrosion inhibitors 5-methyl-benzotriazole (100% FoD) and mercaptobenzothiazole (85% FoD) were determined in concentrations ranging between 1.68 and 62.2  $\text{ng L}^{-1}$ . These compounds are used in various industrial processes and their presence could be attributed to extensive urbanization<sup>55</sup>, as well as industrial activities and maritime traffic occurring in the Saronikos Gulf area. Concentration levels of 5-methyl-benzotriazole increased post-wastewater cleanup from 2.82 to 5.08  $\mu\text{g L}^{-1}$ . In S1-biol seawater, it was determined in concentration levels equal to 3.40  $\text{ng L}^{-1}$ , showing a significant reduction. However, it was determined in the surface layer (2m) of sampling station S1 at 19.9  $\text{ng L}^{-1}$ , while in close to the bottom (18m), a concentration of 1.68  $\text{ng L}^{-1}$  was measured. This increase could be due to maritime scrubber water and industrial runoffs in the wider area of Elefsis Bay, where the largest shipyard in Greece operates, and also attests to additional point sources besides wastewater inputs. This compound has also been detected in corresponding sediments of sampling station S1, in a concentration of 0.877  $\mu\text{g kg}^{-1}$  d.w. Its determination is possibly due to its lipophilicity (log P equal to 1.40) [Table S5] and continuous accumulation in the sediment layer of Elefsis Bay.

A total of 12 PFAS were detected in the tested seawater samples. Legacy PFAS perfluorooctanesulfonic acid (PFOS) was determined at concentration levels ranging between 1.07 and 18.4  $\text{ng L}^{-1}$  in the study area, in comparable levels to similar aquifers<sup>56,57</sup>. Next



generation PFAS were also detected throughout the sampling area, like 4,8-dioxa-3H-perfluorononanoic acid (ADONA), which was determined in concentration levels ranging between 2.82 and 160 ng L<sup>-1</sup>. Its concentration levels close to the sea bottom (69m) of station S7 were equal to 160 ng L<sup>-1</sup>, while near the surface (2m), it was determined at 83.5 ng L<sup>-1</sup>. This could be due to its lipophilicity (log P value equal to 4.10), leading to sorption on suspended organic material and removal from the aqueous seawater compartment. During wastewater cleanup, longer carbon chain PFAS concentrations are generally reduced, while concentrations of smaller carbon chain PFAS are increased. For example, concentration values of PFOS and 6:2 fluorotelomer sulfonic acid (6:2 FTS) are reduced from 375 and 108 ng L<sup>-1</sup> to 133 and 58.7 ng L<sup>-1</sup>, respectively, between influent and effluent wastewater. However, perfluorobutanesulfonic acid (PFBS), perfluoropentanoic acid (PFPeA) and perfluoropentanesulfonic acid (PFPeS) concentrations increased from <LOD values to 1.63, 0.548 and 0.721 ng L<sup>-1</sup>, respectively. This could be due to PFAS carbon chains breaking down during the cleanup processes. Regarding PFOS, its concentration in S1-biol seawater was calculated equal to 18.4 ng L<sup>-1</sup> and is consequently reduced to 0.561 ng L<sup>-1</sup> in seawater close to the surface (2m) of station S1. Particularly 6:2 FTS concentration is continuously decreased between S1-biol seawater and benthic seawater (18m) of station S1, where it was determined at 25.5 and 0.321 ng L<sup>-1</sup>, respectively, whereas its concentration close to the surface (2m) was not determined above LOD values. It seems that the only source of these PFAS in the marine environment of Elefsis Bay is effluent wastewater discharges. The difference in calculated concentrations between legacy and next generation PFAS in the seawater column could be attributed to the efficient removal rates of longer chain PFAS by WWTPs, as well as the possible break-down of the carbon chain due to degradation occurring during the wastewater cleanup process<sup>58–60</sup>.

View Article Online  
DOI: 10.1039/D5VA00287G



Antidiabetic medication metformin and antiepileptics pregabalin and carbamazepine along with the latter's human consumption metabolite, 10-hydroxy-carbamazepine were simultaneously determined in stations S1 and S7, at concentration levels ranging between 0.235 and 78.8 ng L<sup>-1</sup>. These sampling points continuously receive effluent wastewater from the Thriassio and Psyttalia WWTPs, in which aforementioned pharmaceuticals have been previously reported<sup>6,42</sup>, and also confirmed by this study. Interestingly, benzodiazepines bromazepam, citalopram and the latter's metabolite, norcitalopram, along with the non-steroidal anti-inflammatory drugs (NSAIDs) ibuprofen, ketoprofen and salicylic acid, were determined in stations S1 and S7 at concentration levels ranging between 0.363 to 60.1 ng L<sup>-1</sup>, and 2.18 and 76.3 ng L<sup>-1</sup>, respectively. The occurrence of benzodiazepines in the marine environment could be likely attributed to their low rate of metabolism<sup>6,61</sup>, as well as their ineffective removal from the wastewater stream by WWTPs, while the presence of NSAIDs could be linked to different point sources, due to their effective removal ratio by WWTPs reported in this study [Table S8].

Regarding PPPs, a total of 23 substances were detected throughout the sampling area. Substances metolachlor, pentachlorophenol (PCP) and fluometuron were determined in the study area in detection frequencies equaling 100%, 62% and 54%, respectively. Their concentration levels ranged between 0.132 and 35.2 ng L<sup>-1</sup> in the sampling area. Fungicide azoxystrobin was also determined in seawater samples with 23% FoD and its highest measured concentration was determined in seawater close to the surface (2m) of station S11, equal to 1.96 ng L<sup>-1</sup>. Azoxystrobin was determined in the Psyttalia WWTP influent wastewater at 3.56 ng L<sup>-1</sup>, while in effluents, its concentration was calculated at 2.64 ng L<sup>-1</sup>. In seawater from sampling station S7 (50m) its concentration was determined at 1.00 ng L<sup>-1</sup>, while interestingly sediment samples from sampling station S8 contained azoxystrobin in concentration values equal to 0.806 µg kg<sup>-1</sup> d.w., reflecting on its continuous accumulation on the sediment layer in a short





distance from its main point source, possibly due to sea currents and circulation patterns occurring in the Saronikos Gulf. The occurrence of these substances are linked to agricultural activities and have been previously reported in other studied marine substrates<sup>62,63</sup>.

### 5.3. Occurrence and spatial distribution of ECs and PPs in sediment samples

Total results from sediment samples analysis are presented in the Supplement [Table S6]. Determined ECs and PPs per chemical class and sampling depth are visualized as a heatmap [Figure 7]. The most chemically burdened sediments in the sampling area, in terms of cumulative concentration of ECs, were in station S8, where 43 analytes were detected in a total concentration of 1360  $\mu\text{g kg}^{-1}$  d.w. and station S7, where 24 analytes were detected in a total concentration of 675  $\mu\text{g kg}^{-1}$  d.w. Sediments from station S1 contained 42 compounds in cumulative concentration of 281  $\mu\text{g kg}^{-1}$  d.w., while in station S11, a total of 26 compounds were determined at 90.8  $\mu\text{g kg}^{-1}$  d.w. This feature coincides with the consistently lower dissolved oxygen concentrations recorded near the bottom of station S8<sup>37, 38</sup>, and has been linked to the presence and dispersion of the treated wastewater plume in the inner Saronikos Gulf, southwest of Psyttalia, following the prevailing circulation pattern. In this end, the sediments serve as indicators of the long-term environmental condition, rather than capturing transient or short-term fluctuations, and reveal the link of their quality with the hydrographic characteristics of the marine environment.

The highest concentration levels of the stimulant nicotine and its metabolites anabasine and cotinine were determined at sampling station S8, with levels ranging between 12.6  $\mu\text{g kg}^{-1}$  d.w. and 578  $\mu\text{g kg}^{-1}$  d.w., indicating the extensive inhabitation of the surrounding Saronikos Gulf area. These substances were also found in sediment of station S7, but at concentrations lower by an order of magnitude compared to those in S8, suggesting that the prevailing seawater circulation during the last years may transfer Psyttalia WWTP effluents toward the seabed at station S8. This hypothesis is further supported by the presence of the four other most abundant





ECs in sampling station S8 sediments, namely gabapentin ( $680 \mu\text{g kg}^{-1} \text{ d.w.}$ ) valsartan ( $97.5 \mu\text{g kg}^{-1} \text{ d.w.}$ ), apophedrin ( $71.9 \mu\text{g kg}^{-1} \text{ d.w.}$ ) and venlafaxine ( $67.5 \mu\text{g kg}^{-1} \text{ d.w.}$ ), which are closely associated with wastewater discharges, as confirmed by this study's wastewater analysis. The relatively higher concentrations of ECs linked to WWTP discharges in the analyzed sediments underscore the long-term chemical accumulation of these substances at the seabed, contributing to its ongoing quality degradation.

Corrosion inhibitors 5-methyl-benzotriazole and mercaptobenzothiazole were both determined with a frequency of detection surpassing 50% in the sampling area, at concentration levels ranging between  $0.281$  and  $1.84 \mu\text{g kg}^{-1} \text{ d.w.}$  Both compounds were determined simultaneously in sediments from stations S1, where the industrial area of Elefsis Bay is located, and S11, where maritime traffic occurs, due to the Piraeus port operations. Plasticizers di-(2-ethylhexyl) phthalate (DEHP), diethyl phthalate (DEP) and dimethyl phthalate (DMP) were identified with a detection frequency of 100% for DEHP and 50% for both DEP and DMP. Their cumulative concentrations ranged between  $9.65 \mu\text{g kg}^{-1} \text{ d.w.}$  in sampling station S11, where maritime traffic is frequent, and  $84.8 \mu\text{g kg}^{-1} \text{ d.w.}$  in sampling station S8, close to the Psytalia WWTP. The passenger ship traffic in this area is very intense, considering that Piraeus port is one of the largest in the Mediterranean, and also that the biggest shipyard of Greece operates in Elefsis Bay. Reported plasticizer DEHP is regulated under EU legislation for its seawater presence<sup>60,61</sup>, alas limitations are yet to be imposed for its existence in sediment samples.

A total of 9 PFAS were detected in sediment samples throughout the sampling area. Legacy PFAS like PFOS and PFOA were determined at concentrations ranging between  $1.19$  and  $7.85 \mu\text{g kg}^{-1} \text{ d.w.}$  in the sampling area. These PFAS are not completely removed from the wastewater stream during cleanup, as proved by wastewater and the seawater analysis close to the WWTP pipelines, thus determined quantities in the sediment layer could be attributed to



their continuous discharge by local wastewater treatment facilities, as well as diffuse sources as has been previously reported in the literature<sup>57</sup>. Their persistent and toxic nature has led to their gradual replacement by next generation PFAS. Perfluorobutanoic acid (PFBA) was determined to have the highest concentration among PFAS in tested sediment samples, ranging between 27.8 and 28.4  $\mu\text{g kg}^{-1}$  d.w. Interestingly, perfluorohexadecanoic acid (PFHxDA), the longest carbon chain PFAS determined in this study, was solely detected in sediment samples. Its unique presence in the sediment layer could be due to its high lipophilicity (log P value equal to 10.3), leading to its sorption to suspended material and its removal from the aqueous seawater compartment [Table S5].

Antibiotics sulfadiazine and sulfisoxazole from the sulfonamide sub-class were determined at concentration levels equal to 0.378 and 0.293  $\mu\text{g kg}^{-1}$  d.w., respectively, at station S8. On the other hand, antibiotics cinoxacin and norfloxacin, from the quinolone sub-class were determined in concentration levels between 0.126 and 7.65  $\mu\text{g kg}^{-1}$  d.w. in all tested sediment samples. Most sulfonamide antibiotics have been replaced in terms of their prescription pattern by new quinolone antibiotics, while the ones that are still in use are being mainly administered as veterinary medications (see Sect. 5.4: Evidence of shift in pharmaceutical accumulation patterns). It is reported that sulfonamide sulfadiazine is currently used as an antimicrobial agent in aquacultures and not prescribed for humans anymore<sup>62</sup>. Antihypertensive drug valsartan, which is among the most prescribed blood pressure regulators, was determined in 75% of tested sediments. In sampling stations S7 and S8, this substance was determined at concentration levels equal to 69.2 and 97.5  $\mu\text{g kg}^{-1}$  d.w., respectively. These concentration levels could be attributed to the compound's high partition coefficient value (log P value of 4.40), which attests to its hydrophobic nature. Repelling effects from the aqueous seawater compartment, as well as possible precipitation mechanisms could be the key factors in valsartan's transportation and deposition on the sediment layer.



Out of the 18 PPPs and their TPs, 10 of them were determined in sediments of station S1, close to the Thriassio WWTP, which receives urban and hospital wastewater, as well as the industrial and agricultural areas of Elefsis Bay. Their cumulative concentration was measured at 23.8  $\mu\text{g kg}^{-1}$  d.w. Plant growth regulators 4-iodophenoxyacetic acid and maleic hydrazide were the most abundant PPPs in the study area, in terms of concentration levels, which were calculated between 3.51 and 6.70  $\mu\text{g kg}^{-1}$  d.w. The highest detection frequencies for PPPs were observed for the plant growth regulator 4-iodophenoxyacetic acid and the fungicide azoxystrobin, the highest determined concentration of which was 0.806  $\mu\text{g kg}^{-1}$  d.w. in sampling station S8. Azoxystrobin was determined in analyzed effluent wastewater samples in a concentration range of 2.64 and 3.91  $\mu\text{g kg}^{-1}$  d.w. [Table S8].

Interestingly, a total of 30% (n=13) of detected compounds in sampling station S8 sediments were also determined in seawater from sampling station S7. This corroborates a certain sea current direction within the Saronikos gulf, as well as circulation mechanisms and patterns of direct deposition towards the southwest part of the sampling area, where station S8 is located.

#### 5.4. Evidence of shift in pharmaceuticals use

A past study investigating pharmaceuticals<sup>28</sup> is the only one that has been carried out in the wider area of the Saronikos Gulf concerning LC-amenable compounds. This study focused exclusively on a dataset of 156 pharmaceutical compounds and their metabolites in seawater samples collected in 2013, which were analyzed via low resolution mass spectrometry. A direct comparison of the reported results provides evidence of a shift in pharmaceutical compounds sources and precipitation patterns in the study area during sampling.

Specifically, antibiotic compounds like clarithromycin were only detected during the 2013 sampling campaign. This could be possibly attributed to changes in drug prescription and consumption during the last decade, which indicates a significant reduction in clarithromycin



prescription rates around the world<sup>63,64</sup>. This is confirmed by its <LOD concentration values in the influent and effluent samples from the WWTPs of Thriasio and Psytalia and also in the seawater sampled close to the pipelines the two WWTPs, reported in this study [Table S8]. In replacement of these, new types of antibiotics have emerged, specifically quinolones, such as cinoxacin and norfloxacin, which were detected only during the 2023 campaign. Antibiotic compounds sulfamethizole and sulfamethoxazole, which were only detected during the 2013 sampling campaign, belong to the chemical class of sulfonamides. Due to their adverse side effects towards human health, they are now mainly utilized as veterinary drugs<sup>62</sup>, while more substances from the quinolone group are administered in replacement of sulfonamides. Additionally, results from Si-biol and S7-biol stations seawater analysis indicate that sulfamethoxazole concentration levels ranged between 3.34 and 4.75 ng L<sup>-1</sup>, which were significantly lower than those calculated in effluent wastewater samples from the Psytalia WWTP, equal to 725 ng L<sup>-1</sup>.

Moreover, benzodiazepine drug diazepam was identified solely in 2013. Instead, different benzodiazepines were detected during the 2023 sampling campaign, namely bromazepam and citalopram, along with more mild antidepressant substances. Influent wastewater analysis from both WWTPs operating in the area reflects on benzodiazepine consumption. Detected psychoactive substances from the benzodiazepine group include alprazolam, citalopram, midazolam and temazepam. This difference in determination can be attributed to possible changes in prescription patterns by modern psychiatry<sup>65</sup>. Benzodiazepines, however, are being sparingly administered, due to concerns about their dependance and possible adverse effects.

Furthermore, the psychoactive substance venlafaxine, along with its human metabolite O-desmethyl-venlafaxine were detected during the 2023 sampling period. Determination of both parent compound and human metabolite provides additional confidence value to



conducted research. As reported in the literature, venlafaxine is excreted from the human body as 5% unchanged and 29% metabolized to O-desmethyl-venlafaxine<sup>66</sup>. Parent compound determination is difficult in seawater possibly due to its limited concentration levels and extensive dilution occurring in the sea. On the other hand, venlafaxine determination was accomplished in sediments sampled in 2023, where years of deposition have significantly increased its concentration, rendering it detectable via the employed experimental workflow and analytical technique.

The antibiotic substance amoxicillin, which belongs to the group of  $\beta$ -lactams, was uniquely detected during the 2013 sampling period. However, posterior research has proven that due to abiotic mechanisms taking place in the aquatic ecosystem, the  $\beta$ -lactam ring breaks down due to hydrolysis<sup>67,68</sup>. As a result, MS and MS/MS data from amoxicillin determination do not produce mandatory fragment ions and consequently, this antibiotic was not considered as a finding in the present study.

Finally, analytical method constrains should also be considered, e.g., pharmaceuticals like duloxetine and flumequine, and illicit drugs like MDMA were solely determined during the 2013 sampling period at very low concentration levels, lower than the LOD values attained in the 2023 study. These particular compounds were also not determined in concentration levels surpassing LOD values in tested wastewater samples. This could be due to better sensitivity levels achieved by conducting LC-QqQ-MS analysis employed in the 2013 samples<sup>28</sup>, compared to the LC-TIMS-QTOF-MS technique implemented herein. To that end, a complementary LRMS technique is proposed for utilization in future studies regarding illicit drug determination, as their low concentration levels in seawater samples render them difficult to detect in HRMS analysis.

## 5.5. Environmental implications



During the sampling period, the study area is characterized at moderate to poor eutrophication status, based on Chl-a concentrations (Table S2). Moreover, Particulate Organic Carbon (POC) is also high, compared to other coastal areas of Greece<sup>38</sup>. The chemical quality of European coastal waters is regulated by the Water Framework Directive (WFD). Directive 2013/39/EU<sup>64</sup> amended previous legislation annexes 2000/60/EC and 2008/105/EC by adding more regulated chemicals to existing law. In 2022, new proposals were submitted to the European Parliament, under which new ECs were prioritized, and specific environmental quality standards (EQS) were set for several substances<sup>61</sup>. Recently, the Commission has adopted a new watchlist of substances in surface waters, suspected of posing a risk to the environment and human health<sup>17</sup>. The detection and determination of these substances is paramount to evaluate the environmental quality of the study area.

A total of 15 regulated substances were present in seawater samples, including 10 PFAS, 2 pharmaceuticals, 2 plant protection products, and 1 plasticizer. EU legislation suggests that determined PFAS concentrations are multiplied by a relative potency factor (RPF) and presented as  $\Sigma$ PFOA equivalents. Detected priority substances are listed in the Supplement, along with their annual average standards (AA-EQS), maximum allowable concentrations (MAC-EQS) and RPFs for PFAS [Table S9].

Besides seawater close to the sea bottom (18m) of sampling station S1 (sum of PFOA equivalents concentrations equal to 2.90 ng L<sup>-1</sup>), AA-EQS value of 4.40 ng L<sup>-1</sup> <sup>61</sup> was surpassed in every other tested seawater sample. The sum of PFOA equivalents concentration levels in the rest of the study area ranged between 4.53 and 37.3 ng L<sup>-1</sup>. These findings could be attributed to the large scale of industrial activities taking place in the Saronikos Gulf and the Elefsis Bay, as well as the extent of urbanization in the area. Antiepileptic substance carbamazepine, NSAID ibuprofen and plasticizer DEHP did not exceed AA-EQS values during sampling, the maximum concentration values of which were calculated equal to 3.77, 4.78 and



31.6 ng L<sup>-1</sup>, respectively. Insecticide clothianidin was also detected in the sampling area, in maximum concentration levels of 0.829 ng L<sup>-1</sup>, below the proposed AA-EQS levels<sup>65</sup>.

Following the calculation of ECs' concentrations in each tested sample, a prioritization of analytes was conducted by evaluating compounds' bioaccumulative potential, as well as calculating the extent of predicted no effect concentration (PNEC) exceedance in both environmental matrices. Ecotoxicological data are presented in the Supplement [Table S5]<sup>45,47,73</sup>. Analyte information includes lipophilic properties (log P values) and persistent bioaccumulative and toxic (PBT) data, which are reflected by recorded lowest PNEC values. In this study, RQFs for the worst-case scenario were calculated based on the maximum MEC, as suggested in the literature. However, considering the availability of a limited number of tested samples, general risk factors were calculated based on the median and not the average concentration values. Risk evaluations for both tested marine compartments were conducted in this study [Tables 1 & 2].

Seawater analysis suggests that a total of 20 ECs exceeded PNEC values in terms of determined concentration (RQF > 1, high risk), in at least one tested sample [Table 1]. In detail, concentration levels of linear alkylbenzene sulfonates C12-LAS and C13-LAS exceeded seawater PNEC values in 2 tested samples each, with worst-case scenario RQFs equal to 48.4 and 11.4, respectively. Concentration levels of 7 other antiseptic and disinfectant substances exceeded seawater PNEC values in at least one tested seawater sample. In detail, benzododecinium ( $n_{\text{exceedance}}=10$ ), N-methyldodecylamine ( $n_{\text{exceedance}}=3$ ), N,N-dimethyldodecylamine ( $n_{\text{exceedance}}=2$ ), benzyldimethyltetradecylammonium ( $n_{\text{exceedance}}=1$ ), BAC-14 ( $n_{\text{exceedance}}=2$ ), BAC-16 ( $n_{\text{exceedance}}=1$ ) and benzyldimethylhexadecylammonium ( $n_{\text{exceedance}}=1$ ) exhibited maximum RQFs ranging between 1.04 and 11.0 throughout the study area. Fungicide othilinone exceeded PNEC values in terms of calculated concentration in 2 seawater samples and had a worst-case scenario RQF equal to 1.62. Other plant protection





products nicosulfuron, fluometuron and dinoterb also exceeded PNEC values in a single tested seawater each, with maximum RQFs ranging between 1.01 and 30.7. Concentration levels of industrial chemicals 4-nitrophenol ( $n_{\text{exceedance}}=2$ ), bisphenol AF ( $n_{\text{exceedance}}=2$ ), 2-phenylphenol ( $n_{\text{exceedance}}=1$ ) and 1-naphthol ( $n_{\text{exceedance}}=1$ ) exhibited exceedance of their recorded PNEC values in tested seawater. The highest maximum RQF for 4-nitrophenol was measured at 23.6. Finally, veterinary drug fenbendazole ( $n_{\text{exceedance}}=2$ ), NSAID ibuprofen ( $n_{\text{exceedance}}=4$ ) and antifungal substance thiabendazole ( $n_{\text{exceedance}}=1$ ) also exceeded PNEC values in terms of measured concentrations, with their worst-case scenario RQFs ranging between 1.13 and 15.7.

In sediments, a different image is depicted through data inspection, considering the chemical classification of compounds exceeding PNEC values ( $\text{RQF} > 1$ , high risk) in terms of calculated concentration. A total of 12 analytes were determined in  $\text{RQFs} > 1$  in at least one sediment sample [Table 2]. Extremely high RQF values were calculated for PFOS, compared to other analytes. Concentration levels of this particular PFAS exceeded sediment PNEC values by at least 181 times, with a maximum RQF equal to 586. Similar but relatively lower RQFs were calculated for plasticizer DEHP, which presented a risk quotient range between 51.4 and 471. Antifungal medication oclithinone exceeded PNEC values in terms of concentration levels in all tested sediment samples, with a maximum RQF equal to 1.95, whereas pharmaceuticals 2-phenethylamine ( $n_{\text{exceedance}}=3$ ), fenbendazole ( $n_{\text{exceedance}}=2$ ) and miconazole ( $n_{\text{exceedance}}=1$ ) exhibited high-risk RQFs in tested sediments. The psychoactive substance venlafaxine was determined in RQF range between 1.53 and 2.27 in all samples where it was detected. Furthermore, nicotine, along with its human metabolite anabasine, exceeded sediment PNEC values by a maximum of 15.9 and 6.51 times, respectively. Lastly, plant protection substances methyl-pirimiphos and isoproturon were determined in concentration levels higher than their PNEC values in 100% of sediment samples, where they were detected. Their highest RQFs were equal to 9.11 and 1.75, respectively.





## 6. Conclusions

This study's main objective was to assess the presence, distribution and ecological risk of ECs and Priority Pollutants in the Saronikos Gulf and the Elefsis Bay marine environment, in relation to anthropogenic activities, utilizing a novel ion mobility HRMS analytical technique. Reported results reflect the long-term chemical burden on the study area. Due to the continuous treated wastewater discharges by local WWTPs, seawater in the area is susceptible to certain chemical stressors, associated with human-related activities, such as low metabolism percentage pharmaceuticals and plant protection products. Additionally, the prolonged deposition of ECs onto sediments has resulted in the accumulation of high-risk industrial chemicals and PFAS, as well as pharmaceuticals and coffee and tobacco related substances. Comparison of these results with previous studies conducted in the area reveals clear shifts in pharmaceutical consumption patterns by the local population.

Besides chemical identification and prioritization, this study reflects possible circulation patterns occurring in the Saronikos Gulf. Seawater samples from sampling station S7 (close to the Psytalia WWTP pipelines) were chemically encumbered mainly by pharmaceuticals, industrial chemicals and plant protection products. Sediment from sampling station S8 (southwest of the Psytalia WWTP) was determined to be the most chemically encumbered, both in number of detected ECs and in cumulative concentration of chemicals. A total of 30% (n=13) of compounds detected in S8 sediments were also determined in seawater from sampling station S7, indicating circulation mechanisms and patterns of direct deposition.

Overall, the environmental quality status of the study area was not determined to be heavily burdened by anthropogenic chemicals. Risk assessment performed in studied samples showed that only a small fraction of detected substances exceed annual average values of concentrations reported in EU legislation. Regarding PNEC values, a total of 20 compounds




presented high-risk for studied seawater, while 12 compounds presented high-risk for sediments. Among the latter, PFOS and DEHP exhibited the highest extent of PNEC exceedance in considered sediments.

This study is only the first step towards unravelling the pollution status of the Greek coastal and open sea areas. Utilization of HRMS techniques provides the ability of retrospective analysis, by performing wide-scope target screening on existing raw data, through new and enriched databases. Non-target screening protocols will also be developed and applied in the future to expand the chemical domain, which is covered by HRMS analysis of environmental samples. In the future, more seawater, sediment and wastewater samples should be analyzed to clearly recognize pollution patterns and compound degradation procedures. Marine biota samples should be analyzed as well, in the interest of determining possible biological factors of compound degradation and transformation, and also the transportation of determined chemicals through the marine food web.




View Article Online  
DOI: 10.1039/D5VA00287G



765




**Figure 1:** Sampling stations in the Saronikos Gulf and Elefsis Bay, along with detailed anthropogenic pressures in the wider study area.



**Figure 2:** Pie charts presenting the distribution of analytes per chemical class in tested matrices. Abbreviations: CTC: Coffee & tobacco related compounds, FOOD: Food additives, DRG: Illicit drugs, IND: Industrial chemicals, PFAS: Per- and polyfluorinated alkyl substances, PCP: Personal care products, PHR: Pharmaceuticals & TPs, PPP: Plant protection products & TPs, SRF: Surfactants




775

View Article Online  
DOI: 10.1039/D5VA00287G

776

777 **Figure 3:** Venn diagram of analytes determined in different tested matrices of the study area

778




779

780 **Figure 4:** Sampling station chemical encumbrance heatmap per chemical class of ECs and


781 sampling depth for the considered seawater samples

782






**Figure 5:** Concentrations of the main ECs' categories at the surface layer of the study area, starting at 2 m depth. Abbreviations: CTC: coffee and tobacco related compounds, IND: industrial chemicals, PFAS: per- and polyfluorinated alkyl substances, PCP: personal care products, PHR: pharmaceuticals & TPs, PPP: plant protection products & TPs, SRF: surfactants.



**Figure 6:** Concentrations of the main ECs' categories at the near bottom layer of the study area. Abbreviations: CTC: coffee & tobacco related compounds, IND: industrial chemicals, PFAS: per- and polyfluorinated alkyl substances, PCP: personal care products, PHR: pharmaceuticals & TPs, PPP: plant protection products & TPs, SRF: surfactants. Last depth = near bottom depth of each station.



**Figure 7:** Sampling station chemical encumbrance heatmap per chemical class of ECs for the considered sediment samples.

**Table 1:** List of determined analytes, whose concentration levels exceed recorded PNEC values in at least one tested seawater sample, along with their frequency of detection, percentage and number of seawater samples where PNEC was exceeded and RQF range.

SEAWATER SAMPLES					
Analyte name	%FoD	PNEC exceedance	Minimum RQF	Median RQF	Maximum RQF
C12-LAS	15%	15% (n=2)	0	0	48.4
Nicosulfuron	8%	8% (n=1)	0	0	30.7
4-nitrophenol	15%	15% (n=2)	0	0	23.6
Fenbendazole	15%	15% (n=2)	0	0	15.7
C13-LAS	15%	15% (n=2)	0	0	11.4
Benzododecinium	100%	77% (n=10)	0.330	1.17	11.0
N-methyldodecylamine	92%	23% (n=3)	0	0.548	4.35
Ibuprofen	46%	31% (n=4)	0	0	4.34
N,N-dimethyldodecylamine	15%	15% (n=2)	0	0	3.09
Bisphenol AF	15%	15% (n=2)	0	0	2.98
2-phenylphenol	15%	8% (n=1)	0	0	2.42
Benzyldimethyltetradecylammonium	100%	8% (n=1)	0.0589	0.346	2.00
BAC 14	15%	15% (n=2)	0	0	1.96
Octhilinone	15%	15% (n=2)	0	0	1.62
1-naphthol	15%	8% (n=1)	0	0	1.25
Thiabendazole	23%	8% (n=1)	0	0	1.13
Fluometuron	54%	8% (n=1)	0	0.0126	1.10
Benzyldimethylhexadecylammonium	15%	8% (n=1)	0	0	1.08
BAC 16	8%	8% (n=1)	0	0	1.04
Dinoterb	92%	8% (n=1)	0	0.0548	1.01





**Table 2:** List of determined analytes, whose concentration levels exceed recorded PNEC values in at least one tested sediment sample, along with their frequency of detection, percentage and number of sediment samples where PNEC was exceeded and RQF range.

SEDIMENT SAMPLES					
Analyte Name	%FoD	PNEC exceedance	Minimum RQF	Median RQF	Maximum RQF
Perfluorooctanesulfonic acid (PFOS)	100%	100% (n=4)	181	231	586
Di-(2-ethylhexyl) phthalate (DEHP)	100%	100% (n=4)	51.4	267	471
Octhilinone	100%	100% (n=4)	1.07	1.57	1.95
2-phenethylamine	100%	75% (n=3)	0.595	1.42	3.51
Venlafaxine	75%	75% (n=3)	1.53	2.27	2.27
Methyl-pirimiphos	50%	50% (n=2)	8.65	8.88	9.11
Miconazole	75%	50% (n=2)	0.549	1.20	4.39
Nicotine	50%	50% (n=2)	1.39	8.65	15.9
Anabasine	50%	25% (n=1)	0.499	3.50	6.51
Fenbendazole	50%	25% (n=1)	0.211	0.901	1.59
Isoproturon	25%	25% (n=1)	1.75	1.75	1.75
Phorate-sulfone	25%	25% (n=1)	7.89	7.89	7.89



**Authors contribution**

**Loukovois R.** (<https://orcid.org/0009-0005-1852-0412>): Data curation, Formal analysis, Investigation, Methodology, Validation, Writing - original draft, Writing - review & editing.

**Parinos C.** (<https://orcid.org/0000-0002-7700-374X>): Conceptualization, Data curation, Funding Acquisition, Investigation, Methodology, Supervision, Writing - original draft, Writing - review & editing.

**Gkotsis G.** (<https://orcid.org/0000-0002-1744-4228>): Data curation, Investigation, Writing - review & editing.

**Nika M.-C.** (<https://orcid.org/0000-0003-0050-8923>): Data curation, Investigation, Writing - review & editing.

**Hatzianestis I.** (<https://orcid.org/0000-0002-5206-9689>): Conceptualization, Funding Acquisition, Methodology, Resources, Supervision, Writing - review & editing.

**Pavlidou A.** (<https://orcid.org/0000-0002-0031-4726>): Conceptualization, Funding Acquisition, Methodology, Resources, Supervision, Writing - original draft, Writing - review & editing.

**Thomaidis N.** (<https://orcid.org/0000-0002-4624-4735>): Conceptualization, Funding Acquisition, Resources, Supervision, Writing - review & editing.

**Acknowledgements**

This work was financed by the Hellenic Ministry of Environment, Energy and Climate Change - Special Secretariat of Water, under the Monitoring Program for the assessment of ecological quality of rivers, coastal and transitional waters of Greece (Article 8; WFD; 2000/60/EC) co-financed by Greece and the EU, as well as the EU - Horizon Europe under the RHE-MEDiation Lighthouse project (HORIZON-MISS-2022-OCEAN-01-03-101113045).



832 We sincerely thank the officers and crew of R/V Aegaeo (HCMR, Greece) for their assistance  
833 during the sampling campaign, PhD candidate Nikolaos Boinis (<https://orcid.org/0009-0008-6909-6270>) for the development of the data mining tool collecting ECs' physicochemical  
834 properties and ecotoxicological data from online databases and Dr. Reza Aalizadeh  
835 (<https://orcid.org/0000-0001-9167-9060>) for the development of the utilized suspect list and  
836 sharing valuable insights regarding suspect screening through HRMS.  
837  
838

#### 839 **Data availability**

840 All data considered in this study will be available upon request.  
841

#### 842 **Conflict of interest**

843 The authors declare that they have no conflict of interest.  
844

#### 845 **References**

- 846 1. Nilsen E, Smalling KL, Ahrens L, Gros M, Miglioranza KSB, Picó Y, et al. Critical  
847 review: Grand challenges in assessing the adverse effects of contaminants of emerging  
848 concern on aquatic food webs. *Environ Toxicol Chem*. 2018 Oct 8;38(1):46–60.
- 849 2. Das R, Raj D. Sources, distribution, and impacts of emerging contaminants – a critical  
850 review on contamination of landfill leachate. *J Hazard Mater Adv*. 2025 Feb;17:100602.
- 851 3. K'oreje KO, Okoth M, Van Langenhove H, Demeestere K. Occurrence and treatment of  
852 contaminants of emerging concern in the African aquatic environment: Literature  
853 review and a look ahead. *J Environ Manage*. 2020 Jan;254:109752.



- 854 4. Buckley TJ, Egeghy PP, Isaacs K, Richard AM, Ring C, Sayre RR, et al. Cutting-edge  
855 computational chemical exposure research at the U.S. Environmental Protection  
856 Agency. *Environ Int.* 2023 Aug;178:108097.
- 857 5. Rout PR, Zhang TC, Bhunia P, Surampalli RY. Treatment technologies for emerging  
858 contaminants in wastewater treatment plants: A review. *Sci Total Environ.* 2021  
859 Jan;753:141990.
- 860 6. Osooha JO, Anyanwu BO, Ejileugha C. Pharmaceuticals and personal care products as  
861 emerging contaminants: Need for combined treatment strategy. *J Hazard Mater Adv.*  
862 2023 Feb;9:100206.
- 863 7. Wilkinson J, Hooda PS, Barker J, Barton S, Swinden J. Occurrence, fate and  
864 transformation of emerging contaminants in water: An overarching review of the field.  
865 *Environ Pollut.* 2017 Dec;231:954–70.
- 866 8. Kajtazi A, Kajtazi M, Santos Barbeta MF, Bandini E, Eghbali H, Lynen F. Prediction  
867 of Retention Indices in LC-HRMS for Enhanced Structural Identification of Organic  
868 Micropollutants in Water: Selectivity-Based Filtration. *Anal Chem.* 2025 Jan  
869 14;97(1):65–74.
- 870 9. Da Silva AC, Alves De Oliveira LV, Amaral Alexandre L, Rocha Ribas M, Lemos Dal  
871 Pizzol J, Rocha G, et al. Suspect screening and quantitative analysis of 165  
872 contaminants of emerging concern in water, sediments, and biota using LC-MS/MS:  
873 Ecotoxicological and human health risk assessment. *Sci Total Environ.* 2025  
874 Feb;963:178434.



- 875 10. Peng L, Fu D, Qi H, Lan CQ, Yu H, Ge C. Micro- and nano-plastics in marine  
876 environment: Source, distribution and threats — A review. *Sci Total Environ.* 2020  
877 Jan;698:134254. View Article Online  
DOI: 10.1039/D5VA00287G
- 878 11. Gago-Ferrero P, Díaz-Cruz MS, Barceló D. UV filters bioaccumulation in fish from  
879 Iberian river basins. *Sci Total Environ.* 2015 June;518–519:518–25.
- 880 12. Thomaidi VS, Stasinakis AS, Borova VL, Thomaidis NS. Is there a risk for the aquatic  
881 environment due to the existence of emerging organic contaminants in treated domestic  
882 wastewater? Greece as a case-study. *J Hazard Mater.* 2015 Feb;283:740–7.
- 883 13. Pavlidou A, Simboura N, Rousselaki E, Tsapakis M, Pagou K, Drakopoulou P, et al.  
884 Methods of eutrophication assessment in the context of the water framework directive:  
885 Examples from the Eastern Mediterranean coastal areas. *Cont Shelf Res.* 2015  
886 Oct;108:156–68.
- 887 14. E.U. EUR-Lex 2015/495 [Internet]. 2015/495 Mar 20, 2015. Available from: [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32015D0495&qid=1742945848237)  
888 [content/EN/TXT/?uri=CELEX%3A32015D0495&qid=1742945848237](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32015D0495&qid=1742945848237)
- 889 15. E.U. EUR-Lex 2018/840 [Internet]. 2018/840 June 5, 2018. Available from: [https://eur-](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32018D0840&qid=1742945871473)  
890 [lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32018D0840&qid=1742945871473)  
891 [content/EN/TXT/?uri=CELEX%3A32018D0840&qid=1742945871473](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32018D0840&qid=1742945871473)
- 892 16. Cortes G, Cuenca N. Factsheets of the substances for the 4th Watch List under the  
893 Water Framework Directive.  
894



- 895 17. European Commission. Joint Research Centre. Selection of substances for the 5th  
896 Watch List under the Water Framework Directive. [Internet]. LU: Publications Office;  
897 2025 [cited 2025 May 20]. Available from: <https://data.europa.eu/doi/10.2760/956398>
- 898 18. Álvarez-Muñoz D, Rodríguez-Mozaz S, Maulvault AL, Tediosi A, Fernández-Tejedor  
899 M, Van Den Heuvel F, et al. Occurrence of pharmaceuticals and endocrine disrupting  
900 compounds in macroalgae, bivalves, and fish from coastal areas in Europe. *Environ*  
901 *Res.* 2015 Nov;143:56–64.
- 902 19. Vitale D, Picó Y, Spanò N, Torreblanca A, Del Ramo J. Carbamazepine exposure in the  
903 sea anemones *Anemonia sulcata* and *Actinia equina*: Metabolite identification and  
904 physiological responses. *Sci Total Environ.* 2020 Nov;744:140891.
- 905 20. Cervený D, Fick J, Klaminder J, McCallum ES, Bertram MG, Castillo NA, et al. Water  
906 temperature affects the biotransformation and accumulation of a psychoactive  
907 pharmaceutical and its metabolite in aquatic organisms. *Environ Int.* 2021  
908 Oct;155:106705.
- 909 21. Castaño-Ortiz JM, Gil-Solsona R, Ospina-Alvarez N, García-Pimentel MM, León VM,  
910 Santos LHMLM, et al. Bioaccumulation and fate of pharmaceuticals in a Mediterranean  
911 coastal lagoon: Temporal variation and impact of a flash flood event. *Environ Res.* 2023  
912 July;228:115887.
- 913 22. Choi Y, Kim K, Kim D, Moon H bang, Jeon J. Ny-Ålesund-oriented organic pollutants  
914 in sewage effluent and receiving seawater in the Arctic region of Kongsfjorden. *Environ*  
915 *Pollut.* 2020 Mar;258:113792.



- 916 23. Alygizakis N, Giannakopoulos T, Thomaidis NS, Slobodnik J. Detecting the sources of  
917 chemicals in the Black Sea using non-target screening and deep learning convolutional  
918 neural networks. *Sci Total Environ.* 2022 Nov;847:157554.
- 919 24. Gutiérrez-Martín D, Gil-Solsona R, Saaltink MW, Rodellas V, López-Serna R, Folch A,  
920 et al. Chemicals of emerging concern in coastal aquifers: Assessment along the land-  
921 ocean interface. *J Hazard Mater.* 2023 Apr;448:130876.
- 922 25. Kang D, Ahn YY, Moon HB, Kim K, Jeon J. Exploring micropollutants in polar  
923 environments based on non-target analysis using LC-HRMS. *Mar Pollut Bull.* 2024  
924 Dec;209:117083.
- 925 26. Soriano Y, Doñate E, Asins S, Andreu V, Picó Y. Fingerprinting of emerging  
926 contaminants in L'Albufera natural park (Valencia, Spain): Implications for wetland  
927 ecosystem health. *Chemosphere.* 2024 Sept;364:143199.
- 928 27. Nödler K, Voutsas D, Licha T. Polar organic micropollutants in the coastal environment  
929 of different marine systems. *Mar Pollut Bull.* 2014 Aug;85(1):50–9.
- 930 28. Alygizakis NA, Gago-Ferrero P, Borova VL, Pavlidou A, Hatzianestis I, Thomaidis NS.  
931 Occurrence and spatial distribution of 158 pharmaceuticals, drugs of abuse and related  
932 metabolites in offshore seawater. *Sci Total Environ.* 2016 Jan;541:1097–105.
- 933 29. Lougkovois R, Gkotsis G, Parinos C, Hatzianestis I, Nika MC, Pavlidou A, et al. Storm  
934 Daniel Extreme Flood Event in Thessaly, Greece: Assessing the Pollution Status of the  
935 Impacted Coastal Marine Areas through Extended Screening of Emerging Contaminants  
936 Using LC-TIMS-HRMS. *Environ Sci Technol Lett.* 2025 Mar 4;acs.estlett.5c00122.





- 937 30. Nannou C, Efthymiou C, Boti V, Albanis T. Unveiling the pharmaceutical footprint in  
938 freshwater and seawater: HRMS method optimization, validation, and uncertainty  
939 estimation. *Microchem J.* 2025 Mar;210:112933.
- 940 31. Acir IH, Guenther K. Endocrine-disrupting metabolites of alkylphenol ethoxylates – A  
941 critical review of analytical methods, environmental occurrences, toxicity, and  
942 regulation. *Sci Total Environ.* 2018 Sept;635:1530–46.
- 943 32. Ansari AA, Gill SS, editors. *Eutrophication: Causes, Consequences and Control: Volume 2* [Internet]. Dordrecht: Springer Netherlands; 2014 [cited 2025 Mar 26].  
944 Available from: <https://link.springer.com/10.1007/978-94-007-7814-6>
- 946 33. Parinos C, Hatzianestis I, Chourdaki S, Plakidi E, Gogou A. Imprint and short-term fate  
947 of the Agia Zoni II tanker oil spill on the marine ecosystem of Saronikos Gulf. *Sci Total*  
948 *Environ.* 2019 Nov;693:133568.
- 949 34. Parinos C, Hatzianestis I, Chourdaki S, Plakidi E, Gogou A. Dataset on the imprint of  
950 the Agia Zoni II tanker oil spill on the marine ecosystem of Saronikos Gulf. *Data Brief.*  
951 2019 Dec;27:104664.
- 952 35. Pavlidou A, Simboura N, Pagou K, Assimakopoulou G, Gerakaris V, Hatzianestis I, et  
953 al. Using a holistic ecosystem-integrated approach to assess the environmental status of  
954 Saronikos Gulf, Eastern Mediterranean. *Ecol Indic.* 2019 Jan;96:336–50.
- 955 36. Rousselaki E, Michalopoulos P, Pavlidou A, Kaberi H, Prifti E, Dassenakis M. Pore-  
956 water nutrient concentrations variability under different oxygen regimes: A case study  
957 in Elefsis Bay, Greece. *Sci Total Environ.* 2024 Mar;915:169830.



- 958 37. Kontoyiannis H. Observations on the circulation of the Saronikos Gulf: A  
959 Mediterranean embayment sea border of Athens, Greece. *J Geophys Res Oceans*. 2010  
960 June;115(C6):2008JC005026. View Article Online  
DOI: 10.1039/D5VA00287G
- 961 38. Simboura N, Panayotidis P, Papathanassiou E. A synthesis of the biological quality  
962 elements for the implementation of the European Water Framework Directive in the  
963 Mediterranean ecoregion: The case of Saronikos Gulf. *Ecol Indic*. 2005 Aug;5(3):253–  
964 66.
- 965 39. Borova VL, Maragou NC, Gago-Ferrero P, Pistos C, Thomaidis NS. Highly sensitive  
966 determination of 68 psychoactive pharmaceuticals, illicit drugs, and related human  
967 metabolites in wastewater by liquid chromatography–tandem mass spectrometry. *Anal*  
968 *Bioanal Chem*. 2014 July 1;406(17):4273–85.
- 969 40. Gago-Ferrero P, Borova V, Dasenaki ME, Thomaidis NS. Simultaneous determination  
970 of 148 pharmaceuticals and illicit drugs in sewage sludge based on ultrasound-assisted  
971 extraction and liquid chromatography–tandem mass spectrometry. *Anal Bioanal Chem*.  
972 2015 June 1;407(15):4287–97.
- 973 41. Nikolopoulou V, Alygizakis NA, Nika MC, Oswaldova M, Oswald P, Kostakis M, et al.  
974 Screening of legacy and emerging substances in surface water, sediment, biota and  
975 groundwater samples collected in the Siverskyi Donets River Basin employing wide-  
976 scope target and suspect screening. *Sci Total Environ*. 2022 Jan;805:150253.
- 977 42. Gago-Ferrero P, Bletsou AA, Damalas DE, Aalizadeh R, Alygizakis NA, Singer HP, et  
978 al. Wide-scope target screening of >2000 emerging contaminants in wastewater samples  
979 with UPLC-Q-ToF-HRMS/MS and smart evaluation of its performance through the



980 validation of 195 selected representative analytes. *J Hazard Mater.* 2020

981 Apr;387:121712.

982 43. Thomaidis N, Baessmann C, Damalas DE, Diamanti K, Galvin B, Georgakaki M, et al.

983 National and Kapodistrian University of Athens LC-TIMS-HRMS Target List

984 [Internet]. 2024. Available from: <https://zenodo.org/records/13862246>

985 44. Aalizadeh R, Nikolopoulou V, Thomaidis NS. Development of Liquid Chromatographic

986 Retention Index Based on Cocamide Diethanolamine Homologous Series (C( *n* )-DEA).

987 *Anal Chem.* 2022 Nov 22;94(46):15987–96.

988 45. Freeling F, Alygizakis NA, Von Der Ohe PC, Slobodnik J, Oswald P, Aalizadeh R, et

989 al. Occurrence and potential environmental risk of surfactants and their transformation

990 products discharged by wastewater treatment plants. *Sci Total Environ.* 2019

991 Sept;681:475–87.

992 46. Aalizadeh R, Alygizakis NA, Schymanski EL, Krauss M, Schulze T, Ibáñez M, et al.

993 Development and Application of Liquid Chromatographic Retention Time Indices in

994 HRMS-Based Suspect and Nontarget Screening. *Anal Chem.* 2021 Aug

995 24;93(33):11601–11.

996 47. NORMAN network ecotoxicology database [Internet]. [cited 2025 Mar 30]. Available

997 from: <https://www.norman->

998 [network.com/nds/ecotox/lowestPnecsIndex.php?checkSelect=0](https://www.norman-network.com/nds/ecotox/lowestPnecsIndex.php?checkSelect=0)

999 48. Li J, O'Brien JW, Tschärke BJ, Verhagen R, He C, Shimko KM, et al. Occurrence,

1000 Removal, and Risk Assessment of Antimicrobials and Their Transformation Products in

1001 Effluent from Australian Wastewater Treatment Plants. *Environ Sci Technol.* 2025 Apr

1002 8;59(13):6825–38.



- 1003 49. Rauseo J, Spataro F, Pescatore T, Patrolecco L. Multiresidue determination and  
1004 predicted risk assessment of emerging contaminants in sediments from Kongsfjorden,  
1005 Svalbard. *Sci Total Environ.* 2024 Apr;922:171156. View Article Online  
DOI: 10.1039/D5VA00287G
- 1006 50. Pavlidou A, Simboura N, Assimakopoulou G, Gerakaris V, Hatzianestis I, Panayotidis  
1007 P, et al. Using a holistic ecosystem-integrated approach to assess the environmental  
1008 status of Saronikos Gulf, Eastern Mediterranean. 2019;
- 1009 51. López-Pacheco IY, Silva-Núñez A, Salinas-Salazar C, Arévalo-Gallegos A, Lizarazo-  
1010 Holguin LA, Barceló D, et al. Anthropogenic contaminants of high concern: Existence  
1011 in water resources and their adverse effects. *Sci Total Environ.* 2019 Nov;690:1068–88.
- 1012 52. González-Alonso S, Merino LM, Esteban S, López De Alda M, Barceló D, Durán JJ, et  
1013 al. Occurrence of pharmaceutical, recreational and psychotropic drug residues in surface  
1014 water on the northern Antarctic Peninsula region. *Environ Pollut.* 2017 Oct;229:241–54.
- 1015 53. Pisetta AM, Roveri V, Guimarães LL, De Oliveira TMN, Correia AT. First report on the  
1016 occurrence of pharmaceuticals and cocaine in the coastal waters of Santa Catarina,  
1017 Brazil, and its related ecological risk assessment. *Environ Sci Pollut Res.* 2022  
1018 Sept;29(42):63099–111.
- 1019 54. Liu L, Aljathelah NM, Hassan H, Giraldez BW, Leitão A, Bayen S. Targeted and  
1020 suspect screening of contaminants in coastal water and sediment samples in Qatar. *Sci*  
1021 *Total Environ.* 2021 June;774:145043.
- 1022 55. Struk-Sokołowska J, Gwoździej-Mazur J, Jurczyk Ł, Jadwiszczak P, Kotowska U,  
1023 Piekutin J, et al. Environmental risk assessment of low molecule benzotriazoles in urban  
1024 road rainwaters in Poland. *Sci Total Environ.* 2022 Sept;839:156246.



Open Access Article. Published on 05 December 2025. Downloaded on 12/25/2025 12:51:00 AM.  
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



56. Topaz T, Gridish N, Sade T, Zedaka H, Suari Y, Konomi A, et al. Exploring Per- and Polyfluoroalkyl Substances (PFAS) in Microestuaries: Occurrence, Distribution, and Risks. *Environ Sci Technol Lett.* 2024 May 14;11(5):466–71.

57. Gago-Ferrero P, Gros M, Ahrens L, Wiberg K. Impact of on-site, small and large scale wastewater treatment facilities on levels and fate of pharmaceuticals, personal care products, artificial sweeteners, pesticides, and perfluoroalkyl substances in recipient waters. *Sci Total Environ.* 2017 Dec;601–602:1289–97.

58. Shapiro EF, Lin ZW, Cifuentes ES, Barajas-Rodriguez FJ, Gwinn R, Dichtel WR, et al. Removal of PFAS and pharmaceuticals from municipal wastewater using a novel  $\beta$ -cyclodextrin adsorbent over distinct contact times. *Water Res.* 2025 Aug;282:123631.

59. Sanzana S, Fenti A, Iovino P, Panico A. “A review of PFAS remediation: Separation and degradation technologies for water and wastewater treatment.” *J Water Process Eng.* 2025 May;74:107793.

60. Cai P, Chen T, Zhan M, Ma X, Takaoka M, Sun C, et al. Theoretical and experimental insights into the degradation mechanism of PFBS under subcritical hydrothermal conditions. *J Hazard Mater.* 2025 July;491:137935.

61. Greenblatt D, Shader R, Divoll M, Harmatz J. Benzodiazepines: a summary of pharmacokinetic properties. *Br J Clin Pharmacol* [Internet]. 1981 Feb [cited 2025 Apr 8];11(S1). Available from: <https://bpspubs.onlinelibrary.wiley.com/doi/10.1111/j.1365-2125.1981.tb01833.x>

62. Rousis N, Denardou M, Alygizakis N, Galani A, Bletsou A, Damalas D, et al. Assessment of Environmental Pollution and Human Exposure to Pesticides by

Environmental Science: Advances Accepted Manuscript

- 1047 Wastewater Analysis in a Seven-Year Study in Athens, Greece. *Toxics*. 2021 Oct  
1048 11;9(10):260. View Article Online  
DOI: 10.1039/D5VA00287G
- 1049 63. Lee S, Choi Y, Kang D, Jeon J. Proposal for priority emerging pollutants in the  
1050 Nakdong river, Korea: Application of EU watch list mechanisms. *Environ Pollut*. 2024  
1051 Jan;341:122838.
- 1052 64. Directive 2013/39/EU of the European Parliament and of the Council of 12 August 2013  
1053 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the  
1054 field of water policyText with EEA relevance.
- 1055 65. EUR-Proposal for a DIRECTIVE OF THE EUROPEAN PARLIAMENT AND OF  
1056 THE COUNCIL amending Directive 2000/60/EC establishing a framework for  
1057 Community action in the field of water policy, Directive 2006/118/EC on the protection  
1058 of groundwater against pollution and deterioration and Directive 2008/105/EC on  
1059 environmental quality standards in the field of water policy [Internet]. Oct 26, 2022.  
1060 Available from: [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:52022PC0540)  
1061 [content/EN/TXT/HTML/?uri=CELEX:52022PC0540](https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:52022PC0540)
- 1062 66. Shahi MK, Odoi JO, Anuntawirun S, Roongrojmongkhon N, Singchat W, Prasanpan J,  
1063 et al. Antimicrobial resistance in global catfish aquaculture: Trends, control strategies,  
1064 and public health implications.
- 1065 67. ClinCalc.com. Clarithromycin - Total Prescriptions and Patients Per Year (2013 - 2022)  
1066 [Internet]. [cited 2025 Mar 26]. Available from:  
1067 <https://clincalc.com/DrugStats/Drugs/Clarithromycin>





68. Martínez-González NA, Di Gangi S, Pichierri G, Neuner-Jehle S, Senn O, Plate A. Time Trends and Factors Associated with Antibiotic Prescribing in Swiss Primary Care (2008 to 2020). *Antibiotics*. 2020 Nov 23;9(11):837.

69. Antidepressant use in Europe continues to break records [Internet]. [cited 2025 Mar 30]. Available from: [https://www.europeandatajournalism.eu/cp\\_data\\_news/Antidepressant-use-in-Europe-continues-to-break-records/?utm\\_source=chatgpt.com](https://www.europeandatajournalism.eu/cp_data_news/Antidepressant-use-in-Europe-continues-to-break-records/?utm_source=chatgpt.com)

70. DRUGBANK data on venlafaxine metabolism [Internet]. [cited 2025 Mar 30]. Available from: <https://go.drugbank.com/drugs/DB00285>

71. Fabregat-Safont D, Gracia-Marín E, Ibáñez M, Pitarch E, Hernández F. Analytical key issues and challenges in the LC-MS/MS determination of antibiotics in wastewater. *Anal Chim Acta*. 2023 Jan;1239:340739.

72. Felis E, Kalka J, Sochacki A, Kowalska K, Bajkacz S, Harnisz M, et al. Antimicrobial pharmaceuticals in the aquatic environment - occurrence and environmental implications. *Eur J Pharmacol*. 2020 Jan;866:172813.

73. PubChem database [Internet]. [cited 2025 Mar 30]. Available from: <https://pubchem.ncbi.nlm.nih.gov/>

Open Access Article. Published on 05 December 2025. Downloaded on 12/25/2025 12:51:00 AM.  
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Environmental Science: Advances Accepted Manuscript

**Occurrence, distribution and risk assessment of emerging contaminants  
and priority substances in the Saronikos Gulf and Elefsis Bay, Greece,  
utilizing liquid chromatography high-resolution mass spectrometry**

[View Article Online](#)

DOI: 10.1039/D5VA00287G

Rallis Lougkovois<sup>1,2</sup>, Constantine Parinos<sup>1,\*</sup>, Georgios Gkotsis<sup>2</sup>, Maria-Christina Nika<sup>2</sup>,  
Ioannis Hatzianestis<sup>1</sup>, Alexandra Pavlidou<sup>1</sup>, Nikolaos S. Thomaidis<sup>2,\*</sup>

<sup>1</sup> Hellenic Centre for Marine Research, Institute of Oceanography, 46.7 Km Athens-Sounio  
av., Mavro Lithari, 19013, Anavyssos, Attiki, Greece

<sup>2</sup> National and Kapodistrian University of Athens, Department of Chemistry, Laboratory of  
Analytical Chemistry, University Campus, Zografou, 15771, Athens, Greece

**Corresponding authors:**

\* Constantine Parinos, email: [ksparinos@hcmr.gr](mailto:ksparinos@hcmr.gr), tel: +30 2291076372

\* Nikolaos Thomaidis, email: [ntho@chem.uoa.gr](mailto:ntho@chem.uoa.gr), tel: +30 2107274317



**DATA AVAILABILITY STATEMENT**

[View Article Online](#)  
DOI: 10.1039/D5VA00287G

All data considered in this study will be available upon request.

