


# ChemComm

## Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

## COMMUNICATION

## Activation of gaseous PH<sub>3</sub> with low coordinate diaryltetrylene compounds

Cite this: DOI: 10.1039/x0xx00000x

Jonathan W. Dube,<sup>a</sup> Zachary D. Brown,<sup>b</sup> Christine A. Caputo,<sup>b</sup> Philip P. Power,<sup>b</sup> and Paul J. Ragnogna<sup>a</sup>Received 00th January 2012,  
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x


www.rsc.org/

**The reaction of phosphine gas with a low coordinate diarylgermylene or diarylstannylene results in both oxidative addition and arene elimination at the group 14 atom. The products were characterised by <sup>31</sup>P NMR spectroscopy and X-ray crystallography, and represent the first P – H bond activation by a heavy group 14 element compound.**

The activation of small molecule substrates by coordinatively unsaturated main group molecules is a continuously expanding area of research.<sup>1</sup> Of particular intrigue is the fact that these commodity, and typically unreactive, chemicals can be activated directly without the use of transition metals. The activation of dihydrogen by a main group compound under ambient conditions was first achieved by using a digermene (**A**; Figure 1).<sup>2</sup> Subsequent studies have shown that frustrated Lewis pairs (FLPs), ie. **B**),<sup>3</sup> stable singlet carbenes (eg. **C**),<sup>4</sup> diaryltetrylenes (**D**),<sup>5</sup> heavier p-block alkyne analogues,<sup>2,6</sup> silylenes,<sup>7</sup> and group 13 dimetallenes (**E**)<sup>8</sup> can also accomplish the same feat. Several of these unique systems also activate ammonia under ambient conditions, a feature that is not as common for transition metals.<sup>9</sup>

Bertrand *et al.* have shown that unique, stable *N*-heterocyclic carbenes (ie. **C**) can also insert into B – H, Si – H, and P – H bonds.<sup>9</sup> While primary and secondary phosphines were used in work reported previously, reactions involving phosphine (PH<sub>3</sub>) are rare.<sup>11</sup> Driess *et al.* have recently shown that a Si(II) center (**F**)<sup>12</sup> inserts into the P – H bond of phosphine to produce a ligand stabilized Si(H)(PH<sub>2</sub>) fragment.<sup>13</sup>

The primary mode of reactivity for low valent main group centers is to undergo oxidative addition across the X – H (X = H, B, N, Si, P) bond. For the *N*-heterocyclic carbenes and heavier analogues there are few examples to the contrary. The diaryltetrylenes (**D**, EAr<sub>2</sub>; E = Ge, Sn; Ar = C<sub>6</sub>H<sub>3</sub>-2,6-Mes<sub>2</sub>)<sup>13</sup> however, have displayed different reactivity based on the group 14 element. In the reaction with ammonia, germanium exclusively favours the oxidative addition pathway, while tin exclusively favours arene elimination to remain in the +2 oxidation state (Scheme 1).<sup>5</sup> In this context, we report an extension of the reactivity studies on diaryltetrylenes to PH<sub>3</sub>, ammonia's heavier congener. In contrast to ammonia, both oxidative addition and arene elimination products are observed for germanium and tin, and also in different ratios relative to each other.




**Figure 1.** Examples of main group compounds that activate small molecules. Note that Dipp = 2,6-diisopropylphenyl and that Mes = 2,4,6-trimethylphenyl.


The reaction of the diarylgermylene (GeAr<sub>2</sub>; Ar = C<sub>6</sub>H<sub>3</sub>-2,6-Mes<sub>2</sub>) with an excess of PH<sub>3</sub> (80 psi) in a stainless steel pressure reactor for three hours results in the complete

consumption of the purple starting material to give a colourless solution. Analysis of a fraction of the concentrated reaction mixture by  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy revealed the presence of two signals ( $\delta_{\text{P}} = -232$ ;  $-180$ ), which integrate to an approximate 80:20 ratio. The  $^{31}\text{P}$  NMR spectrum was particularly informative revealing the signals to be proton coupled as a triplet of doublets and a triplet ( $\delta_{\text{P}} = -232$ ,  $^1J_{\text{P-H}} = 181$  Hz,  $^2J_{\text{P-H}} = 11.2$  Hz;  $\delta_{\text{P}} = -180$ ,  $^1J_{\text{P-H}} = 183$  Hz), consistent with the formation of both the oxidative addition product,  $\text{Ar}_2\text{E}(\text{H})(\text{PH}_2)$  (**1**), and the arene elimination product,  $\{\text{ArE}(\mu\text{-PH}_2)\}_2$  (**2**) (Scheme 2). The  $^1\text{H}$  NMR spectrum of a recrystallized sample of **1** revealed a doublet of triplets ( $\delta_{\text{H}} = 5.33$ ,  $^2J_{\text{P-H}} = 11.2$  Hz,  $^3J_{\text{H-H}} = 4.3$  Hz), consistent with a germanium hydride adjacent to a  $\text{PH}_2$  functionality. The phosphorus hydrides were also observed as a doublet of doublets in the  $^1\text{H}$  NMR spectrum ( $\delta_{\text{H}} = 1.29$ ,  $^1J_{\text{P-H}} = 181$  Hz,  $^3J_{\text{H-H}} = 4.3$  Hz) and agree nicely to the other coupling constants. The FT-IR spectrum also reveals signals consistent with a  $\text{P-H}$  ( $\nu = 2310$   $\text{cm}^{-1}$ ) and a  $\text{Ge-H}$  stretch ( $\nu = 2069$   $\text{cm}^{-1}$ ). Confirmation of the solid-state structure of **1** was obtained from an X-ray diffraction study on single crystals grown from a saturated hexane solution at  $-35^\circ\text{C}$ . There is a significant presence of  $\text{GeH}_2\text{Ar}_2$  in the crude reaction mixture (40-50%), which cannot be efficiently separated from **1**.<sup>15</sup> While the origin of this species is not clear, its identity was unambiguously confirmed by  $^1\text{H}$  NMR and FT-IR spectroscopies as well as by single crystal X-ray diffraction. The analogous reaction of the diarylstannylene ( $\text{SnAr}_2$ ) with  $\text{PH}_3$  (80 psi) requires longer reaction times (24 hours) to go to completion, as evidenced by the disappearance of the characteristic purple colour. The proton decoupled and coupled  $^{31}\text{P}$  NMR spectra again revealed two signals ( $\delta_{\text{P}} = -249$ ,  $-227$ ), in this case in an approximate 68:32 ratio. The phosphorus – proton coupling constants are comparable to those of the analogous germanium species ( $\delta_{\text{P}} = -249$ ,  $^1J_{\text{P-H}} = 174$  Hz,  $^2J_{\text{P-H}} = 18.6$  Hz;  $\delta_{\text{P}} = -227$ ,  $^1J_{\text{P-H}} = 174$  Hz). Phosphorus – tin coupling is also observed with the values being consistent with a tin – phosphorus covalent bond ( $\delta_{\text{P}} = -249$ ,  $^1J_{\text{Sn-P}} = 496$  Hz,  $^1J_{\text{Sn-P}} = 518$  Hz;  $\delta_{\text{P}} = -227$ ,  $^1J_{\text{Sn-P}} = 597$  Hz,  $^1J_{\text{Sn-P}} = 621$  Hz). The  $^1\text{H}$  NMR spectrum reveals the characteristic doublet of triplets for the  $\text{Sn-H}$  ( $\delta_{\text{H}} = 6.15$ ,  $^2J_{\text{P-H}} = 18.6$  Hz,  $^3J_{\text{H-H}} = 3.6$  Hz) while the FT-IR spectrum displayed resonances consistent the  $\text{P-H}$  and  $\text{Sn-H}$  bond vibrations. The solid-state structure of the oxidative addition product,  $\text{Ar}_2\text{Sn}(\text{H})(\text{PH}_2)$ , **3**, was obtained as structural confirmation, while unfortunately structural verification of the arene elimination product,  $\{\text{ArSn}(\mu\text{-PH}_2)\}_2$ , **4**, has proven elusive despite considerable effort. This is likely a result of the fact that completely separating **3** and **4** was not possible under the conditions employed.


The solid-state structures of the oxidative addition products **1**, and **3** are shown in Figure 3. The structures are nearly identical, and also similar to the ammonia insertion product, with the tetrel center in a distorted tetrahedral geometry. The  $\text{Ge-H}$  and  $\text{Sn-H}$  distances are 1.58(4) and 1.90(5) Å. The  $\text{Ge-C}$  bond lengths are 1.981(3) and 2.003(3) Å, while the  $\text{Ge-P}$  bond length is 2.3194(11) Å. The analogous bond lengths for **3**



**Scheme 2.** The reaction of phosphine with the diaryltetraylenes (E = Ge, Sn) to give both the oxidative addition (**1** or **3**) and arene elimination (**2** or **4**) products. Ratios determined by  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy of the crude reaction mixture.



**Figure 2.** Plots of  $^{31}\text{P}\{^1\text{H}\}$  and  $^{31}\text{P}$  NMR spectra for the reaction of  $\text{EAr}_2$  (E = Ge, Sn) with phosphine. Inset: FT-IR spectrum of **1** focusing on hydride vibrations.



**Figure 3.** The solid-state structure of **1** (left) and **3** (right). Ellipsoids are drawn at 50% probability, hydrogen atoms, with exception to the main group hydrides, and hexane solvate (**1**) are removed for clarity. Selected bond lengths (Å) and angles ( $^\circ$ ): **1** Ge – H 1.58(4), Ge – P 2.3194(11), P – H 1.34(4), P – H 1.02(4), C – Ge – C 112.5(1), P – Ge – H 102.1(13); **3** Sn – H 1.90(5), Ge – P 2.5997(17), P – H 1.391(10), P – H 1.384(10), C – Sn – C 138.1(2), P – Sn – H 96.1(14).

are 2.193(6), 2.181(6), and 2.5997(17) Å, respectively. For both **1** and **3** the phosphorus – hydrogen bond lengths fall within the range of 1.32 and 1.42 Å with the exception of one very short outlier (1.02(4) Å in **1**).<sup>16</sup> The C – E – C bond angle has increased considerably from the free diaryltetraylene<sup>14</sup> for **3** ( $138.1(2)^\circ$  *cf.*  $114.7(2)^\circ$  for  $\text{SnAr}_2$ ) while the same angle in **1** is slightly smaller than in  $\text{GeAr}_2$  ( $112.5(1)^\circ$  *cf.*  $114.4(2)^\circ$  for  $\text{GeAr}_2$ ). The P – E – H bond angles are much smaller in comparison, consistent with the use of unhybridized p-orbitals in the bonding, and also an obvious result of the steric pressure of the terphenyl substituents.

In conclusion, a diarylgermylene and a diarylstannylene react with phosphine gas under ambient conditions to produce both oxidative addition and arene elimination products, the structures of the oxidative addition products being confirmed crystallographically. This study represents a rare example of activation of a P – H bond at a low coordinate main group center, and the first of the heavier group 14 elements (Ge, Sn).

We thank the Natural Science and Engineering Research Counsel (NSERC), the Ontario government, *The University of Western Ontario*, and the U.S. Department of Energy (DE-FG02-07ER46475), for their financial support. The phosphine setup was aided greatly by the support of Cytec industries and the excellent work of the chemistry department electronics shop (J. Vanstone and J. Aukema).

## Notes and references

<sup>a</sup> Department of Chemistry and Center for Advanced Materials and Biomaterials Research (CAMBR), *The University of Western Ontario*, 1151 Richmond St., London, Ontario, Canada, N6A 5B7.

<sup>b</sup> Department of Chemistry, *The University of California*, 1 Shields Ave, Davis, California, USA, 95616.

Electronic Supplementary Information (ESI) available: Selected NMR spectra and ORTEP diagrams. See DOI: 10.1039/c000000x/

†All manipulations were performed in a nitrogen filled MBraun glovebox with solvents stored over 4Å molecular sieves or a potassium mirror. Phosphine (Cytec) was loaded into a stainless steel Parr reactor and excess gas was swept with N<sub>2</sub> and controllably burnt after the desired reaction time. The product distribution was determined from integration of the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of the isolated crude powder in C<sub>6</sub>D<sub>6</sub>.

**Experimental Data:** **1:** <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>, δ): 1.29 (dd, 2H, <sup>1</sup>J<sub>P-H</sub> = 181 Hz, <sup>3</sup>J<sub>H-H</sub> = 4.3 Hz), 1.79 (s, 12H), 1.84 (s, 12H), 2.19 (s, 12H), 5.33 (dt, 1H, <sup>2</sup>J<sub>P-H</sub> = 11.2 Hz, <sup>3</sup>J<sub>H-H</sub> = 4.3 Hz), 6.71 (d, 2H, <sup>3</sup>J<sub>H-H</sub> = 6.4 Hz), 6.75 (s, 4H), 6.83 (s, 4H), 7.02 (t, 1H, <sup>3</sup>J<sub>H-H</sub> = 7.6 Hz); <sup>31</sup>P{<sup>1</sup>H} NMR (161.8 MHz, C<sub>6</sub>D<sub>6</sub>, δ): -232; <sup>31</sup>P NMR (161.8 MHz, C<sub>6</sub>D<sub>6</sub>, δ): -232 (td, <sup>1</sup>J<sub>P-H</sub> = 181 Hz, <sup>2</sup>J<sub>P-H</sub> = 11.2 Hz); FT-IR (cm<sup>-1</sup>): 2069 (Ge-H), 2310 (P-H); **3:** <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>, δ): resonances that could assigned to **3** are listed: 0.84 (dd, 2H, <sup>1</sup>J<sub>P-H</sub> = 174 Hz, <sup>3</sup>J<sub>H-H</sub> = 3.6 Hz, <sup>2</sup>J<sup>117</sup><sub>Sn-H</sub> = 61.8 Hz, <sup>2</sup>J<sup>119</sup><sub>Sn-H</sub> = 68.7 Hz), 6.15 (dt, 1H, <sup>2</sup>J<sub>P-H</sub> = 18.6 Hz, <sup>3</sup>J<sub>H-H</sub> = 3.6 Hz); <sup>31</sup>P{<sup>1</sup>H} NMR (161.8 MHz, C<sub>6</sub>D<sub>6</sub>, δ): -249 (<sup>1</sup>J<sup>117</sup><sub>Sn-P</sub> = 496 Hz, <sup>1</sup>J<sup>119</sup><sub>Sn-P</sub> = 518 Hz); <sup>31</sup>P NMR (161.8 MHz, C<sub>6</sub>D<sub>6</sub>, δ): -249 (td, <sup>1</sup>J<sub>P-H</sub> = 174 Hz, <sup>2</sup>J<sub>P-H</sub> = 18.6 Hz); HRMS: found (calculated) for C<sub>48</sub>H<sub>53</sub>NaPSn 803.2687 (803.2800).

**Crystallographic Details:** **1:** C<sub>48</sub>H<sub>53</sub>GeP, C<sub>6</sub>H<sub>14</sub>; FW = 819.64 g/mol; colourless block; Size: 0.052 x 0.061 x 0.130 mm; Monoclinic, *P2<sub>1</sub>/c*; Unit Cell: a = 14.813(3) Å, b = 19.868(4) Å, c = 16.899(3) Å, β = 111.25(3)°, V = 4635.1(16) Å<sup>3</sup>; Z = 4; ρ = 1.175; T = 150°K; λ = 0.71073 nm (MoKα); F(000) = 1752; μ = 0.729; 2θ<sub>max</sub> = 55.12; 2θ<sub>min</sub> = 3.30; 35818 reflections, 10616 unique (R<sub>int</sub> = 0.0449); parameters = 538; restraints = 5; R<sub>1</sub> = 0.0583; wR<sup>2</sup> = 0.1449; R<sub>1</sub>(all data) = 0.0751; wR<sup>2</sup>(all data) = 0.1563; GooF = 1.017; percent complete: 99.3%; **3:** C<sub>48</sub>H<sub>53</sub>GeP; FW = 779.56 g/mol; colourless block; Size: 0.040 x 0.055 x 0.131 mm; Triclinic, *P2<sub>1</sub>/c*; Unit Cell: a = 10.3576(11) Å, b = 22.461(2) Å, c = 17.5165(19) Å, β = 99.453(3)°, V = 4019.8(7) Å<sup>3</sup>; Z = 4; ρ = 1.288; T = 150°K; λ = 0.71073 nm (MoKα); F(000) = 1624; μ = 0.707; 2θ<sub>max</sub> = 55.0; 2θ<sub>min</sub> = 2.98; 45865 reflections, 9226 unique (R<sub>int</sub> = 0.1763); parameters = 474; restraints = 4; R<sub>1</sub> = 0.0661; wR<sup>2</sup> = 0.1475; R<sub>1</sub>(all data) = 0.1692; wR<sup>2</sup>(all data) = 0.1888; GooF = 1.007; percent complete: 99.9%.

- 1 P. P. Power, *Nature*, 2010, **463**, 171-177.
- 2 G. H. Spikes, J. C. Fettinger, P. P. Power, *J. Am. Chem. Soc.*, 2005, **127**, 12232-12233;
- 3 a) G. C. Welch, R. R. San Juan, J. Masuda, D. W. Stephan, *Science*, 2006, **314**, 1124-1126; b) D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.*, 2010, **49**, 46-76.
- 4 a) G. D. Frey, V. Lavallo, B. Donnadieu, W. W. Schoeller, G. Bertrand, *Science*, 2007, **316**, 439-441; b) D. Martin, M. Soleihavoup, G. Bertrand, *Chem. Sci.*, 2011, **2**, 389-399.
- 5 a) Y. Peng, B. D. Ellis, X. Wang, P. P. Power, *J. Am. Chem. Soc.*, 2008, **130**, 12268-12269; b) Y. Peng, J.-D. Guo, B. D. Ellis, Z. Zhu, J. C. Fettinger, P. P. Power, *J. Am. Chem. Soc.*, 2009, **131**, 16272-16282; c) P. P. Power, *Chem. Rec.*, 2012, **12**, 238-255; d) P. P. Power, *Acc. Chem. Res.*, 2011, **44**, 627-637.
- 6 a) Y. Peng, M. Byrnda, B. D. Ellis, J. C. Fettinger, E. Rivard, P. P. Power, *Chem. Commun.*, 2008, 6042-6044; b) P. P. Power, *Organometallics*, 2007, **26**, 4362-4372; c) J. Li, C. Schenk, C. Goedecke, G. Frenking, C. Jones, *J. Am. Chem. Soc.*, 2011, **133**, 18622-18625; d) R. C. Fischer, P. P. Power, *Chem. Rev.*, 2010, **110**, 3877-3923.
- 7 A. V. Protchenko, K. H. Birjkumar, D. Dange, A. D. Schwarz, D. Vidovic, C. Jones, N. Kaltsayannis, P. Mountford, S. Aldridge, *J. Am. Chem. Soc.*, 2012, **134**, 6500-6503.
- 8 Z. Zhu, X. Wang, Y. Peng, H. Lei, J. C. Fettinger, E. Rivard, P. P. Power, *Angew. Chem. Int. Ed.* **2009**, **48**, 2031.
- 9 J. Zhao, A. S. Goldman, J. F. Hartwig, *Science*, 2006, **307**, 1080-1082.
- 10 a) G. D. Frey, J. D. Masuda, B. Donnadieu, G. Bertrand, *Angew. Chem. Int. Ed.*, 2010, **49**, 9444-9447; While formally not an NHC, a ethynyl dithiocarbamate has also been shown to react with related substrates b) G. Ung, G. D. Frey, W. W. Schoeller, G. Bertrand, *Angew. Chem. Int. Ed.*, 2011, **50**, 9923-9925.
- 11 E. Fluck, *Top. Curr. Chem.* 1971, **35**, 1-64. For a review on transition metal P-H bond activation see: L. Coudray, J.-L. Montchamp, *Eur. J. Org. Chem.* 2008, 3601-3613.
- 12 a) M. Driess, S. Yao, M. Brym, C. van Wüllen, D. Lentz, *J. Am. Chem. Soc.*, 2006, **128**, 9628-9629; b) S. Yao, Y. Xiong, M. Driess, *Organometallics*, 2011, **30**, 1748-1767.
- 13 C. Prasang, M. Stoelzel, S. Inoue, A. Meltzer, M. Driess, *Angew. Chem. Int. Ed.*, 2010, **49**, 10002-11005.
- 14 R. S. Simons, L. Pu, M. M. Olmstead, P. P. Power, *Organometallics*, 1997, **16**, 1920-1925.
- 15 Both products are hexanes soluble and crystallize from hexanes solution at -35°C. Therefore, the products cannot be consistently isolated free of each other in usable quantities.
- 16 There is disorder about the germanium and phosphorus atoms, with the major component refining to 91% occupancy. While the hydride refines to a logical location by bond angle, the bond distance is much shorter than one would expect, potentially a result of this disorder.