


Faraday Discussions

Accepted Manuscript



This manuscript will be presented and discussed at a forthcoming Faraday Discussion meeting. All delegates can contribute to the discussion which will be included in the final volume.

Register now to attend! Full details of all upcoming meetings: <http://rsc.li/fd-upcoming-meetings>



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Free-electron laser induced processes in thin molecular ice

Björn Siemer^a, Sebastian Roling^a, Robert Frigge^a, Tim Hoger^a, Rolf Mitzner^{a,b}, Helmut Zacharias^{a,*}

5 DOI: 10.1039/b000000x [DO NOT ALTER/DELETE THIS TEXT]

Intermolecular reactions in and on icy films on silicate and carbonaceous grains constitute a major route for the formation of new molecular constituents in interstellar molecular clouds. In more diffuse regions and in protoplanetary discs energetic radiation can trigger reaction routes far from thermal equilibrium. As analog of interstellar ice-covered dust grains, highly-oriented pyrolytic graphite (HOPG) covered with D₂O, NO, and H atoms is irradiated by ultrashort XUV pulses and desorbing ionic and neutral products are analysed. Yields of several products show a nonlinear intensity dependence and thus allow to elucidate reaction dynamics by two-pulse correlated desorption.

1 Introduction

Molecular ice layers on grain particles are active environments for the trapping, formation, and release of new and even complex molecular species into the interstellar medium.^{1,2} In dense molecular clouds at low temperatures ($T_s \sim 10$ to 50 K) physisorbed atoms and molecules may undergo exothermic reactions to form new species. In particular the formation of water and of molecular hydrogen is believed to take place under such conditions.^{3,4,5,6,7,8} Inner cores of large molecular clouds in the interstellar medium (ISM) are largely opaque to radiation in the visible, ultraviolet (UV), and vacuum ultraviolet (VUV) spectral range, and only partially transparent for IR radiation.⁹ Although only a low level of VUV and extreme ultraviolet (XUV) radiation is present¹⁰ due to the interaction of cosmic rays with the constituents of the clouds, its impact on the desorption of simple and complex molecules is relevant.

30 At the outer rims of such clouds and particular in the photodissociation regions, the temperatures rise and the densities are considerably lower.¹ In these regions more energetic photons can penetrate into the clouds and reach the surface of ice-covered grains with temperatures up to about 100 K. Such conditions also prevail in the outer regions of approximately solar-mass protoplanetary discs at distances from about 3 to 50 AU from the central object.^{11,12} Typical grains consist of silicate and carbonaceous cores, and show sizes in the range from a few nm for polycyclic aromatic hydrocarbons (PAHs) to 1000 nm for amorphous silicates.^{13,14,15} With graphite and graphene-like surfaces the (photo)chemistry on some of these grains can be simulated in laboratory experiments.




Fig. 1 Illustration of reactive processes on icy grains

Figure 1 shows schematically the typical processes that might occur on such grains when they interact with UV/VUV radiation. These basic processes have been described in various review articles.^{1,2,4,8,12} Besides direct photodesorption of molecules adsorbed on the icy surface, more involved processes can be triggered. Photodissociation of film constituents generates energetic fragments which may directly leave or induce, via collisions, other molecules to leave the grain particle. Furthermore, reactions between dissociation products and constituents of the ice layer or even of the grain core may occur. The high kinetic energy of the primary dissociation products and the solid state density of the film enable reaction cascades very different from equilibrium gas phase chemistry.

The interaction of VUV radiation with condensed molecular films has been studied in recent years extensively by various groups. Early studies involved broad band VUV radiation from hydrogen and helium discharge lamps,^{16,17} while recently tunable radiation from frequency doubled lasers¹⁸ and synchrotron light sources¹⁹ has been employed. Besides simple monomolecular ices like H₂O, CO and N₂,²⁰ ice films of mixed constituents address issues of radical reactions in these films.²¹ A prototypical radical is nitric oxide which has been observed in the gas phase of clouds²² and which, according to simulations, builds up to significant concentrations in ices.²³ It also is the main precursor for nitrogen and oxygen chemical networks.^{24,25} Such reactions become more important after the incorporation of more complex molecules, like benzene,¹⁸ small polycyclic aromatic hydrocarbons²⁶, and C₆₀ fullerenes²⁷ as dopants into water ice.

Important aspects of VUV and XUV induced processes in condensed molecular films are further addressed by low-energy electron beam irradiation of these systems. Such electrons are also liberated in XUV irradiation of the samples. The primary ions, either generated directly by the photons or by secondary photoelectrons, may initiate very effective ion-molecule reactions such as protonation of photoproducts and original constituents. Electron stimulated desorption of various hydrogen and oxygen containing positive ions²⁸ and aromatic molecules²⁹ from water ice has been reported, as well as the interaction with more complex ices.³⁰




Fig. 2a Schematic diagram of the optical layout of the experimental set-up. The spectrum shows in dark colour a single shot spectrum and in white the spectral envelop over 1000 pulses.

In this paper we report on preliminary studies of the interaction of XUV photons with graphite surfaces covered by hydrogen atoms, nitric oxide, or water ice. We make use of the ultrashort duration of XUV pulses from a free-electron laser, on the order of 30 fs, to shed light onto reaction pathways triggered by this radiation. Although the photon energy of these pulses is certainly higher than that from external irradiation sources of molecular clouds due to the atomic hydrogen ionization limit, the basic dynamic features induced by this radiation in the ice films are expected to be similar to the action of VUV photons. Moreover, internal radiation created by cosmic ray collisions produce XUV radiation, even in dense clouds.

2 Experimental

The experiments were performed at beamlines 1 and 3 of the free-electron laser at Hamburg (FLASH), which provides pulsed radiation in the XUV spectral range with fundamental photon energies from about $h\nu = 25$ to 300 eV.³¹ In the present experiments, FLASH operated at photon energies of $h\nu = 38.15$ eV ($\lambda = 32.5$ nm) and 57.1 eV ($\lambda = 21.7$ nm) in a 5 Hz single bunch mode. The spectral energy width averaged to about $h\Delta\nu \sim 0.8$ eV, with pulse energies up to about (13.5 ± 3.5) μ J. The average pulse duration was measured to ca. $t_L \sim 30$ fs.³²

For time-resolved experiments, a wave front dividing beam splitter and pulse split delay unit (SDU) is employed.^{33,34} This SDU provides two jitter-free replica pulses for XUV pump/ XUV probe experiments, see Fig. 2a. The incoming FLASH beam is split into two parts by the first mirror which is partially moved into the beam path. Each partial beam propagates on a separated beam path (orange and green) and both are recombined again into the original direction with the final mirrors of their respective beam paths. Following recombination, the soft x-ray beams are focused by a toroidal mirror to ellipsoidal spot sizes of $220 \mu\text{m} \times 175 \mu\text{m}$ and $235 \mu\text{m} \times 125 \mu\text{m}$ for the variable and fixed beam paths, respectively. The spatial overlap of the two foci is monitored by an x-ray CCD camera in the equivalent focal position via a reflecting quartz plate which is brought into the beam path. In this way the focal diameter has been measured and the stability of the spatial overlap ensured over the whole delay range. The set-up further facilitated a measurement of the different intensities in the two focal spots. The \hat{p} -polarized XUV pulses strike the highly oriented pyrolytic graphite (HOPG) surface at an angle of incidence of $\theta = 67.5^\circ$ relative to the surface normal. At these photon energies of 38.15 eV and 57.1 eV and




Fig. 2b Experimental set-up for the detection of desorption products

incident angle about 42 % and 32 % of the incident radiation is reflected from graphite and about 12 % and 5 % from thick ice films, respectively.^{35,36}

Figure 2b shows schematically the set-up for the detection of desorbing species. The HOPG sample exhibited a mosaic angle of 0.4° and a diameter of 10 mm. It is mounted on a liquid nitrogen containing reservoir attached to a holder on an X-Y-Z stage in an ultra-high vacuum (UHV) chamber. The UHV chamber has been evacuated by an oil-free pumping system to a base pressure of $< 7 \cdot 10^{-10}$ mbar. The HOPG sample was prepared by cleaving with adhesive tape and long-time annealing at 800 K by electron bombardment from the rear side. After sample preparation the temperature of the graphite crystal is held at $T_s = 95$ K, as monitored with a K-type thermocouple attached to the surface.

Molecules are dosed onto the HOPG surface through a tube at normal incidence at a constant chamber pressure for several minutes or via atomic and molecular beams. Hydrogen and deuterium atoms are supplied by a heated tungsten capillary, using a design which avoids ionic species. Hydrogen only sticks as chemisorbed atomic hydrogen after surmounting a considerable adsorption barrier of $E_b \sim 200$ meV,^{37,38} an energy which can be achieved when heating the tube to temperatures above 2000°C. NO molecules are dosed by a doubly differentially pumped pulsed thermal molecular beam. Nitric oxide condenses as dimers in a self-terminating monolayer. At this surface temperature water condenses as microporous amorphous ice. For water ice, coverages of 80 to 200 ML have been prepared and investigated. At XUV photon energies of 38 eV and 57 eV the penetration depth of the radiation into ice amounts to about 9 nm and 15.7 nm, respectively.^{35,36} Therefore, at an incident angle of $\theta = 67.5^\circ$ the first 11 and 18 ML, respectively, are irradiated using a value of $1.15 \cdot 10^{15}$ molecule cm^{-2} for a monolayer (ML) of water on graphite.

Cations desorbing directly after irradiation by the XUV pulse are measured by a Wiley-McLaren type time-of-flight mass spectrometer (ToF) mounted along the surface normal direction. Then the sample holder served as the first repelling electrode. Desorbing neutral molecules are first ionized by tuneable laser radiation (see Fig. 2b) and then detected by the same ToF spectrometer. The output of the micro-channel plates is directly monitored on a digital oscilloscope, gated and forwarded to




Fig. 3 Time-of-flight mass spectrum of species desorbed from ice/graphite by XUV FEL pulses at $h\nu = 38.15$ eV.

a computer. For neutral hydrogen atoms a (2+1) REMPI scheme via the two-photon
 5 excitation $2s \ ^2S_{1/2} \leftarrow 1s \ ^2S_{1/2}$ at $\lambda \sim 243$ nm is chosen, for NO a (1+1) REMPI
 scheme via the $A \ ^2\Sigma^+ \leftarrow X \ ^2\Pi$ γ -band transitions around $\lambda \sim 226$ nm. This UV probe
 laser radiation is provided by a frequency doubled, tuneable dye laser with nanosec-
 ond pulse duration, which is electronically synchronized to the FEL pulses.

For each delay time setting between the desorbing XUV and the detecting tuneable
 10 dye laser pulses the time-of-flight mass spectra are summed over 1000 single shots.
 For each shot the signal and the relative FLASH pulse energy as given by the gas
 monitor detector is stored. Normalization and averaging were performed in the data
 analysis.

3 Results

15

Water ice

XUV photon induced reactions in undoped D₂O water ice multilayers with thick-
 nesses up to about 200 ML are observed. At the photon energies employed, 38.15 eV
 20 and 57.1 eV, the exciting radiation is absorbed in the top most ice layers and does
 not reach the substrate. At respective penetrations depth of 9.0 and 15.7 nm into the
 ice and an angle of incidence of $\theta = 67.5^\circ$ only the top most 11 and 18 ML of the
 ice, respectively, are irradiated. After illuminating the surface with soft x-ray radi-
 ation at $h\nu = 38.15$ eV ($\lambda = 32.1$ nm) with pulse energies up to (13.5 ± 3.5) μ J the
 25 directly desorbing cations are detected with a time-of-flight mass spectrometer.
 When reducing the acceleration voltage in the spectrometer an asymmetric line
 shape for the detected ions arises and thus the kinetic energy distribution of these
 ions can be derived.

A typical mass spectrum is shown in Fig. 3. Besides the expected masses of D₂O⁺,
 30 OD⁺, O⁺, and D⁺ resulting from fragmentation and ionization, products of reactions




Fig 4 Dependence of the O_2^+ ion signal on the pulse energy of the desorbing FEL pulses at $h\nu = 57.1$ eV. The other oxygen containing ions observed (Fig. 5) show similar intensity dependencies.

5 within the ice layers are also observed, in particular D_3O^+ , D_2O_2^+ , and O_2^+ . In addition, water cluster ions $(\text{D}_2\text{O})_n^+$ and deuterated clusters $(\text{D}_2\text{O})_n\text{D}^+$ which are formed by dissociative deuteration are observed at higher masses.³⁹ The yield of these reactively formed products exhibits a nonlinear dependence on the FEL intensity which is exemplarily shown in Fig. 4 for O_2^+ ions. The signals can be fitted by a
 10 power law, $Y \sim I^n$, where $n = 3$, even for unfocussed FEL pulses. In an oversimplified picture, such a power dependence then suggests that three XUV photons are involved in the formation of the product species. Strong pulse-to-pulse fluctuations of the FEL output power causes a broad scatter of single shot data, which leaves some variations in the exponent n .

15 Such a nonlinear intensity dependence allows then to investigate the temporal development of the signal via an XUV pump – XUV probe set-up and thus partially elucidate the reaction dynamics. For the present experiment the SDU is employed with an asymmetric intensity splitting of the FEL pulse. In the overlapping foci of both beams this resulted in an averaged intensity of $(6.5 \pm 1.2) \cdot 10^{10}$ W/cm² and
 20 $(1.1 \pm 0.1) \cdot 10^{11}$ W/cm² for the variable and fixed beam, respectively. Thus, an asymmetric yield is expected as a function of temporal delay between both pulses. For each delay, Δt , the spatial overlap of both beams was controlled by measuring the beam profile by reflecting the beam off a quartz onto a CCD camera positioned at the equivalent focal position plate before entering the experimental chamber, see
 25 Fig. 2a.

Figure 5 shows the yield of various product ions as a function of delay times $\Delta t = -1$ and $+5$ ps between both XUV pulses. The data points are averages of several runs. Besides the expected D_2O^+ ions, several fragment ions - D^+ , O^+ , and OD^+ - and reaction products - D_2O_2^+ , O_2^+ , and D_3O^+ (see Fig. 6) - are also observed. The tem-
 30 poral structure of the two-pulse correlation curves are very similar for all ions shown. Fitting the longer main signal by a Gaussian curve yields a maximum at $\Delta t = 2.2$ ps and a width of the signal of about 2.7 ps (FWHM). In addition, for the oxygen








Fig. 5 Two-pulse correlated XUV desorption of various cation fragments from D₂O ice on graphite. The photon energy of the desorbing laser was $h\nu = 38.15$ eV.

containing products, a second maximum is observed for $\Delta t = 0.6$ ps. Despite the fact
 5 that this relative maximum is supported by just one data point, it was observed in
 every run. The very limited beam time at the FEL prohibited a finer temporal grid. It
 is worth noting that the signal of nearly all ions including the D⁺ ion approach zero
 for longer temporal delays. The D₂O⁺ ion signal, however, shows a large background
 signal of more than 50% of the peak signal which does not change with delay. A
 10 large part of this signal is thus generated by a single photon process, as expected.

From the observation that the maximum of the signals do not occur when both puls-
 es overlap in time it may be concluded that not only pure photonic processes are
 involved in forming these cations. Instead, collisions between fragments within the
 ice layer appear to be important. The first step for the generation of these products
 15 certainly involves the photodissociation of a water molecule:



20




Fig. 6 Two-pulse correlated XUV desorption of D_3O^+ from ice/graphite. The photon energy of the desorbing laser was $h\nu = 38.15$ eV.









where E_{kin} and E_{exc} denote the kinetic energy of the liberated electron and the excess energy available for distribution over internal and translational degrees of freedom, respectively.

Since all products detected in Figs. 5 and 6, except D_2O^+ , show a yield close to zero at large pulse separations in the two-pulse correlation, these products are not generated in a single photon event. The hydroxyl radicals produced in the beam path of the FEL beam penetrating the ice layer may then collide with each other and form hydrogen peroxide:






where also substrate phonons can act as the third collision partner to carry away the excess energy of the reaction.

The temporal behaviour of the two-pulse correlation signals suggests that hydrogen peroxide serves as a reservoir molecule for the other products. A second photon then either ionizes hydrogen peroxide, produces the other fragments, or strips off both hydrogen atoms:



In the gas phase the three product channels show similar reaction probabilities in the

VUV.⁴³ This second step may be followed by the absorption of a third photon which then yields the corresponding atomic and molecular ions, if they are not already produced in the previous step.



The nonlinear yield dependence, which is equal or larger than $n = 3$, suggests that this last step indeed occurs.

In this reaction system process (2), the formation of hydrogen peroxide, is certainly the step which governs the temporal evolution of the signal. Since the excess energy release in processes (1a) through (1a'') is high, with a maximum kinetic energy in the OD radicals of up to $E_{\text{kin}} = 3.3 \text{ eV}$, neighbouring hydroxyl radicals can even react within the temporal overlap of both FEL pulses at $\Delta t = 0 \text{ ps}$. This part then contributes to the small signal at time zero. However, it is evident that it requires some more time to accumulate an optimal concentration of hydrogen peroxide. The two-pulse correlation results (Fig. 5) suggest that this occurs at about $\Delta t = 2.2 \text{ ps}$ for hydroxyl radicals presumably produced in the bulk ice. Surface hydroxyl may react on a faster time scale as is suggested by the peak observed at $\Delta t = 0.6 \text{ ps}$. This mechanism requires at least two XUV photons from the FEL beam, in accordance with the recorded power dependencies of $n \geq 3$. Hydrogen peroxide is thus an essential intermediate precursor for the formation of oxygen containing ions. This conclusion is supported by earlier experiments where ice layers have been irradiated by electron beams with kinetic energies in the same region as the photon energies employed here.^{49,50}

Other oxygen containing cations observed, as shown in Fig. 5, may have additional channels which contribute to the signal height. In particular, D_2O^+ shows a large background of about one half of the peak intensity, respectively, when both pulses are separated in time. This can easily be understood by the direct and dissociative ionization of the ice constituents. It is interesting to note that for D^+ , the fast peak at $\Delta t = 0.6 \text{ ps}$ is missing. A direct D^+ ion abstraction seems not to take place. Moreover, since we observe a significant two-pulse correlation signal, which can only be observed for a nonlinear intensity dependence, a multiphoton process must also be invoked for this product channel.

The hydronium cation, D_3O^+ , which is also observed to show a pronounced temporal dependence in two-pulse correlated desorption (see Fig. 6), is easily formed by deuteration reactions. For this product, a fast correlation peak after ca. $\Delta t = 0.6 \text{ ps}$ is also observed, followed by a major peak at about $\Delta t \sim 2.2 \text{ ps}$. This is again interpreted as a distinction between surface and bulk reactions. The relative integrated yield of the surface contributions is about 0.09.

In this case either a process via the D^+ ion



or a reaction like



is conceivable. The D^+ ions observed (Fig. 5) display only the broad peak at




Fig. 7 (a) Velocity distribution of H^+ after XUV desorption from ice/graphite. Two Maxwellian velocity distribution with $T = 7000\text{ K}$ and 19000 K approximate most of the distribution. Some deviation at low velocities can be noticed. (b) Kinetic energy distribution of H^+ from ice/graphite. The red curve is a fit to a single sided exponential yielding a kinetic temperature of 16245 K . Photon energy: 38.15 eV .

$\Delta t = 2.2\text{ ps}$. The D_3O^+ ion products (Fig. 6) exhibit both peaks in the signal at $\Delta t = 0.6\text{ ps}$ and at 2.2 ps . At longer pulse delays the D_3O^+ signal approaches zero. This last observation argues against reaction (5b), because the generation of the water cation requires only a single XUV photon. This is an efficient process also for widely separated pulses, as is witnessed by the large background in the temporal dependence of the D_2O^+ signal (Fig. 5). It remains to explain the non-appearance of the fast peak at $\Delta t = 0.6\text{ ps}$ in the D^+ ion signal (Fig. 5). It is conceivable that these early D^+ ions react with other products or neutral film constituents, see, e.g., reaction (5a), before they were able to leave the surface. Certainly, additional experiments are required and in particular the kinetic energy release into the different channels to be measured before a definite conclusion can be drawn.

It is important to note that photoinduced processes within the bulk ice lead not only to the formation of new species, in this case hydrogen peroxide, but to the release of these species and dissociation products from the grain surface into the gas phase. Collisions occurring between photodissociation products and ice constituents are central to the formation of new species. Furthermore, even without reactions, such energetic collisions will lead to the desorption of neutral species from the ice layer, as has been observed for vibrational resonant IR-induced desorption from molecular ices, like N_2O ,^{51,52} CD_3F ,⁵³ and methane.⁵⁴

Kinetic energies of directly desorbing ions

As is evident from Fig. 5, the XUV pulses also induce the desorption of hydrogen ions from ice layers. In this case, the temporal behaviour of the two-pulse correlated desorption signal suggests that D^+ ions liberated in the bulk predominantly contribute to the signal observed. However, H^+ and D^+ are also observed to directly desorb from the ice layers. These hydrogen ions show a high kinetic energy as shown in Fig. 7, where both the velocity and kinetic energy distributions of H^+ are plotted. The velocity distribution in Fig. 7a can be fitted by two Maxwellian distributions with respective temperatures of $T = 7000\text{ K}$ and 19000 K and relative integrated yields of 45% and 55%. From this analysis, an average kinetic energy of $E_{\text{kin}} = 1.8\text{ eV}$ for the H^+ ions is obtained. When the measured distribution is transformed into the energy picture (Fig. 7b) a single-sided exponential distribution, $W(E_{\text{kin}})$, is ob-

10 | [journal], [year], [vol], 00–00

This journal is © The Royal Society of Chemistry [year]

tained

$$W(E_{\text{kin}}) \sim a 1/\sqrt{T} \exp\{-E_{\text{kin}} / kT\}. \quad (6)$$

Such distributions are often found in plasma physics to determine kinetic temperatures. In this case a best fit yields a kinetic temperature of $T_{\text{kin}} = 16245$ K, corresponding to an energy of about $E_{\text{kin}} = 2.1$ eV. For O^+ and O_2^+ ions, high kinetic energies of 0.84 eV and 0.39 eV, respectively, have also been observed.⁵⁵

Neutral hydrogen

10 Different from other groups which investigate the reaction dynamics of physisorbed hydrogen on ice,⁴ graphite,⁵⁶ and silicate surfaces^{7,57} we concentrate here on chemisorbed hydrogen atoms on a graphite substrate. Such conditions are reached in warmer parts of the ISM where molecular gases have already been desorbed from the grain particles. Such grain temperatures prevail, e.g., at the outer rims of molecular clouds⁴ and of protoplanetary discs.¹²

The high photon energy of the FEL radiation enables the direct desorption of ionized and neutral hydrogen species. For this experiment on HOPG (0001) we kept, after cleaning, the surface temperature at room temperature to prohibit the adsorption of water from the UHV background gas. Hydrogen atoms are dosed onto the surface from a heated tungsten capillary. In this way the adsorption barrier of ca. 200 meV can be overcome. In this experiment, a hydrogen coverage of about 0.25 ML is adsorbed on the surface.

Neutral H atoms are desorbed by $h\nu = 38.8$ eV radiation using p-polarized FEL pulses with an average pulse energy of 13 μJ , incident under an angle of $\theta = 67.5^\circ$ relative to the surface normal. Slight focusing resulted in an illuminated area of about 0.19 mm^2 , an applied fluence of 6.9 mJ/cm^2 and a fluence absorbed by the substrate of about 3.9 mJ/cm^2 . By changing the temporal delay between the probe and desorption laser pulses an arrival time spectrum of the H atoms in the probe volume is recorded.⁵⁸ After transformation into the velocity space, rather slow velocities for the desorbing hydrogen atoms are observed, see Fig. 8. A theoretical analysis of the desorption process reveals that multiple electron scattering on the H adsorbate causes the desorption, thus a typical DIMET (desorption induced by multiple electronic transitions) process is observed.⁵⁹ These hot electrons are generated by the absorption of the high energy XUV photon by the graphite substrate, producing a dense cloud of hot electrons in the vicinity of the surface with a rather short lifetime of about half a picosecond. Multiple electron scattering leads to a successive population of higher vibrational states of the C-H vibration, and, in a final step, the desorption out of one of these vibrational states. A peak at a velocity of $v = 425$ ms^{-1} , and two shoulders at 255 ms^{-1} and 136 ms^{-1} are apparent. These very slow H atoms show an average kinetic energy of only $E_{\text{kin}} = 0.8$ meV. A second maximum at a velocity of $v = 1000$ ms^{-1} yields an average kinetic energy of $E_{\text{kin}} = 5.3$ meV. The relative integrated intensities of the very slow and the faster peak are about the same.

The theoretical analysis shows good agreement with the experimental observations when the very slow peak is caused by desorption of H atoms out of a hydrogen surface dimer pair which occupies the para position on the underlying graphene lattice.^{38,60,61} The faster peak can be reconciled when one assumes that the desorption




Fig. 8 Velocity distribution of neutral H atoms desorbing from chemisorbed H on graphite in a uniform para adsorbate configuration. This results in a coverage of 0.25 ML H on graphite. Average kinetic energies of $\langle E_{\text{kin}} \rangle = 0.8$ meV and 5.3 meV are deduced for the para pairs and uniform para configurations, respectively.

occurs out of a uniform para configuration of adsorbed H atoms.⁵⁸ In this configuration all H atoms are only adsorbed in a para position relative to each other on the graphene lattice. Thereby additional energy is gained by the adsorbate system, as depicted in Fig. 9. There the adsorption energies gained by both atoms when adsorbing a second hydrogen atom in ortho-, para-, and uniform para-position relative to a first one is shown. Besides a barrier-free adsorption, as for a para pair, the energy gain is even higher by about $\Delta E \sim 0.6$ eV in the uniform para configuration. Thus, instead that both hydrogen atoms show a binding energy of about $E_b \sim 0.8$ eV each as for the adsorption of single isolated H atoms,⁶³ the uniform para pair shows together an adsorption energy of $E_b \sim 2.6$ eV, thus 1.0 eV larger than the sum of two single H atoms on graphene.⁶² This theoretical result is also in good agreement with the experimental observation.

Nitric oxide

The investigation of the XUV light induced desorption process of nitric oxide from graphite is driven by the desire to learn more about the fundamental microscopic desorption dynamics of small molecules. NO can be detected with very high sensitivity and with internal state selection by resonantly enhanced (1+1) REMPI. It are the population distributions over the internal states of a product molecule which allow a deeper understanding of the microscopic dynamics. Figure 10 shows a rotational state resolved REMPI spectrum of the $A^2\Sigma^+ \leftarrow X^2\Pi$ ($v'' = 0, 1$) γ -bands around $\lambda = 226$ nm after excitation of the nitric oxide covered graphite surface by XUV light at $h\nu = 57.1$ eV. It is evident that besides molecules in the vibrational ground state, also vibrationally excited molecules are detected. An analysis of the rotational




Fig. 9 Potential energy diagram for H atoms chemisorbed on graphene for the ortho-, para-, and uniform para configurations.

state distribution yields a non-Boltzmann distribution with an average rotational energy in $v'' = 0$ of $\langle E_{\text{rot}} \rangle = 311 \text{ cm}^{-1}$ or a rotational temperature of about $\langle E_{\text{rot}} \rangle / k = 448 \text{ K}$.⁶³ For the vibrationally excited state the same distribution of the population in the rotational states is observed. The high intensity of the lines belonging to the $\gamma(1-1)$ band suggest already a considerable vibrational excitation. For these two vibrational states a vibrational temperature of about $T_{\text{vib}} \sim 1580 \text{ K}$ is found. Higher vibrational states could not be measured due to the limited beam time at the FEL facility. The translational energy of neutral NO molecules was obtained by changing the temporal delay between the FEL pulse (at $h\nu = 38 \text{ eV}$) and the detection laser pulse. A distribution is observed which can be fitted by the combination of two Maxwellian velocity distributions. They show the temperature parameters of $T_{\text{kin}} = 480 \text{ K}$ and $T_{\text{kin}} = 1717 \text{ K}$ and relative yields of 35% and 65%, respectively. On average, a kinetic energy of $E_{\text{kin}} = 170 \text{ meV}$ is released in the desorption of neutral NO molecules. At this photon energy the desorption cross section was determined to about $\sigma = (1.1 \pm 0.4) \cdot 10^{-17} \text{ cm}^2$.⁶³ The desorption yield of NO from this $(\text{NO})_2/\text{graphite}$ system exhibits a slightly nonlinear dependence on the XUV pulse intensity.⁶³ Exponents between $n = 1.4$ and 3 have been observed in different runs. Such a dependence would then enable the application of two-pulse correlated excitation of the system in order to learn more about the temporal behaviour of the excited system.

At surface temperatures below $T_s \sim 100 \text{ K}$ nitric oxide condenses as cis-dimer $(\text{NO})_2$ on graphite.⁶⁴ The release of NO molecules thus has to be preceded by the dissociation of the NO dimer. Direct dissociative ionisation cross sections are about one order of magnitude lower than the desorption cross section measured in this study.⁶⁵




Fig. 10 Spectrum of the NO $\gamma(0-0)$ and $\gamma(1-1)$ bands around $\lambda = 227$ nm and 224.5 nm, respectively, after desorption from $(\text{NO})_2/\text{graphite}$ by XUV pulses of $h\nu = 57.1$ eV.

Therefore, a direct photoexcitation of the adsorbate is unlikely, and an indirect excitation mechanism is supported. The XUV photons are predominantly absorbed by the graphite substrate, thereby creating a cloud of hot electrons which interacts with the adsorbate. This interaction leads to a temporarily excited $(\text{NO})_2^-$ dimer anion, where one NO molecule lies flat on the surface. For the dimer anion the trans geometry is, however, energetically more favourable than the cis geometry.⁶⁶ Therefore, the temporarily formation of a dimer anion leads to both a strong rotational excitation of the leaving NO fragment due to this transition from a cis to a trans conformation of the dimer while in the excited state, as well as to a high vibrational excitation due to the transient negative ion, as in the case of metal substrates. Both conclusions are supported by the experimental observations.

15 Conclusion

Femtosecond pulsed XUV radiation from free-electron laser sources is a valuable tool to induce reactive processes in molecular ices and elucidate the dynamics of such systems. Preliminary results of such reactions in water ice leading to cationic products have been presented. The high photon energy induces the formation of new species, in this case D_2O_2 , which may act as reservoirs for further reactions. The observation of extremely slow H atoms from the chemisorbed state on graphite is really surprising and may have an influence on the understanding of the dynamics in accretion discs. The extension to other systems, such as doped or molecular ices where the neutral constituents themselves can be detected with internal state selection, will shed light onto the microscopic desorption dynamics. The study of the $(\text{NO})_2$ thin film presented provides a first glimpse of the wealth of information which can be obtained.

However, many interesting questions remain at present unanswered, including the velocity distribution of most of the oxygen containing cation products observed and the relative roles of direct photoprocesses and processes induced by fast electrons liberated by the ionization of ice constituents. Also, the detection of neutral desorbing products, e.g. D atoms or OD radicals would contribute to a deeper understanding. Future experiments on VUV and XUV FEL facilities will allow to address these questions and to contribute to a deeper microscopic understanding of reaction and desorption processes in complex ices.

Acknowledgement

The authors thank the FLASH team at HASYLAB for providing the XUV pulses, D. Gerlich, T. Olsen and J. Thrower for helpful discussions, and S. Eppenhoff and F. Wahlert for extensive technical support. The Bundesministerium für Bildung und Forschung provided partial financial support via grant nos. 05 KS4PMC/8 and 05 KS7PM1 within the research program FSP 301 “FLASH” which is gratefully acknowledged.

References

- ^a *Physikalisches Institut, Westfälische Wilhelms-Universität, 48149 Münster, Germany: E-mail:*
10 *hzach@uni-muenster.de*
- ^b *present address: Helmholtz-Zentrum Berlin für Materialien und Energie, Berlin*
- 1 A.G.G.M. Tielens, *Rev. Mod. Phys.*, 2013, **85**, 1021
 - 2 E. Herbst, E.F. van Dishoeck, *Annu. Rev. Astro. Astrophys.*, 2009, **47**, 427
 - 3 R.J. Gould, E.E. Salpeter, *Astrophys. J.*, 1963, **138**, 393
 - 4 D.A. Williams, E. Herbst, *Surf. Sci.*, 2002, **500**, 823
 - 5 L. Hornekaer, A. Baurichter, V. Petrunin, D. Field, A.C. Luntz, *Science*, 2003, **302**, 1943
 - 6 S.C. Creighan, J.S.A. Perry, S.D. Price, *J. Chem. Phys.*, 2006, **124**, 114701
 - 7 J.L. Lemaire, G. Vidali, S. Baouche, M. Chehrouri, H. Chaabouni, H. Mokrane, *Astrophys. J. Lett.*, 2010, **725**, L156
 - 8 G. Vidali, *J. Low Temp. Phys.*, 2013, **170**, 1
 - 9 J.S. Mathis, P.G. Mezger, N. Panagia, *Astron. Astrophys.*, 1983, **128**, 212
 - 10 S.S. Prasad, S.P. Tarafdar, *Astrophys. J.*, 1983, **267**, 603
 - 11 J. Najita, E.A. Bergin, J.N. Ullom, *Astrophys. J.*, 2001, **561**, 880
 - 12 T. Henning, D. Semenov, *Chem. Rev.*, 2013; dx.doi.org/10.1021/cr400128p
 - 13 J.S. Mathis, W. Rumpl, K.H. Nordsiek, *Astrophys. J.*, 1977, **217**, 425
 - 14 B.T. Draine and H.M. Lee, *Astrophys. J.*, 1984, **285**, 89
 - 15 D. Hollenbach, M.J. Kaufman, E.A. Bergin, G.J. Melnick, *Astrophys. J.*, 2009, **690**, 1497
 - 16 M.S. Westley, R.A. Baragiola, R.E. Johnson, G.A. Baratta, *Nature*, 1995, **373**, 405
 - 17 S.R. Baggott, K.W. Kolasinski, L.M.A. Perdigo, D. Riedel, Q. Guo, R.E. Palmer, *J. Chem. Phys.*, 2002, **117**, 6667
 - 18 J.D. Thrower, D.J. Burke, M.P. Collings, A. Dawes, P.D. Holton, F. Jamme, P. Kendall, W.A. Brown, I.P. Clark, H.J. Fraser, M.R.S. McCoustra, N.J. Mason, A.W. Parker, *Astrophys. J.*, 2008, **673**, 1233
 - 19 E.C. Fayolle, M. Bertin, C. Romanzin, X Michaut, K.I. Öberg, H. Linnartz, J.-H. Fillion, *Astrophys. J. Lett.*, 2011, **739**, L36
 - 20 K.I. Öberg, E.F. van Dishoeck, H. Linnartz, *Astron. Astrophys.*, 2009, **496**, 281
 - 21 K.I. Öberg, E.F. van Dishoeck, H. Linnartz, S. Andersson, *Astrophys. J.*, 2010, **718**, 832
 - 22 D. McGonagle, L.M. Ziurys, W.M. Irvine, Y.C. Minh, *Astrophys. J.*, 1990, **359**, 121
 - 23 S.B. Charnley, S.D. Rogers, P. Ehrenfreund, *Astron. Astrophys.*, 2001, **378**, 1024
 - 24 D.T. Halfen, A.J. Apponi, L.M. Ziurys, *Astrophys. J.*, 2001, **561**, 244
 - 25 E. Congiu, H. Chaabouni, C. Laffon, P. Parent, S. Baouche, F. Dulieu, *J. Chem. Phys.*, 2012, **137**, 054713
 - 26 J. Bouwman, H.M. Cuppen, M. Steglich, L.J. Allamandola, H. Linnartz, *Astron. Astrophys.*, 2011, **529**, A46
 - 27 S.H. Cuyllle, H. Linnartz, J.D. Thrower, *Chem. Phys. Lett.*, 2012, **550**, 79
 - 28 T.M. Orlando, A.B. Alexandrov, J. Herring, *J. Phys. Chem. B*, 2003, **107**, 9370
 - 29 J.D. Thrower, M.P. Collings, F.J.M. Rutten, M.R.S. McCoustra, *Chem. Phys. Lett.*, 2011, **505**, 106
 - 30 A.G.M. Abdulgalil, D. Marchione, J.D. Thrower, M.P. Collings, M.R.S. McCoustra, F. Islam, M.E. Palumbo, E. Congiu, F. Dulieu, *Phil. Trans. R. Soc. A*, 2013, **371**, 20110586
 - 31 K. Tiedtke et al., *New J. Phys.*, 2009, **11**, 023029

- 32 R. Mitzner, A.A. Sorokin, B. Siemer, S. Roling, M. Rutkowski, H. Zacharias, M. Neeb, T. Noll, F. Siewert, W. Eberhardt, M. Richter, P. Juranic, K. Tiedtke, J. Feldhaus, *Phys. Rev. A*, 2009, **80**, 025401
- 33 R. Mitzner, B. Siemer, M. Neeb, T. Noll, F. Siewert, S. Roling, M. Rutkowski, A.A. Sorokin, M. Richter, P. Juranic, K. Tiedtke, J. Feldhaus, W. Eberhardt, H. Zacharias, *Opt. Expr.*, 2008, **16**, 19909
- 34 M. Wöstmann, R. Mitzner, T. Noll, S. Roling, B. Siemer, F. Siewert, S. Eppenhoff, F. Wahlert, H. Zacharias, *J. Phys. B: At. Mol. Opt. Phys.*, 2013, **46**, 164005
- 35 B.L. Henke, E.M. Gullikson, E.M. Davis, *At. Data Nucl. Data Tables*, 1993, **54**, 18
- 36 J.H. Hubell, W.J. Feigele, E.A. Briggs, R.T. Brown, D.T. Cromer, R.J. Howerton, *J. Phys. Chem. Ref. Data*, 1977, **4**, 471
- 37 L. Jeloica, V. Sidis, *Chem. Phys. Lett.*, 1999, **300**, 157
- 38 X. Sha, B. Jackson, *Surf. Sci.*, 2002, **496**, 318
- 39 V. Hermann, B.D. Kay, A.W. Castleman Jr., *Chem. Phys.*, 1982, **72**, 185; F. Dong, S. Heinbuch, J.J. Rocca, E.R. Bernstein, *J. Chem. Phys.*, 2006, **124**, 224319
- 40 J. E. Reutt, L. S. Wang, Y. T. Lee, D. A. Shirley, *J. Chem. Phys.*, 1986, **85**, 6928
- 41 J. H. D. Eland, *Adv. Mass Spectrom.*, 1974, **6**, 917
- 42 F.S. Ashmore A.R. Burgess, *J. Chem. Soc. Faraday Trans. 2*, 1997, **73**, 1247; S. N. Foner, R. L. Hudson, *J. Chem. Phys.*, 1962, **36**, 2676
- 43 R. Atkinson, D. L. Baulch, R. A. Cox, J. N. Crowley, R. F. Hampson, R. G. Hynes, M. E. Jenkin, M. J. Rossi, and J. Troe, *Atmos. Chem. Phys.*, 2004, **4**, 1461
- 44 L.J. Stief, V.J. De Carlo, *J. Chem. Phys.*, 1969, **50**, 3
- 45 R.G. Tonkyn, J.W. Winniczek, M.G. White, *Chem. Phys. Lett.*, 1989, **164**, 137
- 46 *NIST – JANAF Thermochemical Tables, 4th edition*, M.W. Chase, ed., 1998, Monograph 9, ISBN 1-56396-831-2
- 47 R.T. Wiedmann, R.G. Tonkyn, M.G. White, K. Wang, V. McKoy, *J. Chem. Phys.*, 1992, **97**, 768
- 48 R. L. Kelly, *J. Phys. Chem. Ref. Data*, 1987, **16**, suppl. 1
- 49 T.M. Orlando, M.T. Sieger, *Surf. Sci.*, 2003, **528**, 1
- 50 X.N. Pan, A.D. Bass, J.P. Jay-Gerin, L. Sanche, *Icarus*, 2004, **172**, 521
- 51 B. Redlich, H. Zacharias, G. Meijer, G. von Helden, *Surf. Sci.*, 2002, **502 – 503**, 325
- 52 B. Redlich, L. van der Meer, H. Zacharias, G. Meijer, G. von Helden, *Nucl. Instrum. Meth. A*, 2003, **507**, 556
- 53 B. Redlich, H. Zacharias, G. Meijer, G. von Helden, *Phys. Chem. Chem. Phys.*, 2002, **4**, 3448
- 54 B. Redlich, H. Zacharias, G. Meijer, G. von Helden, *J. Chem. Phys.*, 2006, **124**, 044704
- 55 B. Siemer, T. Hoger, M. Rutkowski, R. Treusch, H. Zacharias, *J. Phys. Condens. Matter*, 2010, **22**, 084013
- 56 E.R. Latimer, F. Islam, S.D. Price, *Chem. Phys. Lett.*, 2008, **455**, 174
- 57 L. Gavilan, J.L. Lemaire, G. Vidali, T. Sabri, C. Jäger, *Astrophys. J.*, to be published
- 58 B. Siemer, T. Olsen, T. Hoger, M. Rutkowski, C. Thewes, S. Düsterer, J. Schiötz, H. Zacharias, *Chem. Phys. Lett.*, 2010, **500**, 291
- 59 J.A. Misewich, T.F. Heinz, D.M. Newns, *Phys. Rev. Lett.*, 1992, **68**, 3737
- 60 N. Rougeau, D. Teillet-Billy, V. Sidis, *Chem. Phys. Lett.*, 2006, **431**, 135
- 61 L. Hornekaer, E. Rauls, W. Xu, Z. Slijivancanin, R. Otero, I. Stensgaard, E. Laegsgaard, B. Hammer, F. Besenbacher, *Phys. Rev. Lett.*, 2006, **97**, 186102
- 62 R. Frigge, T. Hoger, B. Siemer, H. Witte, M. Silies, H. Zacharias, T. Olsen, J. Schiötz, *Phys. Rev. Lett.*, 2010, **104**, 256102
- 63 B. Siemer, T. Hoger, M. Rutkowski, S. Düsterer, H. Zacharias, *J. Phys. Chem. A*, 2011, **115**, 7356
- 64 J.P. Coulomb, J. Suzanne, M. Bienfait, M. Matecki, A. Thomy, B. Croset, C. Marti, *J. de Physique*, 1980, **41**, 1155
- 65 J.A.R. Samson, T. Masuoka, P.N. Pareek *J. Chem. Phys.*, 1985, **83**, 11
- 66 B. Urban, A. Strobel, V.E. Bondybey, *J. Chem. Phys.*, 1999, **111**, 8939