



Comparative evaluation of the cardio-metabolic effects of hydroxytyrosol and its lipophilic derivatives (hydroxytyrosyl acetate and ethyl hydroxytyrosyl ether) in hypercholesterolemic rats

Journal:	<i>Food & Function</i>
Manuscript ID:	FO-ART-12-2013-060677.R1
Article Type:	Paper
Date Submitted by the Author:	15-Mar-2014
Complete List of Authors:	<p>Tabernero, Maria; Hospital La Paz Health Research Institute, Experimental Surgery Department Sarria, Beatriz; CSIC, Largo, Carlota; Hospital La Paz Health Research Institute, Experimental Surgery Department Martínez-López, Sara; Institute of Food Science, Technology and Nutrition (CSIC), Metabolism and Nutrition Madrona, Andres; Instituto de Química Médica. CSIC., Espartero, Jose Luis; University of Seville, Bravo, Laura; Institute of Food Science, Technology and Nutrition (CSIC), Metabolism and Nutrition Mateos, Raquel; Institute of Food Science, Technology and Nutrition (CSIC), Metabolism and Nutrition</p>

1 **Comparative evaluation of the metabolic effects of hydroxytyrosol and its**
2 **lipophilic derivatives (hydroxytyrosyl acetate and ethyl hydroxytyrosyl**
3 **ether) in hypercholesterolemic rats**

4
5 María Tabernero[§], Beatriz Sarriá[∞], Carlota Largo[§], Sara Martínez-López[∞], Andrés Madrona[‡],
6 José Luis Espartero[‡], Laura Bravo[∞], Raquel Mateos^{∞*}.

7
8 [§]Hospital La Paz Health Research Institute (IdiPaz); Madrid, Spain. [∞]Instituto de Ciencia y
9 Tecnología de Alimentos y Nutrición (ICTAN-CSIC), Madrid, Spain. [‡]Dpto. Química Orgánica y
10 Farmacéutica. Facultad de Farmacia, Universidad de Sevilla, Seville, Spain.

11
12

13 **Running title:** Metabolic effects of hydroxytyrosol and its lipophilic derivatives

14 * **Corresponding author:** Raquel Mateos. Department of Metabolism and Nutrition,
15 Instituto de Ciencia y Tecnología de Alimentos y Nutrición (ICTAN-CSIC). José Antonio
16 Novais 10, Ciudad Universitaria, 28040, Madrid, Spain. Phone: +34.91.544.56.07; Fax:
17 +34.91.549.36.27; e-mail: raquel.mateos@ictan.csic.es

18

19

20 **Abstract**

21 Hydroxytyrosol (HT), a virgin olive oil phenolic phytochemical with proven health benefits,
22 has been used to generate new lipophilic antioxidants to preserve fats and oils against
23 autoxidation. The aim of this work is to comparatively evaluate the physiological effects of
24 HT and the lipophilic derivatives, hydroxytyrosyl acetate (HT-Ac) and ethyl hydroxytyrosyl
25 ether (HT-Et) in a high-cholesterol fed animals. Male Wistar rats (n=8) were fed a standard
26 diet (*C* group), a cholesterol-rich diet (*Chol* group) or cholesterol-rich diets supplemented
27 with the phenolic compounds (*HT* group, *HT-Ac* group and *HT-Et* group) for 8 weeks. Body
28 and tissue weights, lipid profile, redox status, and biochemical, hormonal, and inflammatory
29 biomarkers were evaluated. Plasma levels of total, LDL-cholesterol, glucose, insulin and
30 leptin, as well as malondialdehyde in serum increased in *Chol* compared to *C* ($p<0.05$). Rats
31 fed the test diets improved their glucose, insulin, leptin and MDA levels and antioxidant
32 capacity status, being HT-Ac the most effective compound. The studied phenolic compounds
33 also modulated TNF- α and IL-1 β plasma levels compared to *Chol*. HT-Ac and HT-Et
34 improved adipose tissue distribution and adipokine production, decreasing MCP-1 and IL-1 β
35 levels. Our results confirm the metabolic effects of HT, which are maintained and even
36 improved by the hydrophobic derivatives, particularly in HT-Ac.

37

38 **Keywords:** Ethyl hydroxytyrosyl ether; hydroxytyrosol; hydroxytyrosyl acetate;
39 hypercholesterolemic rats; metabolic stress.

40

41 1. Introduction

42 The most representative phenolic compound in virgin olive oil, hydroxytyrosol
43 (HT), plays an important role in the prevention of degenerative diseases¹⁻². Explicitly, it
44 is involved in the reduction of LDL oxidation, a well-known cardiovascular risk factor³,
45 and it is transiently associated with LDL lipoproteins⁴. Moreover, HT is involved in the
46 inhibition of lipid and protein oxidation in human plasma⁵, having anti-inflammatory⁶
47 and antiplatelet aggregation activities⁷. Additionally, there is evidence that olive oil
48 phenolic compounds decrease plasma glucose levels in rats⁸⁻⁹. HT can be efficiently
49 recovered from olive by-products, which has prompted the study of the biological and
50 health effects of HT from alperujo^{8,10} or olive leaves.^{9,11}

51 On the other hand, the food industry demands new lipophilic antioxidants to
52 preserve fats and oils against autoxidation. Two series of hydrophobic derivatives of HT,
53 hydroxytyrosyl esters¹² and hydroxytyrosyl ethers¹³, have been synthesized.
54 Hydroxytyrosyl acetate (HT-Ac) merits special interest among hydroxytyrosyl esters,
55 since it is an antioxidant naturally present in virgin olive oil¹⁴ that is transported across
56 the small intestinal epithelial cell barrier more efficiently than HT,¹⁵ showing a higher
57 hepatic bioavailability than HT¹⁶. Moreover, HT-Ac has shown protective effects against
58 oxidative DNA damage in blood cells¹⁷, iron-induced oxidative stress in human cervical
59 cells,¹⁸ and oxidative stress in HepG2 cells¹⁹. Hydroxytyrosyl ethyl ether (HT-Et), with an
60 alkyl chain with the same length as the acyl chain in HT-Ac, was included in this study to
61 assess the influence of the different functional groups (etherification versus esterification)
62 on the biological activity of the two phenolic compounds. HT-Et is absorbed to a higher
63 extent in Caco-2 cells than its precursor HT,²⁰ with absorption rate similar to that of HT-
64 Ac,¹⁵ being also broadly taken-up by HepG2 cells²¹. In addition, HT-Et protects hepatic
65 human HepG2 cells against oxidative stress²² and inhibits platelet activation after oral

66 administration in rats²³. These antecedents justify using this compound although it is not
67 naturally present in virgin olive oil.

68 Both HT-Ac and HT-Et maintain the orthodiphenolic group intact and show
69 higher antioxidant capacity than HT²⁴⁻²⁵, preserving their potential application as
70 antioxidants to stabilize foodstuffs or as functional food ingredients.

71 With these antecedents, the aim of the present study was to comparatively
72 analyse the ability of HT and its lipophilic derivatives, HT-Ac and HT-Et, counteracting
73 the metabolic deregulation derived from consuming a high-cholesterol diet, focusing on
74 the effects on plasma lipids, glucose levels, hormone response, oxidative stress and pro-
75 inflammatory status. In addition, body weight gain, adipose tissue distribution and
76 secretion of adipokines from visceral fat were evaluated.

77

78 **2. Experimental**

79 **2.1. Materials and Chemical**

80 Cholesterol, cholic acid, palmitic acid, 6-hydroxy-2,5,7,8-tetramethylchroman-2-
81 carboxylic acid (Trolox), 1,1,3,3-tetraethoxypropane, fluorescein, 2,2'-azobis(2-
82 amidinopropane) dihydrochloride (AAPH), trizma base, dithiothreitol, 1,1,3,3-
83 tetraethoxypropane and 2,4-dinitrophenylhydrazine were purchased from Sigma-Aldrich
84 Chemical (Madrid, Spain). Sodium hydroxide, sulphuric acid, sodium hydrogen phosphate
85 and potassium dihydrogen phosphate were from Panreac (Madrid, Spain). The Bradford
86 reagent was from BioRad (BioRad Laboratories S.A., Madrid, Spain). Free fatty acids (FFA)
87 colorimetric kit was purchased from Roche (Roche Applied Science, Madrid Spain). Rat
88 adipocyte (RADPCYT-82K, MilliplexMap Rat Adipocyte Panel) and adipokine (RADPK-
89 81K, MilliplexMap Kit Rat Serum Adipokine Panel) Milliplex kits were acquired from

90 Millipore (Millipore, Billerica, MA, US, USA). Other reagents were of analytical or
91 chromatographic quality.

92 HT was isolated with 98% purity from olive oil waste water following a patented
93 procedure²⁶ and further purified by column chromatography.

94 HT-Ac was obtained from HT in ethyl acetate after incubation with *p*-
95 toluenesulfonic acid and purification by column chromatography following a patented
96 procedure¹².

97 HT-Et was obtained from HT by chemical synthesis as described elsewhere¹³.

98

99 **2.2. Diet, animals and experimental design**

100 Forty male Wistar rats (200–225 g body weight) were purchased from an accredited
101 supplier (Charles River Laboratorios España, S.A. Barcelona, Spain) and housed in
102 metabolic cages. They had free access to food and water, and were maintained under a
103 normal light–dark cycle in the Experimental Surgery Service of La Paz University Hospital
104 (registration number: 280790001941). After one week of acclimation, animals were
105 randomly distributed into 5 different experimental groups (8 rats per group). One group
106 received the standard, maintenance rodent diet (A04-SAFE, Augy, France), which is the
107 control (*C* group), and four groups were fed the standard diet supplemented with 2%
108 cholesterol and 0.4% cholic acid in order to generate the hypercholesterolemic model as
109 described elsewhere²⁷. In three of these groups the hypercholesterolemic diet was
110 supplemented with 0.04% of HT (*HT* group), HT-Ac (*HT-Ac* group) and HT-Et (*HT-Et*
111 group), respectively, to assess the potential beneficial effects of these phenolic compounds in
112 the hypercholesterolemic rat model. Nutritional and energetic content of each diet is
113 summarized in Table 1. During the whole intervention, animal health status and dietary
114 tolerance were monitored by veterinary individual observation in addition to weekly control

115 of animals' weight (Table 2). After 8 weeks animals were subjected to complete
116 exsanguination under general anesthesia (isoflurane 2%). Plasma and serum fractions were
117 separated by centrifugation (10 minutes at 2500 g) using pretreated EDTA or Silica Act Cot
118 Activator blood collection tubes (BD Vacutainer, Plymouth, UK) respectively, and stored at -
119 20°C until further analysis. Liver, kidney, heart and adipose tissue from retroperitoneal and
120 epididymal areas were collected, weighed and washed in ice-cold phosphate buffered saline.
121 Samples were immediately frozen in liquid nitrogen and stored at -80°C until analysis. The
122 study protocol was approved by the Institutional Animal Ethics Committee of La Paz
123 University Hospital (Madrid, Spain) and procedures were performed in accordance with
124 Spanish law for the protection of experimental animals and other research purposes (RD
125 53/2013, BOE n° 34 Sec I pg 11370 8th February 2013).

126

127 **2.3. Blood biochemical analyses**

128 Triglycerides, total cholesterol, LDL-cholesterol, HDL-cholesterol, glucose and
129 creatinine in plasma were determined using an automated analyser (Beckman Coulter-Former
130 Olympus Diagnostics AU 5420, Nyon, Switzerland). Plasma free fatty acids (FFA)
131 concentration was determined with a colorimetric commercial assay kit (Free Fatty Acids,
132 Half Micro Test, Roche Applied Science) using palmitic acid as standard.

133

134 **2.4. Determination of oxygen radical scavenging capacity (ORAC) in serum.**

135 Serum antioxidant activity was analysed using the hydrophilic oxygen radical
136 scavenging capacity (ORAC) assay according to the method developed by Huang, Ou,
137 Hampsch-Woodill, Flanagan & Prior²⁸. The fluorescence at 485 and 528nm excitation and
138 emission wavelengths, respectively, was determined in a 96 well microplate reader (Bio-Tek,
139 Winooski, VT, USA). ORAC values were expressed as micromoles of Trolox / mL serum.

140

141 2.5. Determination of Malondialdehyde (MDA) in serum and liver.

142 MDA was determined as its hydrazone by high-performance liquid chromatography
143 using 2,4-dinitrophenylhydrazine for derivatization²⁹. Livers (0.5g) were homogenized (1:5
144 w/v) in ice-cold 0.25M Trizma base buffer pH 7.4 containing 5mM dithiothreitol using an
145 Ultra Turrax (IKA® Works Inc., WilmingtonNC) at 18000 rpm. After centrifugation
146 (11000rpm 30min, 4°C), supernatants were collected for MDA quantification. Serum samples
147 were analysed directly. Standard MDA was prepared by acidic hydrolysis of 1,1,3,3-
148 tetraethoxypropane in 1% sulphuric acid. Concentrations were expressed as nanomoles of
149 MDA per milligram of protein in liver tissue and per millilitre in serum. Protein content in
150 liver homogenates was estimated by the Bradford method using a Bio-Rad protein assay kit.

151

**152 2.6. Determination of cytokines and hormones in plasma and adipose tissue by
153 immunoassay.**

154 A sample of white adipose tissue from the retroperitoneal area (0.2g) was collected
155 and homogenized (1:3.5 w/v) in PBS (pH 7.4) with Triton X-100 and protease inhibitor by
156 low temperature sonication (17 microns, 3 consecutive pulses of 15 seconds at 30 seconds
157 intervals). After centrifugation (5000rpm, 10min, 4°C), the lipid layer was removed and
158 supernatants were collected for adipokine determination. The concentration of leptin,
159 interleukin-1 β (IL-1 β), interleukin-6 (IL-6), tumour necrosis factor- α (TNF- α), monocyte
160 chemoattractant protein-1 (MCP-1) and plasminogen activator inhibitor-1 (PAI-1) in the
161 retroperitoneal adipose tissue were determined using a Rat Adipocyte Multiplex Kits
162 (RADPCYT-82K Milliplex Map Kit, Rat Adipocyte Panel, Millipore, Billerica, MA, US).
163 IL-1 β , IL-6, TNF- α , insulin and leptin concentrations were determined in plasma samples
164 obtained at the end of the intervention using a Rat Adipokine multiplex kits (RADPK-81K,

165 Milliplex Map Kit, Rat Serum Adipokine Panel, Millipore, Billerica, MA, US). Multianalyte
166 profiling panels were used according to manufacturer's instructions and analysed on a
167 LuminexLX200 Analyzer. Data were analysed using the 3.1 xPONENT software (Millipore);
168 high and low concentration quality controls were used with all the biomarkers. Protein
169 content in adipose tissue homogenates was estimated by the Bradford method using a Bio-
170 Rad protein assay kit.

171

172 **2.7. Statistical analysis.**

173 All data were expressed as the mean \pm standard deviation (SD). Non parametric
174 significant test (Kruskal-Wallis one-way analysis of variance and Mann-Whitney *U* test) was
175 performed to compare values. Differences were considered statistically significant when $p <$
176 0.05 and in Mann-Whitney *U* test the significance was adjusted with a Bonferroni test. Data
177 were analysed using the statistical package SPSSv.19.0 software (SPSS, Institute Inc, Cary,
178 NC).

179

180 **3. Results**

181 **3.1. Food intake, body weight gain and tissues weights**

182 All diets provided equivalent amounts of carbohydrates and proteins. However,
183 cholesterol-rich diets led to an excess of lipids resulting in an 8.5 Kcal/100g higher caloric
184 intake. During the 8-week study period, animals in the five experimental groups had similar
185 daily food intakes (approximately 20g/d).

186 Therefore, the groups consuming the cholesterol-rich diets (*Chol*, *HT*, *HT-Ac* and
187 *HT-Et*) presented a daily caloric excess of approximately 2 Kcal/day. However, this did not
188 induce significant differences in the body weight gain compared with the control animals, as
189 can be seen in Table 2.

190 Considering an average daily food intake of 20g and an average body weight of
191 300g, the daily phenolic compound intake was 25mg/kg body weight/day. This is a relatively
192 low dose, comparable with doses used in some other studies⁸⁻¹¹ and within a range proven to
193 elicit no toxic effects in rats. Accordingly, no pathologic alterations or behaviours were
194 observed in the animals during the study. Organs, including kidney or heart among others,
195 presented normal appearance by gross observation. However, livers in animals fed the
196 cholesterol-rich diets showed a whitish appearance compared with the *C* group, suggestive of
197 fat accumulation, being significantly heavier than those of the *C* group (Table 2). Adipose
198 tissue depots were obtained from retroperitoneal and epididymal areas, and when their
199 weights were normalized for the final animal body weight, no differences were observed
200 among different treatments. However, the ratio obtained with the adipose tissue from
201 retroperitoneal area/epididymal area showed significant differences, being higher in the *Chol*
202 and *HT* groups than in the *C* group.

203

204 **3.2. Plasma cholesterol profile**

205 As Figure 1 shows, animals fed the cholesterol-rich diets had a marked
206 hyperlipidaemia, with high LDL-cholesterol and total cholesterol concentrations, although
207 HDL-cholesterol levels were not affected as compared to *C* group. Although not reaching the
208 values of control animals, supplementation of the cholesterol-rich diet with 0.04% HT-Ac
209 significantly decreased the total cholesterol and LDL-cholesterol levels as compared with the
210 *Chol* group ($p<0.05$), which was not attained in the *HT* and *HT-Et* groups.

211

212 **3.3. Biochemical parameters and hormones involved in energy homeostasis**

213 The high-cholesterol diet used in the present study induced a significant increase in
214 glucose levels in the *Chol* group compared to control animals ($p<0.05$). All the tested

215 phenolic compounds counteracted the glucose increase, with a partial effect of HT-Et
216 whereas values in *HT-Ac* and *HT* groups returned to control levels (Table 3).

217 Triglycerides concentrations were similar in all groups, although phenolic
218 compounds supplementation induced a moderate yet non-significant decrease. No differences
219 were observed in plasma concentrations of free fatty acids, whereas creatinine concentrations
220 were statistically higher in *Chol* and *HT* groups, compared to *C*, *HT-Ac* and *HT-Et* groups
221 (Table 3).

222 After consuming the high-cholesterol diet for 8 weeks significant differences in
223 plasma insulin and leptin concentrations were observed in *Chol* compared to *C* group.
224 However, rats consuming diets supplemented with HT and HT-Ac had significantly lower
225 plasma insulin levels than *Chol*, decreasing to control or even lower values as in the HT-Ac
226 group (Table 3). *HT-Et* also presented lower insulin levels than *Chol* group although not
227 statistically different ($p=0.070$), showing intermediate values between *Chol* and *C* groups.
228 Leptin levels were lower in *HT* and *HT-Et* groups than in the *Chol* group, although not
229 reaching values of control animals. Again, *HT-Ac* group had significantly lower plasma
230 leptin levels than *Chol* group and in the range of the control values. In the retroperitoneal
231 adipose tissue, leptin concentration was slightly lower in all cholesterol treated groups
232 compared to the *C* animals, although not significantly different.

233

234 **3.4. Redox status and inflammatory biomarkers in blood and tissue samples**

235 Serum antioxidant activity (Table 4), analysed using the ORAC assay, was lower in
236 *Chol* group compared to the control group without reaching the level of statistical
237 significance. However, rats fed with *HT* and *HT-Ac* diets showed a significantly higher
238 serum antioxidant activity compared with *Chol* group ($p<0.05$).

239 Regarding the biomarker of lipid peroxidation, *Chol* group showed MDA levels
240 significantly higher than *C* animals in serum and liver ($p < 0.05$). Supplementation with any of
241 the studied phenolic compounds decreased MDA levels to control values in serum (Table 4).
242 However, although the three phenolic compounds-supplemented groups showed lower
243 hepatic MDA levels than *Chol* group, differences were statistically significant only in the
244 *HT-Et* group ($p < 0.05$) (Table 4).

245 Consuming the cholesterol-rich diet significantly increased the pro-inflammatory
246 cytokine $\text{TNF}\alpha$ plasma concentrations compared to *C* group, effect that was totally
247 counteracted by the three phenolic compounds (Table 4). A similar response was observed
248 with $\text{IL-1}\beta$, with increased values in the *Chol* group returning to control levels in the *HT*, *HT-*
249 *Ac* and *HT-Et* animals, although differences did not reach the level of statistical significance
250 (Table 4). Plasma IL-6 was not affected by the cholesterol-rich diets or the phenolic
251 supplementation.

252 Cytokine concentrations in the visceral adipose tissue from retroperitoneal area of the
253 rats in the *Chol* group were similar to those in *C* group. $\text{TNF}\alpha$ and IL-6 concentrations were
254 unchanged in all the experimental groups; however, $\text{IL-1}\beta$ concentrations were lower in *HT-*
255 *Ac* and *HT-Et* groups. Accordingly, MCP-1 showed significantly lower concentrations in
256 *HT-Ac* and *HT-Et* groups compared to *C* group (Table 4). PAI-1 concentrations showed
257 similar tendency, but differences did not reach statistical significance.

258

259 **4. Discussion**

260 In the present study the effects of HT were studied in comparison with *HT-Ac* and
261 *HT-Et*, two lipophilic derivatives of HT with a higher intestinal bioavailability than their
262 precursor^{15, 20}. The evaluation of the biological properties of these three compounds was
263 performed focusing on plasma lipids, energy homeostasis and oxidative and inflammatory

264 status in rats fed a cholesterol-rich diet. This study shows that HT and its hydrophobic
265 derivatives induce beneficial metabolic effects in hypercholesterolemic rats. Particularly,
266 HT-Ac showed the highest capacity to counteract the metabolic stress induced by the high-
267 cholesterol diet.

268 Supplementation of the diets with a moderate amount of phenolic compounds
269 (0.04%) resulted in a daily intake of 25mg/kg body weight, which is far beyond any possible
270 toxic level considering a recently published a study where consumption of up to 500
271 mg/kg/day was described as a No Observed Adverse Effects Level (NOAEL)³⁰.

272 As expected, feeding rats with a diet rich in cholesterol resulted in an increase in
273 total cholesterol and LDL cholesterol levels. In this hypercholesterolemic model, HT and
274 HT-Et showed modest plasma LDL-cholesterol lowering effects, in contrast to HT-Ac that
275 significantly reduced total and LDL-cholesterol levels compared to *Chol* group. These results
276 are in line with previous human and animal studies showing that phenolic compounds present
277 in virgin olive oil exert beneficial cardiovascular effects, particularly improving lipid profile⁸,
278 ^{10, 31}. In fact, the European Food Safety Authority has recently issued a positive opinion on
279 olive oil phenolic compounds capacity to protect LDL cholesterol from oxidative damage³².

280 Several markers show that rats in the present study suffered hepatic stress induced
281 by the cholesterol-rich diet, which may have led to an incipient insulin resistance².
282 Comparing the *Chol* and *C* groups, the high-cholesterol diet induced a significant increase in
283 glucose, creatinine, insulin and leptin plasma concentrations. In addition, *Chol*
284 hypercholesterolemic animals presented higher liver weights than *C* animals and the adipose
285 depots were mainly distributed in the retroperitoneal area. Interestingly, the diets
286 supplemented with phenolic compounds counteracted the glucose and insulin increase,
287 particularly HT and HT-Ac. This outcome is in accordance with previous studies showing
288 that HT and other olive oil phenolic compounds reduce plasma glucose concentration in

289 alloxan-diabetic rats⁸⁻⁹ by means of alleviating oxidative stress and free radicals as well as
290 enhancing enzymatic defences. A recent paper evidenced the capacity of oleuropein aglycone
291 in the prevention of cytotoxic amyloid aggregation of human amylin, a hallmark of Type-II
292 diabetes³³. In accordance, the PREDIMED study showed that consuming a virgin olive oil-
293 enriched traditional Mediterranean diet for three months decreased total and LDL cholesterol
294 together with plasma glucose in asymptomatic high cardiovascular-risk patients, although no
295 changes were observed in insulin levels³⁴. In line with these results, a significant decrease of
296 glycaemia related to a long term daily intake of a virgin olive oil rich in phenolic compounds
297 was observed in healthy young subjects³⁵. HT-Ac also maintained plasma leptin at
298 concentrations similar to those in the *C* group, while this effect was more discrete in the case
299 of HT and HT-Et. Similarly, in rats that consumed the diet supplemented with HT-Ac and
300 HT-Et creatinine concentrations were restored to control levels, but not with HT. The renal
301 protective effect of olive phenols reducing creatinine levels has already been described in
302 diabetic rats⁸.

303 The retroperitoneal/epididymal fat ratio provides information about adipose tissue
304 distribution, being directly proportional to the abdominal fat accumulation. Both
305 retroperitoneal and epididymal adipose depots are considered visceral fat associated with
306 higher levels of inflammation and lipolysis than subcutaneous fat, having been related with
307 insulin resistance³⁶. The retroperitoneal adipose depots in rats could be related to abdominal
308 adiposity in humans since they are located inside the peritoneum attached to the dorsal area
309 of the abdomen, whereas epididymal adipose tissue would correspond to gonadal fat, which
310 exists in mice and rats but not in humans. The groups supplemented with the lipophilic
311 compounds (*HT-Ac* and *HT-Et*) moderately improved the retroperitoneal/epididymal fat ratio
312 compared to *Chol* group, whereas the group that consumed HT did not show changes. Our
313 results are in agreement with the reduction in abdominal fat deposition described in a

314 metabolic syndrome animal model that consumed a diet supplemented with an olive leaf
315 extract rich in HT³⁷.

316 Metabolic disturbance which includes hyperlipidaemia, hyperglycaemia and
317 hyperinsulinemia often involves chronic inflammation and oxidative stress^{2, 8, 11}. Therefore,
318 redox status and inflammatory biomarkers were analyzed in plasma in order to understand
319 the potential mechanisms underlying the effects of the studied phenolic compounds.
320 Although the possible health implications related to MDA changes remain unknown, in this
321 study elevated systemic MDA concentrations were observed in the *Chol* group, suggesting a
322 marked oxidative stress in these animals that was counteracted by the consumption of the
323 phenolic compounds. Moreover, the serum antioxidant capacity of animals consuming the
324 phenolic compounds was higher than in the *Chol* group, particularly HT and HT-Ac.
325 However, at the hepatic level phenolic compounds supplementation did not decrease MDA
326 levels to control values although hepatic peroxidation was reduced in all cases, especially in
327 the *HT-Et* group (Table 4). It is noteworthy that the studied phenolic compounds follow
328 different metabolic pathways when ingested. HT-Ac is extensively hydrolysed into free HT,
329 whereas HT-Et remains unaltered, yielding more lipophilic metabolites than those generated
330 after HT and HT-Ac hepatic metabolism^{15, 20}. This would result in higher HT-Et
331 bioaccumulation in hepatic tissue, conferring higher protection against oxidation than HT and
332 HT-Ac metabolites.

333 The protective role of olive oil phenolic compounds against oxidative damage is
334 well established and has been reviewed recently¹. The antioxidant and free radical
335 scavenging capacity of olive oil phenols seem to be related to their anti-inflammatory effects,
336 which have been reported in different animal models^{38, 39}. Gong, Geng, Jiang, Cao,
337 Yoshimura&Zhong⁴⁰ showed the capacity of HT to decrease the pro-inflammatory cytokines
338 IL-1 β and TNF α but not to increase the anti-inflammatory cytokine IL-10 in carrageenan-

339 induced acute inflammation and hyperalgesia in rats. Moreover, the role of olive oil phenols
340 reducing postprandial inflammatory response in obese subjects has been recently attributed to
341 the inhibition of NF- κ B, which is an important link between oxidation and inflammation in
342 the postprandial state⁴¹. In the present study, diets supplemented with HT, HT-Ac and HT-Et
343 had anti-inflammatory effects decreasing plasma TNF α in rats fed cholesterol-rich diet.
344 Similar behaviour was observed with IL-1 β although not reaching statistical significance
345 probably due to the high variability that this parameter presented in the *Chol* group. In
346 adipose tissue, supplementation with HT-Ac and HT-Et decreased MCP-1 and IL-1 β below
347 control levels.

348 Nutrients' mechanisms of action are strongly related with the capacity to modulate
349 gene expression. Llorente-Cortes and co-workers³⁴ showed that consuming a traditional
350 Mediterranean diet enriched with virgin olive oil for three months improved lipid profile and
351 plasma glucose levels, prevented the increased expression of cyclooxygenase-2 (COX2) and
352 LDL receptor-related protein-1 (LRP-1) genes, and reduced the expression of MCP1 gene
353 compared with a traditional Mediterranean diet enriched with nuts or with a low fat diet.
354 COX2 and MCP1 genes are involved in inflammation whereas LRP1 takes part in foam cell
355 formation. Likewise, Konstantinidou and co-workers³¹ observed lower plasma oxidative and
356 inflammatory status in healthy subjects after consuming for three months a Mediterranean
357 diet supplemented with virgin olive oil rich in phenolic compounds compared to the control
358 group. In the study, pro-atherogenic genes related with inflammation (IFN γ , ARHGAP15 and
359 IL7R) and oxidative stress (ADRB2) were down-regulated in peripheral blood mononuclear
360 cells (PBMCs). Changes in gene expression were associated with decreases in lipid oxidative
361 damage and systemic inflammation markers. In vitro studies have established the capacity of
362 HT to modulate adipocyte lipid content and gene expression partially mediated by the
363 reduction of the transcription factors PPAR α and C/EBP α ⁴². In phytochemicals' capacity to

364 modify gene expression in different tissues and cell lines, the chemical structure of the
365 phytochemical plays an important role not only in their direct scavenging of free radicals, but
366 also in the target protein and therefore in the molecular mechanism involved in its biological
367 action. Thus, considering the extensive hydrolysis that HT-Ac undergoes during intestinal
368 absorption yielding HT¹⁵, both compounds could follow similar mechanisms to regulate the
369 unbalance induced by the high-cholesterol rich diet. Our results suggest changes in the
370 expression of redox and inflammatory related genes, based on studies recently published by
371 other research groups; however, further work is required to elucidate the molecular
372 mechanisms underlying the regulatory metabolic effect of HT and its derivatives.

373 A limitation of this study is that a control group fed a standard diet supplemented
374 with HT was not included to assess the effect of this compound within a balanced diet in
375 healthy animals.

376 In summary, HT and its lipophilic derivatives, HT-Ac and HT-Et, were able to
377 reduce the metabolic unbalance induced by a high-cholesterol diet in rats, being HT-Ac, the
378 lipophilic HT derivative naturally present in virgin olive oil, the most effective phenolic
379 compound. Since HT-Ac can be easily obtained from natural HT¹², this compound might be
380 proposed as an interesting bioactive ingredient in the production of functional foods, having
381 the added value of contributing to induce beneficial metabolic properties.

382

383 **List of abbreviations:** **C group:** Control group ; **Chol group:** cholesterol-rich diet group;
384 **AAPH:** 2,2'-azobis (2-amidinopropane) dihydrochloride; **COX 2:** cyclooxygenase-2; **FFA:**
385 Free fatty acids; **HT:** hydroxytyrosol; **HT-Ac:** hydroxytyrosyl acetate; **HT-Et:** ethyl
386 hydroxytyrosyl ether; **IL-1b:** interleukin-1 β ; **IL-6:** interleukin-6; **LRP1:** LDL receptor-
387 related protein-1; **MCP-1:** monocyte chemoattractant protein-1; **ORAC:** oxygen radical
388 scavenging capacity; **PAI-1:** plasminogen activator inhibitor-1; **SD:** standard deviation;
389 **TNF- α :** tumour necrosis factor- α ; **Trolox:** 6-hydroxy-2,5,7,8-tetramethylchroman-2-
390 carboxylic acid

391

392 **Acknowledgements**

393 Study funded by projects RTA2007-00036 (INIA), Excellence Project AGR-5098
394 (Junta de Andalucia) and Consolider-Ingenio (CSD2007-00063). The authors thank the
395 technical support and expertise in blood analysis provided by CORE lab (University Hospital
396 La Paz).SM-L is a JAE predoctoral fellow funded by CSIC and the European Social Fund.

397

398

399 **Conflict of interest statement**

400 The authors have declared no conflict of interest.

401

402

403

404

405

406

407 **References**

- 408 1. M. I. Fernandez-Mar, R. Mateos, M. C. Garcia-Parrilla, B. Puertas, and E. Cantos-
409 Villar, *Food Chem.*, 2012, **130**, 797-813.
- 410 2. A. Dey and J. Lakshmanan, *Food Funct.*, 2013, **4**, 1148-1184.
- 411 3. K. De la Torre-Carbot, J. L. Chávez-Servín, O. Jaúregui, A. I. Castellote, R. M.
412 Lamuela-Raventós, T. Nurmi, H. E. Poulsen, A. V. Gaddi, J. Kaikkonen, H. F.
413 Zunft, H. Kieseewetter, M. Fitó, M. I. Covas, and M. C. López-Sabater. *J. Nutr.*,
414 2010, **140**, 501-508.
- 415 4. M. Gonzalez-Santiago, J. Fonolla, and E. Lopez-Huertas. *Pharm. Res.*, 2010, **61**,
416 364-370.
- 417 5. J. Ruano, J. López-Miranda, F. Fuentes, J. A. Moreno, C. Bellido, P. Perez-
418 Martinez, A. Lozano, P. Gómez, Y. Jiménez, and F. Pérez Jiménez, *J. Am. Coll.*
419 *Cardiol.* 2005, **46**, 1864–1868.
- 420 6. P. Bogani, C. Galli, M. Villa and F. Visioli, *Atherosclerosis.* 2007, **190**, 181–186.
- 421 7. J. Ruano, J. López-Miranda, R. de la Torre, J. Delgado-Lista, J.Fernández, J.
422 Caballero, M. I. Covas, Y. Jiménez, P. Pérez-Martínez, C. Marín, F. Fuentes and F.
423 Pérez-Jiménez. *Am. J. Clin. Nutr.* 2007, **86**, 341–346.
- 424 8. K. Hamden, N. Allouche, M.Damak, and A. Elfeki, *Chem. Biol. Interact.*, 2009,
425 **180**, 421–432.
- 426 9. H. Jemai, A. E. Feki, and S. Sayadi, *J. Agric. Food Chem.* 2009, **57**, 8798–8804.
- 427 10. G. Rodriguez-Gutierrez, G.G. Duthie, S. Wood, P. Morrice, F. Nicol, M. Reid, L.L.
428 Cantlay, T. Kelder, G.W. Horgan, J.F.B. Guzman, and B. de Roos, B. *Mol. Nutr.*
429 *Food Res.* 2012, **56**, 1131-1147.
- 430 11. H. Jemai, I. Fki, M. Bouaziz, Z. Bouallagui, A.E. Feki, H. Isoda, and S.Sayadi, *J.*
431 *Agric. Food Chem.* 2008, **56**, 2630–2636.
- 432 12. F. Alcudia, A. Cert, J.L. Espartero, R. Mateos, and M. Trujillo, PCT
433 WO2004/005237. (2004).

- 434 13. A. Madrona, G. Pereira-Caro, R. Mateos, G. Rodriguez, M. Trujillo, J. Fernandez-
435 Bolanos, and J.L. Espartero, *Molecule*, 2009, **14**, 1762–1772.
- 436 14. R. Mateos, J.L. Espartero, M. Trujillo, J.J. Ríos, M. León-Camacho, F. Alcudia, A.
437 Cert, *J. Agric. Food Chem.* 2001, **49**, 2185–2192.
- 438 15. R. Mateos, G. Pereira-Caro, S. Saha, R. Cert, M. Redondo-Horcajo, L. Bravo and
439 P.A. Kroon, *Food Chem.* 2011, **125**, 865–872.
- 440 16. R. Mateos, L. Goya, and L. Bravo, *J. Agric. Food Chem.* 2005, **53**, 9897–9905.
- 441 17. S. Grasso, L. Siracusa, C. Spatafora, M. Renis and C. Tringali, *Bioorg. Chem.* 2007,
442 **35**, 137–152.
- 443 18. Z. Bouallagui, M. Bouaziz, S. Lassoued, J.M. Engasser, M. Ghoul and S. Sayadi,
444 *Appl. Biochem. Biotechnol.* 2011, **163**, 592–599.
- 445 19. G. Pereira-Caro, R. Mateos, B. Sarria, R. Cert, L. Goya and L. Bravo, *Food Chem.*
446 2012, **131**, 869–878.
- 447 20. G. Pereira-Caro, R. Mateos, S. Saha, A. Madrona, J.L. Espartero, L. Bravo, L. and
448 P.A. Kroon, *J. Agric. Food Chem.* 2010, **58**, 11501–11509.
- 449 21. G. Pereira-Caro, L. Bravo, A. Madrona, J.L. Espartero and R. Mateos, *J. Agric.*
450 *Food Chem.* 2010, **58**, 798–806.
- 451 22. G. Pereira-Caro, B. Sarria, A. Madrona, J.L. Espartero, L. Goya, L. Bravo and R.
452 Mateos, *J. Agric. Food Chem.* 2011, **59**, 5964–5976.
- 453 23. J. Munoz-Marin, J.P. De La Cruz, J.J. Reyes, J.A. Lopez-Villodres, A. Guerrero, I.
454 Lopez-Leiva, J.L. Espartero, M.T. Labajos, and J.A. Gonzalez-Correa, *Food Chem.*
455 *Toxicol.*, 2013, **58**, 295–300.
- 456 24. G. Pereira-Caro, A. Madrona, L. Bravo, J.L. Espartero, F. Alcudia, A. Cert, and R.
457 Mateos, *Food Chem.* 2009, **115**, 86–91.
- 458 25. R. Mateos, M. Trujillo, G. Pereira-Caro, A. Madrona, A. Cert, and J.L. Espartero, *J.*
459 *Agric. Food Chem.* 2008, **56**, 10960–10966.

- 460 26. J. Fernandez-Bolaños, A. Heredia, G. Rodríguez, R. Rodríguez, A. Jiménez and R.
461 Guillen, US 6,849,770 B2. 2005.
- 462 27. S.G. Sayago-Ayerdi, R. Mateos, R.I. Ortiz-Basurto, C. Largo, J. Serrano, A.B.
463 Granado-Serrano, B. Sarria, L. Bravo, M. Tabernero, *Food Chem.* 2014, **148**, 54-59.
- 464 28. D. Huang, B. Ou, M. Hampsch-Woodill, J.A. Flanagan and R.L. Prior, *J. Agric.*
465 *Food Chem.* 2002, **50**, 4437-4444.
- 466 29. R. Mateos, E. Lecumberri, S. Ramos, L. Goya and L. Bravo, *J. Chromatogr. B.*
467 *Analyt. Technol. Biomed. Life Sci.* 2005, **827**, 76–82.
- 468 30. D. Aunon-Calles, L. Canut, F. Visioli, *Food Chem. Toxicol.* 2013, **55**, 498-504.
- 469 31. V. Konstantinidou, M.I. Covas, D. Munoz-Aguayo, O. Khymenets, R. De la Torre,
470 G. Saez, M.C. Tormos, E. Toledo, A. Marti, V. Ruiz-Gutiérrez, M.V. Ruiz-Mendez
471 and M. Fitó, *Faseb. J.* 2010, **24**, 2546-2557.
- 472 32. EFSA Panel on Dietetic Products, Nutrition and Allergies, *EFSA Journal*, 2011, **9**,
473 2033-2058.
- 474 33. S. Rigacci, V. Guidotti, M. Bucciattini, M. Parri, C. Nediani, E. Cerbai, M. Stefani,
475 and A. Berti, *J. Nutr. Biochem.* 2010, **21**, 726-735.
- 476 34. V. Llorente-Cortes, R. Estruch, M.P. Mena, E. Ros, M.A. González, M. Fitó, R.M.
477 Lamuela-Raventós, and L. Badimon, *Atherosclerosis* 2010, **208**, 442–450.
- 478 35. M.J. Oliveras-Lopez, M. Innocenti, F. Martín-Bermudo, H. Lopez-Garcia de la
479 Serrana and N. Mulinacci, *Eur. J. LipidSci. Technol.* 2012, **114**, 999-1006.
- 480 36. S. Yamashita, T. Nakamura, I. Shimomura, M. Nishida, S. Yoshida, K. Kotani, K.
481 Kameda-Takemuara, K. Tokunaga and Y. Matsuzawa, *Diabetes Care*, 1996, **19**,
482 287-291.
- 483 37. H. Poudyal, F. Campbell and L. Brown, *J. Nutr.* 2010, **140**, 946-953.
- 484 38. R. De la Puerta, E. Martínez-Domínguez and V Ruiz-Gutiérrez, *Z Naturforsch C.*
485 2000, **55**, 814–819.

- 486 39. C. Puel, J. Mardon, A. Agalias, M.J. Davicco, P. Lebecque, A. Mazur, M.N.
487 Horcajada, A.L. Skaltsounis and V. Coxam, *J Agric Food Chem.* 2008, **56**, 9417–
488 9422.
- 489 40. D. Gong, C. Geng, L. Jiang, J. Cao, H. Yoshimura, and L.E. *Phytother. Res.* 2009,
490 **23**, 646–650.
- 491 41. A. Perez-Herrera, J. Delgado-Lista, L.A. Torres-Sanchez, O.A., Rangel-Zuñiga, A.
492 Camargo, J.M. Moreno-Navarrete, B. Garcia-Olid, G.M. Quintana-Navarro, J.F.
493 Alcala-Diaz, C. Muñoz-Lopez, F. Lopez-Segura, J.M. Fernandez-Real, M.D. Luque
494 de Castro, J. Lopez-Miranda and F. Perez-Jimenez, *Mol. Nutr. Food Res.* 2012, **56**,
495 510-514.
- 496 42. I. Warnke, R. Goralczyk, E. Fuhrer and J. Schwager, *Nutr. Metab.* 2011, **8**, 1-13.
497
498
499
500
501
502
503
504
505
506
507
508

509

510 **Figure Legends**

511

512 **Figure 1.** Plasma lipid profile (total-cholesterol, LDL- and HDL-Chol) of rats fed for 8
513 weeks with standard diet (*C* group), cholesterol-rich diet (*Chol* group) and cholesterol-rich
514 diet supplemented with hydroxytyrosol (*HT* group), hydroxytyrosyl acetate (*HT-Ac*) and
515 ethyl hydroxytyrosyl ether (*HT-Et*). Data represent the means of 8 determination \pm SD. Bars
516 without a common letter differ, $p < 0.05$.

517

518

Figure 1.




Table 1. Dietary composition of the experimental diets.

Component (g/100g dry weight)	<i>C</i>	<i>Chol</i>	<i>HT</i>	<i>HT-Ac</i>	<i>HT-Et</i>
Carbohydrates	65.93	64.20	64.17	64.17	64.17
Protein	17.58	17.12	17.11	17.11	17.11
Lipids (others than cholesterol)	3.30	3.21	3.21	3.21	3.21
Cholesterol	0.00	2.19	2.19	2.19	2.19
Cholic acid	0.00	0.44	0.44	0.44	0.44
HT and derivatives	0.00	0.00	0.04	0.04	0.04
Caloric content (Kcal/100g)	219.2	227.7	227.7	227.7	227.7

1 **Table 2.** Body and tissue weights of rats in the control group (*C*), the group consuming the non-supplemented cholesterol-rich diet (*Chol*) or the
 2 cholesterol-rich diet supplemented with hydroxytyrosol (*HT*), hydroxytyrosyl acetate (*HT-Ac*) and ethyl hydroxytyrosyl ether (*HT-Et*). Data
 3 represents the mean of 8 determinations \pm SD*.

	<i>Standard diet</i>		<i>Hypercholesterolemic diet</i>		
	<i>C</i>	<i>Chol</i>	<i>HT</i>	<i>HT-Ac</i>	<i>HT-Et</i>
Body weight (g)					
Initial	242 + 6	244 + 8	244 + 7	239 + 11	243 + 4
Final	336 + 18	326 + 30	331 + 16	317 + 10	329 + 14
Body weight gain (g/56d)	94 + 15	82 + 32	87 + 12	78 + 11	86 + 12
Tissue weights (g)					
Epididymal fat/ body weight	0.016 + 0.003	0.016 + 0.002	0.015 + 0.001	0.017 + 0.003	0.015 + 0.004
Retroperitoneal fat/ body weight	0.018 + 0.004	0.015 + 0.002	0.014 + 0.002	0.017 + 0.001	0.016 + 0.005
Ratio Retroperitoneal/epididymal	0.88 + 0.09 ^a	1.07 + 0.08 ^b	1.07 + 0.08 ^b	1.00 + 0.05 ^{ab}	0.93 + 0.07 ^{ab}
Liver / body weight	0.024 + 0.002 ^a	0.038 + 0.003 ^b	0.029 + 0.018 ^{ab}	0.036 + 0.002 ^b	0.037 + 0.002 ^b
Kidney / body weight	0.0053 + 0.0022	0.0057 + 0.0005	0.0058 + 0.0008	0.0061 + 0.0005	0.0059 + 0.0006
Heart / body weight	0.0026 + 0.0003	0.0028 + 0.0005	0.0025 + 0.0001	0.0029 + 0.0005	0.0028 + 0.0003

4 * Mean values in a row with unlike superscripts are significantly different according to Kruskal-Wallis one-way analysis of variance and Mann-Whitney
 5 U test ($p < 0.05$).

6

7

8 **Table 3.** Metabolic biomarkers in plasma and adipose tissue of rats in the control group (*C*), the group consuming the un-supplemented
 9 cholesterol-rich diet (*Chol*) or the group fed the cholesterol-rich diet supplemented with hydroxytyrosol (*HT*), hydroxytyrosyl acetate (*HT-Ac*)
 10 and ethyl hydroxytyrosyl ether (*HT-Et*). Data represents the means of eight determinations \pm SD*.
 11

Plasma nutrients		Standard diet	Hypercholesterolemic diet			
		<i>C</i>	<i>Chol</i>	<i>HT</i>	<i>HT-Ac</i>	<i>HT-Et</i>
Glucose (mg/dL)		140.50 \pm 15.23 ^a	182.29 \pm 18.17 ^b	143.25 \pm 19.83 ^a	154.75 \pm 23.36 ^a	160.00 \pm 22.53 ^{ab}
Triglycerides (mg/dL)		106.71 \pm 21.69	100.29 \pm 18.33	83.00 \pm 29.50	90.50 \pm 19.49	84.57 \pm 13.90
Free Fatty Acids (mM equivalents palmitic acid)		0.33 \pm 0.07	0.30 \pm 0.13	0.34 \pm 0.07	0.31 \pm 0.08	0.36 \pm 0.07
Creatinine (mg/dL)		0.52 \pm 0.02 ^a	0.58 \pm 0.05 ^b	0.55 \pm 0.02 ^b	0.53 \pm 0.03 ^a	0.53 \pm 0.03 ^a
Hormones						
Plasma	Insulin (ng/mL)	972 \pm 341 ^a	1560.2 \pm 997.9 ^b	934.1 \pm 390 ^a	608.1 \pm 301.6 ^a	1121.3 \pm 617.1 ^{ab}
	Leptin (ng/mL)	2786 \pm 908 ^a	4174 \pm 1116 ^b	3609 \pm 1022 ^{ab}	2970 \pm 1150 ^a	3468.1 \pm 1291 ^{ab}
Adipose tissue	Leptin (ng/ μ g pr)	682 \pm 189	501 \pm 178	503 \pm 137	503 \pm 148	671 \pm 283
	Adiponectin (ng/ μ g pr)	726415 \pm 139506	658737 \pm 145887	593934 \pm 67860	693282 \pm 139463	653668 \pm 66473

12
 13 * Mean values in a row with unlike superscripts are significantly different according to Kruskal-Wallis one-way analysis of variance and Mann-Whitney
 14 U test ($p < 0.05$).

15
 16

17 **Table 4.** Redox status and inflammatory biomarkers in plasma and adipose tissue of rats fed the control diet (*C*), the un-supplemented
 18 cholesterol-rich diet (*Chol*) and the cholesterol-rich diet supplemented with hydroxytyrosol (*HT*), hydroxytyrosyl acetate (*HT-Ac*) and ethyl
 19 hydroxytyrosyl ether (*HT-Et*). Data represents the mean of 8 determinations \pm SD*.
 20

			<i>Standard diet</i>	<i>Hypercholesterolemic diet</i>			
			<i>C</i>	<i>Chol</i>	<i>HT</i>	<i>HT-Ac</i>	<i>HT-Et</i>
Antioxidant capacity	Serum ORAC (mM T.eq)		27.4 \pm 1.8 ^{ab}	25.6 \pm 2.3 ^a	28.4 \pm 1.6 ^b	28.0 \pm 1.2 ^b	26.2 \pm 1.1 ^a
Lipid oxidation	Serum (nmol/mL)		1.97 \pm 0.75 ^a	3.54 \pm 0.24 ^b	2.36 \pm 0.23 ^a	2.19 \pm 0.39 ^a	2.36 \pm 0.30 ^a
	Liver (nmol/mg pr)		0.98 \pm 0.33 ^a	3.33 \pm 1.14 ^b	2.52 \pm 0.92 ^{bc}	2.35 \pm 0.81 ^{bc}	2.11 \pm 0.53 ^c
Inflammatory mediators	Plasma	TNF α (ng/mL)	3.0 \pm 0.7 ^a	4.2 \pm 0.7 ^b	3.7 \pm 0.8 ^a	2.7 \pm 0.6 ^a	3.4 \pm 0.7 ^{ab}
		IL-1 β (ng/mL)	11.6 \pm 6.3	26.50 \pm 21.5	13.8 \pm 14.6	11.5 \pm 7.1	8.0 \pm 4.9
		IL-6 (ng/mL)	4.9 \pm 4.4	3.66 \pm 1.63	6.0 \pm 3.3	5.7 \pm 3.7	5.9 \pm 2.4
	Adipose tissue	TNF α (ng/ μ g pr)	2.5 \pm 0.6	2.2 \pm 0.5	2.2 \pm 0.5	2.0 \pm 0.4	1.9 \pm 0.3
		IL-1 β (ng/ μ g pr)	3.9 \pm 0.9 ^a	3.7 \pm 0.8 ^a	3.8 \pm 0.6 ^a	2.9 \pm 0.4 ^b	3.1 \pm 0.2 ^b
		IL-6 (ng/ μ g pr)	24.6 \pm 7.0	23.7 \pm 6.1	25.7 \pm 7.4	20 \pm 8	23.4 \pm 6.5
		MCP-1 (ng/ μ g pr)	32.3 \pm 7.4 ^a	27.6 \pm 3.9 ^a	28.5 \pm 3.9 ^a	26 \pm 4 ^b	25.6 \pm 3.4 ^b
		PAI-1 (ng/ μ g pr)	87.6 \pm 22.9	98.9 \pm 32.9	92.4 \pm 15.4	73.3 \pm 18.0	81.5 \pm 12.4

21

22 * Mean values in a row with unlike superscripts are significantly different according to Kruskal-Wallis one-way analysis of variance and Mann-Whitney

23 U test ($p < 0.05$).

