



**Probing the substrate specificity of *Trypanosoma brucei*
GlcNAc-PI de-N-acetylase with synthetic substrate
analogues.**

Journal:	<i>Organic & Biomolecular Chemistry</i>
Manuscript ID:	OB-ART-10-2013-042164.R1
Article Type:	Paper
Date Submitted by the Author:	30-Jan-2014
Complete List of Authors:	Capes, Amy; University of Dundee, BCDD Crossman, Arthur; University of Dundee, BCDD Urbaniak, Michael; University of Dundee, BCDD Gilbert, Sophie; University of Dundee, BCDD Ferguson, Michael; University of Dundee, BCDD Gilbert, Ian; University of Dundee, BCDD

Probing the substrate specificity of *Trypanosoma brucei* GlcNAc-PI de-*N*-acetylase with synthetic substrate analogues.

Amy S. Capes,[†] Arthur Crossman,[†] Michael D. Urbaniak, Sophie H. Gilbert, Michael A. J. Ferguson,^{*} Ian H. Gilbert^{*}

Division of Biological Chemistry and Drug Discovery, College of Life Sciences, University of Dundee, Dow Street, Dundee, DD1 5EH, UK.

[†]These authors contributed equally to the work.


^{*}Co-corresponding authors. Phone +44 (0) 1382 386240, Email m.a.j.ferguson@dundee.ac.uk and i.h.gilbert@dundee.ac.uk

Table of Contents Entry

Probing the substrate specificity of *Trypanosoma brucei* GlcNAc-PI de-*N*-acetylase with synthetic substrate analogues.

Amy S. Capes, Arthur Crossman, Michael D. Urbaniak, Sophie H. Gilbert, Michael A. J. Ferguson,
Ian H. Gilbert

A series of synthetic analogues of 1-D-(2-amino-2-deoxy- α -D-glucopyranosyl)-*myo*-inositol 1-(1,2-di-*O*-hexadecanoyl-*sn*-glycerol 3-phosphate), consisting of 7 variants of either the D-*myo*-inositol, D-GlcpN or the phospholipid components, were prepared and following *N*-acylation were tested as substrates and inhibitors of GlcNAc-PI de-*N*-acetylase, a genetically validated drug target in *Trypanosoma brucei*, the causative agent of human African trypanosomiasis.



Abstract

A series of synthetic analogues of 1-D-(2-amino-2-deoxy- α -D-glucopyranosyl)-*myo*-inositol 1-(1,2-di-*O*-hexadecanoyl-*sn*-glycerol 3-phosphate), consisting of 7 variants of either the D-*myo*-inositol, D-GlcpN or the phospholipid components, were prepared and tested as substrates and inhibitors of GlcNAc-PI de-*N*-acetylase, a genetically validated drug target enzyme responsible for the second step in the glycosylphosphatidylinositol (GPI) biosynthetic pathway of *Trypanosoma brucei*. The D-*myo*-inositol in the physiological substrate was successfully replaced by cyclohexanediol and is still a substrate for *T. brucei* GlcNAc-PI de-*N*-acetylase. However, this compound became sensitive to the stereochemistry of the glycoside linkage (the β -anomer was neither substrate or inhibitor) and the structure of the lipid moiety (the hexadecyl derivatives were inhibitors). Chemistry was successfully developed to replace the phosphate with a sulphonamide, but the compound was neither a substrate or an inhibitor, confirming the importance of the phosphate for molecular recognition. We also replaced the glucosamine by an acyclic analogue, but this also was inactive, both as a substrate and inhibitor. These findings add significantly to our understanding of substrate and inhibitor binding to the GlcNAc-PI de-*N*-acetylase enzyme and will have a bearing on the design of future inhibitors.

Introduction

The enzymes of the glycosylphosphatidylinositol (GPI) biosynthetic pathway are located in the endoplasmic reticulum, contain between one and thirteen predicted trans-membrane domains and are mostly present as components of multi-subunit complexes.¹ No high-resolution structural data are available on any of these enzymes and our research group has been probing the specificities of several of the enzymes in the GPI pathway of the protozoan parasite *Trypanosoma brucei*, the causative agent of African sleeping sickness in humans and the related disease Nagana in cattle, using synthetic substrate analogues *in vitro*.²⁻⁸ One of the key enzymes of interest is an amidase, the GlcNAc-PI de-*N*-acetylase (EC [3.5.1.89](#)) that de-*N*-acetylates 1-D-(2-acetamido-2-deoxy- α -D-glucopyranosyl)-*myo*-inositol 1-(1,2-di-*O*-hexadecanoyl-*sn*-glycerol 3-phosphate) (**1**, α -D-Glc_pNAc-PI) to 1-D-(2-amino-2-deoxy- α -D-glucopyranosyl)-*myo*-inositol 1-(1,2-di-*O*-hexadecanoyl-*sn*-glycerol 3-phosphate) (**2**, α -D-Glc_pN-PI), Figure 1.




Figure 1 Some previously prepared GPI analogues

This enzyme catalyses the second step in the *T. brucei* GPI biosynthetic pathway, which is a prerequisite for all subsequent steps in the pathway.⁹ In earlier studies, we showed that *T. brucei* GlcNAc-PI de-*N*-acetylase is a zinc-dependent metalloenzyme¹⁰ and demonstrated, by construction of a condition-null mutant cell line, that it is essential for the bloodstream form of the parasite and, therefore, a genetically validated drug target.¹¹ Previous studies with other substrate analogues showed that the phosphate, 2'-NHAc and 3'-OH groups of the natural substrate α -D-Glc_pNAc-PI (**1**) are critical for recognition by the *T. brucei* GlcNAc-PI de-*N*-acetylase.²⁻⁴ In contrast, the

diacylglycerol moiety is not strictly required and may be efficiently replaced with an octadecyl chain,⁴ as shown in analogues **3** and **4**. In the case of the *T. brucei* enzyme, we had hypothesised that one or more of the inositol 2, 3, 4 and 5-OH groups is/are not required.²⁻⁴ We came to this hypothesis from the ability of the enzyme to recognise and process both α -D-GlcpNAc-[L]-PI (**5**) and β -D-GlcpNAc-PI (**6**). Molecular dynamics simulations, showed that α and β anomers can adopt conformations in which the phosphate, the 2'-amide and the 3'-OH overlay. Given the 2' amide is where the reaction occurs, and evidence suggests that the 3'-OH and phosphate are important for recognition of the substrate with the enzyme, these conformations are likely to be the active enzyme-bound conformations. In these conformations the inositol 2, 3, 4 and 5- hydroxyls are in different orientations for the two α and β anomers, implying the hydroxyls are not critical for interaction with the enzyme. Further evidence was obtained from a compound in which the inositol 2-hydroxyl was alkylated. This was also a substrate for the *T. brucei* enzyme. One of the goals of the study described in this paper was to investigate the hypothesis that the inositol 2, 3, 4 and 5-OH groups are not required.

Bearing in mind these key structural features, we have synthesised a variety of analogues to further probe the requirements for substrate recognition by the *T. brucei* GlcNAc-PI de-*N*-acetylase and, specifically, to test:

1. The hypothesis that the inositol 2, 3, 4 and 5-OH groups are not required for enzyme recognition, a series of pseudodisaccharides (**7** – **10**, Figure 2) containing a cyclohexanediol moiety in place of the inositol aglycone were prepared.
2. Whether the phosphate group can be replaced by more cell-permeable sulphonamide isosteres,¹² compounds **11** and **12** (Figure 2) were prepared.
3. Whether, given the essentiality of the 2'-NHR and 3'-OH groups but non-essentiality of the 4'- and 6'-OH groups for substrate recognition, the glucosamine residue might be simplified to a simple acyclic structure, as in compound **13**.




Figure 2 Target molecules


The *N*-acetylated derivatives of the above analogues required for biological studies with the de-*N*-acetylase were prepared from the corresponding amines by standard procedures.¹³ All the analogues were examined for their recognition and processing by the *T. brucei* GlcNAc-PI de-*N*-acetylase.

Results and discussion

Synthesis of analogues 7 – 13

The synthesis of the required α and β -glucosaminy (1'→1) cyclohexandiol building blocks **16** and **17**, respectively, began by reacting the known¹⁴ trichloroacetimidate **14** and the commercially available 1*R*,2*R*-*trans*-cyclohexandiol **15**, Scheme 1, with a catalytic amount of trimethylsilyl trifluoromethanesulfonate (TMSOTf). The separation of anomers was achievable at this stage and these anomers were vital in providing the glucosamine-phosphodiester target analogues discussed herein. For the sake of brevity, we have chosen to describe in the main text the formation of the α -anomers, while details for the corresponding β -anomers appears in the Supplementary Data Section. Therefore, the pseudodisaccharide **16** was coupled to the hydrogen

phosphonate **18**¹⁵, and the ensuing mixture of diastereoisomeric phosphonic diesters, was oxidised with iodine in pyridine–water¹⁶ to give the corresponding phosphodiester **19**. The diester was transformed into the triol **20** after conventional *O*-deacylation of the latter compound with 0.05 M methanolic NaOMe in CH₂Cl₂–MeOH.




Scheme 1 Synthesis of **7**

Our first attempt at hydrogenolysis of the triethylammonium (TEA) salt **20** over Pd(OH)₂/C gave, surprisingly, the alkyl alcohol secondary amine **21**. Initially, the azide of **20** was reduced to the primary amine, however, if there are peroxides present in the THF then the formation of THF hydroperoxide is, apparently, possible which could then lead to an amine-THF coupling *via* a free-radical-based mechanism.¹⁷ This intermediate is susceptible to a Pd-mediated THF ring opening reaction that gives an imine which is then further hydrogenated to an aminobutanol.¹⁷ Consequently, after purchasing a fresh bottle of anhydrous stabilised THF, the second hydrogenolysis attempt at **20** → **7** proceeded without incident.

The preparation of the dipalmitoyl glycerol pseudodisaccharide **9** was accomplished from the triacetate **16**, Scheme 2. However, the acetate protecting groups in **16** are unsuitable because if they were left in place and removed by base at the final step of the synthesis, then those requisite


esters of the lipid fragment would likewise be saponified. Therefore, the acetates of **16** needed to be swapped to a more appropriate protecting group but first, the temporary *tert*-butyldimethylsilyl (TBDMS) protection of the 2-OH, **16** \rightarrow **22**, was performed and then followed by conventional *O*-deacylation, as previously described for **19** \rightarrow **20**, furnished the triol **23**.



Scheme 2 Synthesis of **9**

The benzyl group was chosen as the 3', 4' and 6'-OH protecting group because, from our past experiences synthesising GPI analogues, the benzyl group has been a very reliable protecting group *via* ease of installation and removal. Thus, the triol **23** was benzylated with benzyl bromide in the presence of NaH, as the base, to afford compound **24**. We next turned our attention towards the reduction of the azide in **24** and because of the issues with Pd(OH)₂ catalysed reduction in the presence of peroxidic THF discussed earlier, we chose to reduce the azide *via* the Staudinger reaction¹⁸ to give the amine **25** which was subsequently *tert*-butyl carbamate (Boc) protected to furnish **26**. Desilylation of **26** using HF-pyridine conditions afforded the alcohol **27**. The known hydrogen phosphonate **28**¹⁹ was coupled, as already described, to the 2-OH of **27** which, after oxidation, was isolated and characterised as the TEA salt **29**. Hydrogenolysis of **29** over Pd(OH)₂/C gave the Boc protected derivative **30** and subsequent cleavage of the Boc group produced the deprotected target analogue **9**.


Sulphonamides are potential isosteres for the phosphate group; they have the same tetrahedral shape and polar oxygen atoms. The synthesis of the sulphonamides **11** and **12**, Scheme 3, was accomplished by reacting commercially available 1*R*,2*R*-1-amino-2-benzyloxycyclohexane **31** with 1-octadecanesulfonyl chloride in the presence of triethylamine and CH₂Cl₂ to give the benzyl derivative **32**. The benzyl group was removed by hydrogenolysis over Pd(OH)₂/C to furnish the alcohol **33**. Coupling of **33** with the known trichloroacetimidate **34**²⁰ resulted in an inseparable mixture of the α,β-anomers **35** in the ratio of ~1:1, as determined by ¹H NMR spectroscopy. Finally, hydrogenolysis of the aforementioned anomers **35** over Pd(OH)₂/C and subsequent silica gel column chromatography (9:1 CH₂Cl₂ – MeOH) gave first the β-anomer **11** and then the α-anomer **12**.



Scheme 3 Synthesis of sulphonamides **11** and **12**

We were also interested in seeing if we could replace the glucose ring with an acyclic moiety. From our knowledge of the SAR, retaining the 2-amino and 3-hydroxy groups are important for activity. The synthesis of the amino-phosphate **13**, Scheme 4, began by epoxide ring-opening of cyclohexene oxide **36** with *p*-methoxybenzyl alcohol (PMBOH), using Cu(BF₄)₂·*n*H₂O as a catalyst²¹ to give the known racemic PMB monoprotected cyclohexanediol²² contaminated with unreacted PMBOH. Chromatographic separation of this PMB cyclohexyl derivative from the excess of PMBOH was not achievable, in our hands, and so the entire PMB reaction residue was

acetylated with acetic anhydride in the presence of pyridine and a catalytic amount of 4-(dimethylamino)pyridine DMAP to furnish the acetate **37**, which was easily separated from the acetate of *p*-methoxybenzyl alcohol by silica gel column chromatography.²³ After deacetylation, the resulting alcohol²² residue was alkylated using sodium hydride and allyl bromide to give the allyl derivative **38**. The epoxide **39** was prepared upon reacting **38** with 3-chloroperbenzoic acid (*m*CPBA), and the subsequent hydrolysis of epoxide **39** with DMSO, H₂O and a catalytic amount of KOH,²⁴ worked smoothly to furnish diol **40**. The primary alcohol of **40** was protected using *tert*-butyl(chloro)diphenylsilane (TBDPSCI) and DMAP to give compound **41**. The azido group of **43** was satisfactorily installed (51% yield over two steps) *via* the mesylate **42** obtained by reacting the secondary alcohol **41** with methanesulfonyl chloride in the presence of pyridine, followed by treatment of **42** with sodium azide under forcing conditions. The crude mesylate **42** was used directly in the displacement reaction but a small portion of **42** was purified for a full characterisation of this intermediate. The *p*-methoxybenzyl protecting group of **43** was removed with mild acid to give alcohol **44** and then it was phosphorylated, as previously described, to give the phosphoric diester **45**. Thereafter, the removal of the silyl protecting group of **45** with 1.0 M tetrabutylammonium fluoride (TBAF) in THF proceeded smoothly to give **46** which, after hydrogenolysis over Pd(OH)₂/C, provided the TEA salt **13**.

Scheme 4 Synthesis of racemic **13**

Biological results

Substrate analogues:

The ability of the *T. brucei* GlcNAc-PI de-*N*-acetylase to recognise and process the synthetic pseudodisaccharides **7** – **13** was tested in *T. brucei* cell-free system using an LC-MS/MS assay.^{8,10} The *N*-acetylated analogues of these compounds were prepared as previously described to give compounds **47** – **53** (Figure 3).¹³




Figure 3 *N*-Acetylated analogues

Using LC-MS/MS, multiple reaction monitoring of characteristic transitions for the *N*-acetylated and corresponding amine form of each compound was used to directly measure the rate of conversion of the *N*-acetylated compound to the free amine (Table 1). As no suitable transition was identified for the amine form of **13**, the enzymatic turnover was accessed by reacting any free amine formed with *d*₆-Ac₂O, and measuring the formation of the *d*₃-*N*-acetylated form by LC-MS/MS.

Over the range of enzyme concentration that gave a linear turnover for α -D-GlcpNAc-PI (**1**) the substrate analogue α -D-GlcpNAc-IPC₁₈ (**3**) was de-*N*-acetylated at 450% the rate of α -D-

GlcNAc-PI (**1**). The increased turnover is most likely due to improved accessibility of the compound, conferred by the single alkyl chain, to the membranes that contain the de-*N*-acetylase enzyme in the cell-free system. Of the synthetic pseudodisaccharides (**47** – **50**) tested, only the α -anomer of the dipalmitoylated compound **49** showed any appreciable turnover at 22% the rate of α -D-GlcNAc-PI (**1**).

Table 1 Recognition of synthetic analogues by *T. brucei* GlcNAc-PI de-*N*-acetylase.

Compound	<i>m/z</i> transition for <i>NH</i> ₂	<i>m/z</i> transition for <i>NHAc</i>	Fragment assignment	Turnover / <i>pMol</i> / 10 ⁶ cells equiv.	Relative turnover ^a
1	1012 > 241	972 > 241	C ₆ H ₁₀ O ₈ P	6.1 ± 0.9	100%
3	673 > 223	715 > 223	C ₆ H ₈ O ₇ P	27.0 ± 6.0	450%
47	608 > 100	650 > 100	C ₆ H ₁₂ O	ND	-
48	608 > 100	650 > 100	C ₆ H ₁₂ O	ND	-
49	906 > 255	948 > 255	O ₂ CC ₁₅ H ₃₁	1.3 ± 0.3	22%
50	906 > 255	948 > 255	O ₂ CC ₁₅ H ₃₁	ND	-
51	633 > 332	592 > 332	NHSO ₂ C ₁₈ H ₃₇	ND	-
52	633 > 332	592 > 332	NHSO ₂ C ₁₈ H ₃₇	ND	-
53^b		563 > 447	C ₂₄ H ₄₇ O ₅ P	ND	-

^a turnover relative to α -D-GlcNAc-PI (**1**), ^b No suitable MRM, ND – turnover not detected. The multiple reaction monitoring (MRM) transition is shown as [Parent ion *m/z*] > [Daughter ion *m/z*].

Inhibitors:

Since the majority of the compounds were not processed by the *T. brucei* GlcNAc-PI de-*N*-acetylase, we tested their ability to inhibit the turnover of the α -D-GlcNAc-IPC₁₈ (**3**) substrate by the *T. brucei* GlcNAc-PI de-*N*-acetylase in the LC-MS/MS assay. Most compounds showed no

inhibitory activity at 100 μM . However, compounds **47** and **48** showed significant inhibition, with IC_{50} values of $11 \pm 4 \mu\text{M}$ and $37 \pm 20 \mu\text{M}$, respectively, indicating that they can be recognised but not processed by the *T. brucei* GlcNAc-PI de-*N*-acetylase. Interestingly, the potency of **47** and **48** is comparable to that observed with the, hydroxamic acid pseudodisaccharide analogue **54** (Figure 4), where the *N*-acetyl group is replaced with the zinc-chelating hydroxamic acid ($\text{IC}_{50} = 19 \pm 0.5 \mu\text{M}$)⁸ and suggests that the zinc-chelating group may not be driving the potency of the latter compound.




Figure 4 Hydroxamic acid

The ability of **49** and **48** to act as a substrate and inhibitor, respectively, of the *T. brucei* GPI pathway was confirmed using the trypanosome cell-free system with [³H]-mannose labelling (Figure 5). Priming the cell-free system with **49** produced three bands corresponding to the addition of 1-3 mannose residues (Figure 5A), and, consistent with this assignment, the bands were sensitive to jackbean α -mannosidase. As these mannosylated compounds lack the inositol 2-OH group they cannot undergo inositol acylation, a prerequisite for the transfer of ethanolamine, and thus are not processed past the Man_3 -species.²⁵ Priming the cell-free system with **3** (α -D-GlcNAc-IPC₁₈ at 10 μM) was efficiently prevented by incubation with **48** at 100 μM (Figure 5B), confirming that inhibition of the GlcNAc-PI de-*N*-acetylase is sufficient to prevent the formation of downstream GPI precursors.




Figure 5 The trypanosome GPI biosynthesis in the cell-free system. A. The *T. brucei* cell-free system was incubated without exogenous substrate, with **1**, α -D-GlcpNAc-PI (10 μ M), or with **49** (100 μ M) in the presence of GDP-[3 H]Mannose to stimulate the production of radiolabelled mannosylated GPI intermediates. B. Inhibition of the turnover of **3**, α -D-GlcpNAc-IPC₁₈ (10 μ M), in the presence of **48** (100 μ M). Glycolipid products were extracted, separated by high-performance thin-layer chromatography, and visualised by fluorography. DPM – dolichol-phosphate-mannose, M1 – Man₁ species, M2 – Man₂ species, M3 – Man₃ species, A' – EtNPMann₃ species, where the identity and migration of the species depends on the glycolipid substrate employed.

Previous studies (see the introduction for more detail) have shown that both α -D-GlcpNAc-PI (**1**) and β -D-GlcpNAc-PI (**6**) are recognised and processed by the *T. brucei* GlcNAc-PI de-*N*-acetylase, leading to the hypothesis that one or more of the inositol 2, 3, 4, and 5-OH groups is/are not required. Our data supports this hypothesis with an important caveat; when these hydroxyl groups are removed, substrate recognition and turnover is dependent on both the stereochemistry of the glycosidic linkage and the lipid composition. With the diacylglycerol lipid containing compounds **49** and **50**, only the natural α -anomer **49** is both recognised and processed, whereas the β -anomer **50** is neither a substrate nor an inhibitor. However, neither the α nor β -anomer of the octadecyl lipid containing compounds **47** and **48** is processed, although both appear to be recognised and act as inhibitors. These sets of diastereoisomers differ only in the identity of their lipid component, with the more flexible diacylglycerol moiety allowing the glycan to be recognised and processed. Thus, it appears that the requirement for the presence of inositol 2, 3, 4, and 5-OH groups for recognition by the *T. brucei* GlcNAc-PI de-*N*-acetylase is nuanced and may depend on the conformational flexibility of the substrate analogue.

The inability of the sulphonamide-containing compounds, **51** and **52**, to act as substrates or inhibitors confirms the importance of the phosphate group in substrate recognition. It may be that the presence of the negatively charged phosphate is essential for binding at the enzyme active site.

The inactivity of compound **53** is difficult to interpret. It may be that removal of the inositol 2, 3, 4, and 5-OH groups is not compatible with having a modified glucosamine moiety, or that the entire glucosamine ring is required. Having said this, the glucosamine “replacement” in compound **53** is likely to have a considerable degree of conformational flexibility, which could allow it to take up multiple orientations within the active site.

Conclusions

In summary, we have prepared a series of compounds to probe the substrate specificity and inhibition of enzymes involved at an early stage of GPI biosynthesis. The enzyme of interest to us, GlcNAc-PI de-*N*-acetylase, proved to be fastidious in its processing of variants of α -D-Glc β N-PI. We conclude that the glucosamine and the phospholipid moieties are essential for binding and that, while the *D*-*myo*-inositol residue is the preferred aglycone for recognition by the enzyme, the dispensing of it entirely with a cyclohexanediol group is tolerated but should be used with caution. Further work on this enzyme should focus on using the emerging structure activity relationship data to develop less synthetically complex, cell permeable analogues, which will be valuable chemical tools and may serve as leads for a drug discovery programme.

Experimental

Synthesis general methods

^1H , ^{13}C , ^{31}P NMR spectra were recorded on a Bruker AVANCE 500 MHz spectrometer using tetramethylsilane or the residual solvent as the internal standard. High resolution electrospray ionisation mass spectra [HRMS (ESI)] were recorded with a Bruker microTof spectrometer.

Melting points were determined on a Reichert hot-plate apparatus and are uncorrected. Optical rotations were measured with a Perkin-Elmer 343 polarimeter. Thin layer chromatography (TLC) was performed on Kieselgel 60 F₂₅₄ (Merck) plates with various solvent systems as developers, followed by detection under UV light or by charring with sulfuric acid-water-ethanol (15:85:5). Column chromatography was performed on Kieselgel 60 (0.040 – 0.063 mm) (Merck). Radial-band chromatography (RBC) was performed using a Chromatotron (model 7924T, TC Research UK) with silica gel F₂₅₄ TLC standard grade as the adsorbent. Iatrobeds (6RS-8060) were purchased from SES Analyssysteme. All reactions were carried out under argon in commercially available dry solvents, unless otherwise stated.

1R,2R-1-O-(2-Azido-3,4,6-tri-O-acetyl-2-deoxy- α -D-glucopyranosyl)-cyclohexanediol **16 and the β -anomer **17****

After drying overnight over P₂O₅ in a vacuum desiccator, the glycosyl donor¹⁴ **14** (731 mg, 1.54 mmol) and the acceptor **15** (Sigma-Aldrich) were dissolved in 1:1 Et₂O – CH₂Cl₂ (10 mL). To this solution was added activated 4Å molecular sieves (1g) and TMSOTf (5.4 μ L, 0.03 mmol) at rt under argon. The reaction mixture was stirred at rt overnight, whereafter it was neutralised with TEA, percholated through a short column of silica gel (further elution with EtOAc) and the subsequent eluent was concentrated under reduced pressure. RBC [elution first with PE (40 – 60°) and then with 3:2 PE (40 – 60°) – EtOAc] of the residue gave first the α -linked pseudodisaccharide **16** (255 mg, 39%) as a waxy solid; $[\alpha]_D^{25} +83.7^\circ$ (*c* 2.37, CHCl₃); δ_H (500 MHz, CDCl₃) 5.42 (dd, 1H, $J_{2',3'} = J_{3',4'} = 10.3$ Hz, H-3'), 5.09 (d, 1H, $J_{1',2'} = 3.7$ Hz, H-1'), 4.95 (dd, 1H, $J_{3',4'} = J_{4',5'} = 10.3$ Hz, H-4'), 4.18 (dd, 1H, $J_{5',6'a} = 4.8$, $J_{6'a,6'b} = 12.2$ Hz, H-6'a), 4.08 (m, 1H, H-5'), 4.01 (dd, 1H, $J_{5',6'b} = 2.2$, $J_{6'a,6'b} = 12.2$ Hz, H-6'b), 3.58 (dd, 1H, H-2'), 3.44 (m, 1H, H-2), 3.23 (m, 1H, H-1), 3.05 (s, 1H, 2-OH), 2.50 – 1.90 (m, 11H, 3 x CH₃, H-3a and 6a), 1.64 (m, 2H, H-4a and 5a), 1.35 – 1.15 (m, 4H, H-3b, 4b, 5b and 6b); δ_C (125 MHz, CDCl₃) 170.6 – 169.8 (3 x COCH₃), 99.3 (C-1'), 87.5 (C-

1), 74.1 (C-2), 71.7 (C-3'), 68.6 (C-4'), 67.8 (C-5'), 62.1 (C-2'), 61.9 (C-6'), 32.0, 31.7, 24.4, 23.8, 20.7 (COCH₃) 20.6 (COCH₃); HRMS (ESI) calcd. for C₁₈H₂₇N₃NaO₉ [M + Na]⁺ 452.1640, found 452.1622 and then the β -anomer **17** (202 mg, 31%) as a white solid; mp 120 – 122 °C; [α]_D²⁵ +21.5° (c 1.15, CHCl₃); δ _H (500 MHz, CDCl₃) 4.91 (m, 2H, H-3' and 4'), 4.43 (d, 1H, $J_{1',2'} = 8.1$ Hz, H-1'), 4.12 (m, 2H, H6'a and 6'b), 3.67 (m, 1H, H-5'), 3.45 (m, 1H, H-2'), 3.40 – 3.30 (m, 2H, H-1 and 2), 2.50 – 1.93 (m, 11H, 3 x CH₃ and 2H-cyclitol), 1.70 – 1.60 (m, 2H, cyclitol) 1.35 (m, 1H, cyclitol), 1.17 (m, 3H, cyclitol); δ _C (125 MHz, CDCl₃) 170.6 – 169.6 (3 x COCH₃), 102.1 (C-1'), 88.2, 73.2, 72.1, 71.8 (C-5'), 68.3, 63.7 (C-2'), 61.8 (C-6'), 32.2, 30.9, 24.2, 23.6, 20.7 (COCH₃), 20.6 (COCH₃); HRMS (ESI) calcd. for C₁₈H₂₈N₃O₉ [M + H]⁺ 430.1820, found 430.1831.

Triethylammonium 1R,2R-1-O-(2-azido-3,4,6-tri-O-acetyl-2-deoxy- α -D-glucopyranosyl)-cyclohexanediol 2-(n-octadecylphosphate) **19**

Each of the compounds **16** (117mg, 0.27 mmol) and **18**¹⁵ (237 mg, 0.54 mmol) were dried overnight over P₂O₅ in a vacuum desiccator, whereafter anhyd pyridine was evaporated therefrom. They were then dissolved in dry pyridine (10 mL), pivaloyl chloride (216 μ L, 1.76 mmol) was added and the resulting solution was stirred under argon at rt for 1 h. A freshly prepared solution of iodine (274 mg, 1.08 mmol) in 9:1 pyridine – water was then added and stirring of the reaction mixture was continued for 45 min. After the addition of CH₂Cl₂ (20 mL), the organic solution was washed successively with 5% aq NaHSO₃ (25 mL), water (25 mL), 1M TEAB buffer solution (3 x 15 mL), dried (MgSO₄) and concentrated under reduced pressure. RBC of the residue (elution first with CH₂Cl₂ and then with 9:1 CH₂Cl₂ – MeOH) afforded the TEA phosphate derivative **19** (140 mg, 60%); [α]_D²⁵ +68.6° (c 1.07, CHCl₃); δ _H (500 MHz, CDCl₃) 5.40 (dd, 1H, $J_{3',4'} = 9.2$ Hz, H-3'), 5.27 (d, 1H, $J_{1',2'} = 3.7$ Hz, H-1'), 4.94 (t, 1H, $J_{4',5'} = 9.6$ Hz, H-4'), 4.24 – 4.13 (m, 2H, H6'a and H-1 or 2), 4.07 – 3.98 (m, 2H, H-5' and 6'b), 3.84 – 3.76 (m, 3H, OCH₂ and H-1 or 2), 3.17 (dd, 1H, $J_{2',3'} = 10.6$ Hz, H-2'), 2.83 (q, 6H, $J = 6.8$ Hz, 3 x CH₂CH₃), 2.05 – 1.95 (m, 10H, 3 x COCH₃ and 1H-cyclitol), 1.87 (m, 1H, cyclitol), 1.62 – 1.45 (m, 6H, OCH₂CH₂ and 4H-cyclitol), 1.33 – 1.14

(41H, [CH₂]₁₅, 3x CH₂CH₃ and 2H-cyclitol), 0.81 (t, 3H, *J* = 6.8 Hz, CH₂CH₃); δ_C (125 MHz, CDCl₃) 169.5 – 168.7 (3 x COCH₃), 96.9 (C-1'), 76.4 (C-1 or C-2), 73.9 (C-1 or C-2), 69.4 (C-3'), 67.9 (C-4'), 66.7 (C-5'), 65.2 (OCH₂), 61.1 (C-6'), 60.0 (C-2'), 44.4 [N(CH₂CH₃)₃], 30.9, 29.7, 28.7 – 28.1, 24.8, 21.7, 21.0, 20.3, 19.6, 13.1 (CH₂CH₃), 7.5 [N(CH₂CH₃)₃]; δ_P (202 MHz, CDCl₃) –0.05 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₃₆H₆₃N₃O₁₂P [M – NEt₃ – H][–] 760.4155, found 760.4154.

Triethylammonium 1*R*,2*R*-1-*O*-(2-azido-2-deoxy-α-*D*-glucopyranosyl)-cyclohexanediol 2-(*n*-octadecylphosphate) 20

To a solution of compound **19** (78 mg, 0.09 mmol) in 1:1 CH₂Cl₂ – MeOH (10 mL) was added 5.4 M NaOMe in MeOH (0.10 mL). The mixture was kept for 3 h at rt and was then neutralised with Amberlite IR-120 (H⁺) ion-exchange resin, filtered and the filtrate concentrated under reduced pressure. Column chromatography (elution first with 3:1 CH₂Cl₂ – MeOH and then with 2:1 → 1:1) of the residue furnished the TEA salt **20** (45 mg, 67%) as a waxy solid; [α]_D²⁵ +51.6° (*c* 4.5, 1:1 THF – MeOH); δ_H (500 MHz, 1:1 CDCl₃ – MeOH-*d*₄) 5.18 (d, 1H, *J*_{1',2'} = 3.5 Hz, H-1'), 4.25 (m, 1H, H-1 or 2), 3.96 – 3.65 (7H, OCH₂, H-3', 5', 6'a,b and H-1 or 2) 3.40 (t, 1H, *J*_{3',4'} = *J*_{4',5'} = 9.6 Hz, H-4'), 3.13 (q, 6H, *J* = 7.3 Hz, 3 x CH₂CH₃), 3.06 (dd, 1H, *J*_{2',3'} = 10.5 Hz, H-2'), 2.03 (m, 1H, cyclitol), 1.90 (m, 1H, cyclitol), 1.64 (m, 6H, OCH₂CH₂ and 4H-cyclitol), 1.43 – 1.22 (41H, [CH₂]₁₅, 3x CH₂CH₃ and 2H-cyclitol), 0.88 (t, 3H, *J* = 6.8 Hz, CH₂CH₃); δ_C (125 MHz, 1:1 CDCl₃ – MeOH-*d*₄) 98.7 (C-1'), 76.3 (C-1 or C-2), 74.2 (C-1 or C-2), 73.8 72.4, 72.3, 66.9 (OCH₂), 64.5 (C-2'), 62.7, 47.4 [N(CH₂CH₃)₃], 32.0, 31.0 – 30.6, 29.8, 29.5, 27.1, 23.9, 22.5, 22.1, 15.0 (CH₂CH₃), 9.6 [N(CH₂CH₃)₃]; δ_P (202 MHz, 1:1 CDCl₃ – MeOH-*d*₄) –0.47 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₃₀H₅₇N₃O₉P [M – NEt₃ – H][–] 634.3838, found 634.3850.

1*R*,2*R*-1-*O*-[2-(4-Hydroxybutyl)amino-2-deoxy-α-*D*-glucopyranosyl]-cyclohexanediol 2-(*n*-octadecylphosphate) 21

A solution of the azido compound **20** (45 mg, 0.06 mmol) in 1:1 THF – MeOH (5 mL) containing 10-20% Pd(OH)₂ on carbon (15 mg) was stirred under a hydrogen atmosphere at rt for 30 min before it was percolated through a short column of Chelex 100 on a bed of Celite (further elution with 1:1 THF – MeOH). The eluent was concentrated under reduced pressure and the ensuing residue was purified by column chromatography (elution gradient 6:1 → 4:1 CH₂Cl₂ – MeOH) to give the hydroxybutylamino compound **21** (15 mg, 38%); $[\alpha]_D^{25} +34.0^\circ$ (c 1.5, 1:1 CHCl₃ – MeOH); δ_H (500 MHz, 1:1 CDCl₃ – MeOH-d₄) 5.43 (d, 1H, $J_{1',2'} = 3.6$ Hz, H-1'), 4.05 (m, 1H, H-1 or 2), 3.93 (dd, 1H, $J_{3',4'} = 10.4$ Hz, H-3'), 3.90 – 3.50 (8H, OCH₂ butyl, POCH₂, H-5', 6'a,b and H-1 or 2), 3.38 (m, 1H, H-4'), 3.17 (t, 2H, $J = 7.6$ Hz, NCH₂), 3.00 (dd, 1H, $J_{2',3'} = 10.4$ Hz, H-2'), 2.18 (m, 1H, cyclitol), 2.08 (m, 1H, cyclitol), 1.86 (m, 2H, CH₂ butyl), 1.76 – 1.57 (6H, POCH₂CH₂, CH₂ butyl and 2H-cyclitol), 1.45 – 1.20 (34H, [CH₂]₁₅ and 4H-cyclitol), 0.89 (t, 3H, $J = 6.8$ Hz, CH₂CH₃); δ_C (125 MHz, 1:1 CDCl₃ – MeOH-d₄) 97.3 (C-1'), 83.2 (C-1 or C-2), 79.3 (C-1 or C-2), 74.0, 71.8, 71.7, 67.1, 62.3, 61.6 (C-2'), 54.8, 50.6, 48.7 (NCH₂), 34.2, 33.6, 31.9, 30.9 – 30.6, 27.0, 25.3, 25.2, 15.0 (CH₂CH₃); δ_P (202 MHz, 1:1 CDCl₃ – MeOH-d₄) 0.66 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₃₄H₆₇NO₁₀P [M – H][–] 680.4508, found 680.4554.

1R,2R-1-O-(2-Amino-2-deoxy- α -D-glucopyranosyl)-cyclohexanediol 2-(*n*-octadecylphosphate)

7

A solution of the TEA salt **20** (52 mg, 0.07 mmol) in 1:1 stabilised THF – MeOH (5 mL) containing 10-20% Pd(OH)₂ on carbon (5 mg) was stirred under a hydrogen atmosphere at rt for 1 h. Work-up as described for the derivative **21** gave, after column chromatography (5:1 CH₂Cl₂ – MeOH), the amino compound **7** (33 mg, 77%); $[\alpha]_D^{25} +57.5^\circ$ (c 3.3, 1:1 CHCl₃ – MeOH); δ_H (500 MHz, 1:1 CDCl₃ – MeOH-d₄) 5.42 (d, 1H, $J_{1',2'} = 3.9$ Hz, H-1'), 4.05 (m, 1H, H-1 or 2), 3.90 – 3.65 (6H, OCH₂, H-3', 5', 6'a,b), 3.58 (m, 1H, H-1 or 2), 3.32 (m, 3H, H-4' and MeOH-d₄), 3.05 (dd, 1H, $J_{2',3'} = 10.5$ Hz, H-2'), 2.10 (m, 2H, 2H-cyclitol), 1.74 – 1.57 (4H, OCH₂CH₂ and 2H-cyclitol), 1.44 – 1.21 (34H, [CH₂]₁₅ and 4H-cyclitol), 0.90 (t, 3H, $J = 6.8$ Hz, CH₂CH₃); δ_C (125 MHz, 1:1 CDCl₃

– MeOH- d_4) 97.9 (C-1'), 81.8 (C-1 or C-2), 79.9 (C-1 or C-2), 74.5, 71.8, 71.7, 66.7 (OCH₂), 62.4 (C-6'), 55.8 (C-2'), 34.0, 33.4, 33.1, 31.9, 31.8, 30.8, 30.5, 26.9, 25.2, 25.1, 23.8, 14.5 (CH₂CH₃); δ_P (202 MHz, 1:1 CDCl₃ – MeOH- d_4) 0.54 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₃₀H₆₁NO₉P [M + H]⁺ 610.4078, found 610.4050.

1*R*,2*R*-1-*O*-(2-Azido-3,4,6-tri-*O*-acetyl-2-deoxy- α -D-glucopyranosyl)-2-*O*-(*tert*-butyldimethylsilyl)-cyclohexanediol 22

The alcohol **16** (284 mg, 0.66 mmol) was dried overnight in a desiccator over P₂O₅ under high vacuum and then dissolved in anhyd. CH₂Cl₂ (10 mL). To this solution, at room temperature, was added 2,6-lutidine (154 μ L, 1.32 mmol) and *tert*-butyldimethylsilyl trifluoromethane sulfonate (228 μ L, 0.99 mmol). After 30 min, CH₂Cl₂ (25 mL) and brine (25 mL) were added and the organic layer separated. The aqueous layer was re-extracted with CH₂Cl₂ (25 mL) and the combined organic layers were washed with brine (2 x 25 mL), dried (MgSO₄) and concentrated under reduced pressure. RBC [elution first with PE (40-60°) and then with 1:1 PE(40-60°) – Et₂O] of the residue gave the azide **22** (284 mg, 79%) as an oil; $[\alpha]_D^{25} +90.2^\circ$ (*c* 1.52, CHCl₃); δ_H (500 MHz, CDCl₃) 5.40 (t, 1H, $J_{2',3'} = J_{3',4'} = 9.7$ Hz, H-3'), 5.35 (d, 1H, $J_{1',2'} = 3.3$ Hz, H-1'), 5.20 (t, 1H, $J_{3',4'} = J_{4',5'} = 9.7$ Hz, H-4'), 4.26 (dd, 1H, $J_{5',6'a} = 4.7$, $J_{6'a,6'b} = 12.1$ Hz, H-6'a), 4.10 (m, 2H, H-5' and 6'b), 3.75 (m, 1H, H-1 or 2), 3.57 (m, 1H, H-1 or 2), 3.17 (dd, 1H, H-2'), 2.10 – 2.03 (3 x s, 9H, 3 x COCH₃), 1.89 (m, 2H-cyclitol), 1.70 – 1.25 (6H-cyclitol), 0.89 (s, 9H, 3 x CH₃), 0.12 – 0.08 (2 x s, 6H, 2 x CH₃); δ_C (125 MHz, CDCl₃) 170.6 – 169.7 (3 x COCH₃), 97.9 (C-1'), 79.7 (C-1 or 2), 72.6 (C-1 or 2), 70.4 (C-3'), 68.7 (C-4'), 67.6 (C-5'), 62.1 (C-6'), 61.0 (C-2'), 32.5, 29.8, 25.9, 22.1, 20.7 (COCH₃), 20.6 (COCH₃), 18.0, -4.1, -4.9; HRMS (ESI) calcd. for C₂₄H₄₂N₃O₉Si [M + H]⁺ 544.2685, found 544.2698.

1*R*,2*R*-1-*O*-(2-Azido-2-deoxy- α -D-glucopyranosyl)-2-*O*-(*tert*-butyldimethylsilyl)-cyclohexanediol 23

To a solution of the triacetate **22** (185 mg, 0.34 mmol) in 1:1 CH₂Cl₂ – MeOH (92 mL) was added 5.4 M NaOMe in MeOH (230 μ L). The mixture was kept for 30 min at rt and was then neutralised with Amberlite IR-120 (H⁺) ion-exchange resin, filtered and the filtrate concentrated under reduced pressure. The residue, so obtained, was percolated through a short silica-gel column (further elution with EtOAc) and the eluent was concentrated under reduced pressure to afford the triol **23** (136 mg, 96%) as a white solid, mp 122 – 123 °C (from 10:1 hexane – Et₂O); [α]_D²⁵ +84.1° (*c* 1.63, CHCl₃); δ_{H} (500 MHz, CDCl₃) 5.22 (d, 1H, $J_{1',2'} = 3.4$ Hz, H-1'), 4.02 (t, 1H, $J_{3',4'} = 9.4$ Hz, H-3'), 3.90 (dd, 1H, $J_{5',6'a} = 2.3$, $J_{6'a,6'b} = 11.6$ Hz, H-6'a), 3.81 (dd, 1H, $J_{5',6'b} = 2.2$, $J_{6'a,6'b} = 11.6$ Hz, H-6'b), 3.73 (m, 2H, H-5' and H-1 or 2), 3.65 (t, 1H, $J_{4',5'} = 9.4$ Hz, H-4'), 3.59 (m, 1H, H-1 or 2), 3.15 (dd, 1H, $J_{2',3'} = 10.4$ Hz H-2'), 1.85 (m, 2H, cyclitol), 1.60 (m, 2H, cyclitol) 1.50 – 1.20 (4H, cyclitol), 0.89 (s, 9H, 3 x CH₃), 0.12 – 0.08 (2 x s, 6H, 2 x CH₃); δ_{C} (125 MHz, CDCl₃) 98.2 (C-1'), 78.9 (C-1 or 2), 71.5, 71.4, 70.5 (C-4'), 62.9 (C-2'), 61.6 (C-6'), 31.9, 29.4, 25.9, 22.5, 21.7, 18.0, -4.3, -4.9; HRMS (ESI) calcd. for C₁₈H₃₆N₃O₆Si [M + H]⁺ 418.2368, found 418.2365.

1R,2R-1-O-(2-Azido-3,4,6-tri-O-benzyl-2-deoxy- α -D-glucopyranosyl)-2-O-(tert-butyltrimethylsilyl)-cyclohexanediol **24**

To a stirred and cooled (0 °C) solution of the triol **23** (70 mg, 0.17 mmol) in DMF (10 mL) under argon was added NaH (19 mg, 0.78 mmol) and the solution was stirred for 15 min before benzyl bromide (93 μ L, 0.78 mmol) was added dropwise. The reaction mixture was stirred at rt overnight and then poured slowly and carefully into ice-cold water (50 mL). After dilution with EtOAc (50 mL), the EtOAc solution was washed with brine (25 mL), dried (Na₂SO₄) and concentrated under reduced pressure. RBC [elution gradient PE (40-60°) → 10:1 → 7:1 → 4:1 PE (40-60°) – Et₂O] of the residue yielded the fully protected compound **24** (86 mg, 74%); [α]_D²⁵ +57.9° (*c* 1.78, CHCl₃); δ_{H} (500 MHz, CDCl₃) 7.33 – 7.03 (15H, 3 x Ph), 5.17 (d, 1H, $J_{1',2'} = 3.6$ Hz, H-1'), 4.84 – 4.37 (6H, 3 x CH₂Ar), 3.93 (t, 1H, $J_{3',4'} = 9.0$ Hz, H-3'), 3.83 (m, 1H, H-5'), 3.72 – 3.60 (m, 3H, H-4', 6'a and 1 or 2), 3.57 (dd, 1H, $J_{5',6'b} = 2.0$, $J_{6'a,6'b} = 10.7$ Hz, H-6'b), 3.48 (m, 1H, H-1 or 2),

3.26 (dd, 1H, $J_{2',3'} = 10.3$ Hz, H-2'), 1.80 (m, 2H, cyclitol), 1.32 (m, 2H, cyclitol), 1.40 – 1.10 (4H, cyclitol), 0.82 (s, 9H, 3 x CH₃), 0.02 – 0.00 (2 x s, 6H, 2 x CH₃); δ_C (125 MHz, CDCl₃) 137.0 – 136.8 (Ph), 127.4 – 126.7 (Ph), 97.0 (C-1'), 79.2 (C-3'), 77.9 (C-1 or 2), 77.4, 74.3, 74.1, 72.5, 71.3, 69.8 (C-5'), 67.3 (C-6'), 62.6 (C-2'), 31.3, 28.7, 24.5, 21.7, 21.0, 18.4, -5.2, -5.8; HRMS (ESI) calcd. for C₃₉H₅₄N₃O₆Si [M + H]⁺ 688.3776, found 688.3780.

1R,2R-1-O-(2-Amino-3,4,6-tri-O-benzyl-2-deoxy- α -D-glucopyranosyl)-2-O-(tert-butyltrimethylsilyl)-cyclohexanediol 25

To a stirred solution of **24** (86 mg, 0.12 mmol) in 10:1 THF – water (5 mL) at 60 °C was added Ph₃P (98 mg, 0.38 mmol). After 3 h, TLC showed the complete disappearance of the starting material. The reaction was then cooled to rt, poured into water (25 mL) and extracted with CH₂Cl₂ (3 x 25 mL). The combined organics were washed successively with water (25 mL), brine (25 mL), dried (MgSO₄) and concentrated under reduced pressure. RBC [elution gradient hexane → 7:1 → 3:1 → 1:1 → 1:3 hexane – EtOAc] of the residue afforded the amino derivative **25** (53 mg, 62%); $[\alpha]_D^{25} +65.6^\circ$ (c 1.09, CHCl₃); δ_H (500 MHz, CDCl₃) 7.35 – 7.09 (15H, 3 x Ph), 4.98 – 4.44 (7H, H-1' and 3 x CH₂Ar), 3.87 (m, 1H, H-5'), 3.75 (dd, 1H, $J_{5',6'a} = 3.6$, $J_{6'a,6'b} = 10.6$ Hz, H-6'a), 3.61 (m, 3H, H-4', 6'b and 1 or 2), 3.55 (t, 1H, $J_{3',4'} = 9.2$ Hz, H-3'), 3.45 (m, 1H, H-1 or 2), 2.77 (dd, 1H, $J_{1',2'} = 3.7$, $J_{2',3'} = 9.7$ Hz, H-2'), 1.98 (m, 1H, cyclitol), 1.70 (m, 1H, cyclitol), 1.63 – 1.16 (6H, cyclitol), 0.84 (s, 9H, 3 x CH₃), 0.00 (2 x s, 6H, 2 x CH₃); δ_C (125 MHz, CDCl₃) 138.7 – 138.0 (Ph), 128.5 – 127.7 (Ph), 100.2 (C-1'), 84.1 (C-3'), 79.9 (C-1 or 2), 78.9, 75.6, 74.9, 73.5, 71.4 (C-5'), 71.0, 68.7 (C-6'), 56.4 (C-2'), 31.9, 29.4, 25.9, 22.4, 21.8, 18.0, -4.3, -4.4; HRMS (ESI) calcd. for C₃₉H₅₆NO₆Si [M + H]⁺ 662.3871, found 662.3874.

1R,2R-1-O-[2-N-(tert-Butoxycarbonyl)amino-3,4,6-tri-O-benzyl-2-deoxy- α -D-glucopyranosyl]-2-O-(tert-butyltrimethylsilyl)-cyclohexanediol 26

The amine **25** (147 mg, 0.22 mmol) was dissolved in EtOAc (10 mL) at rt. Di-tert-butylidicarbonate (58 mg, 0.26 mmol) was then added and the mixture was stirred overnight at rt.

Afterwards, the reaction mixture was diluted with EtOAc (25 mL) and then washed successively with water (25 mL), brine (25 mL), dried (Na₂SO₄) and concentrated under reduced pressure. RBC [elution gradient hexane → 7:1 → 5:1 hexane – EtOAc] of the residue afforded the Boc protected derivative **26** (132 mg, 79%); $[\alpha]_{\text{D}}^{25} +39.0^{\circ}$ (*c* 1.06, CHCl₃); δ_{H} (500 MHz, CDCl₃) 7.30 – 7.05 (15H, 3 x Ph), 4.95 (d, 1H, $J_{1',2'} = 3.3$ Hz, H-1'), 4.76 – 4.38 (7H, NH and 3 x CH₂Ar), 3.90 (m, 1H, H-2') 3.83 (m, 1H, H-5'), 3.69 (dd, 1H, $J_{5',6'a} = 4.1$, $J_{6'a,6'b} = 10.7$ Hz, H-6'a), 3.61 (m, 3H, H-3', 4', and 6'b), 3.48 (m, 1H, H-1 or 2), 3.37 (m, 1H, H-1 or 2), 2.01 (m, 1H, cyclitol), 1.74 (m, 1H, cyclitol), 1.52 (m, 2H, cyclitol), 1.35 (s, 9H, 3 x BocCH₃), 1.30 – 1.10 (4H, cyclitol), 0.83 (s, 9H, 3 x CH₃), 0.00 (2 x s, 6H, 2 x CH₃); δ_{C} (125 MHz, CDCl₃) 155.3 (C=O), 138.6 – 138.2 (Ph), 128.4 – 127.5 (Ph), 99.1 (C-1'), 81.5, 81.3 (C-1 or 2), 79.5, 78.5, 75.3, 75.1, 73.4, 73.2 (C-1 or 2), 71.4 (C-5'), 68.8 (C-6'), 54.6 (C-2'), 33.4, 30.8, 28.5, 26.1, 23.3, 22.9, 18.1, -3.9, -4.3; HRMS (ESI) calcd. for C₄₄H₆₄NO₈Si [M + H]⁺ 762.4396, found 762.4393.

1R,2R-1-O-[2-N-(*tert*-Butoxycarbonyl)amino-3,4,6-tri-O-benzyl-2-deoxy- α -D-glucopyranosyl]-cyclohexanediol **27**

To a stirred solution of the silyl derivative **26** (114 mg, 0.15 mmol) in THF (10 mL) at 0° C was added ~70% HF-pyridine (90 μ L). The solution was stirred overnight at rt whereafter a further aliquot of ~70% HF-pyridine (90 μ L) was added and the solution was left to stir overnight; this process was continued on day 3. On day 4, TLC revealed the complete disappearance of the starting material, whereafter satd NaHCO₃ (1 mL) was added dropwise to quench the reaction and the resulting solution was poured into brine (25 mL) and extracted with EtOAc (3 x 25 mL). The EtOAc extracts were combined and washed with brine (25 mL), dried (MgSO₄) and concentrated under reduced pressure. RBC [elution gradient hexane → 2:1 hexane – EtOAc] of the residue furnished the alcohol **27** (97 mg, 49%); $[\alpha]_{\text{D}}^{25} +27.2^{\circ}$ (*c* 1.08, CHCl₃); δ_{H} (500 MHz, CDCl₃) 7.36 – 7.10 (15H, 3 x Ph), 5.12 (d, 1H, $J_{1',2'} = 3.6$ Hz, H-1'), 4.86 – 4.44 (7H, NH and 3 x CH₂Ar), 3.95 (dd, 1H, $J_{3',4'} = J_{4',5'} = 9.7$ Hz, H-4'), 3.86 (m, 1H, H-2'), 3.75 – 3.61 (4H, H-3', 5' and 6'a,b), 3.47 (m, 1H,

H-1 or 2), 3.32 (m, 1H, H-1 or 2), 2.10 – 1.91 (2H, cyclitol), 1.65 (m, 2H, cyclitol), 1.43 (s, 9H, 3 x CH₃), 1.33 – 1.16 (4H, cyclitol); δ_C (125 MHz, CDCl₃) 155.6 (C=O), 138.4 – 138.0 (Ph), 128.4 – 127.7 (Ph), 99.5 (C-1'), 85.0 (C-1 or 2), 80.5, 79.7, 78.5, 75.0, 74.0 (C-1 or 2), 73.4, 71.2 (C-4'), 68.7 (C-6'), 54.9 (C-2'), 32.9, 31.7, 28.4, 24.3, 23.9; HRMS (ESI) calcd. for C₃₈H₅₀NO₈ [M + H]⁺ 648.3531, found 648.3527.

Triethylammonium 1*R*,2*R*-1-*O*-[2-*N*-(*tert*-butoxycarbonyl)amino-3,4,6-tri-*O*-benzyl-2-deoxy- α -D-glucopyranosyl]-cyclohexanediol 2-(1,2-di-*O*-hexadecanoyl-*sn*-glycerol 3-phosphate) 29

This compound was obtained from the alcohol **27** (55.7 mg, 0.086 mmol) and 1,2-di-*O*-hexadecanoyl-*sn*-glycerol 3-hydrogenphosphate TEA salt **28**¹⁹ (126 mg, 0.17 mmol) in the presence of pivaloyl chloride (69 μ L, 0.56 mmol) essentially as described for the 2-(*n*-octadecyl phosphate) **19**. After the oxidation with iodine (87 mg, 0.34 mmol) in 9:1 pyridine – water followed by the same aqueous workup as described for **19**, RBC (elution first with CH₂Cl₂ and then with 20:1 \rightarrow 15:1 CH₂Cl₂ – MeOH) afforded the TEA phosphate derivative **29** (57 mg, 52%) as an opaque oil; $[\alpha]_D^{25} +27.7^\circ$ (*c* 1.08, CHCl₃); δ_H (500 MHz, CDCl₃) 12.5 (brs, 1H, NH TEA salt), 7.36 – 7.10 (15H, 3 x Ph), 5.24 (m, 1H, H-2 glycerol), 4.96 (s, 1H, H-1'), 4.84 – 4.44 (6H, 3 x CH₂Ar), 4.39 (m, 1H, 1- or 3-CHa glycerol), 4.17 (dd, 1H, *J* = 6.6, *J* = 12.0 Hz, 1- or 3-CHb glycerol), 4.13 – 3.97 (m, 4H, H-2', 1 or 2 cyclitol and 1- or 3-CH₂ glycerol), 3.95 (m, 1H, H-4'), 3.85 (t, *J*_{2',3'} = *J*_{3',4'} = 9.9 Hz, H-3'), 3.74 (dd, 1H, *J*_{5',6'a} = 4.1, *J*_{6'a,6'b} = 10.7 Hz, H-6'a), 3.66 (m, 2H, H-5' and 6'b), 3.50 (m, 1H, H-1 or 2), 2.97 (q, 6H, *J* = 7.1 Hz, 3 x CH₂CH₃), 2.27 (m, 4H, 2 x COCH₂), 2.21 – 1.95 (2H, cyclitol), 1.72 – 1.50 (m, 6H, 2 x COCH₂CH₂ and 2H cyclitol), 1.44 (s, 9H, 3 x CH₃), 1.33 – 1.18 (61H, 3 x CH₂CH₃, 2 x [CH₂]₁₂ and 4H cyclitol), 0.88 (t, 6H, *J* = 7.3 Hz, 2 x CH₂CH₃); δ_C (125 MHz, CDCl₃) 172.4 (C=O), 171.9 (C=O), 155.4 (C=O), 138.9 – 137.4 (Ph), 127.3 – 126.3 (Ph), 98.9 (C-1'), 80.0, 76.8, 73.7, 72.7, 72.3, 70.3, 69.8, 69.5, 68.1, 67.6, 62.4, 61.8, 61.4, 53.5, 44.2 [N(CH₂CH₃)₃], 33.3, 33.1, 30.9, 30.2, 28.7 – 28.1, 27.5, 23.4, 21.7, 13.1, 7.40 [N(CH₂CH₃)₃];

δ_P (202 MHz, $CDCl_3$) 0.04 (with heteronuclear decoupling); HRMS (ESI) calcd. for $C_{73}H_{115}NO_{15}P$ $[M - NEt_3 - H]^-$ 1276.8010, found 1276.8015.

1R,2R-1-O-[2-N-(*tert*-Butoxycarbonyl)amino-2-deoxy- α -D-glucopyranosyl]-cyclohexanediol 2-(1,2-di-O-hexadecanoyl-*sn*-glycerol 3-phosphate) 30

A solution of the benzylated compound **29** (57 mg, 0.041 mmol) in 1:1 THF – *n*-propanol (10 mL) containing 10-20% $Pd(OH)_2$ on carbon (15 mg) was stirred under 3 atm of hydrogen for 3 h before it was percolated through a short column of Chelex 100 on a bed of Celite (further elution with 1:1 THF – *n*-propanol). The eluent was concentrated under reduced pressure and the ensuing residue was purified by column chromatography (elution gradient 7:1 \rightarrow 4:1 CH_2Cl_2 – MeOH) to give the Boc protected derivative **30** (30 mg, 73%); $[\alpha]_D^{25} +39.1^\circ$ (c 3.00, 1:1 CH_2Cl_2 – MeOH); δ_H (500 MHz, 1:1 $CDCl_3$ – MeOH- d_4) 5.25 (m, 1H, H-2 glycerol), 4.90 (s, 1H, H-1'), 4.45 (m, 1H, 1- or 3-CHa glycerol), 4.20 (dd, 1H, $J = 6.7$, $J = 11.7$ Hz, 1- or 3-CHb glycerol), 4.00 (m, 2H, 1- or 3-CH₂ glycerol), 3.75 (m, 1H, H-3' and 4'), 3.65 (m, 1H, H-1 or 2), 3.55 (dd, 1H, $J_{1',2'} = 3.0$, $J_{2',3'} = 10.5$ Hz, H-2'), 3.47 (m, 1H, H-1 or 2), 3.40 (m, 3H, H-5' and 6'a,b), 2.40 – 2.00 (m, 6H, 2 x COCH₂ and 2H cyclitol), 1.74 – 1.57 (m, 6H, 2 x COCH₂CH₂ and 2H cyclitol), 1.47 (s, 9H, 3 x CH₃), 1.40 – 1.20 (52H, 2 x $[CH_2]_{12}$ and 4H cyclitol), 0.89 (t, 6H, $J = 7.1$ Hz, 2 x CH₂CH₃); δ_C (125 MHz, 1:1 $CDCl_3$ – MeOH- d_4) 174.6 (C=O), 174.1 (C=O), 158.2 (C=O), 98.9 (C-1'), 82.0, 78.0, 73.3, 73.0, 72.0, 71.1, 64.0, 63.1, 62.0, 57.1, 56.4, 34.7, 34.5, 33.0 – 29.5, 28.5, 25.4, 24.3, 23.5, 23.1, 14.3; δ_P (202 MHz, 1:1 $CDCl_3$ – MeOH- d_4) –0.28 (with heteronuclear decoupling); HRMS (ESI) calcd. for $C_{52}H_{97}NO_{15}P$ $[M - H]^-$ 1006.6601, found 1006.6635.

1R,2R-1-O-(2-Amino-2-deoxy- α -D-glucopyranosyl)-cyclohexanediol 2-(1,2-di-O-hexadecanoyl-*sn*-glycerol 3-phosphate) 9

To a solution of the *tert*-butoxycarbonyl protected compound **30** (30 mg, 0.030 mmol) in 1:1 CH_2Cl_2 – MeOH (1 mL) was added 9:1 trifluoroacetic acid (TFA) – water (5 mL). After stirring 3 h at rt, toluene (5 mL) was added and the solvents were removed under reduced pressure. Toluene

(2 x 5 mL) was evaporated off twice from the residue (to remove traces of TFA and water) to give the pseudodisaccharide phosphate derivative **9** (25 mg, 93%) which did not require any further purification; $[\alpha]_D^{25} +28.0^\circ$ (*c* 2.50, 1:1 CHCl₃ – MeOH); δ_H (500 MHz, 1:1 CDCl₃ – MeOH-d₄) 5.28 (d, 1H, $J_{1,2'} = 3.7$ Hz, H-1'), 5.14 (m, 1H, H-2 glycerol), 4.34 (dd, 1H, $J = 3.2$, $J = 12.0$ Hz, 1- or 3-CHa glycerol), 4.10 (dd, 1H, $J = 6.6$, $J = 12.0$ Hz, 1- or 3-CHb glycerol), 3.96 (m, 1H, H-1 or 2), 3.88 (m, 2H, 1- or 3-CH₂ glycerol), 3.71 (m, 3H, H-3' and 6'a,b), 3.62 (dd, 1H, $J_{4',5'} = 9.4$, $J_{5',6'} = 3.9$ Hz, H-5'), 3.45 (m, 1H, H-1 or 2), 3.33 (t, 1H, $J_{3',4'} = 9.4$ Hz, H-4'), 2.96 (dd, 1H, $J_{2',3'} = 10.5$ Hz, H-2'), 2.24 (m, 4H, 2 x COCH₂), 2.0 (m, 2H, cyclitol), 1.68 – 1.47 (m, 6H, 2 x COCH₂CH₂ and 2H cyclitol), 1.35 – 1.14 (52H, 2 x [CH₂]₁₂ and 4H cyclitol), 0.79 (t, 6H, $J = 7.1$ Hz, 2 x CH₂CH₃); δ_C (125 MHz, 1:1 CDCl₃ – MeOH-d₄) 175.3 (C=O), 174.8 (C=O), 98.0 (C-1'), 82.5 (C-1 or 2), 80.2 (C-1 or 2), 74.2 (C-5'), 71.6, 71.5, 71.4, 64.8 (CH₂ glycerol), 63.8 (CH₂ glycerol), 62.3 (C-6'), 55.7 (C-2'), 35.5, 35.4, 34.0, 33.5, 33.2, 31.0 – 30.4, 26.2, 25.3, 23.9, 15.1; δ_P (202 MHz, 1:1 CDCl₃ – MeOH-d₄) 0.10 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₅₂H₉₇NO₁₅P [M – H][–] 906.6077, found 906.6094.

N*-[(1*R*,2*R*)-2-(Benzyloxy)cyclohexyl]octadecane-1-sulphonamide **32*

To a solution of CH₂Cl₂ (10 mL) and triethylamine (2.1 mL) under argon was added (1*R*, 2*R*)-1-amino-2-benzyloxycyclohexane **31** (1.0 g, 4.87 mmol) and 1-octadecanesulfonyl chloride (2.1 g, 5.95 mmol), purchased from Sigma-Aldrich and Alfa Aesar, respectively. The reaction mixture was stirred at rt overnight, whereafter it was diluted with CH₂Cl₂ (40 mL), washed successively with water (25 mL), brine (25 mL), dried (MgSO₄) and concentrated under reduced pressure. RBC (elution first with hexane and then with a gradient of 5:1 → 3:1 hexane – EtOAc) gave the sulphonamide **32** (1.9 g, 76%) as white needles; mp 63 – 64 °C; $[\alpha]_D^{25} -35.6^\circ$ (*c* 1.00, CHCl₃); δ_H (500 MHz, CDCl₃) 7.40 – 7.20 (m, 5H, Ph), 4.68 (d, 1H, $J = 1.4$ Hz, OCH₂), 4.46 (d, 1H, $J = 5.0$ Hz, NH), 3.23 – 3.11 (m, 2H, H1 and 2), 3.03 – 2.90 (m, 2H, SO₂CH₂), 2.23 (m, 2H, H3a and 6a), 1.80 – 1.60 (m, 4H, SO₂CH₂CH₂, H4a and 5a), 1.35 – 1.11 (m, 34H, 15 x [CH₂]₁₅, H3b, 4b,

5b and 6b), 0.88 (t, 3H, $J = 6.8$ Hz, CH_2CH_3); δ_{C} (125 MHz, CDCl_3) 138.1, 128.5, 127.8, 127.7, 80.1 (C1 or 2), 70.6 (OCH_2), 57.6 (C1 or 2), 53.2 (SO_2CH_2), 33.2 (C3 or 6), 31.9, 30.1 (C3 or 6), 29.7, 29.5, 29.4, 29.1, 28.3, 24.2, 23.8, 23.5, 22.7, 14.1 (CH_2CH_3); HRMS (ESI) calcd. for $\text{C}_{31}\text{H}_{56}\text{NO}_3\text{S}$ $[\text{M} + \text{H}]^+$ 522.3975, found 522.3981.

N*-[(1*R*,2*R*)-2-Hydroxycyclohexyl]octadecane-1-sulphonamide **33*

A solution of the benzyloxysulphonamide **32** (200 mg, 0.38 mmol) in 5:1 THF – AcOH (6 mL) containing 10 – 20% $\text{Pd}(\text{OH})_2$ on carbon (50 mg) was stirred under a slight over pressure of hydrogen at room temperature for 2 h before it was percolated through a short column of Celite on a bed of silica gel (further elution with EtOAc). The eluent was concentrated under reduced pressure to give the deprotected alcohol **33** (140 mg, 85%) as a white solid which was used without any further purification; mp 101 – 102 °C; $[\alpha]_{\text{D}}^{25} -7.3$ (c 3.40, CHCl_3); δ_{H} (500 MHz, CDCl_3) 4.54 (d, 1H, $J = 7.4$ Hz, NH), 3.32 (m, 1H, H2), 3.15 – 3.02 (m, 3H, SO_2CH_2 and H1), 2.54 (brs, 1H, OH), 2.12 – 2.03 (m, 2H, H3a and 6a), 1.90 – 1.63 (m, 4H, $\text{SO}_2\text{CH}_2\text{CH}_2$, H4a and 5a), 1.50 – 1.10 (m, 34H, $[\text{CH}_2]_{15}$, H3b, 4b, 5b and 6b), 0.88 (t, 3H, $J = 6.8$ Hz, CH_2CH_3); δ_{C} (125 MHz, CDCl_3) 73.8 (C2), 59.8 (C1), 53.5 (SO_2CH_2), 34.1 (C3 or 6), 31.9, 29.7, 29.6, 29.5, 29.3, 29.1, 28.3, 24.8 (C4 or 5), 24.0 (C4 or 5), 23.7, 14.1 (CH_2CH_3); HRMS (ESI) calcd. for $\text{C}_{24}\text{H}_{50}\text{NO}_3\text{S}$ $[\text{M} + \text{H}]^+$ 432.3506, found 432.3487.

N*-(1*R*,2*R*)-2-*O*-(2-Azido-3,4,6-tri-*O*-benzyl-2-deoxy-*D*-glucopyranosyl)-cyclohexyloctadecane-1-sulphonamide **35*

A mixture of trichloroacetimidate **34**²⁰ (130 mg, 0.21 mmol), acceptor **33** (109 mg, 0.25 mmol) and activated 4 Å molecular sieves (200mg) in dry CH_2Cl_2 (15 mL) was stirred under argon at room temperature for 15 min. Then TMSOTf (5.3 μL , 0.029 mmol) was added and the solution was stirred at room temperature for an additional 2 h. It was then percolated through a short column of silica gel (elution with EtOAc) and the eluent was concentrated under reduced pressure. RBC (elution gradient 1:5 \rightarrow 1:3 EtOAc – hexane) of the residue gave the pseudodisaccharide **35**

(140 mg, 75%) as an oily mixture of α , β anomers in the ratio of $\sim 1:1$, as determined by ^1H NMR spectroscopy; δ_{H} (500 MHz, CDCl_3) 7.40 – 7.10 (30H, 6 x Ph α and β), 5.62 (d, 1H, $J = 2.5$ Hz, NH α or β), 5.46 (s, 1H, NH α or β), 4.98 (d, 1H, $J_{1',2'} = 3.6$ Hz, H1' α), 4.90 – 4.44 (m, 12H, 6 x CH_2Ph α and β), 4.30 (d, 1H, $J_{1',2'} = 8.0$ Hz, H1' β), 3.91 (m, 2H), 3.77 – 3.57 (m, 7H), 3.47 (m, 1H), 3.42 – 3.27 (m, 4H), 3.17 (m, 1H, H1 α/β or 2 α/β), 3.10 – 2.96 (m, 4H, 2 x SO_2CH_2), 2.37 (m, 2H, cyclitol), 2.13 (m, 2H, cyclitol), 1.90 – 1.18 (α and β $\text{SO}_2\text{CH}_2\text{CH}_2$, $[\text{CH}_2]_{15}$ and H's cyclitol), 0.88 (t, 6H, $J = 7.1$ Hz, 2 x CH_2CH_3 α and β); δ_{C} (125 MHz, CDCl_3) 137.9, 137.8, 137.6, 128.5 – 127.7 (Ph), 101.2 (C1' β), 99.6 (C1' α), 83.7, 82.7, 82.3, 81.4, 78.1, 77.5, 75.8, 75.6, 75.3, 75.1, 74.9, 73.6, 73.5, 71.4, 68.6, 68.2, 66.2, 64.7, 57.9 (C1 α/β or C2 α/β), 57.5, 53.6, 51.9, 34.0, 32.3, 32.0, 31.9, 31.4, 29.7, 29.5, 29.4, 29.3, 28.6, 28.5, 24.2, 23.8, 23.5, 23.4, 22.7, 14.2 (CH_2CH_3 α and β); HRMS (ESI) calcd. for $\text{C}_{51}\text{H}_{76}\text{N}_4\text{O}_7\text{SNa}$ $[\text{M} + \text{Na}]^+$ 911.5372, found 911.5282.

***N*-(1*R*,2*R*)-2-*O*-(2-Amino-2-deoxy- β -D-glucopyranosyl)-cyclohexyloctadecane-1-sulphonamide 11 and the α -anomer 12**

A solution of the anomeric mixture **35** (108 mg, 0.12 mmol) in 5:1 THF – AcOH (6 mL) containing 10 – 20% $\text{Pd}(\text{OH})_2$ on carbon (25 mg) was stirred under a slight over pressure of hydrogen at room temperature for 24 h before it was filtered through a bed of Celite. The catalyst was further washed with 1:1 THF – MeOH (2 x 10 mL) and the washings were combined and concentrated under reduced pressure. Column chromatography (9:1 CH_2Cl_2 – MeOH) gave first the β anomer **11** (9.8 mg, 14%) as a waxy solid; $[\alpha]_{\text{D}}^{25} +2.5$ (c 0.98, MeOH); δ_{H} (500 MHz, MeOH-d_4) 4.44 (d, 1H, $J_{1',2'} = 8.1$ Hz, H1'), 3.94 (dd, 1H, $J_{5',6'a} = 2.1$, $J_{6a',6'b} = 11.7$ Hz, H6'a), 3.65 – 3.50 (m, 2H, H1 or 2 and 6'b), 3.34 (m, 2H, H3' and 5'), 3.22 (q, 1H, $J_{3',4'} = J_{4',5'} = 9.2$ Hz, H4'), 3.09 (m, 3H, H1 or 2 and SO_2CH_2), 2.63 (t, 1H, $J_{2',3'} = 9.0$ Hz, H2'), 2.13 (m, 2H, cyclitol), 1.90 – 1.20 (38H, $\text{SO}_2\text{CH}_2\text{CH}_2$, $[\text{CH}_2]_{15}$ and 6H cyclitol), 0.89 (t, 3H, $J = 7.0$ Hz, CH_2CH_3); δ_{C} (125 MHz, MeOH-d_4) 100.7 (C1'), 80.5 (C1 or 2), 78.5 (C3' or 5'), 72.2 (C4'), 62.9 (C6'), 58.1 (C1 or 2 or 2'), 58.0 (C1 or 2 or 2'), 54.1 (SO_2CH_2), 35.4, 32.2, 30.8, 30.6, 30.5, 25.4, 25.0, 24.7, 14.5 (CH_2CH_3); HRMS (ESI)

calcd. for $C_{30}H_{61}N_2O_7S$ $[M + H]^+$ 593.4194, found 593.4178. Continued elution gave the α anomer **12** (14.5 mg, 20%) as an oil; $[\alpha]_D^{25} +66.5$ (c 1.45, MeOH); δ_H (500 MHz, MeOH- d_4) 5.28 (d, 1H, $J_{1',2'} = 3.7$ Hz, H1'), 3.81 (m, 1H, H6'a), 3.71 (m, 3H, H3', 5' and 6'b), 3.42 (m, 1H, H1 or 2), 3.34 (m, 1H, H4'), 3.23 (m, H1 or 2), 3.08 (m, 1H, H2' and SO_2CH_2), 2.24 (m, 1H, cyclitol), 1.96 (m, 1H, cyclitol), 1.84 – 1.67 (m, 4H, $SO_2CH_2CH_2$ and 2H cyclitol), 1.52 – 1.21 (34H, $[CH_2]_{15}$ and 4H cyclitol), 0.90 (t, 3H, $J = 7.0$ Hz, CH_2CH_3); δ_C (125 MHz, MeOH- d_4) 97.3 (C1'), 81.4 (C1 or 2), 73.2 (C3' or 5'), 70.4 (C3' or 4' or 5'), 70.3 (C3' or 4' or 5'), 60.8 (C6'), 56.6 (C1 or 2), 54.9 (C2'), 52.6 (SO_2CH_2), 32.7, 32.3, 31.6, 29.4, 29.3, 29.2, 29.0, 28.8, 27.8, 24.2, 23.5, 23.3, 22.3, 13.0 (CH_2CH_3); HRMS (ESI) calcd. for $C_{30}H_{61}N_2O_7S$ $[M + H]^+$ 593.4194, found 593.4181.

trans*-2-(4-Methoxybenzyloxy)cyclohexyl acetate **37*

$Cu(BF_4)_2 \cdot nH_2O$ (42 mg, 0.18 mmol) was dissolved in CH_2Cl_2 (20 mL) and cyclohexene oxide **36** (1.8 mL, 17.8 mmol) and 4-methoxybenzyl alcohol (10 mL, 80.2 mmol) were added. The reaction mixture was stirred for 24 h, diluted with water (20 mL), and the aqueous layer extracted with CH_2Cl_2 (3 x 30 mL). The combined organic extracts were washed with brine, dried over $MgSO_4$, filtered and the solvent was removed *in vacuo*. The resulting crude monoprotected PMB cyclohexanediol²² was used with no further purification in the next step, whereby it was dissolved in pyridine (7.5 mL), cooled to 0 °C, before DMAP (3 mg, 0.26 mmol) and acetic anhydride (4.5 mL, 48.0 mmol) were added. The reaction mixture was stirred overnight at rt, diluted with water (100 mL) and extracted with EtOAc (3 x 100 mL). The combined organic extracts were washed with water (100 mL), brine (100 mL), dried over $MgSO_4$, filtered and the solvent was removed under reduced pressure. Column chromatography (5:1 hexane – Et_2O) of the ensuing residue afforded the oily product **37** (1.37 g, 55%); δ_H (500 MHz, $CDCl_3$) 7.24 (d, 2H, $J = 8.7$ Hz, Ph), 6.86 (d, 2H, $J = 8.6$ Hz, Ph), 4.83 – 4.79 (m, 1H, H-1), 4.56 – 4.49 (2 x d, 2H, $J = 11.7$ Hz, CH_2Ar), 3.80 (s, 3H, OCH_3), 3.38 – 3.33 (m, 1H, H-2), 2.04 (s, 3H, CH_3), 2.02 – 1.98 (m, 2H, H-3a and 6a), 1.70 – 1.62 (m, 2H, H-4a and 5a), 1.43 – 1.18 (m, 4H, H-3b, 4b, 5b and 6b); δ_C (125 MHz, $CDCl_3$) 170.5

(C=O), 159.1, 131.0, 129.0, 114.0, 113.7, 78.4 (C-2), 75.3 (C-1), 71.0 (CH₂Ar.), 55.3 (OCH₃), 30.0, 29.9, 23.6, 23.3, 21.4 (CH₃); HRMS (ESI) calcd. for C₁₆H₂₂NaO₄ [M + Na]⁺ 301.1410, found 301.1397.

1-*trans*-2-(Allyloxy)cyclohexyloxy)methyl]-4-methoxybenzene **38**

The acetate **37** (938 mg, 3.37 mmol) was dissolved in MeOH (10 mL) and NaOMe (5.4 M in MeOH, 150 μ L) was added and the solution stirred for 1 h at rt. Afterwards, TLC revealed that there was still the presence of **37** and, thus, a further aliquot of NaOMe (5.4 M in MeOH, 100 μ L) was added and the reaction mixture was stirred for an additional 24 h. After which, the reaction was neutralised with Amberlite IR-120 (H⁺) ion-exchange resin, filtered and the crude solution was passed down a short plug of silica gel (elution with EtOAc) to afford the known 2-PMB protected alcohol²² as a pale yellow oil which was used without further purification in the next step. To a stirred and cooled (0 °C) solution of the alcohol²² (886 mg, 3.75 mmol,) in DMF (40 mL) was added NaH (60% dispersion in mineral oil, 750 mg, 18.7 mmol) and the solution was stirred for 30 min before allyl bromide (2.92 mL, 22.8 mmol) was added dropwise. The reaction mixture was stirred under argon for a further 18 h at rt, quenched with MeOH (50 mL), H₂O (250 mL) was added and then the resulting solution was extracted with EtOAc (3 x 250 mL). The combined organic extracts were washed with H₂O (3 x 250 mL), brine (250 mL), dried over MgSO₄, filtered and the solvent was removed under reduced pressure. The resulting residue was passed down a short plug of silica gel (elution with EtOAc) and, after evaporation to dryness, purified by RBC (elution gradient hexane \rightarrow 5:1 Et₂O – hexane) to afford the allyl product **38** (712 mg, 69%) as a clear oil; δ_{H} (500 MHz, CDCl₃) 7.32 (d, 2H, J = 8.6 Hz, Ph), 6.90 (d, 2H, J = 8.7 Hz, Ph), 6.02 – 5.94 (m, 1H, CH₂CH=CH₂), 5.32 – 5.17 (4 x m, 2H, CH₂CH=CH₂), 4.63 (dd, 2H, J = 11.4 Hz, CH₂Ar), 4.17 (m, 2H, CH₂CH=CH₂), 3.83 (s, OCH₃), 3.37 – 3.29 (m, 2H, H-1 and 2), 2.03–2.00 (m, 2H, cyclitol), 1.69–1.67 (m, 2H, cyclitol), 1.39–1.20 (m, 4H, cyclitol); δ_{C} (125 MHz, CDCl₃) 159.0, 135.8 (CH₂CH=CH₂), 131.5, 129.1, 116.1 (CH₂CH=CH₂), 113.7, 81.0 (C-1 or 2), 80.8 (C-1 or 2),

71.6 (CH₂CH=CH₂), 71.0 (CH₂Ar), 55.3 (OCH₃), 30.33, 30.31, 23.59, 23.58; HRMS (ESI) calcd. for C₁₇H₂₄NaO₃ [M + Na]⁺ 299.1618, found 299.1607.

2-**{[trans-2-((4-Methoxybenzyl)oxy)cyclohexyloxy]methyl} oxirane 39**

Compound **38** (2.22 g, 8.02 mmol) was dissolved in CH₂Cl₂ (30 mL) and *m*CPBA (4.15 g, 24.1 mmol) was added and the reaction mixture stirred for 18 h at rt. Afterwards, the reaction mixture was washed successively with 10% aq sodium sulfite (80 mL), water (100 mL), 10% aq NaOH (80 mL) and brine (80 mL). The organic phase was then filtered through cotton wool and the solvent was removed *in vacuo*. The crude material was passed down a short plug of silica gel (further elution with EtOAc) and evaporated to dryness under reduced pressure. RBC (elution gradient 1:1 → 2:1 Et₂O – hexane) furnished the epoxide **39** (1.50 g, 64%) as a clear oil; δ_H (500 MHz, CDCl₃) 7.20 (dd, 2H, *J* = 8.7 Hz, Ph), 6.77 (d, 2H, *J* = 8.7 Hz, Ph), 4.50 (s, 2H, CH₂Ar) 3.76 – 3.40 (5H, OCH₃ and 1- or 3-CH₂ propyl), 3.25 (m, 2H, H-1 and 2), 3.05 (m, 1H, H-2 propyl), 2.67 – 2.52 (2H, 1- or 3-CH₂ propyl), 1.90 (m, 2H, cyclitol), 1.57 (m, 2H, cyclitol), 1.22-1.06 (m, 4H, cyclitol); δ_C (125 MHz, CDCl₃) 159.0, 131.4, 131.3, 129.14, 129.08, 113.7, 82.3 (C-1 or 2), 82.1 (C-1 or 2), 80.8 (C-1 or 2), 71.5 (CH₂Ar), 71.2 (1- or 3-CH₂ propyl), 70.4 (1- or 3-CH₂ propyl), 55.2 (OCH₃), 51.3 (C-2 propyl), 51.1 (C-2 propyl), 44.44 (1- or 3-CH₂ propyl), 44.38 (1- or 3-CH₂ propyl), 30.3, 30.20, 30.16, 23.6, 23.5; HRMS (ESI) calcd. for C₁₇H₂₄NaO₄ [M + Na]⁺ 315.1567, found 315.1553.

3-**{[trans-2-((4-Methoxybenzyl)oxy)cyclohexyl]oxy}propane-1,2-diol 40**

To a solution of **39** (1.50 g, 5.13 mmol) in DMSO (56.4 mL) was added water (10.8 mL) and aq 0.3 M KOH (2.4 mL). The reaction mixture was heated to 100 °C for 18 h, and then diluted with water (200 mL) followed by extraction with CH₂Cl₂ (3 x 200 mL). The combined organic extracts were washed with water (100 mL), brine (100 mL), dried over MgSO₄, filtered and the solvent was removed under reduced pressure. The residue, so obtained, was percolated through a short column of silica gel (further elution with EtOAc) and the subsequent eluent was concentrated under reduced pressure. RBC (elution first with hexane → 6:1 EtOAc – hexane) of the residue

afforded the diol **40** (942 mg, 65) as a clear oil; δ_{H} (500 MHz, CDCl_3) 7.28 (d, 2H, $J = 8.6$ Hz, Ph), 6.88 (dd, 2H, CH, $J = 8.6$, Ph), 4.59 – 4.50 (2H, CH_2Ar), 3.82 – 3.47 (8H, OCH_3 , H-2 propyl, 1- and 3- CH_2 propyl), 3.25 (m, 2H, H-1 and 2), 2.50 (bs, 1H, OH), 2.40 (bs, 1H, OH), 2.11 – 2.01 (m, 2H, cyclitol), 1.68 (m, 2H, cyclitol), 1.23 (m, 4H, cyclitol); δ_{C} (125 MHz, CDCl_3) 159.3, 159.2, 130.5, 130.4, 129.5, 129.4, 113.84, 113.83, 83.40 (C-1 or 2), 82.39 (CH, C-1 or 2), 80.86 (C-1 or 2), 80.78 (C-1 or 2), 72.31 (1- or 3- CH_2 propyl), 71.37 (C-2 propyl), 71.03 (1- or 3- CH_2 propyl), 70.81 (1- or 3- CH_2 propyl), 70.51 (C-2 propyl), 64.20 (1- or 3- CH_2 propyl), 63.83 (1- or 3- CH_2 propyl), 55.24 (OCH_3), 30.75, 30.54, 29.99, 29.93, 23.78, 23.71, 23.69; HRMS (ESI) calcd. for $\text{C}_{17}\text{H}_{26}\text{NaO}_5$ $[\text{M} + \text{Na}]^+$ 333.1672, found 333.1678.

1-[(*tert*-Butyldiphenylsilyl)oxy]-3-[[*trans*-2-((4-methoxybenzyl)oxy)cyclohexyl]oxy]propan-2-ol **41**

To a solution of the primary alcohol **40** (942 mg, 3.34 mmol) and DIPA (5.8 mL, 3.75 mmol) in CH_2Cl_2 (5 mL) was added TBDPSCl (1.04 mL, 4.00 mmol) dropwise followed by DMAP (5 mg, 0.038 mmol) and the reaction stirred for 24 h at rt. Afterwards, TLC revealed the presence of the starting material **40**; thus an additional aliquot of TBDPSCl (0.521 mL, 2.00 mmol) was added and the reaction mixture was stirred for a further 3 h, whereafter it was quenched with water (60 mL) and then extracted with CH_2Cl_2 (3 x 60 mL). The combined organic extracts were washed successively with water (100 mL), brine (50 mL), filtered through cotton wool and the solvent was removed under reduced pressure. The residue so obtained was percolated through a short column of silica gel (further elution with EtOAc) and the subsequent eluent was concentrated under reduced pressure. RBC (elution first with hexane \rightarrow 1:2 EtOAc – hexane) of the residue afforded the silyl protected product **41** (1.33 g, 76%) as a pale yellow oil; δ_{H} (500 MHz, CDCl_3) 7.68 – 6.79 (14H, Ph), 4.56-4.44 (m, 2H, CH_2Ar), 3.90 – 3.51 (8H, OCH_3 , H-2 propyl, 1- and 3- CH_2 propyl), 3.25 (m, 2H, H-1 and 2), 3.12 (bs, 1H, OH), 1.97 (m, 2H, cyclitol), 1.66 (m, 2H, cyclitol), 1.30-1.15 (m, 4H, cyclitol), 1.05 (s, 9H, 3 x CH_3); δ_{C} (125 MHz, CDCl_3) 159.1, 135.6, 135.6, 134.7, 133.5, 133.49, 133.42, 130.9, 130.8, 129.73, 129.72, 129.3, 129.2, 127.74, 127.71, 113.79, 113.77, 82.9 (C-1 or 2),

82.3 (C-1 or 2), 80.7 (C-1 or 2), 80.6 (C-1 or 2), 71.7, 71.6, 71.2, 71.1, 71.0, 70.4, 65.0, 64.8, 55.3, 55.2, 30.6, 30.4, 30.11, 30.06, 29.7, 26.9, 23.69, 23.66, 19.29, 19.28; HRMS (ESI) calcd. for $C_{33}H_{44}NaO_5Si$ $[M + Na]^+$ 571.2856, found 571.2860.

1-[(*tert*-Butyldiphenylsilyl)oxy]-3-[(*trans*-2-[(4-methoxybenzyl)oxy]cyclohexyl)oxy]propan-2-yl methanesulfonate **42**

To the secondary alcohol **41** (1.33 g, 2.54 mmol) in pyridine (5 mL) was added mesyl chloride (0.63 mL, 8.14 mmol) dropwise. The reaction mixture was stirred at room temperature for 24 h and then quenched with saturated $NaHCO_3$ (10 mL) followed by extraction with CH_2Cl_2 (3 x 20 mL). The combined organic extracts were washed with water (60 mL), brine (60 mL), filtered through cotton wool and then evaporated to dryness; whereafter toluene (5 mL) was added and evaporated therefrom. The resulting oil was passed through a short column of silica gel (elution with EtOAc) to give a clear yellow oil after evaporation to dryness under reduced pressure. This oil was used in the following step without further purification. However, a small sample of the product (100 mg) was purified by RBC (1:1 Et_2O – hexane,) to afford a clear, colourless oil of **42** for analytical analyses; δ_H (500 MHz, $CDCl_3$) 7.60 – 6.73 (14H, Ph), 4.70 (m, 1H, H-2 propyl), 4.61–4.35 (m, 2H, CH_2Ar), 3.83 – 3.69 (m, 7H, OCH_3 , 1- and 3- CH_2 propyl), 3.18 (m, 2H, H-1 and 2), 2.90 (s, 3H, SO_2CH_3), 1.90 (m, 2H, cyclitol), 1.57 (m, 2H, cyclitol), 1.24 – 1.08 (m, 4H, cyclitol), 0.98 (s, 9H, 3 x CH_3); δ_C (125 MHz, $CDCl_3$) 159.1, 135.6, 135.54, 135.52, 132.94, 132.89, 132.8, 131.1, 131.0, 129.92, 129.90, 129.2, 129.1, 127.9, 113.78, 113.77, 82.5 (CH, CH propyl), 82.4 (CH, C-1 or 2), 82.1 (CH, C-1 or 2), 81.9 (CH, C-1 or 2), 80.5 (CH, C9), 71.2, 71.0, 68.9, 68.8, 63.5, 55.3 (OCH_3), 38.5 (SO_2CH_3), 38.4, 30.0, 30.0, 29.8, 26.8, 23.5, 23.4, 19.2; HRMS (ESI) calcd. for $C_{34}H_{46}NaO_7SSi$ $[M + Na]^+$ 649.2626, found 649.2628.

{2-Azido-3-[(*trans*-2-[(4-methoxybenzyl)oxy]cyclohexyl)oxy] propoxy} (*tert*-butyl)diphenylsilane **43**

A solution of the mesylate **42** in DMF (10 mL) containing sodium azide (496 mg, 7.64 mmol) was heated and stirred at 125 °C for 24 h, cooled and then poured into water (40 mL). The

resulting aqueous solution was extracted with CH_2Cl_2 (3 x 40 mL) and the combined organic extracts were washed successively with water (100 mL), brine (100 mL), filtered through cotton wool and concentrated under reduced pressure. A solution of the residue in EtOAc was percolated through a short column of silica gel (elution with EtOAc) and the eluent concentrated under reduced pressure. RBC of the residue (elution first with hexane \rightarrow 1:1 Et_2O – hexane) gave the azide **43** (1.30 g, 51%) as a clear oil; δ_{H} (500 MHz, CDCl_3) 7.62 – 6.73 (14H, Ph), 4.44 (s, 2H, CH_2Ar), 3.71 – 3.32 (8H, OCH_3 , H-2 propyl, 1- and 3- CH_2 propyl), 3.22 (m, 2H, H-1 and 2), 1.86 (m, 2H, cyclitol), 1.55 (m, 2H, cyclitol), 1.25 – 1.04 (m, 4H, cyclitol), 0.99 (s, 9H, 3 x CH_3); δ_{C} (125 MHz, CDCl_3) 159.0, 135.9, 135.8, 135.6, 135.3, 134.8, 133.10, 133.06, 131.31, 131.29, 129.8, 129.7, 129.2, 129.12, 129.08, 129.0, 127.8, 127.73, 127.67, 113.74, 113.70, 82.2 (C-1 or 2), 82.0 (C-1 or 2), 80.5 (C-1 or 2), 80.4 (C-1 or 2), 71.5, 71.4, 71.19, 69.5, 69.0, 64.08, 64.05, 63.2, 63.0, 55.3 (OCH_3), 30.0, 29.9, 26.9, 26.8, 26.6, 23.43, 23.40, 23.35, 19.2; HRMS (ESI) calcd. for $\text{C}_{33}\text{H}_{43}\text{N}_3\text{NaO}_4\text{Si}$ $[\text{M} + \text{Na}]^+$ 596.2915, found 596.2926.

trans*-2-{2-Azido-3-[(*tert*-butyldiphenylsilyl)oxy]propoxy}cyclohexanol **44*

The PMB derivative **43** (131 mg, 0.240 mmol) was dissolved in a solution of 1% TFA in CH_2Cl_2 (8.62 mL). The reaction mixture was stirred at room temperature for 18 h; whereafter an additional aliquot of TFA (43 μL) was added because TLC indicated the presence of the PMB protected starting material (**43**). After a further 4 h, TLC revealed the absence of any starting material (**43**) and the reaction mixture was diluted with CH_2Cl_2 (40 mL) and washed with saturated NaHCO_3 (40 mL), water (40 mL), brine (40 mL) and then filtered through cotton wool. The CH_2Cl_2 extract was concentrated under reduced pressure and the residue was percolated through a short column of silica gel (elution with EtOAc) and concentrated to dryness under reduced pressure. RBC purification of the residue (elution first with hexane \rightarrow 1:1 Et_2O – hexane) afforded the alcohol **44** (67 mg, 64%) as a clear oil; δ_{H} (500 MHz, CDCl_3) 7.60 – 7.31 (10H, Ph), 3.74 – 3.11 (6H, H-1 or 2, H-2 propyl, 1- and 3- CH_2 propyl), 2.99 (m, 1H, H-1 or 2), 2.60 (bd, 1H, OH), 1.96 (m, 2H, cyclitol), 1.65 (m, 2H, cyclitol), 1.23-1.03 (m, 4H, cyclitol), 1.00 (s, 9H, 3 x CH_3); δ_{C} (125

MHz, CDCl₃) 134.75, 134.58, 134.56, 132.2, 131.91, 131.87, 128.9, 126.81, 126.79, 83.6 (C-1 or 2), 83.5 (C-1 or 2), 72.8, 72.7, 67.3, 67.0, 62.8, 62.7, 62.0, 61.7, 31.02, 30.98, 28.7, 28.1, 25.9, 25.7, 23.1, 22.9, 18.2; HRMS (ESI) calcd. for C₂₅H₃₅N₃NaO₃Si [M + Na]⁺ 476.2340, found 476.2351.

Triethylammonium *trans*-2-{2-azido-3-[(*tert*-butyldiphenylsilyl)oxy]propoxy}cyclohexyl *n*-octadecyl phosphate 45

This compound was obtained from the alcohol **44** (280 mg, 0.62 mmol) and the hydrogenphosphonate TEA salt **18**¹⁵ (537 mg, 1.23 mmol) in the presence of pivaloyl chloride (0.48 mL, 3.86 mmol) essentially as described for the TEA salt **19**. After oxidation with iodine (623 mg, 2.47 mmol) in 9:1 pyridine – water followed by the same aqueous workup as described for **19**, column chromatography (CH₂Cl₂ → 8:1 CH₂Cl₂ – MeOH) of the residue afforded the octadecyl phosphate TEA salt **45** (276 mg, 50%) as a yellow paste; δ_H (500 MHz, CDCl₃) 7.63 – 7.30 (10H, Ph), 4.10 (m, 1H, H-1 or 2), 3.92 – 3.33 (7H, OCH₂, H-2 propyl, 1- and 3-CH₂ propyl), 3.28 (m, 1H, H-1 or 2), 3.00 (m, 6H, 3 x CH₂CH₃), 1.95 – 1.73 (m, 2H, cyclitol), 1.58 – 1.46 (m, 4H, OCH₂CH₂ and 2H cyclitol), 1.35 – 1.14 (43H, [CH₂]₁₅, 3 x CH₂CH₃ and 4H cyclitol), 1.73 (s, 9H, 3 x CH₃), 0.80 (t, 3H, CH₃, *J* = 6.8 Hz, CH₂CH₃); δ_C (125 MHz, CDCl₃) 134.9, 134.8, 134.6, 132.6, 132.4, 132.1, 132.0, 128.9, 128.8, 126.8, 126.7, 78.8 (C-1 or 2), 78.6 (C-1 or 2), 76.6 (C-1 or 2), 70.8, 70.1, 68.2, 67.7, 65.7, 65.6, 65.5, 63.0, 62.9, 44.5 [N(CH₂CH₃)₃], 30.9, 29.7, 29.6, 29.5, 29.0, 28.70, 28.65, 28.4, 27.2, 26.0, 25.9, 25.7, 24.8, 24.7, 21.7, 21.1, 21.0, 18.2, 14.3 (CH₂CH₃), 8.5 [N(CH₂CH₃)₃]; δ_P (202 MHz, CDCl₃) –1.2 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₄₃H₇₁N₃O₆PSi [M – NEt₃ – H][–] 784.4885, found 784.4759.

Triethylammonium *trans*-2-(2-azido-3-hydroxypropoxy)cyclohexyl *n*-octadecyl phosphate 46

Compound **45** (68 mg, 0.077 mmol) was dissolved in THF (1 mL) and 1.0 M TBAF in THF (153 μL, 0.15 mmol) was added. The reaction mixture was stirred at rt for 16 h and then diluted with water (25 mL), extracted CH₂Cl₂ (3 x 25 mL) and the combined organic extracts were washed with aq 1.0 M TEAB (2 x 10 mL). The organic phase was filtered through cotton wool, the solvent was removed under reduced pressure and the resulting residue was purified by column

chromatography (8:1 CH₂Cl₂ – MeOH) to afford the alcohol **46** (50 mg, 100%) as a white paste; δ_{H} (500 MHz, CDCl₃) 4.08 – 3.33 (8H, OCH₂, H-1 or 2, H-2 propyl, 1- and 3-CH₂ propyl), 3.25 (m, 1H, H-1 or 2), 3.09 (m, 6H, 3 x CH₂CH₃), 2.14 – 1.99 (m, 2H, cyclitol), 1.66-1.59 (m, 4H, OCH₂CH₂ and 2H cyclitol), 1.30-1.19 (43H, [CH₂]₁₅, 3 x CH₂CH₃ and 4H cyclitol), 0.88 (t, 3H, CH₃, J = 6.5 Hz, CH₂CH₃); δ_{C} (125 MHz, CDCl₃) 81.6 (C-1 or 2), 78.6 (C-1 or 2), 68.6, 68.5, 66.03, 65.99, 62.4, 61.7, 60.9, 60.6, 45.4 [N(CH₂CH₃)₃], 32.2, 32.0, 31.9, 30.8, 30.74, 30.68, 29.73, 29.67, 29.42, 29.38, 25.8, 23.9, 23.8, 23.8, 23.7, 22.7, 14.1 (CH₂CH₃), 8.5 [N(CH₂CH₃)₃]; δ_{P} (202 MHz, CDCl₃) –1.02 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₂₇H₅₃N₃O₆P [M – NEt₃ – H][–] 546.3677, found 546.3673.

Triethylammonium *trans*-2-(2-amino-3-hydroxypropoxy)cyclohexyl *n*-octadecyl phosphate **13**

Pearlman's catalyst [10 – 20% Pd(OH)₂ on carbon, 15 mg] was added to a solution of the azide **46** (45 mg, 0.069 mmol) in 1:1 THF – MeOH (10 mL) and the mixture was stirred under a hydrogen atmosphere at rt for 2 h. Processing as described for **21** gave the amino TEA salt **13** (29 mg, 44%) as a white paste, which did not require any chromatographic purification; δ_{H} (500 MHz, CDCl₃) 4.00 – 3.31 (8H, OCH₂, H-1 or 2, H-2 propyl, 1- and 3-CH₂ propyl), 3.10 (m, 1H, H-1 or 2), 3.02 (q, 6H, J = 7.4 Hz, 3 x CH₂CH₃), 2.04 – 1.93 (m, 2H, cyclitol), 1.63 – 1.50 (m, 4H, OCH₂CH₂ and 2H cyclitol), 1.33-1.05 (43H, [CH₂]₁₅, 3 x CH₂CH₃ and 4H cyclitol), 0.81 (t, 3H, CH₃, J = 6.7 Hz, CH₂CH₃); δ_{C} (125 MHz, CDCl₃) 81.6 (C-1 or 2), 77.9 (C-1 or 2), 67.1, 65.9, 64.8, 53.2, 44.5 [N(CH₂CH₃)₃], 33.0, 31.8, 30.9, 29.7, 29.6, 29.33, 29.29, 29.0, 28.70, 28.65, 28.43, 28.35, 27.9, 24.9, 24.8, 24.8, 24.7, 23.2, 23.1, 22.8, 22.7, 13.1 (CH₂CH₃), 7.5 [N(CH₂CH₃)₃]; δ_{P} (202 MHz, CDCl₃) –0.23 (with heteronuclear decoupling); HRMS (ESI) calcd. for C₂₇H₅₅NO₆P [M – NEt₃ – H][–] 520.3772, found 520.3747.

Biological assays

Materials

The synthesis of 1-D-6-*O*-(2-amino-2-deoxy- α -D-glucopyranosyl)-*myo*-inositol 1-(octadecyl phosphate), (**4**, α -D-GlcpNH₂-IPC₁₈),¹⁵ has been described previously. The corresponding *N*-acetyl derivate α -D-GlcpNAc-IPC₁₈ (**3**) was prepared by treatment with acetic anhydride¹³, and the concentration of stock solutions determined by measurement of the inositol content by selected ion-monitoring GC-MS.⁸ Bloodstream form *Trypanosome brucei* (variant MITat1.4) were isolated and membranes (cell-free system) prepared as described previously and stored at -80 °C.²⁶

Activity assays

Substrate recognition assays were performed using 500 pmol of α -D-GlcpNAc-PI (**1**) in incorporation buffer (25 mM Tris pH 8.0, 50 mM KCl, 50 mM MnCl₂) and varying amounts of trypanosome cell-free system (0 - 15×10^6 cell equivalents per assay) in 96-well plates containing 100 μ L final volume, and incubated at 37 °C for 1 h. The reaction was quenched and the glycolipids enriched and analyzed by LC-MS/MS as described below.

Inhibition assays were performed in 96-well plates in 100 μ L final volume, with 1% v/v DMSO with or without inhibitor. Trypanosome cell-free system (2.5×10^6 cell equivalents per assay) in incorporation buffer were added to wells containing 500 pmol α -D-GlcpNAc-IPC₁₈ (**3**) with or without inhibitor and incubated at 37 °C for 1h. The reaction was quenched and the glycolipids enriched and analyzed by LC-MS/MS as described below.

Glycolipid enrichment

Enrichment of glycolipids was performed in a 96-well plate format. Reactions were quenched by addition of 200 μ L of 5% propan-1-ol, 5 mM NH₄OAc, and the glycolipids were bound to C₁₈ resin (50 mg Isolute Array cartridge), washed three times with 200 μ L 5% propan-1-ol, 5 mM NH₄OAc and eluted with 100 μ L 40% propan-1-ol, 5 mM NH₄OAc into a 96-well collection plate.

Prior to subsequent analysis, compound **13** was dried under nitrogen, resuspended in MeOH (100 μ L) and any free amine reacted with excess d_6 -Ac₂O (1.5 μ L) in the presence of pyridine (10 μ L) for 15 min. The reaction was quenched with water (50 μ L), dried under nitrogen and resuspended in 100 μ L 40% propan-1-ol, 5 mM NH₄OAc.

Liquid chromatography - tandem mass spectrometry of glycolipids

Glycolipids were analyzed by liquid chromatography coupled to an electrospray tandem mass spectrometer (LC-MS/MS). Samples (40 μ L) were injected directly from a 96-well plate onto a 10 \times 1 mm C₁₈ column (ACE, 5 μ M) and then eluted using a binary gradient of 5 - 80% propan-1-ol in 5 mM NH₄OAc (Dionex Ultimate 3000). The gradient consisted of 2 min 0% B, 2 - 4 min 0-100% B, 4 - 8 min 100%, 8 - 9 min 100-0% B, 9 - 10 min 0% B where buffer A consisted of 5% propan-1-ol, 5 mM NH₄OAc and buffer B 80% propan-1-ol, 5 mM NH₄OAc. The glycolipids were analysed on an electrospray triple quadrupole mass spectrometer (Micromass Quattro Ultima) in multiple reaction monitoring mode.

For each pseudodisaccharide analogue (**7** – **12**), standards of the *N*-acetylated compound and corresponding free amine were analyzed separately in order to identify unique transitions for use in subsequent multiple reaction monitoring experiments (Table 1). The ratio of the integrals for these transitions was used to calculate the percentage of substrate conversion to product in a given sample. For compound **13**, standards of the *N*-acetylated compound and the d_3 -*N*-acetylated form were analyzed separately and found to produce a common fragment for use in subsequent multiple reaction monitoring experiments.

For inhibition assays, the turnover of the substrate α -D-GlcpNAc-IPC₁₈ (**3**) was used to calculate the percentage of substrate conversion to product in a given sample.¹⁰ Inhibitor IC₅₀ values were calculated using a four-parameter fit of eight-point potency curves derived from three independent experiments, and are quoted with a standard deviation.

Trypanosome cell-free system assays

The formation of GPI precursors is monitored by following the incorporation of [³H]-mannose and then they were analysed using high-performance liquid chromatography and fluorography as described previously.⁸

Acknowledgements

We would like to acknowledge the Wellcome Trust (programme grant 085622 and strategic awards 08348 and 100476) and the MRC (studentship to ASC) for financial support.

Supplementary material

Supplementary data (additional experimental procedures and characterisation data for the β -anomers **8** and **10** plus ¹H and ¹³C NMR spectra of all the compounds) associated with this article can be found, in the online version, at XXX.

References

1. M. A. J. Ferguson, *J. Cell Sci.*, 1999, **112**, 2799-2809.
2. D. K. Sharma, T. K. Smith, C. T. Weller, A. Crossman, J. S. Brimacombe and M. A. J. Ferguson, *Glycobiology*, 1999, **9**, 415-422.
3. T. K. Smith, M. J. Patterson, A. Crossman, J. S. Brimacombe and M. A. J. Ferguson, *Biochemistry*, 2000, **39**, 11801-11807.
4. T. K. Smith, A. Crossman, C. N. Borrisow, M. J. Paterson, A. Dix, J. S. Brimacombe and M. A. J. Ferguson, *EMBO J.*, 2001, **20**, 3322-3332.

5. M. D. Urbaniak, A. Crossman and M. A. J. Ferguson, *Chem. Biol. Drug Des.*, 2008, **72**, 127-132.
6. M. D. Urbaniak, D. V. Yashunsky, A. Crossman, A. V. Nikolaev and M. A. J. Ferguson, *ACS Chem. Biol.*, 2008, **3**, 625-634.
7. N. Z. Abdelwahab, A. T. Crossman, M. D. Urbaniak and M. A. J. Ferguson, *Carbohydr. Res.*, 2011, **346**, 708-714.
8. N. Z. Abdelwahab, A. T. Crossman, L. Sullivan, M. A. J. Ferguson and M. D. Urbaniak, *Chem. Biol. Drug Des.*, 2012, **79**, 270-278.
9. D. K. Sharma, T. K. Smith, A. Crossman, J. S. Brimacombe and M. A. J. Ferguson, *Biochem. J.*, 1997, **328**, 171-177.
10. M. D. Urbaniak, A. Crossman, T. Chang, T. K. Smith, D. M. F. v. Aalten and M. A. J. Ferguson, *J. Biol. Chem.*, 2005, **280**, 22831-22838.
11. T. Chang, K. G. Milne, M. L. S. Guthrie, T. K. Smith and M. A. J. Ferguson, *J. Biol. Chem.*, 2002, **277**, 50176-50182.
12. N. A. Meanwell, *J. Med. Chem.*, 2011, **54**, 2529-2591.
13. T. K. Smith, S. Cottaz, J. S. Brimacombe and M. A. J. Ferguson, *J. Biol. Chem.*, 1996, **271**, 6476-6482.
14. G. Grundler and R. R. Schmidt, *Liebigs Ann. Chem.*, 1984, 1826-1847.
15. A. Crossman, M. J. Patterson, M. A. J. Ferguson, T. K. Smith and J. S. Brimacombe, *Carbohydr. Res.*, 2002, **337**, 2049-2059.
16. A. V. Nikolaev, I. A. Ivanova, V. N. Shibaev and N. K. Kochetkov, *Carbohydr. Res.*, 1990, **204**, 65-78.
17. H. F. Russell, J. B. Bremner, J. Bushelle-Edghill, M. R. Lewis, S. R. Thomas and F. Bates, II, *Tetrahedron Lett.*, 2007, **48**, 1637-1639.
18. Y. G. Gololobov, I. N. Zhmurova, L. F. Kasukhin, *Tetrahedron*, 1981, **37**, 437-472.
19. I. Lindh and J. Stawinski, *J. Org. Chem.*, 1989, **54**, 1338-1342.

20. S. Cottaz, J. S. Brimacombe and M. A. J. Ferguson, *Carbohydr. Res.*, 1995, **270**, 85-91.
21. J. Barluenga, H. Vázquez-Villa, A. Ballesteros, J. M. Gonzalez, *Org. Lett.* 2002, **4**, 2817-2819.
22. A. Tschöp, A. Marx, A. R. Sreekanth, C. Schneider, *Eur. J. Org. Chem.* 2007, **14**, 2318-2327.
23. A. S. Capes, A. T. Crossman, L. A. Webster, M. A. J. Ferguson and I. H. Gilbert, *Tetrahedron Lett.*, 2011, **52**, 7091-7094.
24. G. Berti, B. Macchia, F. Macchia, *Tet. Lett.* 1965, **38**, 3421-3427.
25. M. L. S. Guthrie and M. A. J. Ferguson, *EMBO J.*, 1995, **14**, 3080-3093.
26. W. J. Masterson, T. L. Doering, G. W. Hart and P.W.Englund, *Cell*, 1989, **62**, 73-80.