


Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

ARTICLE

Thionoglycine as a multifunctional spectroscopic reporter of screw-sense preference in helical foldamers

Cite this: DOI: 10.1039/x0xx00000x

Matteo De Poli^a and Jonathan Clayden^{a*}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

A single thionoglycine (glycine thioamide, $-\text{HNCH}_2\text{C}(=\text{S})-$) residue inserted into a peptide foldamer provides both a pair of germinal protons for use as a ^1H NMR stereochemical probe and a chromophore giving rise to a well defined Cotton effect in CD. Comparison of the response of these two features to a local helically chiral environment validates them as independent methods for quantifying the conformational screw-sense preference of a helical oligomer, in this case a peptide made of repeated Aib units. The sign of the Cotton effect provides a measure of the sign of the screw-sense preference, while both the chemical shift separation of the anisochronous signals of the glycine CH_2 group and the magnitude of the Cotton effect give an estimate of the helicity excess of the oligomer. The thionoglycine unit is readily introduced synthetically by a thionation of a BocGlyAibOMe dipeptide.

Introduction

Foldamers are extended molecular structures with well defined global conformations.¹⁻⁴ Many foldamers are helical,^{4, 5} and such structures fall into two conceptual classes. Oligomeric helical foldamers built from *chiral* monomers typically prefer a single screw sense because the two screw-sense conformers are diastereoisomers, one being of higher energy than the other. By contrast the oligomers of *achiral* monomers, though they may adopt a helical conformation, must necessarily populate equally left (*M*) and right (*P*) handed screw-sense conformers. In some cases the screw senses isomers may interconvert slowly, even to the point of being separable,⁶⁻⁸ while in others the racemisation rate is fast,⁹ even on the NMR timescale.¹⁰

The equilibrium between these screw-sense conformers of achiral helical foldamers may be perturbed in a number of ways – by incorporation of a chiral residue in the achiral chain, for example, or by covalent or non-covalent interactions of chiral ligands with the body or terminus of the chain. We have used such strategies as a means of communicating information through foldamers by a mechanism analogous to that exploited by the G-protein coupled receptor, or for achieving remote stereocontrol over up to 60 bonds.¹¹

Classically, screw-sense preference has been analysed by circular dichroism (CD), which is a particularly powerful method for assigning screw sense as left or right handed.¹² CD may also be quantitative, provided a maximum value of molar ellipticity characteristic of a compound with a complete preference for a single screw sense may be determined.^{13, 14} However, in many cases this value is difficult to estimate accurately, and CD spectra may be difficult to interpret quantitatively if other absorption bands, for example from chiral residues, overlap with those characteristic of the helix itself.¹⁵

We have reported an alternative, practically simple, method for quantifying screw-sense preference which relies on the behaviour of a pair of diastereotopic groups located within the foldamer chain under conditions where the rate of interconversion of the screw-sense conformers is either slow or fast on the NMR timescale.¹⁶⁻¹⁹ At slow exchange, the diastereotopic groups appear as a pair of anisochronous signals by virtue of their location in the chiral environment of the helix. At fast exchange, the diastereotopic groups may still appear anisochronous, but only if the local chiral environment is maintained by the existence of an imbalance between the helical screw-sense conformers. We proposed that the ratio of the anisochronicity of the two signals at slow and fast exchange is directly proportional to this screw-sense preference, expressed as a 'helical excess'. Using the NMR signals arising from ^{13}C -labelled Me groups, we explored this hypothesis by using line shape analysis at a range of temperatures, and extracted from these data thermodynamic and kinetic parameters for screw-sense inversion.¹⁷ Importantly, the value of the equilibrium constant *K* extracted by this method matched closely the value obtained simply from the quotient of the anisochronicities at fast and slow exchange.

More convenient than relying on a specifically synthesised and incorporated ^{13}C - (or ^{19}F -)¹⁹ labelled residue is the use of a pair of diastereotopic protons present in a methylene group – for example the protons of a glycine residue.¹⁸ On the assumption that these diastereotopic signals behave in a comparable manner to those of the ^{13}C -labelled Aib residue, we have used the AB system arising from the coupled anisochronous signals of the CH_2 group of a glycine residue to quantify the different levels of screw-sense control from the N- and C-terminals of a peptide helix,¹⁶ and to compare different controllers of helicity with one another.¹⁸ We have also used the CH_2 groups of related C-terminal amino alcohol residues to


assess the decay of conformational preference over extended foldamer lengths.²⁰

However, while these studies gave values that were self-consistent,¹⁹ we have thus far never demonstrated that the degree of screw-sense control measured by NMR may be correlated directly with the degree of screw-sense control observed by circular dichroism. In this paper we do this by making use of a thionated glycine residue ($-\text{NHCH}_2(\text{C}=\text{S})-$) as a reporter of screw-sense preference. Among the possible peptide bond modifications, thioamides are one of the closest mimics of a peptide linkage. These 'endothiopeptides' have received much attention because of their interesting physical and chemical properties. For example: (1) the thioamide C–N bond undergoes reversible *cis/trans* photoisomerization upon irradiation at 260 nm;^{21, 22} (2) the S atom of a thioamide is a weaker hydrogen bond acceptor than the O atom of an amide, while the more acidic adjacent NH is a stronger hydrogen bond donor;²³ (3) endothiopeptides have been used in structure-activity relationship studies to alter biological activities;^{24, 25} (4) thioamides can act as isosteric peptide bond fluorescence quenchers.²⁶

Thionoglycine contains both a chromophore suitable as a CD reporter in the form of a thioamide group and a pair of potentially diastereotopic ¹H nuclei for use as an NMR reporter, directly comparable with the ¹H NMR spectra of the related glycinamides.¹⁸ By using a thioamide, we ensure that the Cotton effect arising from this chromophore is well separated in the CD spectrum from the bands associated with the amide carbonyls themselves. Typical applications of CD measure helicity in the foldamer *structure as a whole*, in contrast with our NMR methods, which measure conformational preference at a single site.¹⁷ Here we use the thioamide function to achieve a localised CD response at a particular site in the foldamer chain.²⁷ The thionoglycine was located one residue from the C-terminus of the peptide chain, sufficiently far from any chiral residue for its chiral environment to be affected only by the local helicity of the chain, but still incorporated into the hydrogen-bonded network forming the helix.


Design and synthesis of the endothiopeptides

Aib-rich, helical^{28–30} endothiopeptides **1–6** bearing either an N-terminal Cbz group or an N-terminal acetamide (Figure 1) were made by the route shown in Scheme 1. BocGlyAibOMe **7** was thionated regioselectively at the amide carbonyl group using Lawesson's reagent^{31–33} to give the thioamide **8**. The Boc group of **8** was removed under acidic conditions^{31, 33} to provide a common precursor to **1–6**, HCl-H-Gly-ψ[CSNH]AibOMe **9** (Scheme 1),³⁴ which was coupled with range of Aib-containing oligomers by opening of their C-terminal azlactones. Thus **11**, **12** and N₃Aib₃OH gave, respectively, **1**, **2** and **13**. Cbz-protected endothiopeptides **3–6** were obtained by Staudinger reduction of azide **13** followed by N-terminal capping with a range of protected L-amino acids (L-Xxx).



compound	R ¹	R ²	R ³	L-Xxx
1	Me	H	<i>i</i> Pr	AcVal
2	Me	Me	<i>i</i> Pr	Ac(αMe)Val
3	OBn	H	<i>i</i> Pr	CbzVal
4	OBn	H	Bn	CbzPhe
5	OBn	H	Me	CbzAla
6	OBn	Me	<i>i</i> Pr	Cbz(αMe)Val

Figure 1. The endothiopeptides used in this study. A right-handed helix is shown for convenience only; see text for further explanation.



Scheme 1. Reagents and conditions: (a) Lawesson's reagent, THF, 60 °C; (b) HCl/Et₂O 2M, HSCH₂CH₂OH, anisole; (c) TFA, CH₂Cl₂; (d) EDC, MeCN; (e) **9**, *i*Pr₂NEt, MeCN; (f) PMe₃, THF, H₂O; (g) Cbz-L-AA-OH, HOAt, EDC, *i*Pr₂NEt.

NMR and CD analysis

¹H NMR spectra of the endothiopeptides **1–6** in CD₃OD³⁵ at 23 °C showed characteristic AB signals for the methylene protons, centred around 4.25 ppm. Table 1 shows the measured anisochronicity Δδ³⁶ for each Aib-containing oligomer **1–6** XxxAib₄Gly-ψ[CSNH]AibOMe carrying an N-terminal controller of screw-sense. Plotting these values against the anisochronicities previously measured in non-thionated peptides XxxAib₄GlyNH₂,¹⁸ XxxAib₄Aib**Aib₄Or-Bu,¹⁷ and XxxAib₄FibOMe¹⁹ bearing identical N-terminal protected chiral residues Xxx (Figure 2) shows that the response of the thionoglycine probe is directly proportional to that of other NMR reporters of helical screw sense, validating its applicability in this context. This proportionality also strongly suggests that the thioamide-containing foldamers adopt, at least in their Aib₄ portion, the same 3₁₀-helical conformation as that adopted by related all-amide oligomers.^{28–30}

compound	L-Xxx	marker ^a	$\Delta\delta$ Gly(CSNH)
1	AcVal	●	204
2	Ac(α Me)Val	■	195
3	CbzVal	○	135
4	CbzPhe	▼	190
5	CbzAla	▲	104
6	Cbz(α Me)Val	□	252

Table 1: Measured anisochronicities ($\Delta\delta$) in the $CH_2H_B C(=S)NH$ methylene groups of oligomers 1-6. (a) see Figure 2.




Figure 2. Comparing the performance of the glycine thioamide as a 1H NMR probe with other NMR reporters of helical screw-sense; ^afor ^{13}C Aib probe (blue), $\Delta\delta/10$; ^bmeasured anisochronicity of the signals from highlighted nuclei in the 1H , ^{13}C and ^{19}F NMR spectrum.

CD spectra of the thionated peptides 1-6 were recorded in MeOH solution, in order to keep the experimental conditions consistent with those used in the NMR experiments. The thionated peptides display a distinct Cotton effect at about 268 nm, clearly separated from the rest of the peptide bond absorptions, which fall below 240 nm (Figure 3). This strong band is assigned to the π - π^* transition of the thioamide chromophore;^{22, 37-45} it is interesting that the n - π^* band (usually lying at around 340 nm³² is hardly visible. The positions of these two bands match those observed in the UV spectra in the same solvent (see Supporting Information). Analysis of the far-UV portion of the CD spectrum, which usually reveals valuable information on helical conformations, is complicated by the contribution of the thioamide chromophore transitions at lower wavelengths.^{12, 46}




Figure 3. CD spectra of the endothiopeptides 1-6 in methanol. Inset: the region of the spectrum arising from the π - π^* transition of the thioamide chromophore.

The ability to identify the contribution of bonding at a single residue to an overall circular dichroism spectrum is a valuable result of the thionation of the glycine peptide bond. Given the spatial separation of this thioamide chromophore from the nearest stereogenic centre, the sign of the thioamide π - π^* absorption band centred at 268 nm (Figure 3, inset) must result solely from the screw-sense preference experienced in the vicinity of the thionoglycine probe. Despite the common L-configuration of all the endothiopeptides at their N-terminal stereocontrolling residue, 2 and 6, which are capped with a C $^{\alpha}$ -tetrasubstituted amino acid (Ac- or Cbz-(α Me)Val) show a positive Cotton effect, while 1 and 3-5, which are capped with C $^{\alpha}$ -tertiary, proteinogenic amino acids (Ac- and Cbz- protected Val, Phe and Ala), show a negative Cotton effect. This observation is in agreement with the reported behaviour of peptides of the sequence L-XxxAib_n, which adopt a left-handed (*M*) helical screw sense if Xxx is a *tertiary* L-amino acid and right-handed (*P*) helical screw sense if Xxx is a *quaternary* L-amino acid.^{18, 47} Previous studies of this behaviour by circular dichroism, initially identified by using an enantiomerically enriched ^{13}C isotopic label, had been frustrated by the lack of separation between the characteristic chromophores of the helical chain and those of the chiral N-terminal residue.

Circular dichroism is a powerful technique for identifying the sign of screw-sense preference (*M* or *P*), but NMR has been of more use for quantifying the magnitude of the preference because (at least for ^{13}C NMR) maximum peak separation (corresponding to 100% helical excess) may be evaluated from low temperature NMR spectra, in which screw-sense conformers are in slow exchange.⁴⁸ However, the reported values of screw-sense preference induced by each controller determined by this method can now be correlated with the absolute molar ellipticities ($[\theta]_M$) measured at 268 nm (the CSNH π - π^* absorption) (Figure 4). In confirmation that the sign and value of $[\theta]_M^{268}$ is a reporter of both sign and magnitude of screw-sense preference, these two values are proportional, with $[\theta]_M^{268} = 10.6 \times \Delta\delta$ (in ppb).¹⁷




Figure 4. Relationship between the measured absolute molar ellipticities at 268 nm and the h.e. values for peptides 1-6.

As seen in earlier work on related compounds,¹⁸ bulkier and C ^{β} -branched amino acids induce a greater level of screw-sense control, while the role played by protecting groups (amide vs. carbamate) depends on whether the stereocontrolling residue is C ^{α} -tri- or tetrasubstituted, probably because of the different type of β -turn⁴⁷ adopted at the N-terminus of the helix.

X-ray crystal structure analysis

Endothiopeptides **3** and **5**, having respectively Cbz-Val and Cbz-Ala at their N-terminus, formed crystals suitable for X-ray analysis (Figure 5). Interestingly, the unit cell contains equal numbers of peptides in *M* and *P* screw-sense conformations, indicating that both helical conformers are of similar energy in the solid state as well as in solution. Furthermore, each helix is initiated by either one of the two types of β -turn (a distorted Type II or Type III) proposed⁴⁷ to account for the induction of screw-sense preference from the helix N-terminus. It is likely that the asymmetric environment experienced by the thioamide chromophore in solution results from the population of an unequal mixture of these two diastereoisomeric conformers.




Figure 5. X-ray crystal structures of endothiopeptides **3** (top) and **5** (bottom) in their *M* (left) and *P* (right) conformations as observed in the unit cell.[‡] The *M* conformations display an N-terminal Type II β -turn while the *P* conformations display a Type III β -turn.⁴⁷ Cbz phenyl rings have been removed for clarity.

[‡] X-ray data for **3** and **5** have been deposited with the Cambridge crystallographic data centre. Deposition numbers **3**: 969434; **5**: 969435.

Conclusions

A thionoglycine residue inserted into a peptide provides a probe of the magnitude and sign of local helical screw sense, providing a qualitative and quantitative reporter of conformational preference. The versatile probe is readily introduced by selective thionation, and is responsive towards both magnetic and optical spectroscopies, with the helical excess values derived by NMR correlating closely those obtained by CD. The absolute measurement of a single thioamide chromophore contribution to the CD spectrum is made possible by virtue of its well-separated absorption wavelength, which may also permit its use in solvents that absorb in the far-uv region of the CD spectrum. Thioamides may prove useful in the future as tools for the conformational analysis of foldamers and peptide mimics embedded into membranes.

Experimental

BocGlyAibOMe⁴⁹ **7**. Isobutyl chloroformate (0.74 mL, 5.7 mmol) was added dropwise to a cooled (-15 °C) solution of BocGlyOH (1g, 5.7 mmol) and *N*-methylmorpholine (0.66 mL, 6 mmol) in dry THF (15 mL). The suspension was stirred at -15 °C for 15 mins, and a mixture of H-AibOMe-HCl⁵⁰ (1.75 g, 11.4 mmol) and *N*-methylmorpholine (1.3 mL, 11.4 mmol) in dry THF (20 mL) was added. The mixture was warmed to room temperature and left stirring overnight. After solvent evaporation, the residue was taken up in EtOAc (100 mL) and washed with KHSO₄ 5% (3 x 10 mL), sat. NaHCO₃ (3 x 10 mL) and brine (10 mL). The organic phase was dried (MgSO₄) and the solvent removed under reduced pressure. The crude product was purified by column chromatography (CHCl₃/MeOH, 97:3) to give pure BocGlyAibOMe **7** (1.44 g, 92%) as a white solid. *R*_f 0.35 (CHCl₃/MeOH, 97:3); mp = 94-96 °C; IR (ATR, cm⁻¹): 3340, 3275, 1716, 1667, 1534, 1303, 1279, 1244, 1157; ¹H NMR (300 MHz, CDCl₃) δ _H 6.72 (s, 1H), 5.21 (s, 1H), 3.74 (d, *J* = 6.0 Hz, 1H), 3.73 (s, 3H), 1.54 (s, 6H), 1.45 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ _C 175.0, 168.9, 156.3, 80.5, 56.8, 52.9, 44.9, 44.8, 44.8, 28.5, 24.9; MS (ES⁺, MeOH): 275 ([M + H]⁺, 55%), 297 ([M + Na]⁺, 100%); HRMS (ES⁺, MeOH): Calcd for C₁₂H₂₂N₂O₅Na [M + Na]⁺ 297.1426 found 297.1437.

BocGly- ψ [CSNH]AibOMe **8**. A stirred solution of BocGlyAibOMe **7** (1.410 g, 5.1 mmol) and Lawesson's reagent (2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-dithione, 1.041 g, 2.55 mmol) in dry THF (15 mL) was heated to 60 °C for 4 hours. The mixture was cooled to room temperature and the volatiles evaporated under reduced pressure. After purification by column chromatography (EtOAc/PE 2:3) endothiopeptide BocGly- ψ [CSNH]AibOMe **8** (0.87 g, 57%) was obtained as an off-white waxy solid. *R*_f 0.45 (EtOAc/PE, 2:3); mp = 96-98 °C; IR (ATR, cm⁻¹): 3324, 1741, 1667, 1520, 1427, 1409, 1286, 1256, 1153; ¹H NMR (300 MHz, CDCl₃) δ _H 8.63 (s, 1H), 5.39 (s, 1H), 4.08 (d, *J* = 6.4 Hz, 2H), 3.71 (s, 3H), 1.67 (s, 6H), 1.45 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ _C 199.0, 173.4, 156.8, 80.9, 60.0, 53.2, 52.7, 28.2, 23.9; MS (ES⁺, MeOH): (292 [M + H]⁺, 100%); HRMS

(ES⁺, MeOH): Calcd for C₁₂H₂₃N₂O₄S [M + H]⁺ 291.1373, found 291.1372.

HCl·H-Gly-ψ[CSNH]AibOMe 9. The method of Jensen *et al.* was used.³³ Anisole (1.1 mL, 1 mmol) and 2-mercaptoethanol (0.4 mL, 0.6 mmol) were added to a stirred solution of BocGly-ψ[CSNH]AibOMe **8** (565 mg, 1.95 mmol) in dry dioxane (4 mL), followed by HCl (2M solution in diethyl ether, 10 mL, 20 mmol) at room temperature. The reaction was monitored by TLC. A precipitate was observed after 45 mins. Stirring was continued for 2 h and the product was collected by filtration to give HCl·H-Gly-ψ[CSNH]AibOMe as a white solid (241 mg, 65%). *R*_f 0.05 (EtOAc/PE, 2:3); mp = 206-207 °C; IR (ATR, cm⁻¹): 3223, 3045, 2928, 2864, 1741, 1542, 1438, 1428, 1352, 1276, 1181, 1153; ¹H NMR (500 MHz, MeOD) δ_H 3.83 (s, 2H), 3.66 (s, 3H), 1.63 (s, 6H); ¹³C NMR (101 MHz, MeOD) δ_C 194.9, 174.6, 61.3, 52.9, 46.9, 24.4; MS (ES⁺, MeOH): 191 ([M + H]⁺, 100%); HRMS (ES⁺, MeOH): Calcd for C₇H₁₃N₂O₂S [M + Na]⁺ 191.0849, found 191.0848.

AcValAib₄OH 11. To a solution of AcValAib₄OrBu⁴⁷ (100 mg, 0.18 mmol) in dry CH₂Cl₂ (2 mL) TFA was added (2 mL) dropwise. The progress of the reaction was followed by TLC. After reaction completion (4 hs) the mixture was concentrated *in vacuo*. Et₂O was added to the residue (5 mL) and the solvent evaporated. This process was repeated 5 times. No further purification was required. AcValAib₄OH was obtained as a white solid. Yield: 86 mg, 96%. *R*_f 0.10 (CH₂Cl₂/MeOH, 95:5); mp = 242-245 °C (dec.). [α]_D²⁵ = +19.2; IR (ATR, cm⁻¹): 3296, 1648, 1528, 1386, 1364, 1290, 1169; ¹H NMR (300 MHz, MeOD) δ_H 8.57 (s, 1H), 8.27 (d, *J* = 5.7 Hz, 1H), 7.70 (s, 1H), 7.65 (s, 1H), 7.51 (s, 1H), 3.90 – 3.81 (m, 1H), 2.08 – 1.98 (m, 1H), 2.00 (s, 3H), 1.53 – 1.39 (m, 24H), 1.04 (d, *J* = 6.7 Hz, 3H), 1.00 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (126 MHz, MeOD) δ_C 178.4, 177.0, 176.8, 176.8, 174.6, 174.5, 174.1, 62.0, 58.1, 58.0, 58.0, 57.9, 57.3, 31.0, 27.0 - 26.8 (overlapping signals), 26.5, 26.0, 25.9, 25.0, 24.9, 24.8, 24.7, 24.4, 24.4, 24.2, 24.1, 22.5, 19.7; MS (ES⁺, MeOH): 500 ([M + H]⁺, 40%), 522 ([M + Na]⁺, 100%); HRMS (ES⁺, MeOH): Calcd for C₂₃H₄₁N₅O₇Na [M + Na]⁺ 522.2904, found 522.2906.

AcValAib₄Gly-ψ[CSNH]AibOMe 1. To a mixture of AcValAib₄OH **11** (60 mg, 0.12 mmol) and HOAt (1-hydroxy-7-azabenzotriazole, 18 mg, 0.13 mmol) in a mixture of dry CH₂Cl₂ (3 mL) and dry DMF (0.4 mL), was added EDC·HCl (25 mg, 0.11 mmol). After stirring for 20 mins, a solution of HCl·H-Gly-ψ[CSNH]AibOMe **9** (28 mg, 0.12 mmol) and *N,N*-diisopropylethylamine (0.061 mL, 0.35 mmol) in dry CH₂Cl₂ (2 mL) was added. The mixture was left stirring at room temperature for 6 days, after which the solvent was evaporated. The residue taken up in EtOAc (10 mL) and washed with KHSO₄ 5% (3 x 2 mL), sat. NaHCO₃ (3 x 2 mL) and brine (2 mL). The organic phase was dried (MgSO₄) and the solvent removed *in vacuo*. The crude product was purified by column chromatography (CHCl₃/MeOH, 95:5) to give pure AcValAib₄Gly-ψ[CSNH]AibOMe (45 mg, 52%) as a white

solid. *R*_f 0.35 (CH₂Cl₂/MeOH, 95:5); mp = 116-117 °C; [α]_D²⁵ = -29.6; IR (ATR, cm⁻¹): 3273, 1650, 1530, 1416, 1362, 1226, 1156; ¹H NMR (500 MHz, MeOD) δ_H 7.82 (s, 1H), 4.31 (d, A of AB, *J* = 17.7 Hz, 1H), 4.11 (d, B of AB, *J* = 17.7 Hz, 1H), 3.87 (d, *J* = 8.1 Hz, 1H), 3.63 (s, 3H), 2.06 – 1.98 (m, 1H), 2.00 (s, 3H), 1.68 (s, 3H), 1.67 (s, 3H), 1.53 – 1.34 (m, 24H), 1.04 (d, *J* = 6.7 Hz, 3H), 1.01 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (101 MHz, MeOD) δ_C 199.9, 177.9, 177.9, 177.9, 176.8, 175.5, 174.6, 174.1, 61.9, 61.4, 58.3, 58.1, 57.9, 57.9, 52.9, 52.3, 31.0, 30.9, 26.7, 26.6, 26.5, 26.4, 25.0 - 24.8 (overlapping signals), 24.3, 24.2, 22.5, 19.7, 19.6; MS (ES⁺, MeOH): 694 ([M + Na]⁺, 100%); HRMS (ES⁺, MeOH) *m/z* calcd. for C₃₀H₅₃N₇O₈NaS [M + Na]⁺ 694.3569, found 694.3555.

Ac(αMe)ValAib₄Gly-ψ[CSNH]AibOMe 2. To a mixture of Ac(αMe)ValAib₄OH⁴⁷ (74 mg, 0.14 mmol) and HOAt (21 mg, 0.15 mmol) in a mixture of dry CH₂Cl₂ (3 mL) and dry DMF (0.4 mL), EDC·HCl (*N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride, 30 mg, 0.16 mmol) was added. After stirring for 20 mins, a solution of HCl·H-Gly-ψ[CSNH]AibOMe **9** (33 mg, 0.14 mmol) and *N,N*-diisopropylethylamine (0.080 mL, 0.46 mmol) in dry CH₂Cl₂ (2 mL) was added. The mixture was left stirring at room temperature for 7 days, after which the solvent was evaporated. The residue taken up in EtOAc (10 mL) and washed with KHSO₄ 5% (3 x 2 mL), sat. NaHCO₃ (3 x 2 mL) and brine (2 mL). The organic phase was dried (MgSO₄) and the solvent removed *in vacuo*. The crude product was purified by column chromatography (CHCl₃/MeOH, 97:3 → 95:5) to give pure Ac(αMe)ValAib₄Gly-ψ[CSNH]AibOMe (54 mg, 55%) as a white solid. *R*_f 0.7 (CH₂Cl₂/MeOH, 9:1); mp = 191-193 °C; [α]_D²⁵ = +31.2; IR (ATR, cm⁻¹): 3283, 1654, 1530, 1418, 1286; ¹H NMR (400 MHz, MeOD) δ_H 7.76 (s, 1H), 4.22 (d, A of AB, *J* = 17.7 Hz, 1H), 4.02 (d, B of AB, *J* = 17.7 Hz, 1H), 3.53 (s, 3H), 1.98 – 1.89 (m, 1H), 1.93 (s, 3H), 1.58 (s, 3H), 1.57 (s, 3H), 1.45 – 1.26 (m, 24H), 0.92 (d, *J* = 6.8 Hz, 3H), 0.88 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (126 MHz, MeOD) δ_C 199.8, 178.2, 177.8, 177.7, 177.1, 175.3, 175.1, 173.7, 63.9, 61.3, 58.2, 58.0, 57.9, 57.8, 52.7, 52.2, 36.2, 26.5, 26.4, 26.4, 24.9, 24.7, 24.6 - 24.2 (overlapping signals), 23.3, 18.6, 17.9, 17.7; MS (ES⁺, MeOH): 708 ([M + Na]⁺, 100%). HRMS (ES⁺, MeOH) *m/z* calcd. for C₃₁H₅₅N₇O₈NaS [M + Na]⁺ 708.3725, found 708.3712.

N₃Aib₄Gly-ψ[CSNH]AibOMe 13.

1) *Azlactone formation.* N₃Aib₄OH (500 mg, 1.3 mmol) was dissolved in dry CH₂Cl₂ (5 mL) and EDC (0.252 mL, 1.43 mmol) was added. The resulting solution was left stirring for 3 h at room temperature. The solvent was evaporated *in vacuo* and the residue taken up in EtOAc (20 mL) and washed with KHSO₄ 5% (3 x 5 mL). The organic phase was dried (MgSO₄) and concentrated to give the crude azlactone product, which was used directly in the next step.

2) *Azlactone coupling.* The crude azlactone was dissolved in dry CH₂Cl₂ (5 mL) and HCl·H-Gly-ψ[CSNH]AibOMe **9** (230 mg, 1.01 mmol) was added, together with *N,N*-

diisopropylethylamine (0.232 mL, 1.33 mmol). The mixture was left stirring at room temperature for 7 days, after which the solvent was evaporated under reduced pressure. The residue was taken up in EtOAc (10 mL) and washed with KHSO₄ 5% (3 x 2 mL), sat. NaHCO₃ (3 x 2 mL) and brine (2 mL). The organic phase was dried (MgSO₄) and the solvent removed *in vacuo*. The crude product was purified by column chromatography (CH₂Cl₂/MeOH, 97:3 → 95:5) to give pure N₃Aib₄Gly-ψ[CSNH]AibOME (250 mg, 44%) as a colourless solid. *R*_f 0.75 (CH₂Cl₂/MeOH, 95:5); mp = 190-192 °C (dec.); IR (ATR, cm⁻¹): 3277, 2112, 1745, 1657, 1530, 1382, 1275, 1152; ¹H NMR (500 MHz, CDCl₃) δ_H 8.68 (s, 1H), 7.76 (t, *J* = 6.2 Hz, 1H), 7.36 (s, 1H), 6.93 (s, 1H), 6.28 (s, 1H), 4.31 (d, *J* = 6.2 Hz, 2H), 3.64 (s, 3H), 1.67 (s, 6H), 1.49 (s, 12H), 1.44 (s, 6H), 1.40 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ_C 198.6, 175.0, 174.2, 173.9, 173.7, 173.3, 64.0, 60.3, 57.3, 57.0, 56.9, 52.4, 51.7, 25.5 - 25.4 (overlapping signals), 24.9, 24.4; MS (ES⁺, MeOH): 579 ([M + Na]⁺, 100%); HRMS (ES⁺, MeOH) *m/z* calcd. for C₂₃H₄₀N₈O₆NaS [M + Na]⁺ 579.2689, found 579.2697.

H-Aib₄Gly-ψ[CSNH]AibOME 14. N₃Aib₄Gly-ψ[CSNH]AibOME **13** (112 mg, 0.2 mmol) was dissolved in distilled THF (2 mL). Water (0.018 mL, 1 mmol) was added, followed by PMe₃ (1M solution in THF, 2 mL, 2 mmol). The reaction was stirred overnight at room temperature. The volatiles were removed under vacuum and the residue purified by column chromatography (CH₂Cl₂/MeOH, 9:1 → 8:2) to give H-Aib₄Gly-ψ[CSNH]AibOME (85 mg, 81%). *R*_f 0.10 (CH₂Cl₂/MeOH, 95:5); mp = 212-213 °C; IR (ATR, cm⁻¹): 3278, 1730, 1651, 1530, 1383, 1362, 1282, 1221, 1155; ¹H NMR (500 MHz, CDCl₃) δ_H 8.73 (s, 1H), 8.22 (s, 1H), 7.81 (t, *J* = 6.2 Hz, 1H), 7.69 (s, 1H), 6.36 (s, 1H), 4.31 (d, *J* = 6.2 Hz, 2H), 3.64 (s, 3H), 1.75 (brs, 2H), 1.68 (s, 6H), 1.50 (s, 6H), 1.42 (s, 6H), 1.39 (s, 6H), 1.32 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ_C 198.7, 175.4, 174.8, 174.4, 174.1, 174.1, 60.4, 57.3, 56.7, 56.6, 55.5, 52.5, 51.8, 28.0, 25.5, 25.0, 24.4, 18.7, 17.7; MS (ES⁺, MeOH): 553 ([M + Na]⁺, 100%); HRMS (ES⁺, MeOH) *m/z* calcd. for C₂₃H₄₂N₆O₆NaS [M + Na]⁺ 553.2784, found 553.2775.

CbzValAib₄Gly-ψ[CSNH]AibOME 3. To a mixture of CbzValOH (31 mg, 0.12 mmol), HOAt (17 mg, 0.12 mmol) and H-Aib₄Gly-ψ[CSNH]AibOME **14** (33 mg, 0.06 mmol) in dry CH₂Cl₂ (3 mL) was added EDC (0.022 mL, 0.12 mmol). The mixture was left stirring at room temperature for 3 days, then the solvent was evaporated. The residue was taken up in EtOAc (5 mL) and washed with KHSO₄ 5% (3 x 1 mL), sat. NaHCO₃ (3 x 1 mL) and brine (1 mL). The organic phase was dried (MgSO₄) and the solvent removed *in vacuo*. The crude product was purified by column chromatography (EtOAc/PE, 4:1) to give pure CbzValAib₄Gly-ψ[CSNH]AibOME (35 mg, 74%) as a white solid. *R*_f 0.65 (EtOAc/PE, 4:1); mp = 198-200 °C; [α]_D²⁵ = -22.4; IR (ATR, cm⁻¹): 3279, 2983, 1651, 1529, 1284, 1226, 1157; ¹H NMR (500 MHz, MeOD) δ_H 7.47 - 7.23 (m, 5H), 5.11 (s, 2H), 4.28 (d, *J* = 17.7 Hz, 1H), 4.14 (d, *J* =

17.6 Hz, 1H), 3.75 (d, *J* = 7.4 Hz, 1H), 3.63 (s, 3H), 2.08 - 1.97 (m, 1H), 1.67 (s, 6H), 1.52 - 1.32 (m, 24H), 1.02 (d, *J* = 7.0 Hz, 3H), 1.00 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (126 MHz, MeOD) δ_C 199.8, 177.7, 177.7, 177.7, 176.6, 175.3, 174.6, 159.0, 138.2, 129.6, 129.1, 128.7, 67.7, 62.8, 61.3, 58.1, 58.0, 57.7, 57.7, 52.7, 52.2, 31.2, 26.2, 26.2, 26.0, 25.9, 25.1 - 24.7 (overlapping signals), 24.5, 24.3, 19.6, 19.4, 19.3 ppm; MS (ES⁺, MeOH): 786 ([M + Na]⁺, 100%); HRMS (ES⁺, MeOH) *m/z* calcd. for C₃₆H₅₇N₇O₉NaS [M + Na]⁺ 786.3831, found 786.3817.

CbzPheAib₄Gly-ψ[CSNH]AibOME 4. To a mixture of CbzPheOH (37 mg, 0.12 mmol), HOAt (17 mg, 0.12 mmol) and H-Aib₄Gly-ψ[CSNH]AibOME **14** (33 mg, 0.06 mmol) in dry CH₂Cl₂ (3 mL) was added EDC (0.022 mL, 0.12 mmol), followed by N,N-diisopropylethylamine (0.05 mL, 0.03 mmol). The mixture was left stirring at room temperature for 3 days, then the solvent was evaporated. The residue was taken up in EtOAc (5 mL) and washed with KHSO₄ 5% (3 x 1 mL), sat. NaHCO₃ (3 x 1 mL) and brine (1 mL). The organic phase was dried (MgSO₄) and the solvent removed *in vacuo*. The crude product was purified by column chromatography (CH₂Cl₂/MeOH, 95:5) to give pure CbzPheAib₄Gly-ψ[CSNH]AibOME (39 mg, 78%) as a white solid. *R*_f 0.5 (CH₂Cl₂/MeOH, 95:5); mp = 110-113 °C; [α]_D²⁵ = -19.2; IR (ATR, cm⁻¹): 3291, 2984, 1657, 1529, 1383, 1226, 1155; ¹H NMR (500 MHz, MeOD) δ_H 7.90 (s, 1H), 7.81 (s, 2H), 7.39 - 7.16 (m, 1H), 5.10 (d, A of AB, *J* = 12.6 Hz, 1H), 5.06 (d, B of AB, 12.6 Hz, 1H), 4.30 (d, A of AB, *J* = 17.7 Hz, 1H), 4.23 (t, *J* = 7.8 Hz, 1H), 4.11 (d, B of AB, *J* = 17.7 Hz, 1H), 3.63 (s, 3H), 3.00 (ddd, *J* = 31.6, 13.6, 7.8 Hz, 2H), 1.67 (d, *J* = 2.5 Hz, 6H), 1.50 (s, 3H), 1.49 (s, 6H), 1.42 (s, 3H), 1.38 (s, 3H), 1.34 (s, 3H), 1.28 (s, 3H), 1.24 (s, 3H). ¹³C NMR (101 MHz, MeOD) δ_C 199.9, 177.9, 177.9, 177.8, 176.7, 175.5, 174.5, 158.7, 138.3, 138.2, 130.7, 129.7, 129.7, 129.2, 128.7, 128.1, 67.8, 61.4, 58.4, 58.3, 58.1, 57.9, 57.7, 52.8, 52.4, 38.4, 30.9, 26.6 - 26.3 (overlapping signals), 25.1, 24.9, 24.8, 24.3, 24.1; MS (ES⁺, MeOH): 834 ([M + Na]⁺, 100%); HRMS (ES⁺, MeOH) *m/z* calcd. for C₄₀H₅₇N₇O₉NaS [M + Na]⁺ 834.3831, found 834.3818.

CbzAlaAib₄Gly-ψ[CSNH]AibOME 5. To a mixture of CbzAla-OH (28 mg, 0.12 mmol), HOAt (17 mg, 0.12 mmol) and H-Aib₄Gly-ψ[CSNH]AibOME **14** (33 mg, 0.06 mmol) in dry CH₂Cl₂ (3 mL) was added EDC (0.023 mL, 0.12 mmol), followed by N,N-diisopropylethylamine (0.05 mL, 0.03 mmol). The mixture was left stirring at room temperature for 3 days, then the solvent was evaporated. The residue was taken up in EtOAc (5 mL) and washed with KHSO₄ 5% (3 x 1 mL), sat. NaHCO₃ (3 x 1 mL) and brine (1 mL). The organic phase was dried (MgSO₄) and the solvent removed *in vacuo*. The crude product was purified by column chromatography (CH₂Cl₂/MeOH, 97:3) to give pure CbzAlaAib₄Gly-ψ[CSNH]Aib-OME (35 mg, 77%) as a white solid. *R*_f 0.4 (CH₂Cl₂/MeOH, 97:3); mp = 204-205 °C; [α]_D²⁵ = -13.2 IR (ATR, cm⁻¹) 3287, 2984, 1703, 1643, 1529, 1420, 1361, 1208, 1171; ¹H NMR (500 MHz, MeOD) δ_H 7.90 (s, 1H), 7.81 (s,

1H), 7.38 – 7.27 (m, 5H), 5.12 (d, A of AB, $J = 12.7$ Hz, 1H), 5.09 (d, B of AB, $J = 12.7$ Hz, 1H), 4.27 (d, A of AB, $J = 17.7$ Hz, 1H), 4.16 (d, B of AB, $J = 17.7$ Hz, 1H), 4.00 (q, $J = 7.1$ Hz, 1H), 3.63 (s, 3H), 1.67 (s, 6H), 1.50 (s, 6H), 1.47 – 1.36 (m, 18H), 1.34 (d, $J = 7.1$ Hz, 3H); ^{13}C NMR (126 MHz, MeOD) δ_{C} 199.9, 177.9, 177.9, 177.8, 176.8, 176.1, 175.5, 158.8, 138.4, 129.7, 129.3, 128.8, 79.6, 67.8, 61.4, 58.2, 58.1, 57.9, 57.6, 52.8, 52.7, 52.4, 26.2, 25.9, 25.8, 25.4, 25.2 - 24.8 (overlapping signals), 24.7, 17.3; MS (ES^+ , MeOH): 758 ($[\text{M} + \text{Na}]^+$, 100%); HRMS (ES^+ , MeOH) m/z calcd. for $\text{C}_{34}\text{H}_{53}\text{N}_7\text{O}_9\text{NaS} [\text{M} + \text{Na}]^+$ 758.3518, found 758.3505.

Cbz(α Me)ValAib₄Gly- ψ [CSNH]AibOME 6.

*Acyl fluoride formation.*⁵¹ A solution of Cbz(α Me)Val-OH (58 mg, 0.22 mmol) in dry CH_2Cl_2 (2 mL), was treated with pyridine (0.018 mL, 0.22 mmol) and TFFH (tetramethylfluoroformamidinium hexafluorophosphate, 87 mg, 0.33 mmol). The mixture was left stirring at room temperature for 4 h, after which ice-cold water was added (2 mL). The organic phase was separated and washed again with ice-cold water (2 mL), dried (MgSO_4) and concentrated to give crude Cbz(α Me)Val-F as a colorless solid.

Acyl fluoride coupling. The freshly prepared amino acyl fluoride was dissolved in dry CH_2Cl_2 (2 mL) and the resulting solution was added to H-Aib₄Gly- ψ [CSNH]AibOME 14 (39 mg, 0.073 mmol) in dry CH_2Cl_2 (3 mL). N,N -diisopropylethylamine (0.134 mL, 0.77 mmol) was added and the mixture was left stirring at room temperature for 5 days, after which the solvent was evaporated. The residue was taken up in EtOAc (5 mL) and washed with KHSO_4 5% (3 x 1 mL), sat. NaHCO_3 (3 x 1 mL) and brine (1 mL). The organic phase was dried (MgSO_4) and the solvent removed *in vacuo*. The crude product was purified by column chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$, 95:5) to give pure Cbz(α Me)ValAib₄Gly- ψ [CSNH]Aib-OME (51 mg, 70%) as a white solid. R_f 0.25 ($\text{CH}_2\text{Cl}_2/\text{MeOH}$, 95:5); mp = 203-205 °C; $[\alpha]_{\text{D}}^{25} = +34.7$; IR (ATR, cm^{-1}): 3281, 2983, 1707, 1651, 1530, 1454, 1382, 1361, 1287, 1163; ^1H NMR (400 MHz, MeOD) δ_{H} 7.91 (s, 1H), 7.83 (s, 1H), 7.43 – 7.27 (m, 5H), 5.20 (d, A of AB, $J = 12.7$ Hz, 1H), 5.07 (d, B of AB, $J = 12.7$ Hz, 1H), 4.34 (d, A of AB, $J = 17.7$ Hz, 1H), 4.09 (d, B of AB, $J = 17.7$ Hz, 1H), 3.63 (s, 3H), 2.02 (m, 1H), 1.68 (s, 3H), 1.67 (s, 3H), 1.53 – 1.33 (m, 24H), 1.27 (s, 3H), 1.00 (d, $J = 6.8$ Hz, 3H), 0.94 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (101 MHz, MeOD) δ_{C} 199.8, 178.2, 177.8, 177.7, 176.9, 175.9, 175.4, 158.2, 138.7, 129.6, 129.1, 128.7, 79.5, 73.4, 67.7, 63.9, 61.3, 58.2, 58.1, 58.0, 57.8, 57.7, 52.7, 52.2, 36.0, 26.7, 26.6, 24.8, 24.7 - 24.4 (overlapping signals), 23.9, 18.0, 17.7; MS (ES^+ , MeOH): 800 ($[\text{M} + \text{Na}]^+$, 100%); HRMS (ES^+ , MeOH) m/z calcd. for $\text{C}_{37}\text{H}_{63}\text{N}_8\text{O}_9\text{NaS} [\text{M} + \text{NH}_4]^+$ 795.4439, found 795.4433.

Acknowledgements

This work was funded by the European Research Council (ERC Advanced Grant ROCOCO) and was presented at the 23rd American Peptide Society Symposium in Hawaii, 20-27 June

2013. Jonathan Clayden is a recipient of a Royal Society Wolfson Research Merit Award.

Notes and references

^aSchool of Chemistry, University of Manchester, Oxford Road, Manchester M13 9PL, UK.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

References

- S. H. Gellman, *Acc. Chem. Res.*, 1998, **31**, 173.
- G. Guichard and I. Huc, *Chem. Commun.*, 2011, 5933.
- S. Hecht and I. Huc, *Foldamers*, Wiley-VCH, Weinheim, 2007.
- D. J. Hill, M. J. Mio, R. B. Prince, T. S. Hughes and J. S. Moore, *Chem. Rev.*, 2001, **101**, 3893.
- E. Yashima, K. Maeda, H. Iida, Y. Furusho and K. Nagai, *Chem. Rev.*, 2009, **109**, 6102.
- P. N. Shah, J. Min, H.-J. Kim, S.-Y. Park and J.-S. Lee, *Macromolecules*, **44**, 7917.
- P. N. Shah, J. Min, C.-G. Chae, N. Nishikawa, D. Suemasa, T. Kakuchi, T. Satoh and J.-S. Lee, *Macromolecules*, **45**, 8961.
- N. Ousaka, Y. Inai and R. Kuroda, *J. Am. Chem. Soc.*, 2008, **130**, 12266.
- N. Delsuc, T. Kawanami, J. Lefeuvre, A. Shundo, H. Ihara, M. Takafuji and I. Huc, *ChemPhysChem*, 2008, **9**, 1882.
- R. P. Hummel, C. Toniolo and G. Jung, *Angew. Chem. Int. Ed.*, 1987, **99**, 1180.
- L. Byrne, J. Solà, T. Boddaert, T. Marcelli, R. W. Adams, G. A. Morris and J. Clayden, *Angew. Chem. Int. Ed.*, 2013, in press.
- N. Sreerama and R. W. Woody, *Circular Dichroism of Peptides and Proteins in "Circular Dichroism: Principles and Applications"*, Wiley, Eds. Berova, N., Nakanishi, K., Woody, R.W., New York, NY, 2000.
- M. Sugimoto, S. Collet and Y. Ito, *Org. Lett.*, 2002, **4**, 351.
- Y. Ito, T. Miyake, S. Hatano, R. Shima, T. Ohara and M. Sugimoto, *J. Am. Chem. Soc.*, 1998, **120**, 11880.
- Such a difficulty has led to misassignment of screw sense: see ref. 18.
- J. Solà, M. Helliwell and J. Clayden, *J. Am. Chem. Soc.*, 2010, **132**, 4548.
- J. Solà, G. A. Morris and J. Clayden, *J. Am. Chem. Soc.*, 2011, **133**, 3712.
- R. A. Brown, T. Marcelli, M. De Poli, J. Solà and J. Clayden, *Angew. Chem. Int. Ed.*, 2012, **51**, 1395.
- S. J. Pike, M. De Poli, W. Zawodny, J. Raftery, S. J. Webb and J. Clayden, *Org. Biomol. Chem.*, 2013, **11**, 3168.
- J. Solà, S. P. Fletcher, A. Castellanos and J. Clayden, *Angew. Chem. Int. Ed.*, 2010, **49**, 6836.
- J. Helbing, H. Bregy, J. Bredenbeck, R. Pfister, P. Hamm, R. Huber, J. Wachtveitl, L. De Vico and M. Olivucci, *J. Am. Chem. Soc.*, 2004, **126**, 8823.
- D. Wildemann, C. Schiene-Fischer, T. Aumüller, A. Bachmann, T. Kiefhaber, C. Lücke and G. Fischer, *J. Am. Chem. Soc.*, 2007, **129**, 4910.
- E. P. Dudek and G. Dudek, *J. Org. Chem.*, 1967, **32**, 823.

24. S. Yao, R. Zutshi and J. Chmielewski, *Bioorg. Med. Chem. Lett.*, 1998, **8**, 699.
25. J. Lehmann, A. Linden and H. Heimgartner, *Tetrahedron*, 1998, **98**, 8721.
26. J. M. Goldberg and R. L. Baldwin, *Biochemistry*, 1998, **37**, 2546.
27. Inai (N. Ousaka and Y. Inai, *J. Org. Chem.*, 2009, **74**, 1429) has used dehydrophenylalanine derivatives containing different chromophores to distinguish different locations in an extended helix.
28. C. Toniolo, *CRC Crit. Rev. Biochem.*, 1980, **9**, 1.
29. C. Toniolo and E. Benedetti, *Trends Biochem. Sci.*, 1991, **16**, 350.
30. C. Toniolo, M. Crisma, F. Formaggio and C. Peggion, *Biopolymers (Pept. Sci.)*, 2001, **60**, 396.
31. K. Clausen, M. Thorsen and S.-O. Lawesson, *Tetrahedron*, 1981, **37**, 3635.
32. M. De Zotti, B. Biondi, C. Peggion, M. De Poli, H. Fathi, S. Oancea, C. Toniolo and F. Formaggio, *Beilstein J. Org. Chem.*, 2012, **8**, 1161.
33. O. E. Jensen, S.-O. Lawesson, R. Bardi, A. M. Piazzesi and C. Toniolo, *Tetrahedron*, 1985, **41**, 5595.
34. $\psi[X]$ indicates replacement of a peptide CONH bond by X: see Spatola, A. F. in *Chemistry and Biochemistry of Amino Acids, Peptides and Proteins*, Dekker, Ed. B. Weinstein, New York, 1983.
35. Studies of similar Aib-containing oligomers have shown that NMR spectra obtained in CD₃OD remained identical over a four-fold range of concentration. Spectra obtained in less polar solvents were broad and concentration-dependent, suggesting aggregation. See J. Clayden, A. Castellanos, J. Solà and G. A. Morris, *Angew. Chem. Int. Ed.* 2009, **48**, 5962; S. J. Pike, V. Diemer, J. Raftery, S. J. Webb and J. Clayden, manuscript in preparation
36. $\Delta\delta$ for an AB system is given by $\nu_0\Delta\delta = [(f_1 - f_3)^2 - J_{AB}^2]^{1/2} = [(f_2 - f_4)^2 - J_{AB}^2]^{1/2} = [(f_1 - f_4)(f_2 - f_3)]^{1/2}$ where $f_{1,2,3,4}$ are the observed resonance frequencies in order of the four lines comprising the AB multiplet, J_{AB} is the coupling constant and ν_0 is the spectrometer frequency.
37. J. H. Miwa, L. Pallivathucal, S. Gowda and K. E. Lee, *Org. Lett.*, 2002, **4**, 4655.
38. J. Zhao, J.-C. Micheau, C. Vargas and C. Schiene-Fischer, *Chem. Eur. J.*, 2004, **10**, 6093.
39. J. H. Miwa, A. K. Patel, N. Vivatrat, S. M. Popek and A. M. Meyer, *Org. Lett.*, 2001, **3**, 3373.
40. M. Kajtár, M. Hollósi, J. Kajtár, Z. Majer and K. E. Kövér, *Tetrahedron*, 1986, **42**, 3931.
41. M. Hollósi, E. Kollát, J. Kajtár, M. Kajtár and G. D. Fasman, *Biopolymers*, 1990, **30**, 1061.
42. T. T. Sifferlen, M. Rueping, K. Gademann, B. Jaun and D. Seebach, *Helv. Chim. Acta*, 1999, **82**, 2067.
43. A. Bachmann, D. Wildemann, F. Praetorius, G. Fischer and T. Kiefhaber, *Proc. Natl. Acad. Sci. U.S.A.*, 2011, **108**, 3952.
44. D. F. Mierke, A. Geyer and H. Kessler, *Int. J. Pept. Protein Res.*, 1994, **44**, 325.
45. Y. Otani, T. Hori, M. Kawahata, K. Yamagouchi and T. Ohwada, *Tetrahedron*, 2012, **68**, 4418.
46. M. Goodman and C. Toniolo, *Biopolymers*, 1968, **6**, 1673.
47. M. De Poli, M. De Zotti, J. Raftery, J. A. Aguilar, G. A. Morris and J. Clayden, *J. Org. Chem.*, 2013, **78**, 2248.
48. ¹³C NMR may be used to determine absolute screw sense preferences by employing an enantiomerically enriched probe incorporating an isotopic label. See J. Solà, S. P. Fletcher, A. Castellanos and J. Clayden, *Angew. Chem. Int. Ed.*, 2010, **49**, 6836; S. P. Fletcher, J. Solà, D. Holt, R. A. Brown and J. Clayden, *Beilstein J. Org. Chem.* 2011, **7**, 1304.
49. P. Jana, S. Maity, S. K. Maity and D. Halder, *Chem. Comm.*, **47**, 2092.
50. C. Kolano, K. Gomann and W. Sander, *Eur. J. Org. Chem.*, 2004, 4167.
51. L. A. Carpino and A. El-Faham, *J. Am. Chem. Soc.*, 1995, **117**, 5401.