


Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

Catalytic and mechanistic studies into the epoxidation of styrenes using manganese complexes of structurally similar polyamine ligands.

Gennadiy Ilyashenko, Giorgio De Faveri, Thomas Follier, Rawan Al-Safadi, Majid Motevalli and Michael Watkinson*

The Joseph Priestley Building, School of Biological and Chemical Sciences, Queen Mary University of London, Mile End Road, London, E1 4NS, UK.

Email m.watkinson@qmul.ac.uk

Abstract:

Two structurally similar polyamine ligands (**7** and **8**) have been prepared, which differ only by the presence of either a secondary or tertiary nitrogen donor within their N₅ donor set. The ligands, in combination with iron and manganese salts, have been screened for their efficacy as catalysts for the epoxidation of styrene, using both hydrogen peroxide and peracetic acid as oxidants. Clear differences in activity between the two systems were observed, with **7** proving most effective in the presence of MnSO₄ with H₂O₂, whereas ligand **8** proved to be effective with Mn(OTf)₂, MnCl₂ and Mn(ClO₄)₂ using peracetic acid as the oxidant. A Hammett analysis of the initial rate kinetics of the optimal systems, combined with analysis by UV-vis spectroscopy, indicates that the small structural differences in the ligands elicit profound changes in the nature of the active species formed.

Keywords Epoxidation; Polyamine ligands; Manganese; Hydrogen peroxide; Kinetics.


Introduction

The oxidation of organic substrates is fundamental to contemporary synthetic chemistry and there continues to be a demand for new and efficient catalytic methods to effect regio- and stereoselective oxidative transformations.¹ Within the very broad class of catalytic oxidative transformations, asymmetric epoxidation (AE) is an area which has received considerable attention due to the fact that a synthetically useful epoxide is generated containing up to two stereogenic centres from widely available alkene substrates.² Since the seminal reports by Sharpless of the AE of allylic alcohols,³ and Jacobsen⁴ and Katsuki⁵ of the AE of certain

unfunctionalised alkene substrates with manganese-salen complexes, a range of other highly effective and elegant metal-based and organocatalytic systems have been developed which are extremely effective for a wide range of substrates.⁶ Although some of these systems satisfy many of the increasingly demanding environmental and economic constraints of industry, some limitations persist. These include: the need for relatively high catalyst loadings; the environmentally non-benign solvents and oxidants used in some cases; difficulties in ligand synthesis and the ease of their modification in order to fine-tune catalyst activity and stereoselectivity, and relatively low turnover numbers and frequencies when compared to other catalytic systems (e.g. TONs for Noyori's asymmetric hydrogenation systems reach up to 2 400 000, with TOFs similarly impressive⁷). Furthermore, there is no generic catalyst that is capable of the AE of all alkene substrates. As a result there continues to be considerable interest in the development of new and improved catalytic systems.

Owing to their benign environmental character, low toxicity and cost, good atom economy and operational ease, much of the recent effort in this area has been focused on the activation of hydrogen peroxide or peracetic acid with complexes of both iron and manganese.^{6b} Considerable success has been achieved with a range of aminopyridine-based ligands (**1-6**, Scheme 1), as a result of their ability to form catalytically active complexes with both manganese and iron.⁸ For example, manganese complexes of chiral ligands **1** and **2** have already shown promising activity in the epoxidation of the challenging bench-mark substrate styrene (83% yield, 15% *ee* and 85% yield, 43% *ee* respectively).^{8c} However, unwanted and competing decomposition of the hydrogen peroxide with these metals, *via* catalase-like pathways, can often be problematic and the enantioselectivity of these complexes is still modest; their further improvement being contingent on the chemist's ability to readily modify the ligand structure in order to allow additional tuning and optimisation of activity.


We became particularly interested in developing ligands which combine the chiral backbone derived from *trans*-diaminocyclohexane and the coordination environment of ligand **3**, the complexes of which have already exhibited excellent reactivity in oxidation reactions.^{8f} We postulated that ligands **7** and **8** (Scheme 1) would stabilise the putative high-valent manganese species expected to be the active oxidant, whilst also providing the potential to induce asymmetry as a result of the privileged nature of the diamine backbone. In addition the facile inclusion of secondary and tertiary amines in closely related ligands would allow their effects to be assessed. Herein we report our investigations and the unexpected results obtained during the catalytic screening and subsequent preliminary kinetic studies.



Scheme 1: Structure of previously synthesised aminopyridine ligands **1** – **6** and ligands **7** and **8**.

Results and Discussions


The desired ligands **7** and **8** were prepared in six steps in high enantiopurity and excellent yields (**Scheme 2**) by adapting reported methods.⁹



Scheme 2: Synthesis of ligands **7** and **8**. Reagents and conditions: (i) *L*-tartaric acid, acetic acid, water; (ii) pyridine-2-carboxylaldehyde, K_2CO_3 , EtOH, 90%; (iii) $NaBH_4$, MeOH, 88%; (iv) pyridine-2-carboxylaldehyde, toluene, Dean-Stark, reflux, 88%; (v) $NaBH_3(CN)$, MeOH, 87%; (vi) HCO_2H , CH_2O , H_2O , 93%.

Amine **10** was readily obtained after the resolution of *trans*-1,2-diaminocyclohexane into diastereoisomeric tartrate salts, which underwent facile condensation with pyridine-2-carboxylaldehyde followed by reduction of the imines with $NaBH_4$ in excellent yield. With **10** in hand, addition of a further equivalent of pyridine-2-carboxylaldehyde gave aiminal **11** which was reduced with $NaBH_3(CN)$ to obtain ligand **7**. Although it would be preferable to effect this transformation with other reducing agents, such as $NaBH_4$, due to the potential toxicity and high cost of $NaBH_3(CN)$, all of our efforts lead to either a substantial degradation of **11** or the reaction not proceeding to completion. Methylation of **7** was readily achieved using standard Eschweiler-Clarke conditions to produce ligand **8**.

With the ligands in hand we were able to carry out a preliminary screen for catalytic activity in the epoxidation of the challenging bench-mark substrate styrene, using both H₂O₂ and PAA, in combination with various metal salts (**Scheme 3, Table 1**). This was carried out according to previously developed procedures within our group,^{8i,10} wherein both ligand and metal salt are mixed together for 30 minutes in MeCN at room temperature prior to the addition of the substrate. In all cases the complete disappearance of the ligand was confirmed by TLC prior to the addition of oxidant. The reaction was then initiated by the addition of the oxidant: (hydrogen peroxide in acetone was added to the reaction mixture at room temperature as has been widely used in related systems;^{6b,10} PAA was added to the reaction mixture which was cooled on ice to 3-4 °C). Substrate conversion was monitored by collecting aliquots and analysing them by HPLC.



Scheme 3: Epoxidation of styrenes using *in situ* conditions.

Table 1: Epoxidation of styrene under various reaction conditions using ligands **7** and **8**.

Entry	Ligand (1 mol%)	Metal Salt (1mol%)	Oxidant	Time (h) ^a	Conversion % ^b	Yield % ^{b,c}	TON	TOF (min ⁻¹)
1 ^d	7	FeX _{2/3}	H ₂ O ₂ /PAA	8	-	-	-	-
2 ^d	8	FeX _{2/3}	H ₂ O ₂ /PAA	8	-	-	-	-
3	7	MnSO ₄	PAA	5	-	-	-	-
4	7	Mn(OTf) ₂	PAA	5	-	-	-	-
5	7	MnCl ₂	PAA	5	-	-	-	-
6	7	Mn(ClO ₄) ₂	PAA	5	-	-	-	-
7	7	MnSO ₄	H ₂ O ₂	5	100	78	100	0.33
8	7	Mn(OTf) ₂	H ₂ O ₂	8	51	13	51	0.11
9	7	MnCl ₂	H ₂ O ₂	8	65	5	65	0.13
10	7	Mn(ClO ₄) ₂	H ₂ O ₂	8	60	4	50	0.12
11	8	MnSO ₄	H ₂ O ₂	8	25	14	25	0.052
12	8	Mn(OTf) ₂	H ₂ O ₂	8	40	27	40	0.083
13	8	MnCl ₂	H ₂ O ₂	8	41	1	41	0.083
14	8	Mn(ClO ₄) ₂	H ₂ O ₂	8	37	2	37	0.077
15	8	Mn(OTf) ₂	PAA	1.5	100	93	100	1.11
16	8	MnCl ₂	PAA	1.5	100	74	100	1.11
17	8	Mn(ClO ₄) ₂	PAA	1.5	100	90	100	1.11
18	8	MnSO ₄	PAA	5	29	7	29	0.097

a) Minimum time required for the reaction to reach its maximum conversion; b) the conversions and yields were calculated using HPLC; c) in cases when the yields were low

compared to conversions a number of over-oxidation products were also observed, including 1-phenylethyl-1,2-diol and benzaldehyde; d) Fe-salts used: FeCl₃ (anhydrous), Fe(ClO₄)₃, Fe(OTf)₂, Fe(OAc)₂, FeCl₂.

Despite a number of attempts, we did not observe any catalytic activity with the iron salts tested in combination with either ligand or oxidant (Table 1, entries 1 and 2) in contrast to other recently reported results.^{8j} Also the combination of PAA as the oxidant with ligand **7** and a number of the manganese salts did not yield any activity (Table 1, entries 3-6). The combination of either ligands **7** or **8** with a range of manganese salts and H₂O₂ resulted in reasonable substrate conversions, although poor yields were observed in all cases with the exception of ligand **7**, MnSO₄ and H₂O₂ (Table 1, entry 7). Indeed, the catalytic activity of this combination (TON = 100, TOF = 0.33 min⁻¹) is higher than that observed for Katsuki's salalen-Ti catalyst (TON = 100, TOF = 1.16x10⁻³ min⁻¹),¹¹ yet is lower than that of manganese complexes derived from triazacyclononane (TON = 100, TOF = 2.22 min⁻¹) under similar reaction conditions.¹⁰ The considerable difference between the conversions and yields, particularly prevalent for both the perchlorate and chloride salts (Table 1, Entries 9, 10, 13 and 14) prompted a further analysis by HPLC and the ¹H NMR spectroscopy of the crude reaction mixtures. We identified that the low yields were a result of the over-oxidation of the styrene substrate, principally to form the 1-phenylethyl-1,2-diol and benzaldehyde.

Combination of ligand **8** with Mn(OTf)₂ MnCl₂ and Mn(ClO₄)₂ and PAA (Table 1, entries 15-17), resulted in complete conversion of styrene in 1.5 hours (e.g. for Mn(OTf)₂: TON = 100, TOF = 1.11 min⁻¹) and produced the epoxide in excellent yields with only traces of over-oxidation products being observed (Table 1, entries 15-17). This catalytic activity exhibited by this system is comparable to that observed with manganese complexes of ligand **2** under similar reaction conditions (TON = 100, TOF = 1.11 min⁻¹).^{8e} In contrast, MnSO₄ proved relatively inactive in combination with **8** using PAA as the oxidant (Table 1, entry 18). Disappointingly, despite the high catalytic activity exhibited in several cases, only the combination of MnSO₄ and **7** (system 1) and Mn(OTf)₂ and **8** (system 2) ever produced non-racemic product, albeit with very low levels of asymmetric induction (11 and 15% respectively). When compared to the state-of-art, systems such as Katsuki's salalen-Ti and ligand **5** derived catalysts which give yields of 93%¹¹ and 43%^{8h} respectively, the present system clearly exhibits an inferior level of enantioselectivity.

This moved us to investigate the reaction further and we carried out a series of blank experiments which have clearly shown that in order for the epoxidation to occur the reaction

mixture must contain ligand, manganese salt, oxidant and the correct solvent ratio. Upon removal of the ligand from the reaction mixture only a very slow background oxidation was observed, which resulted in less than 20% conversion of substrate over a 48 h period, giving an intractable mixture of products.

A recent report by Browne *et al.* has shown that a decomposition of structurally similar ligand results in the formation of picolinic acid, which then react with manganese to form an effective epoxidation catalyst.¹² Given the structural similarity of **7** and **8** to the ligands used by Browne *et al.*, we were keen to eliminate such a possibility in our study. Fortunately, under the reaction conditions we have employed for our systems, picolinic acid proved to be inactive in the presence of the manganese salts which showed activity. Moreover, the exposure of **7** and **8** to hydrogen peroxide or PAA in acetone-d₆ and subsequent analysis by ¹H NMR spectroscopy showed that within the catalytic reaction times (up to 8 h) both ligands were stable and only after 18 h was some ligand degradation observed.

Finally we prepared authentic metal complexes of both ligands (*vide infra*) and showed that their activity was the same as the *in situ* formed complexes. We are thus confident that during the timeframe of the reaction that the conversions and yields observed are due to the ligand-metal complexes, rather than being due to decomposition products or free metal species.^{2b}

The synthesis of the authentic complexes was carried out by reacting the ligand with the desired manganese salt in acetonitrile according to a previously reported procedure,⁸ⁱ and these were isolated in good to excellent yields as pale yellow or off-white powders (Table 2).

Table 2: Summary of manganese complexes synthesised and their magnetic moments measured by the Evan's NMR method.¹³

Entry	Complex	Yield (%)	$\chi_c/10^{-6}\text{cm}^3\text{g}^{-1}$	$\mu_{\text{eff}} (\mu_{\text{B}})$
1	Mn(7)SO ₄	59	26.6	5.84
2	Mn(7)(ClO ₄) ₂	82	22.5	5.87
3	Mn(7)OAc ₂	79	25.7	5.86
4	[Mn(7)Cl]Cl·H ₂ O	91	28.3	5.88
5	Mn(7)OTf ₂	85	19.3	5.85
6	Mn(8)Cl ₂	82	27.7	5.90
7	Mn(8)OTf ₂	38	19.0	5.84

The complexes were characterized by mass spectrometry, UV/vis and IR spectroscopies. As expected a slight shift in IR-bands attributable to the pyridine moieties were observed between 1650 and 1550 cm⁻¹ which is indicative of coordination to the manganese centre. The UV/vis

spectra showed absorbance bands due to the ligands and, as expected for the manganese(II) centres formulated in Table 2, no d-d transitions. The formation of the complexes was supported by the HRMS with $[\text{MnL}]^+$ ions being observed for all complexes. The Evans' NMR method¹³ was used to determine the magnetic susceptibilities of the complexes and the data obtained from these measurements (Table 2) was also consistent with the spin-only value expected for magnetically dilute manganese(II)-d⁵ complexes.

The reaction between ligand **7** and $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ afforded an off-white solid. Dissolution of this material in a mixture of MeCN/DCM followed by slow evaporation gave pale yellow crystals suitable for single crystal X-ray diffraction. The crystal structure of the complex is shown in **Figure 2** and relevant bond lengths and angles are recorded in **Table 5**. This revealed the manganese centre to display a distorted octahedral geometry with one chloride ion bound directly to the manganese atom and the other chloride counter ion outside the primary coordination sphere. The Mn-N bond lengths are comparable to those previously reported for similar structures.¹⁴




Figure 2: ORTEP plot showing the single crystal X-ray structure of complex $[\text{Mn}(\mathbf{7})\text{Cl}]\text{Cl} \cdot \text{H}_2\text{O}$ with 50% thermal ellipsoids with hydrogen atoms omitted for clarity.

Table 5: Selected bond lengths and angles for $[\text{Mn}(\mathbf{7})\text{Cl}]\text{Cl} \cdot \text{H}_2\text{O}$ complex with esd's.

Bond length (Å)		Bond angle (°)	
Mn(1)-N(1)	2.2441(17)	N(1)-Mn(1)-N(2)	72.87(6)
Mn(1)-N(2)	2.3236 (17)	N(1)-Mn(1)-N(3)	77.57(6)
Mn(1)-N(3)	2.3440 (17)	N(1)-Mn(1)-N(5)	100.28(6)
Mn(1)-N(4)	2.2371 (17)	N(1)-Mn(1)-Cl(1)	116.82(5)
Mn(1)-N(5)	2.2598 (17)	N(2)-Mn(1)-Cl(1)	91.99(5)
Mn(1)-Cl(1)	2.4122(6)	N(3)-Mn(1)-Cl(1)	163.42(4)
		N(4)-Mn(1)-Cl(1)	99.10(4)
		N(5)-Mn(1)-N(3)	72.95(6)

 | N(5)-Mn(1)-Cl(1) 95.33(5)

Despite the lack of the enantioselectivity we decided to investigate the two most active systems with each oxidant more thoroughly i.e. ligand **7**/ MnSO₄/ H₂O₂ (System 1) and ligand **8**/ Mn(OTf)₂/ PAA (System 2) and screened them for activity with a range of *p*-substituted styrenes containing both electron donating and electron withdrawing groups (**Table 2**)

Table 2: Epoxidation of a range of *p*-substituted styrene substrates using the two catalytic systems.

Entry	Catalytic System	Substrate	Time (h) ^a	Conversion %	Yield %	TON ^b	TOF (min ⁻¹)
1	1	<i>p</i> -Methoxystyrene	8	21	-	21	0.044
2	1	<i>p</i> -Methylstyrene	8	89	43	89	0.18
3	1	Styrene	5	100	78	100	0.33
4	1	<i>p</i> -Fluorostyrene	5	100	79	100	0.33
5	1	<i>p</i> -Chlorostyrene	5	100	75	100	0.33
6	1	<i>p</i> -Bromostyrene	6	100	39	100	0.27
7	1	<i>p</i> -Nitrostyrene	6	90	63	90	0.25
8	1	<i>p</i> -Nitrilestyrene	4	100	73	100	0.38
9	2	<i>p</i> -Methoxystyrene	1	100	10	100	1.67
10	2	<i>p</i> -Methylstyrene	2	100	45	100	0.83
11	2	Styrene	1.5	100	93	100	1.11
12	2	<i>p</i> -Fluorostyrene	1.5	100	98	100	1.11
13	2	<i>p</i> -Chlorostyrene	1.5	100	98	100	1.11
14	2	<i>p</i> -Bromostyrene	1.5	100	72	100	1.11
15	2	<i>p</i> -Nitrostyrene	1	100	43	100	1.67
16	2	<i>p</i> -Nitrilestyrene	0.25	100	58	100	6.67

a) Minimum time required for the reaction to reach its maximum conversion; b) based on substrate conversions

The influence of the *para*-substituent on the reaction catalysed with System **1** quickly became apparent, with electron rich styrenes having the lowest conversions and yields (Table 2, entries 1 and 2), whilst other more electron deficient substrates deficient derivatives underwent complete conversion and gave the target epoxide in good yields, with the exception of *p*-bromostyrene (Table 2, entries 3 – 8).

The reactions using System 2 produced broadly similar results. Although the overall conversion of styrenes was excellent in all cases, the yields of the epoxide products differed. Styrene, its *para*-fluoro- and chloro-derivatives were epoxidised in the highest yields (Table

2, entries 11-13), whereas *para*-methyl-, nitro- and nitrile-styrene epoxides were formed in more modest yields, despite complete substrate conversion. Substituted styrene epoxides containing electron donating groups, such as methyl and methoxy are known to undergo hydrolysis in the presence of acids,¹⁵ hence while their decomposition under the reaction conditions is disappointing, it is not unexpected. It is also known that *p*-nitrostyrene oxide undergoes benzylic C-O bond cleavage in the presence of acids, even though the electron-withdrawing nature of nitro- group should, in fact, decrease the rate of the reaction that results from the formation of positive charge at the benzylic position.^{15,16} Therefore it is reasonable to assume similar behaviour from *p*-nitrilestyrene oxide, hence explaining the lower than expected yields of the product. Analysis of the crude reaction mixtures by ¹H NMR-spectroscopy confirmed the formation of diols, further supporting this hypothesis. Finally, for reasons, which are not obvious at this stage, although *p*- bromostyrene substrate was completely consumed by system 1, relatively little epoxide was observed.

In order to gain a further insight into the reaction pathway and understand if the oxygen transfer is a step-wise or concerted process we then investigated the epoxidation of *cis*-stilbene. After subjecting the alkene to the oxidation under both sets of general conditions the crude reaction mixture was analysed by ¹H NMR spectroscopy, which clearly showed that the epoxide obtained had a *trans*-configuration. We therefore decided to investigate the initial rate kinetics of the epoxidation reactions using a Hammett analysis¹⁷ between the rate and *p*-substituents on styrene.¹⁰

Since at the start of the reaction the catalyst is present at the lowest concentration when compared to either the substrate or the oxidant and assuming that the oxygen transfer is the rate-determining step, it is appropriate to use a pseudo-first order approximation and obtain initial rates and initial rate constants (equation 1).

$$R_i = k_i [\text{catalyst}] \text{ where } k_i = k[\text{substrate}] \text{ (Equation 1)}$$

The initial rate constant, k_i , is proportional to the real rate constant, k , hence the Hammett plots obtained with k_i should be representative of those obtained using k . Therefore, we focussed our efforts on obtaining the initial rates and k_i values for both systems. All of the reactions were monitored by HPLC using anisole (and 1,2-dichlorobenzene for *p*-nitrilestyrene due to overlap in the HPLC trace) as the internal standard, which was calibrated simultaneously for each of the styrenes and its corresponding epoxide (see ESI for calibrations and representative reaction profiles). The reactions were run in triplicate in order to produce

accurate and reproducible results and the data are presented in **Table 3**. The k_i -values were used to obtain Hammett plots, which graphically show the influence of the *para*-substituents on the reaction (**Figure 1**).

Table 3: Initial rates and k_i -values for each substrate catalysed by both systems.

Entry	Catalytic System	Substrate	k_i value (s^{-1})	Rate ($mol\ dm^{-3}\ s^{-1}$)
1	1	<i>p</i> -Methoxystyrene	$7.89 \times 10^{-5} \pm 3.57 \times 10^{-6}$	$1.46 \times 10^{-8} \pm 6.06 \times 10^{-10}$
2	1	<i>p</i> -Methylstyrene	$9.29 \times 10^{-5} \pm 2.59 \times 10^{-6}$	$1.72 \times 10^{-8} \pm 4.79 \times 10^{-10}$
3	1	Styrene	$1.17 \times 10^{-4} \pm 4.07 \times 10^{-6}$	$2.16 \times 10^{-8} \pm 7.51 \times 10^{-9}$
4	1	<i>p</i> -Fluorostyrene	$1.32 \times 10^{-4} \pm 6.05 \times 10^{-5}$	$2.44 \times 10^{-8} \pm 1.11 \times 10^{-9}$
5	1	<i>p</i> -Chlorostyrene	$1.40 \times 10^{-4} \pm 5.63 \times 10^{-6}$	$2.59 \times 10^{-8} \pm 1.04 \times 10^{-9}$
6	1	<i>p</i> -Bromostyrene	$1.52 \times 10^{-4} \pm 1.25 \times 10^{-5}$	$2.81 \times 10^{-8} \pm 2.31 \times 10^{-9}$
7	1	<i>p</i> -Nitrostyrene	$1.86 \times 10^{-4} \pm 1.76 \times 10^{-5}$	$3.44 \times 10^{-8} \pm 3.26 \times 10^{-9}$
8	1	<i>p</i> -Nitrilestyrene	$3.29 \times 10^{-4} \pm 8.08 \times 10^{-6}$	$6.09 \times 10^{-8} \pm 1.50 \times 10^{-9}$
9	2	<i>p</i> -Methoxystyrene	$1.85 \times 10^{-3} \pm 1.12 \times 10^{-4}$	$3.41 \times 10^{-7} \pm 2.07 \times 10^{-8}$
10	2	<i>p</i> -Methylstyrene	$1.08 \times 10^{-3} \pm 7.08 \times 10^{-5}$	$1.99 \times 10^{-7} \pm 1.44 \times 10^{-8}$
11	2	Styrene	$7.79 \times 10^{-4} \pm 2.13 \times 10^{-5}$	$1.44 \times 10^{-7} \pm 3.93 \times 10^{-9}$
12	2	<i>p</i> -Fluorostyrene	$3.80 \times 10^{-4} \pm 1.94 \times 10^{-5}$	$7.03 \times 10^{-8} \pm 3.58 \times 10^{-9}$
13	2	<i>p</i> -Chlorostyrene	$3.04 \times 10^{-4} \pm 2.73 \times 10^{-5}$	$5.62 \times 10^{-8} \pm 5.05 \times 10^{-9}$
14	2	<i>p</i> -Bromostyrene	$5.81 \times 10^{-4} \pm 1.75 \times 10^{-5}$	$1.07 \times 10^{-7} \pm 3.24 \times 10^{-9}$
15	2	<i>p</i> -Nitrostyrene	$0.0708 \pm 2.92 \times 10^{-3}$	$1.31 \times 10^{-5} \pm 5.40 \times 10^{-6}$
16	2	<i>p</i> -Nitrilestyrene	0.515 ± 0.0209	$9.52 \times 10^{-5} \pm 3.87 \times 10^{-6}$




Figure 1: Hammett plots for both systems investigated.

Both systems exhibit unusual Hammett plots for manganese-containing epoxidation catalysts.¹⁰ The positive ρ -value (+0.413) obtained in the case of ligand **7** and $MnSO_4$ system indicates that the oxygen is delivered to styrene by the catalyst in a nucleophilic fashion. Thus, it seems unlikely that electrophilic $Mn^V=O$ species, which are often invoked as key intermediates in the oxidation of organic substrates, are involved. The non-linearity in the Hammett plot obtained for ligand **8** and $Mn(OTf)_2$ system is indicative of a fundamental

change.¹⁸ With electron donating groups the catalyst exhibits typical electrophilic behaviour having significant ρ -value of -1.25. However, with electron withdrawing groups the nucleophilic nature of the catalytic species takes over and dramatically influences the ρ -value (+3.64).




Figure 2: UV/vis spectra of catalytically active Systems 1 and 2 in the presence of H₂O₂ and PAA and substrate at various time intervals. a) Mn(7)SO₄ with H₂O₂; b) Mn(7)SO₄ with PAA; c) Mn(8)(OTf)₂ with PAA; d) Mn(8)(OTf)₂ with H₂O₂; e) expansion of Figure 2c showing the weak absorption band at 1090 nm for Mn(8)(OTf)₂ with PAA; f) time-resolved UV-Vis at 1090 nm for Mn(8)(OTf)₂ with PAA.

A brief examination of the active Systems 1 and 2 using UV-vis spectroscopy further highlighted differences between them. The stoichiometric reaction between Mn(7)SO₄ and H₂O₂ resulted in the build-up of a putative manganese(IV) species (**Figure 2a**).^{19,8f} The

addition of 1 equivalent of styrene resulted in the rapid reduction in this species within 5 minutes and within 5 hours only manganese(III) species could be observed by UV-vis spectroscopy. Identical treatment of Mn(7)SO_4 with PAA also initially produced a similar band in the UV-vis spectrum (**Figure 2b**), however this combination of reagents is inactive in the epoxidation of styrene, and it also decomposed within 10 minutes. Any further addition of PAA to the reaction mixture resulted in rapid decomposition of the oxidant, with no change in the UV-vis spectra being observed.

Monitoring of the reaction between Mn(8)(OTf)_2 and PAA using UV-vis spectroscopy did not provide any supporting evidence for the significant formation of the apparent manganese(IV) species observed with Mn(7)SO_4 (**Figure 2c**), although addition of the substrate to the species generated at 20 minutes resulted in very similar spectra. Interestingly, a weak absorption band at 1090 nm consistently appeared within the first minute of the addition of PAA and then rapidly disappeared. Similar bands have been previously reported in related complexes.²⁰ In contrast, treating Mn(8)(OTf)_2 with H_2O_2 (**Figure 2d**) appears to produce a similar manganese(IV) species to those observed with Mn(7)SO_4 . Additionally another absorbance band at 966 nm is also present,¹⁹ however, the addition of styrene to the reaction mixture did not produce any distinct changes in the spectra. Figures 2b and 2d clearly show that while high oxidation state manganese-species can be obtained using both oxidants, they are not the same as the catalytically active species and appear to be unreactive.

In order to gain further insight into the different oxidative and catalytic behaviour exhibited by the manganese complexes of the two ligands, we decided to investigate the electrochemistry of the manganese centres. Due to the difference in solubility of Mn(7)SO_4 and Mn(8)(OTf)_2 and possible counter ion exchange in the presence of supporting electrolytes with poorly coordinating anions we elected to use Mn(7)Cl_2 and Mn(8)Cl_2 complexes.




Figure 3: Cyclic voltammograms of Mn(**7**)Cl₂ and Mn(**8**)Cl₂ in MeCN using (tBu₄N)(BF₄) as electrolyte, Pt-mesh counter electrode and Ag/AgCl as reference electrode.

As expected, both complexes Mn(**7**)Cl₂ and Mn(**8**)Cl₂ exhibit one electron oxidations at $E^{\circ} = +1.11$ vs Ag/AgCl, which we assign to a manganese(II/III) redox process, Figure 3.²¹ The complex Mn(**7**)Cl₂ displays a further clear and reversible oxidative process $E^{\circ} = +1.37$ vs Ag/AgCl with $\Delta E = 78$ mV, which we tentatively assign to a manganese(III/IV) redox couple. Whilst a similar process also appears to occur for Mn(**8**)Cl₂ with $E^{\circ} = +1.13$ vs Ag/AgCl with $\Delta E = 64$ mV, the wave is much broader and rather ill-defined. Nonetheless, this indicates that overall the formation of higher valent manganese species appears to be a less demanding process for Mn(**8**)Cl₂.

Whilst high valent Mn^V=O have long been recognised as active catalysts,^{22,1} recently alternative high valent Mn^{IV}=O species have also emerged as catalytically competent species.^{8f} Studies have revealed subtle differences in reactivity of these species not only in their rates of reaction, but also in the way the actual oxidation reaction occurs i.e. concerted oxygen transfer vs electron transfer. In the present study, which of these possible reactive species is present in the reaction is greatly influenced by the ligand. It well known that *N*-alkylation of secondary amine ligands affects the properties of their corresponding metal complexes through the combination of a number of effects such as: (i) decreasing the outer sphere solvation energy of their complexes; (ii) elimination of the M-N-H⁺⋯O hydrogen bonds, thus affecting the σ -donor power of nitrogen; (iii) elongation of M-N bonds due to steric hindrance and (iv) making nitrogen a better σ -donor due to increased electron donation due to hyperconjugation.²³ In the present case, the oxidation mechanism by which two similar systems **1** and **2** operate appears to be fundamentally different and highlights further how a simple change from N-H to N-Me within a ligand structure alters the overall reactivity. The secondary amine on ligand **7** appears to create an active high valent manganese(IV) species, which could be observed by UV-vis spectroscopy and is nucleophilic in nature. In contrast, the coordination of **8** results in a manganese centre which is easier to oxidise and which appears to form a reactive intermediate with electron rich styrenes that is electrophilic in nature. However, for electron deficient styrenes our observations of non-linearity in the Hammett plot appears to signify a change in the rate determining step and presumably the nature of the reactive intermediate.

Conclusion

We have synthesised chiral ligands and prepared their manganese(II) complexes, which showed efficacy in the epoxidation of styrenes using either hydrogen peroxide or peracetic acid as the oxidant. Despite the inclusion of *trans*-diaminocyclohexane in the ligand backbone, no asymmetric induction was observed in the final products. Nonetheless, some interesting mechanistic observations were made, which reveal that small changes in ligand architectures from **7** to **8** can induce profound changes in both reactivity and the nature of the active oxidant.

Acknowledgments

We are grateful to the EPSRC for financial support to GI (EP/H034838/1) and for the provision of the National Mass Spectrometry Service, University of Wales, Swansea. We are also grateful to Dr Aleksandra Pinczewska for assistance with the cyclic voltammetry.

Experimental:

All reagents were purchased from Aldrich, Acros, Merck or Fluka and were used without further purification unless otherwise stated. The hydrogen peroxide used in the epoxidation procedures was bought from Sigma Aldrich as a 30% solution in water. Acetonitrile was refluxed over CaH₂ for minimum of 6 h and distilled prior to use. All other solvents that were required to be anhydrous were obtained from solvent purifying system MBRAUN MB SPS-800. All water was obtained from an Elga Purelab Option distillation system. ¹H-NMR and ¹³C-NMR spectra were recorded on a 400 MHz and at 100.2 MHz on a Bruker AV400 or a Bruker AMX400 and referenced to the signal of tetramethylsilane (TMS) or residual protic solvent. IR spectra were recorded on a Perkin Elmer Spectrum 65 IR spectrometer equipped with ATR accessory unless otherwise stated. UV-Vis spectra were obtained on a HP 8453 spectrophotometer, absorption maxima (λ_{max}) are expressed in nm, the molar extinction coefficients (ϵ) are expressed in L mol⁻¹cm⁻¹. Electrospray ionisation mass spectrometry was obtained from the EPSRC National Mass Spectrometry Service, University of Wales, Swansea on a Thermofisher LTQ Orbitrap XL. Cyclic voltammetry was performed on an EG&G instruments Versastat II workstation, using tetrabutylammonium hexaphosphate as the support electrolyte, a platinum working electrode, a graphite counter electrode and a silver reference electrode in MeCN. Melting points were measured on a Stuart SMP3 melting point apparatus and are uncorrected. High Performance Liquid Chromatography was performed on a Perkin Elmer Series 200 instrument equipped with UV-Vis detector using an Eclips C18 reverse phase

column and Chiralcel OJ or Chiralcel OD normal phase chiral columns. The calibration procedures and curves, together with selected data and synthesis of styrenes and corresponding epoxides are provided in the ESI.

Resolution of 1,2-diaminocyclohexane was carried out according to previously reported procedure.²⁴ Compounds **10** and **11** were synthesised according to previously reported procedures.²⁵ 4-Nitrostyrene and 4-nitrilestyrene were prepared according to a modified procedure.²⁶ PAA was prepared according to previously reported procedure.^{8a-c}

(1R, 2R)-N¹, N¹, N²-tris(Pyridin-2-ylmethyl)cyclohexane-1,2-diamine 7.⁹

NaBH₃(CN) (0.620 g, 19.4 mmol, 2.00 equiv.) was added portion wise to a solution of (**R, R**)-**11** (2.00 g, 5.19 mmol) in dry MeOH (100 mL) under atmosphere of nitrogen. This was followed by a drop-wise addition of trifluoroacetic acid (0.800 mL, 10.4 mmol, 2.00 equiv.) and reaction was allowed to stir at r.t. for 16 h. NaOH (aq. 20% w/v; 30 mL) was then added to the reaction mixture very slowly and the mixture was extracted with DCM (3 x 100 mL). The combined organic phases were washed with brine (100 mL), dried over MgSO₄ and concentrated *in vacuo* to give **7** (1.98 g, 99%) as a yellow oil: $[\alpha]_D^{25} = -29.8$ (c = 0.66, CHCl₃), (an identical procedure was followed for ent-**7** (1*S*, 2*S*) isomer $[\alpha]_D^{25} = 29.2$ (c = 1, CHCl₃)). (1*R*, 2*R*) isomer; IR (neat, $\nu_{\max}/\text{cm}^{-1}$): 3049, 3006, 2927, 2855, 1590, 1568, 1473, 1432, 755; ¹H-NMR (400 MHz, δ -CDCl₃): 0.84-1.31 (4H,m, CH₂CH₂CHN, CH₂CH₂CHN), 1.46-1.59 (1H, m, CH₂CH₂CHN), 1.61-1.75 (1H, m, CH₂CH₂CHN), 1.95-2.11 (2H, m, CH₂CH₂CHN, CH₂CH₂CHN), 2.34-2.52 (2H, m, CH₂CH₂CHN), 3.67 (3H, d, $J_1 = 13.9$ Hz, NCH₂Py), 3.77 (2H, d, $J_1 = 14.2$ Hz, NCH₂Py), 3.89 (1H, d, $J_1 = 13.9$ Hz, NCH₂Py), 4.00 (1H, broad s, NH) 6.92-7.03 (3H, m, CHPy), 7.17 (1H, d, $J_1 = 7.6$ Hz, CHPy), 7.41-7.52 (5H, m, CHPy), 8.30-8.35 (3H, m, CHPy); ¹³C-NMR (100.2 MHz, δ -CDCl₃): 22.9, 24.5, 25.4, 31.5, 52.2, 56.1, 57.7, 63.9, 121.6, 121.7, 122.1, 123.0, 136.1, 136.3, 148.6, 148.9, 159.8, 160.1; *m/z* (ESI): 197.8 (5 %), 311.7 (5), 388.2 (10), 410.2 (100); HRMS (ESI): [C₂₄H₂₉N₅+Na]⁺ requires 410.2302 found 410.2306.

(1R, 2R)-N¹-Methyl-N¹, N², N²-tris(pyridin-2-ylmethyl)cyclohexane-1,2-diamine 8.

(**R, R**)-**7** (1.60 g, 3.90 mmol) was dissolved in mixture of formaldehyde (30% solution in water, 10mL) and formic acid (15 mL). The mixture was heated at 85 °C for 48 h and cooled to r.t. The mixture was washed with Et₂O (50 mL) and the aqueous phase was cooled on ice. The

mixture was then basified (pH = 14) with NaOH (aq, 40% w/v) and NaCl was added until the solution became saturated. The product was extracted with Et₂O (6 x 30 mL). The combined extracts were dried over MgSO₄, and solvent was removed *in vacuo*. The residue (1.5 g) was re-dissolved in anhydrous Et₂O (30 mL) under atmosphere of nitrogen and filtered to remove any insoluble material. The clear yellow solution was then treated with dry HCl (2M ether solution, 1.5 mL) and copious amounts of precipitate was formed. The solid was collected, washed with anhydrous Et₂O (2 x 10 mL) and re-dissolved in water. The mixture was then basified (pH = 14) with NaOH (aq, 40% w/v) and extracted with DCM (3 x 50 mL). The extracts were combined, dried over MgSO₄ and solvent was removed *in vacuo* to give **8** (1.23 g, 75%) as yellow oil; $[\alpha]_D^{25} = -6.6$ (c = 0.5, CHCl₃), (an identical procedure was used to prepare ent-**8** (1*S*, 2*S*) isomer, $[\alpha]_D^{25} = 7.2$ (c = 0.5, CHCl₃)). (1*R*, 2*R*) isomer; IR (neat, $\nu_{\max}/\text{cm}^{-1}$): 2927, 2852, 1589, 1567, 1472, 1432, 1355, 756; ¹H-NMR (400 MHz, δ -CDCl₃): 0.94-1.33 (4H, m, CH₂CH₂CHN, CH₂CH₂CHN), 1.60-1.70 (2H, m, CH₂CH₂CHN, CH₂CH₂CHN), 1.86-1.94 (1H, m, CH₂CH₂CHN), 2.04 (3H, s, CH₃), 2.08-2.13 (1H, m, CH₂CH₂CHN), 2.56 (1H, dt, $J_1 = 11.0$ Hz, $J_2 = 3.5$ Hz, CH₂CH₂CHN), 2.71 (1H, dt, $J_1 = 10.5$ Hz, $J_2 = 3.2$ Hz, CH₂CH₂CHN), 3.68 (2H, d, $J_1 = 2.2$ Hz, NCH₂Py), 3.81 (2H, d, $J_1 = 14.7$ Hz, NCH₂Py), 4.01 (2H, d, $J_1 = 14.7$ Hz, NCH₂Py), 6.96-7.09 (3H, m, CHPy), 7.41-7.50 (4H, m, CHPy), 7.53 (2H, d, $J_1 = 7.8$ Hz, CHPy), 8.35-8.44 (3H, m, CHPy); ¹³C-NMR (100.2 MHz, δ -CDCl₃): 23.8, 25.6, 25.9, 27.8, 36.2, 56.5, 60.0, 61.0, 65.4, 121.60, 121.67, 123.1, 123.2, 136.0, 136.2, 148.5, 148.6, 160.6, 161.3; m/z (ESI): 203.1 (15 %), 280.1 (5), 402.2 (100); HRMS (ESI): [C₂₅H₃₁N₅ + H]⁺ requires 402.2657 found 402.2655.

General procedure for the synthesis of manganese(II) complexes

The relevant manganese salt (1.0 equiv.) and ligand (1.0 equiv.) were dissolved in acetonitrile (10 mL). The reaction mixture was heated at reflux for 24 h. After cooling to room temperature diethyl ether was added to the mixture until copious amounts of precipitate was formed, which was collected by filtration. In some cases a thick oil formed from which the solvent was decanted and the oil re-dissolved in DCM (5 mL). The resultant solution was filtered through celite and the solvent was removed *in vacuo* to give the product.

[(1*R*, 2*R*)-*N*¹, *N*¹, *N*²-tris(Pyridin-2-ylmethyl)cyclohexane-1,2-diamino]manganese(II) sulfate: an off white solid, 59%: m.p. >220 °C; $[\alpha]_D^{25} = 88.7$ (c = 0.1, chloroform), $\mu_{\text{eff}} = 5.84$ μ_B ; IR (neat, $\nu_{\max}/\text{cm}^{-1}$): 2934, 2856, 1668, 1637, 1594, 1570, 1476, 1433, 1372, 1278, 1258,

1151, 1120, 1094, 1050, 999, 875, 765, 751, 637; m/z (ESI): 449 (100%), 441 (35), 424 (25), 366 (10); HRMS (ESI): $C_{24}H_{28}MnN_5$, requires 442.1749 found 442.1736; λ_{max} (DCM)/nm (ϵ /dm³mol⁻¹cm⁻¹) 258 (10.5×10^3), 263 (10.9×10^3), 268 (9.30×10^3), 322 (1.07×10^3).

[(1*R*, 2*R*)-*N*¹, *N*¹, *N*²-tris(Pyridin-2-ylmethyl)cyclohexane-1,2-diamino]manganese(II) chloride: an off white solid, 91%; m.p. >220 °C; $[\alpha]_D^{25} = 134.8$ ($c = 0.1$, chloroform); $\mu_{eff} = 5.89 \mu_B$; IR (neat, ν_{max}/cm^{-1}): 3050, 2932, 2959, 2414, 1647, 1603, 1590, 1569, 1474, 1434, 1265, 1150, 1081, 995, 760, 729, 698, 637, 620; m/z (ESI) 477 (100%), 388 (45), 297 (45), 221 (17), 181 (12); HRMS (ESI): $C_{24}H_{29}MnN_5Cl$ requires 477.1486 found 477.1484; λ_{max} (DCM)/nm (ϵ /dm³mol⁻¹cm⁻¹) 259 (9.47×10^3), 263 (10.0×10^3), 269 (7.08×10^3). Dissolution of the solid in a mixture of MeCN/DCM followed by slow evaporation yielded crystals suitable for single crystal X-ray diffraction. $C_{24}H_{31}N_5OCl_2Mn$, $M = 531.38$, monoclinic, $a = 9.5670(9)$ Å, $b = 15.0540(15)$ Å, $c = 9.7671(13)$ Å, $\beta = 119.292(2)^\circ$, $V = 1226.8(2)$ Å³, space group P1 2₁ 1, $Z = 2$, $T = 100(2)$ K, $m = 0.783$ mm⁻¹, MoK α ($\lambda = 0.71073$ Å), reflections collected 5802, independent reflections 5174, final R indices $[I > 2\sigma I]$ $R_1 = 0.0392$; $wR_2 = 0.0545$ (all data) $R_1 = 0.0309$, $wR_2 = 0.0524$. Data have been deposited at the Cambridge Crystallographic Database CCDC967858.

[(1*R*, 2*R*)-*N*¹, *N*¹, *N*²-tris(Pyridin-2-ylmethyl)cyclohexane-1,2-diamino]manganese(II) triflate: an off white solid, 85%; m.p. >220 °C; $[\alpha]_D^{25} = 84.4$ ($c = 0.1$, chloroform); $\mu_{eff} = 5.85 \mu_B$; IR (neat, ν_{max}/cm^{-1}): 2937, 1606, 1527, 1485, 1443, 1275, 1258, 1238, 1157, 1100, 1080, 1055, 1029, 930, 763, 636, 574; m/z (ESI) 402 (3%), 442 (25), 591 (100); HRMS (ESI): $C_{24}H_{28}MnN_5$ requires 441.1720 found 441.1720; λ_{max} (DCM)/nm (ϵ /dm³mol⁻¹cm⁻¹) 259 (13.7×10^3), 263 (14.3×10^3), 269 (10.5×10^3), 301 (1.11×10^3).

[(1*R*, 2*R*)-*N*¹, *N*¹, *N*²-tris(Pyridin-2-ylmethyl)cyclohexane-1,2-diamino]manganese(II) acetate: a brown solid, 79%; m.p. >220 °C; $[\alpha]_D^{25} =$ ($c = 0.1$, chloroform); $\mu_{eff} = 5.86 \mu_B$; IR (neat, ν_{max}/cm^{-1}): 3211, 2936, 2859, 1634 (v. broad), 1604, 1566, 1479, 1437, 1376, 1308, 1261, 1154, 1097, 1083, 1053, 1013, 996, 933, 879, 842, 764, 701, 637; m/z (ESI) 501 (100%, $[M - CH_3CO_2]^+$), 441 (15, $[M - (CH_3CO_2)_2 - H]^+$), 366 (15), 348 (5); HRMS (ESI): $C_{24}H_{28}MnN_5$ requires 441.1720 found 441.1719; λ_{max} (DCM)/nm (ϵ /dm³mol⁻¹cm⁻¹) 259 (13.1×10^3), 263 (13.6×10^3), 269 (11.0×10^3), 296 (2.46×10^3).

[(1R, 2R)-N¹, N¹, N²-tris(Pyridin-2-ylmethyl)cyclohexane-1,2-diamino]manganese(II) perchlorate: a pale pink solid, 82%; m.p. >220 °C; $[\alpha]_D^{25} = -5.4$ (c = 0.1, chloroform); $\mu_{\text{eff}} = 5.88 \mu_{\text{B}}$; IR (neat, $\nu_{\text{max}}/\text{cm}^{-1}$): 2931, 2860, 1726, 1605, 1571, 1484, 1441, 1262, 1155, 1086, 1016, 969, 930, 763, 734, 701, 622; m/z (ESI) 541 (100%, $[\text{M} - \text{ClO}_4]^+$); 501 (20, $[\text{M} - (\text{ClO}_4)_2 - \text{H} + \text{CH}_3\text{CO}_2]^+$), 477 (60), 441 (20, $[\text{M} - (\text{ClO}_4)_2]^+$), 402 (25), 384 (30), 366 (25), 294 (15), 221 (75); HRMS (ESI): C₂₄H₂₉MnN₅O₄Cl requires 541.1289 found 541.1276; λ_{max} (DCM)/nm ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$) 258 (10.4 x 10³), 263 (10.9 x 10³), 269 (8.46 x 10³), 302 (1.44 x 10³).

[(1R, 2R)-N¹-Methyl-N¹, N², N²-tris(pyridin-2-ylmethyl)cyclohexane-1,2-diamino]manganese(II) chloride: a light-yellow solid, 82%; m.p. >220 °C; $[\alpha]_D^{25} = 79.7$ (c = 0.1, chloroform); $\mu_{\text{eff}} = 5.90 \mu_{\text{B}}$; IR (neat, $\nu_{\text{max}}/\text{cm}^{-1}$): 2932, 2859, 1663, 1604, 1570, 1479, 1441, 1306, 1260, 1155, 1098, 1052, 1015, 870, 765, 638, 578; m/z (ESI) 491 (100%, $[\text{M} - \text{Cl}]^+$), 414 (20), 339 (10), 228 (15); HRMS (ESI): C₂₅H₃₁MnN₅Cl requires 491.1643 found 491.1636; λ_{max} (DCM)/nm ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$) 258 (8.41 x 10³), 263 (7.65 x 10³), 269 (5.88 x 10³), 325 (0.656 x 10³).

[(1R, 2R)-N¹-Methyl-N¹, N², N²-tris(pyridin-2-ylmethyl)cyclohexane-1,2-diamino]manganese(II) triflate: a light-yellow solid, 38%; m.p. >220 °C; $[\alpha]_D^{25} = 10.6$ (c = 0.1, chloroform); $\mu_{\text{eff}} = 5.85 \mu_{\text{B}}$; IR (neat, $\nu_{\text{max}}/\text{cm}^{-1}$): 2938, 2857, 1661, 1607, 1573, 1484, 1445, 1275, 1260, 1238, 1223, 1158, 1029, 949, 764, 637, 574; m/z (ESI) 605 (100%, $[\text{M} - (\text{CF}_3\text{SO}_3)]^+$), 528 (7), 491 (25, $[\text{M} - (\text{CF}_3\text{SO}_3)_2 + \text{H}_2\text{O} + \text{OH}]^+$); HRMS (ESI): C₂₅H₃₁F₃MnN₅O₃S requires 605.1480 found 605.1464; λ_{max} (DCM)/nm ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$) 258 (9.45 x 10³), 263 (9.77 x 10³), 269 (6.86 x 10³).

General epoxidation procedure using ligand 7 and H₂O₂. A solution of ligand 1 (1.0 mol%) in MeCN (1.0 mL) was delivered to a capped vial containing MnSO₄·H₂O (1.0 mol%) and the mixture was allowed to stir at r.t. for 30 min. Solutions of styrene in MeCN (0.50 M, 1000 μL, 1.0 equiv.) and anisole (0.10 M, 500 μL, see **note 1**) were then delivered to the reaction vial and mixture allowed to equilibrate for 15 min at room temperature. A solution of H₂O₂ (30% aq. 855 μL, 10 equiv.) in acetone (2500 μL) was then added to the reaction in one portion and aliquots (50 μL each) were sampled at specific time intervals starting from the addition of peroxide. The aliquots were filtered through a short pad of silica and MgSO₄ using THF as eluent to remove residual H₂O, peroxides and manganese-salts. The aliquots were then

analysed using HPLC and reaction profiles, yields, conversions and ee were recorded as appropriate.

Epoxidation procedure using ligand 8 and peracetic acid.

A solution of ligand **2** (1.0 mol%) in MeCN (1.0 mL) was delivered to a capped vial containing Mn(OTf)₂ (1.0 mol%) and the mixture was allowed to stir at r.t. for 15 min. Solutions of styrene in MeCN (0.50 M, 1000 μL, 1.0 equiv.) and anisole (0.10 M, 500 μL, **note 1**) were then delivered to the reaction vial followed by the addition of MeCN (2500 μL) and mixture allowed to equilibrate for 15 min at room temperature. The reaction mixture was then cooled on ice and PAA (1.2 equiv.) was added in one portion. The aliquots (50 μL each) were sampled at specific time intervals starting from the addition of oxidant. The aliquots were filtered through a short pad of silica and MgSO₄ using THF as eluent to remove residual PAA and Mn-salts. The aliquots were then analysed using HPLC and reaction profiles, yields, conversions and e.e. were recorded as appropriate.

Note 1: An equivalent amount of 1,2-dichlorobenzene was used as internal standard during the epoxidation of 4-nitrilestyrene due to the overlap of signals on HPLC trace.

References.

1. (a) R. G. Bergman, *Nature*, 2007, **446**, 391-393; (b) P. Saisaha, J. W. de Boer and W. R. Browne, *Chem. Soc. Rev.*, 2013, **42**, 2059-2074; (c) E. P. Talsi and K. P. Bryliakov, *Coord. Chem. Rev.*, 2012, **256**, 1418-1434; (d) D. J. Covell and M. C. White, *Angew. Chem. Int. Ed.*, 2008, **47**, 6448-6451.
2. (a) I. W. C. E. Arends, *Angew. Chem., Int. Ed.*, 2006, **45**, 6250-6252; (b) P. Saisaha, J.W. de Boer and W.R. Browne, *Chem. Soc.Rev.*, 2013, **42**, 2059-2074; (c) G. Shul'pin, *Dalton Trans.*, 2013, **42**, 12794-12818.
3. (a) T. Katsuki and K. B. Sharpless, *J. Am. Chem. Soc.*, 1980, **102**, 5974-5976; (b) R. A. Johnson and K. B. Sharpless, *Catalytic Asymmetric Synthesis*, Ed: I. Ojima, VCH, New York, 1993, chap. 4.1.
4. (a) W. Zhang, J. L. Leobach, S. R. Wilson and E. N. Jacobsen, *J. Am. Chem. Soc.*, 1990, **112**, 2801-2803; (b) E. N. Jacobsen, *Catalytic Asymmetric Synthesis*, Ed: I. Ojima, VCH, New York, 1993, chap. 4.2.
5. (a) R. Irie, K. Noda, Y. Ito, N. Matsumoto and T. Katsuki, *Tetrahedron Lett.*, 1990, **31**, 7345-7348; (b) T. Katsuki, *Coord. Chem. Rev.*, 1995, **140**, 189-214; (c) T. Katsuki, *Adv. Synth. Catal.*, 2002, **344**, 131-147.

6. for example: (a) M. Shibasaki, M. Kanai, S. Matsunaga and N. Kumagai, *Acc. Chem. Res.*, 2009, **42**, 1117-1127; (b) G. D. Faveri, G. Ilyashenko and M. Watkinson, *Chem. Soc. Rev.*, 2011, **40**, 1722-1760.
7. H. Doucet, T. Ohkuma, K. Murata, T. Yokozawa, M. Kozawa, E. Katayama, A. F. England, T. Ikariya, R. Noyori, *Angew. Chem. Int. Ed.*, 1998, **37**, 1703-1707.
8. (a) A. Murphy, G. Dubois and T. D. P. Stack, *J. Am. Chem. Soc.*, 2003, **125**, 5250-5251; (b) A. Murphy, A. Pace and T. D. P. Stack, *Org. Lett.*, 2004, **6**, 3119-3122; (c) A. Murphy and T. D. P. Stack, *J. Mol. Catal. A: Chem.*, 2006, **251**, 78-88; (d) L. Gomez, I. Garcia-Bosch, A. Company, X. Sala, X. Fontrodona, X. Ribas and M. Costas, *Dalton Trans.*, 2007, 5539-5545; (e) M. Wu, B. Wang, S. F. Wang, C. G. Xia and W. Sun, *Org. Lett.*, 2009, **11**, 3622-3625; (f) X. Wu, M. S. Seo, K. M. Davis, Y.-M. Lee, J. Chen, K.-B. Cho, Y. N. Pushkar., and W. Nam, *J. Am. Chem. Soc.*, 2011, **133**, 20088-20091; (g) T. W.-S. Chow, Y. Liu and C.-M. Che, *Chem. Commun.*, 2011, **47**, 11204-11206; (h) L. Gomez, I. Garcia-Bosch, A. Company, J. Benet-Buchholz, A. Polo, X. Sala, X. Ribas and M. Costas, *Angew. Chem., Int. Ed.*, 2009, **48**, 5720-5723; (i) G. Ilyashenko, D. Sale, M. Motevalli and M. Watkinson, *J. Mol. Catal. A: Chem.*, 2008, **296**, 1-8; (j) O. Cussó, I. Garcia-Bosch, X. Ribas, J. Lloret-Fillol and M. Costas, *J. Am. Chem. Soc.*, 2013, **135**, 14871-14878.
9. M. M. Hammoud, J. J. McKamie, M. J. Heeg and J. J. Kodanko, *Dalton Trans.*, 2008, 4843-4845.
10. G. Ilyashenko, G. D. Faveri, S. Masoudi, R. Al-Safadi and M. Watkinson, *Org. Biomol. Chem.*, 2013, **11**, 1942-1951.
11. K. Matsumoto, Y. Sawada, B. Saito, K. Sakai and T. Katsuki, *Angew. Chem., Int. Ed.*, 2005, **44**, 4935-4939.
12. D. Pijper, P. Saisaha, J. W. de Boer, R. Hoen, C. Smit, A. Meetsma, R. Hage, R. P. van Summeren, P. L. Alsters, B. L. Feringa, and W. R. Browne, *Dalton Trans.*, 2010, 10375-10381.
13. D. F. Evans, *J. Chem. Soc.* 1959, 2003-2005.
14. a) G. Chelucci, *Coord. Chem. Rev.*, 2013, **257**, 1887-1932; b) R. Hage, *Recl. Trav. Chim. Pays Bas*, 1996, **115**, 385-195; J. Glerup, P.A. Goodson, D.J. Hodgson, K. Michelsen, K.M. Nielsen, and H. Weihe, *Inorg. Chem.*, 1992, **31**, 4611-4616.
15. J. J. Blumenstein, V. C. Ukachukwu, R. S. Mohan and D. L. Whalen, *J. Org. Chem.*, 1993, **58**, 924-932.
16. H. E. Audier, J. F. Dupin and J. Jullien, *J. Bull. Soc. Chim. Fr.*, 1968, **9**, 3850.
17. (a) L. P. Hammett, *J. Am. Chem. Soc.*, 1937, **59**, 96-103; (b) C. Hansch, A. Leo and R. W.

Taft, *Chem. Rev.*, 1991, **91**, 165-195.

18 (a) P. R. Young and W. P. Jencks, *J. Am. Chem. Soc.*, 1979, **101**, 3288-3294; (b) A. R. Stein, M. Tencer, E. A. Moffatt, R. Dawe and J. Sweet, *J. Org. Chem.*, 1980, **45**, 3539-3540.

19. (a) S. C. Sawant, X. Wu, J. Cho, K.-B. Cho, S. H. Kim, M. S. Seo, Y.-M. Lee, M. Kubo, T. Ogura, S. Shaik, W. Nam, *Angew. Chem. Int. Ed.*, 2010, **49**, 8190–8194; (b) T. H. Parsell, R. K. Behan, M. T. Green, M. P. Hendrich and A. S. Borovik, *J. Am. Chem. Soc.*, 2006, **128**, 8728-8729; (c) I. Garcia-Bosch, A. Company, C. W. Cady, S. Styring, W. R. Browne, X. Ribas and M. Costas, *Angew. Chem. Int. Ed.*, 2011, **50**, 5648-5653.

20. X. Wu, M. S. Seo, K. M. Davis, Y.-M. Lee, J. Chen, K.-B. Cho, Y. N. Pushkar, and W. Nam, *J. Am. Chem. Soc.*, 2011, **133**, 20088–20091.

21. T. J. Hubin, J. M. McCormick, S. R. Collinson, M. Buchalova, C. M. Perkins, N. W. Alcock, P. M. Kahol, A. Raghunathan and D. H. Busch, *J. Am. Chem. Soc.*, 2000, **122**, 2512-2522.

22. for example: E. M. McGarrigle and D. G. Gilheany, *Chem. Rev.*, 2005, **105**, 1565–1602.

23. D. Meyerstein, *Coord. Chem. Rev.*, 1999, **185-186**, 141-147.

24. H.-J. Schanz, M. A. Linseis and D. G. Gilheany, *Tetrahedron: Asymmetry*, 2003, **14**, 2763-2769.

25. (a) H.-J. Cristau, P. P. Cellier, S. Hamada, J.-F. Spindler and M. Taillefer, *Org. Lett.*, 2004, **6**, 913-916; (b) W. Park, M. H. Shin, J. H. Chung, J. Park, M. S. Lah and D. Lim, *Tetrahedron Lett.*, 2006, **47**, 8841-8845.

26. G. A. Molander and A. R. Brown, *J. Org. Chem.*, 2006, **71**, 9681-9686.