

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

NHC-Catalyzed Oxidative Cyclization Reaction for the Synthesis of 3-Substituted Phthalides

Cite this: DOI: 10.1039/x0xx00000x

So Won Youn,* Hyoung Sub Song and Jong Hyub Park

Received 00th January 2012,

Accepted 00th January 2012


DOI: 10.1039/x0xx00000x

www.rsc.org/

An efficient NHC-catalyzed domino oxidation/oxa-Michael addition reaction of 2-alkenylbenzaldehydes has been developed to afford 3-substituted phthalides bearing a C₃-stereogenic center with a broad substrate scope and wide functional group tolerance. The preliminary results of asymmetric process have been provided as well.

Introduction

Phthalides, in particular, containing a C₃-stereogenic center, are versatile building blocks and pervasive motifs in many bioactive natural products (Fig. 1).¹ Consequently, a number of synthetic strategies have been developed for their construction, generally involving the use of chiral auxiliaries and resolutions, chiral organometallics, and transition metal catalysts.² However, organocatalytic synthesis of phthalides are very rare in the literature.³ Wang and Duan demonstrated an aldol-lactonization reaction between 2-formylbenzoates and ketones/aldehydes for the enantioselective synthesis of 3-substituted phthalides using proline-based secondary amine organocatalyst.^{3a} While N-heterocyclic carbenes (NHCs)⁴ have rapidly emerged as an important and powerful class of organocatalysts in various synthetic transformations, their uses in the synthesis of phthalides are very limited.^{3b-c} Scheidt and Chan showed only a single example, synthesis of 3-phenylphthalide, as an intramolecular variant in an NHC-catalyzed hydroacylation of activated ketones.^{3b} Recently, we reported an NHC-catalyzed oxidative cyclization of 2-alkynylbenzaldehydes under aerobic conditions that enabled the easy preparation of a diverse array of phthalides and isocoumarins (Scheme 1, above).^{3c} While it



Scheme 1

can be a more general method utilizing NHCs as an organocatalyst for the synthesis of 3-substituted phthalides, inseparable mixtures of two isomers, phthalides and isocoumarins, were obtained in most cases and, most of all, there is no stereogenic center at C₃ position of 3-substituted phthalides due to its *sp*²-carbon character. In view of the biological importance of the chiral 3-substituted phthalides, various benefits of organocatalysis as an attractive alternative to metal catalysis,⁵ and the potential of application in enantioselective oxidative NHC organocatalysis,^{4b-c} we were interested in developing a new, efficient method catalyzed by NHC for the facile construction of 3-substituted phthalide motif bearing a C₃-stereogenic center. In light of our recent success in NHC-catalyzed oxidative cyclization of 2-alkynylbenzaldehydes,^{3c} we naturally envisaged the use of 2-alkenylbenzaldehydes as a substrate for such a reaction (Scheme 1). On the other hand, NHC-catalyzed reactions of 2-alkenylbenzaldehydes under an inert atmosphere were reported




Fig. 1 Selected examples of naturally occurring 3-substituted phthalide derivatives.

to provide either spiro bis-indanes or dibenzo[8]annulenes via a domino Stetter-aldol-Michael or double Stetter reaction, respectively.⁶ Furthermore, there is another example to use the same substrates under NHC catalysis which underwent a domino aza-benzoin/intramolecular oxa-Michael addition reaction with nitrosoarenes to afford 2,3-benzoxazin-4-ones, which could be further transformed into a phthalide (R' = H, EWG = CO₂Me).⁷ Given the biological importance of the 3-substituted phthalides and the ever increasing necessity of new atom and step economical, sustainable, and eco-friendly methods for the synthesis of valuable compounds, an expedient and efficient synthetic protocol under atmospheric oxygen can lead to a facile and desirable route to the intriguing phthalide motif.

Herein we disclose the realization of this proposal. Noteworthy is that this process shows a broad substrate scope and wide functional group tolerance. Oxidation-cyclization sequence involving a 2-alkenylbenzoic acid intermediate was proposed and the preliminary results of asymmetric process have been provided.

Results and discussion

We began our studies on the proposed reaction using **1a** as the test substrate and examined the reaction parameters to identify optimal conditions (Table 1). In contrast to our previous work,^{3c}

Table 1 Optimization studies

1a

catalyst (20 mol%), base (40 mol%)
solvent (0.2 M), air, 80 °C, 12 h

2a

Legend for catalysts:

- R = (2,4,6-Me)₃Ph, X = Cl (A)
- R = Ph, X = Cl (B)
- R = C₆F₅, X = BF₄ (C)
- R = Bn, X = Cl (D)
- R = Et, X = Br (E)
- R = (2,4,6-Me)₃Ph (F)
- R = (2,6-*i*-Pr)₂Ph (H)
- R = (2,4,6-Me)₃Ph (G)
- R = (2,6-*i*-Pr)₂Ph (I)

| Entry | Catalyst | Base | Solvent | Yield (%) ^a |
|-----------------|----------------------|---|-----------------------------|------------------------|
| 1 | A | DBU | toluene | 60 |
| 2 | B or C | DBU | toluene | 34-38 |
| 3 | D or E | DBU | toluene | 16-17 |
| 4 | F | DBU | toluene | 36 |
| 5 | G | DBU | toluene | 55 |
| 6 | H or I | DBU | toluene | 24-26 |
| 7 | A | NEt₃ or DMAP | toluene | 96-100 |
| 8 | A | <i>i</i> Pr ₂ NEt | toluene | 84 |
| 9 | A | NaOAc or KHMDS | toluene | 71-73 |
| 10 | A | Na ₂ CO ₃ , K ₂ CO ₃ , Cs ₂ CO ₃ , KOH, or K ₃ PO ₄ | toluene | 26-61 |
| 11 | A | NaH or <i>t</i> BuOK | toluene | 20-34 |
| 12 | A | NEt ₃ | MeCN, dioxane, or DMF | 66-79 |
| 13 | A | NEt ₃ | <i>t</i> BuOH, DMSO, or DCE | 41-53 |
| 14 | A | NEt ₃ | THF, acetone, or EtOH | 13-28 |
| 15 ^b | A | NEt ₃ | toluene | 80-85 |
| 16 ^c | A | NEt ₃ | toluene | 72 |
| 17 ^d | A | NEt ₃ | toluene | 89 |

^a Yields were determined by ¹H NMR using trichloroethylene as an internal standard. ^b At 60 °C under either air or O₂ (1 atm). ^c With 10 mol% **A** and 20 mol% NEt₃ for 22 h. ^d In toluene (0.1 M).


the combination of triazolium salt **A** and NEt₃ or DMAP in toluene proved the most superior for this reaction among a variety of heterazolium salts, bases, and solvents examined. Finally, the optimal result was obtained in the presence of 20 mol % **A** and 40 mol % NEt₃ in toluene (0.2 M) at 80 °C, providing **2a** in 96% yield (by ¹H NMR).

Table 2 NHC-catalyzed oxidative cyclization reaction of 2-alkenylbenzaldehydes

| Entry | R ¹ | R ² | R ³ | R ⁴ | Time (h) | Yield (%) ^a |
|------------------|----------------|--------------------------------------|----------------|---|----------|------------------------|
| 1 | H | H | H | CO ₂ Et (1a) | 12 | 86 (2a) |
| 2 ^{b,c} | H | H | H | CO ₂ Et (1a') | 9 | 47 (2a) |
| 3 | H | H | H | CO ₂ Me (1b) | 24 | 90 (2b) |
| 4 | H | H | H | CO ₂ <i>n</i> Bu (1c) | 24 | 88 (2c) |
| 5 | H | H | H | CO ₂ <i>t</i> Bu (1d) | 8 | 64 (2d) |
| 6 | H | H | H | CONMe ₂ (1e) | 24 | 80 (2e) |
| 7 | H | H | H | CN (1f) | 22 | 72 (2f) |
| 8 ^d | H | H | H | COMe (1g) | 24 | 79 (2g) |
| 9 ^d | H | H | H | COEt (1h) | 24 | 72 (2h) |
| 10 ^e | H | H | H | COPh (1i) | 10 | 52 (2i) |
| 11 ^c | H | H | H | P(=O)(OEt) ₂ (1j) | 48 | 84 (2j) |
| 12 | H | H | Me | CO ₂ Et (1k) | 7 | 79 (2k) |
| 13 | H | OMe | H | CO ₂ Et (1l) | 6 | 77 (2l) |
| 14 | H | OMe | H | CO ₂ <i>t</i> Bu (1m) | 72 | 92 (2m) |
| 15 | H | OMe | OMe | CO ₂ Et (1n) | 51 | 82 (2n) |
| 16 ^c | OMe | H | OMe | CO ₂ Me (1o) | 48 | 86 (2o) |
| 17 | H | -OCH ₂ CH ₂ O- | H | CO ₂ Et (1p) | 24 | 80 (2p) |
| 18 | H | Cl | H | CO ₂ Et (1q) | 6 | 69 (2q) |
| 19 | H | F | H | CO ₂ Et (1r) | 7 | 65 (2r) |

Reaction conditions: **1** (1 equiv), **A** (20 mol%), and NEt₃ (40 mol%) in toluene (0.2 M) at 80 °C under aerobic conditions, unless otherwise noted. ^a Isolated yield. ^b (*Z*)-Ethyl 3-(2-formylphenyl)acrylate (**1a'**) was used as a substrate. ^c At 100 °C. ^d At 60 °C. ^e At 25 °C.


With the optimized reaction conditions in hand, we set out to explore the substrate scope of this process. A variety of electron-withdrawing substituents at the alkene moiety were well tolerated in this reaction, thereby generating the corresponding phthalides **2** in moderate to good yields (Table 2, entries 1–11). α,β -Unsaturated ketones displayed higher reactivity than others, requiring lower reaction temperature (25–60 °C, Table 2, entries 8–10). Noteworthy is the fact that (*Z*)-isomer (**1a'**) required more forcing conditions for the reaction to take place and led to **2a** in only moderate yield even at elevated temperature (100 °C, Table 2, entry 2). Next, we proceeded to examine the effect of substituents on the aromatic ring of 2-alkenylbenzaldehydes (Table 2, entries 12–19). The yields remained equally good with both electron-donating and -withdrawing substituents on the aromatic ring. Electron-donating and -withdrawing substituents *para* to the alkene moiety (R²) had no significant effect on the reactivity. However, the *ortho* substituents (R¹) and electron-donating substituents *para* to the aldehyde moiety (R³) appeared to retard the reaction considerably, presumably as a consequence of steric hindrance and the reduced electrophilicity of the aldehyde carbon toward reaction with the NHC catalyst, respectively (Table 2, entries 15–17). Equally noteworthy is that this process can tolerate a

Table 3. NHC-Catalyzed Oxidative Cyclization Reaction for the Synthesis of Various Phthalides

Reaction conditions: **3** (1 equiv), **A** (20 mol%), and NEt_3 (40 mol%) in toluene (0.2 M) at 100 °C under aerobic conditions, unless otherwise noted. Isolated yields are given. ^a Using **B** instead of **A**. ^b At 120 °C. ^c At 80 °C. ^d Diastereomeric ratios (dr) = ~1:1. ^e Under O_2 atmosphere (10 atm).

variety of functional groups including methoxy, halogen, ester, amide, nitrile, ketone, and phosphonate moieties.


Subsequently, we explored the reaction of trisubstituted alkene substrates. To our delight, the oxidative cyclization reactions of **3** also proceeded successfully to afford the corresponding phthalides in moderate to good yields (Table 3), although they generally required more forcing conditions (higher reaction temperature or higher pressure of O_2) and/or longer reaction times for the satisfactory conversion, presumably due to steric and electronic effects. In addition, aryl-substituted alkene substrates (**1s-t**) afforded the related uncyclized products (**5s-t**) as the sole or major product, and attempts to induce further cyclization were unsuccessful (eqn (1)). These findings implicate that 2-alkenylbenzoic acids could be generated from an NHC-catalyzed preoxidation of 2-alkenylbenzaldehydes under aerobic conditions, which could be a plausible intermediate in this oxidative cyclization reaction, in close analogy to our previous report.^{3c, 4b-c, 8} Furthermore, these results also indicate the requirement of carbonyl or related electron-withdrawing functional groups having heteroatoms with lone pair electrons as an alkene's substituent for the subsequent cyclization reaction to take place. Carbonyl groups can be coordinated by the conjugate acid of the base, leading to a more electrophilic alkene moiety. In this regard, a conjugate acid could be generated from the deprotonation process either between excess NEt_3 and $-\text{CO}_2\text{H}$ group of the speculated intermediate (Scheme 3a) or between triazolium salt **A** and NEt_3 for the preparation of an NHC species⁹ (Scheme 3b) (*vide*



infra).


Next, we proceeded to the application of this reaction system to the formation of 6- and 7-membered ring products using **6** and **8**, respectively (eqn (2)). Instead of the expected lactone derivatives, however, indanone **7** and chromanone **9** were obtained exclusively via the Stetter reaction.^{4, 10}

Encouraged by the successful synthesis of various 3-substituted phthalides, we turned our attention to the enantioselective version of this oxidative cyclization reaction, which would be highly useful and sought after. We examined various chiral triazolium salts and reaction parameters for the asymmetric reaction of **1a** (Scheme 2). Very disappointingly, however, no or only modest stereoselectivities were obtained. Low reactivity of the reactions using a preformed carbene in the absence of additional base (condition B) indicates the requirement of the excess amount of base for efficiency of the reaction.

**Scheme 2** Preliminary results on NHC-catalyzed asymmetric oxidative cyclization reaction.

Based on our findings and by analogy with the mechanism proposed for the related NHC-catalyzed reactions under aerobic conditions,^{3c, 4b-c, 8, 11} two plausible mechanisms for this oxidative cyclization reaction can be conceived as follows (Scheme 3): 1) Oxidation-cyclization and 2) cyclization-oxidation sequence. In the oxidation-cyclization sequence, the Breslow intermediate **II** formed by an initial nucleophilic addition of an in situ generated NHC species to the aldehyde functional group is incorporated with electrophilic O_2 , followed by the intramolecular oxa-Michael reactions of the resulting 2-alkenylbenzoic acid intermediate **5a** (Scheme 3a). Alternatively, a cyclization-oxidation mechanism involving the intramolecular cyclization of 2-alkenylbenzaldehydes through a concerted or stepwise pathway prior to oxidation with O_2 could also be proposed (Scheme 3b).^{11a, 12} In the latter case wherein NHC is incorporated in the cyclization step, better asymmetric induction by chiral NHCs could be expected than our results shown in Scheme 2. Furthermore, observation of both no lactone formation in the reaction of **6** and **8** (eqn (2)) and very low reactivity in the absence of excess of base (condition B in Scheme 2) is contrary to a cyclization-oxidation pathway. The former results suggest that the Stetter reaction of **6** and **8** via a **II**-type intermediate is much faster than aldehyde oxidation reaction. In stark contrast, unfavorable 4-membered ring formation with significant ring strain seemingly prevented **1** and **3** from the Stetter reaction.¹³ Instead, oxidation precedes

preferentially the cyclization reaction, resulting in 5-membered ring formation.



Scheme 3 Proposed mechanism.

Therefore, although a mechanism remains elusive at this juncture, the oxidation-cyclization pathway seems to work in this protocol, promoting sequential oxidation/oxa-Michael addition reaction and involving 2-alkenylbenzoic acid intermediates (Scheme 3c). In addition, in a series of detailed ^1H NMR analyses during the reaction progress, the putative intermediate **5a** (and possibly other species such as **II** or **III**) could be observed (for details, see Supporting Information). In the cases of the reactions under condition B in Scheme 2, deprotonation of intermediate **5a** could be achieved by a chiral NHC (i.e., B: = NHC in cyclization catalytic cycle in Scheme 3a),^{9a, 14} whose conjugate acid coordinates to a carbonyl oxygen atom for the subsequent cyclization reaction to take place. No stereoinduction and low reactivity under this condition suggest that a tighter chiral pocket and excess base would be needed for better stereocontrol and reactivity, respectively.

Conclusions

In summary, we have developed an efficient NHC-catalyzed domino oxidation/oxa-Michael addition reaction of 2-alkenylbenzaldehydes. This protocol represents an atom/step economical, sustainable, and eco-friendly route for the facile construction of biologically important 3-substituted phthalides bearing a C3-stereogenic center. Equally noteworthy is that this process can tolerate a variety of functional groups.

The success of this sequential process could be achieved both by exploiting atmospheric oxygen as an oxygen atom source and by introducing an electron-deficient alkene moiety bearing heteroatoms with lone pair electrons activated by the in situ generated conjugate acid of the base. Noteworthy is the fact that molecular oxygen in air plays an essential role as the source of the "O" atom in this transformation, and the reaction atmosphere has a pronounced effect on the reaction outcomes since, as mentioned earlier, the similar NHC-catalyzed reactions of the same substrates under inert atmosphere afforded very different products through different reaction pathways (Scheme 1).⁶

Mechanistically, this protocol might follow the oxidation-cyclization sequence, involving 2-alkenylbenzoic acid intermediates. In addition, the preliminary results of our investigations into an enantioselective NHC-catalyzed oxidative cyclization reaction demonstrate that both a tight chiral pocket and excess base may be required for good stereocontrol and reactivity, respectively. These new findings may present new ideas and possibilities to develop an effective multistep catalysis with the combination of NHC and chiral cocatalyst. Further investigations along this direction will be reported in due course.

Experimental

General procedure for the NHC-catalyzed oxidative cyclization reactions of 2-alkenylbenzaldehydes

To a solution of the substrate **1**, **3**, **6**, or **8** (0.100 mmol, 1 equiv) in toluene (0.5 mL, 0.2 M) were added 2-mesityl-2,5,6,7-tetrahydropyrrolo[2,1-c][1,2,4]triazol-4-ium chloride (**A**) (5.4 mg, 0.020 mmol, 20 mol%) and NEt_3 (6 μL , 0.040 mmol, 40 mol%). The resulting mixture was stirred at the reported temperature for the reported time under aerobic conditions. After the reaction was completed, the reaction mixture was cooled to room temperature, diluted with distilled water, extracted with CH_2Cl_2 (three times), dried over MgSO_4 , and concentrate in vacuo. The residue was purified by column chromatography on silica gel to give the corresponding product.

Acknowledgements

This work was supported by both Basic Science Research Program and Nano-Material Technology Department Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology

(Nos. 2010-0007737, 2012R1A1A2041471, and 2012M3A7B4049654).

Notes and references

Department of Chemistry and Research Institute for Natural Sciences, Hanyang University, Seoul 133-791, Korea

† Electronic Supplementary Information (ESI) available: [Full experimental details, characterization data, and detailed results of screening studies]. See DOI: 10.1039/b000000x/

- (a) T. K. Devon and A. I. Scott, *Handbook of Naturally Occurring Compounds*; Academic Press: New York, 1975; Vol. 1, p 249; (b) R. C. Elderfield, *Heterocyclic Compounds*; Wiley: New York, 1951; Vol. 2, Chapter 2; (c) L. Ge, S.-K. S. Chan, H.-S. Chung and S.-L. Li, *Studies in Natural Products Chemistry*; A. Rahman, Ed.; Elsevier: Amsterdam, 2005; Vol. 32, p 611; (d) J. J. Beck and S.-C. Chou, *J. Nat. Prod.*, 2007, **70**, 891; (e) G. Lin, S. S.-K. Chan, H.-S. Chung and S. L. Li, *Stud. Nat. Prod. Chem.*, 2005, 611.
- (a) M. J. Xiaoang and Z. H. Li, *Curr. Org. Chem.*, 2007, **11**, 833. Chiral auxiliaries and resolutions: (b) R. Pedrosa, S. Sayalero and M. Vicente, *Tetrahedron*, 2006, **62**, 10400; (c) P. J. Choi, J. Sperry and M. A. Brimble, *J. Org. Chem.*, 2010, **75**, 7388 and references therein. Chiral organometallics: (d) P. V. Ramachandran, G.-M. Chen and H. C. Brown, *Tetrahedron Lett.*, 1996, **37**, 2205; (e) K. Knepper, R. E. Ziegert and S. Bräse, *Tetrahedron*, 2004, **60**, 8591; (f) H.-T. Chang, M. Jeganmohan and C.-H. Cheng, *Chem.—Eur. J.*, 2007, **13**, 4356; (g) B. M. Trost and A. H. Weiss, *Angew. Chem., Int. Ed.*, 2007, **46**, 7664. Transition metal catalysts: (h) M. C. Willis, *Angew. Chem., Int. Ed.*, 2010, **49**, 6026; (i) F. Luo, C. Pan and J. Cheng, *Synlett*, 2012, 357; (j) Y. Yamamoto, H. Nishiyama and K. Itoh, *J. Am. Chem. Soc.*, 2005, **127**, 9625; (k) D. H. T. Phan, B. Kim and V. M. Dong, *J. Am. Chem. Soc.*, 2009, **131**, 15608; (l) G. Lv, G. Huang, G. Zhang, C. Pan, F. Chen and J. Cheng, *Tetrahedron*, 2011, **67**, 4879; (m) L. Ackermann and J. Pospech, *Org. Lett.*, 2011, **13**, 4153.
- (a) H. Zhang, S. Zhang, L. Liu, G. Luo, W. Duan and W. Wang, *J. Org. Chem.*, 2010, **75**, 368; (b) A. Chan and K. A. Scheidt, *J. Am. Chem. Soc.*, 2006, **128**, 4558; (c) J. H. Park, S. V. Bhilare and S. W. Youn, *Org. Lett.*, 2011, **13**, 2228.
- For recent selected reviews, see: (a) S. J. Ryan, L. Candish and D. W. Lupton, *Chem. Soc. Rev.*, 2013, **42**, 4906; (b) S. De Sarkar, A. Biswas, R. C. Samanta and A. Studer, *Chem.—Eur. J.*, 2013, **19**, 4664; (c) C. E. I. Knappke, A. Imami and A. J. von Wangelin, *ChemCatChem*, 2012, **4**, 937; (d) H. U. Vora, P. Wheeler and T. Rovis, *Adv. Synth. Catal.*, 2012, **354**, 1617; (e) J. Douglas, G. Churchill and A. D. Smith, *Synthesis*, 2012, 2295; (f) J. Izquierdo, G. E. Hutson, D. T. Cohen and K. A. Scheidt, *Angew. Chem., Int. Ed.*, 2012, **51**, 11686; (g) X. Bugaut and F. Glorius, *Chem. Soc. Rev.*, 2012, **41**, 3511; (h) A. Grossmann and D. Enders, *Angew. Chem., Int. Ed.*, 2012, **51**, 314; (i) V. Nair, R. S. Menon, A. T. Biju, C. R. Sinu, R. R. Paul, A. Jose and V. Sreekumar, *Chem. Soc. Rev.*, 2011, **40**, 5336; (j) D. Enders, O. Niemeier and A. Henseler, *Chem. Rev.*, 2007, **107**, 5606; (k) N. Marion, S. Díaz-González and S. P. Nolan, *Angew. Chem., Int. Ed.*, 2007, **46**, 2988.
- For selected reviews, see: (a) Special issue on Organocatalysis: *Chem. Rev.*, 2007, **107**, 5413–5883; (b) P. I. Dalko and L. Moisan, *Angew. Chem., Int. Ed.*, 2001, **40**, 3726; (c) P. I. Dalko and L. Moisan, *Angew. Chem., Int. Ed.*, 2004, **43**, 5138; (d) C. F. Barbas III, *Angew. Chem., Int. Ed.*, 2007, **47**, 42; (e) D. W. C. MacMillan, *Nature*, 2008, **455**, 304; (f) A. Dondoni and A. Massi, *Angew. Chem., Int. Ed.*, 2008, **47**, 4638; (g) P. Melchiorre, M. Marigo, A. Carlone and G. Bartoli, *Angew. Chem., Int. Ed.*, 2008, **47**, 6138; (h) A. Moyano and R. Rios, *Chem. Rev.*, 2011, **111**, 4703; (i) F. Giacalone, M. Gruttadauria, P. Agrigento and R. Noto, *Chem. Soc. Rev.*, 2012, **41**, 2406.
- (a) E. Sánchez-Larios, J. M. Holmes, C. L. Daschner and M. Gravel, *Org. Lett.*, 2010, **12**, 5772; (b) E. Sánchez-Larios, J. M. Holmes, C. L. Daschner and M. Gravel, *Synthesis*, 2011, 1896.
- Z.-X. Sun and Y. Cheng, *Org. Biomol. Chem.*, 2012, **10**, 4088.
- For examples of NHC-catalyzed oxidation of aldehydes to the corresponding carboxylic acids, see: (a) J. Castells, H. Llitjós and M. Moreno-Mañas, *Tetrahedron Lett.*, 1977, **18**, 205; (b) M. Yoshida, Y. Katagiri, W.-B. Zhu and K. Shishido, *Org. Biomol. Chem.*, 2009, **7**, 4062; (c) L. Gu and Y. Zhang, *J. Am. Chem. Soc.*, 2010, **132**, 914; (d) V. Nair, V. Varghese, R. R. Paul, A. Jose, C. R. Sinu and R. S. Menon, *Org. Lett.*, 2010, **12**, 2653; (e) P.-C. Chiang and J. W. Bode, *Org. Lett.*, 2011, **13**, 2422; (f) J. Zhao, C. Mück-Lichtenfeld and A. Studer, *Adv. Synth. Catal.*, 2013, **355**, 1098 and refs 3c and 4b-c.
- For examples using the conjugate acid of the base used to generate the carbene to activate a basic functionality of the electrophilic reaction counterpart, see: (a) X. Zhao, D. A. DiRocco and T. Rovis, *J. Am. Chem. Soc.*, 2011, **133**, 12466; (b) Ref 3c.
- For reviews, see: (a) J. R. de Alaniz and T. Rovis, *Synlett*, 2009, 1189; (b) M. Christmann, *Angew. Chem., Int. Ed.*, 2005, **44**, 2632.
- For examples of NHC-catalyzed reactions involving esterification under aerobic conditions, see: (a) Y.-K. Liu, R. Li, L. Yue, B.-J. Li, Y.-C. Chen, Y. Wu and L.-S. Ding, *Org. Lett.*, 2006, **8**, 1521; (b) L. Lin, Y. Li, W. Du and W.-P. Deng, *Tetrahedron Lett.*, 2010, **51**, 3571; (c) B. Maji, S. Vedachalan, X. Ge, S. Cai and X.-W. Liu, *J. Org. Chem.*, 2011, **76**, 3016; (d) Y.-C. Xin, S.-H. Shi, D.-D. Xie, X.-P. Hui and P.-F. Xu, *Eur. J. Org. Chem.*, 2011, 6527; (e) J. J. Meng, M. Gao, Y. P. Wei and W. Q. Zhang, *Chem. Asian J.*, 2012, **7**, 872; (f) Y. Li, W. T. Du and W. P. Deng, *Tetrahedron*, 2012, **68**, 3611; (g) E. G. Delany, C.-L. Fagan, S. Gundala, A. Mari, T. Broja, K. Zeitler and S. J. Connon, *Chem. Commun.*, 2013, **49**, 6510; (h) E. G. Delany, C.-L. Fagan, S. Gundala, K. Zeitler and S. J. Connon, *Chem. Commun.*, 2013, **49**, 6513; (i) I. N. C. Kiran, K. Lalwani and A. Sudalai, *RSC Adv.*, 2013, **3**, 1695; (j) J. Zhao, C. Mück-Lichtenfeld and A. Studer, *Adv. Synth. Catal.*, 2013, **355**, 1098 and refs 3c, 4b-c and 8f.
- T. Soeta, Y. Tabatake, S. Fujinami and Y. Ukaji, *Org. Lett.*, 2013, **15**, 2088.
- It cannot be completely excluded that the Stetter reaction of **1** and **3** to form 4-membered rings occurs reversibly.
- For the examples using NHCs as a Brønsted base catalysts, see: (a) E. M. Phillips, M. Riedrich and K. A. Scheidt, *J. Am. Chem. Soc.*, 2010, **132**, 13179; (b) K. A. Jo, M. Maheswara, E. Yoon, Y. Y. Lee, H. Yun and E. J. Kang, *J. Org. Chem.*, 2012, **77**, 2924; (c) T. Boddaert, Y. Coquerel and J. Rodriguez, *Chem. Eur. J.*, 2011, **17**, 2266; (d) S. Wei, X.-G. Wei, X. Su, J. You and Y. Ren, *Chem. Eur. J.*, 2011, **17**, 5965; (e) C. E. I. Knappke, A. J. Arduengo, III, H. Jiao, J.-M. Neudörfl and A. J. von Wangelin, *Synthesis*, 2011, 3784; (f) G. A. Grasa, R. Singh, N. M. Scott, E. D. Stevens and S. P. Nolan, *Chem. Commun.*, 2004, 2890.