

Analyst

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Haptens, bioconjugates, and antibodies for penthiopyrad immunosensing

Cite this: DOI: 10.1039/x0xx00000x

Eric Ceballos-Alcantarilla,^b Antonio Abad-Fuentes,^a Vincenzo Aloisio,^b Consuelo Agulló,^b Antonio Abad-Somovilla,^b Josep V. Mercader^{a,*}

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Haptens, bioconjugates, and antibodies for highly sensitive immunochemical analysis of the new-generation fungicide penthiopyrad are described. Two haptens with equivalent carboxylated linkers were prepared, and the purified active esters were efficiently coupled to proteins. Results revealed a slightly different antibody-eliciting capacity of the two synthetic derivatives. All of the produced antibodies were specific for penthiopyrad, and showed affinity values in the nanomolar range.

Penthiopyrad (Fig. 1) is a new-generation pyrazole carboxamide fungicide that interferes with fungal cell respiration in complex II of the cytochrome system by inhibiting the succinate dehydrogenase activity (Group 7).¹ Penthiopyrad is active against many diseases caused by ascomycetes, deuteromycetes, and basidiomycetes,² and due to its unique mode and site of action, it has shown great potential as complementary active principle in order to fight against fungi resistance to other frequently used fungicides such as strobilurins



Penthiopyrad: R¹ = Me; R² = CH(Me)CH₂CHMe₂Hapten PPa: R¹ = (CH₂)₄CO₂H; R² = CH(Me)CH₂CHMe₂Hapten PPb: R¹ = Me; R² = (CH₂)₅CO₂H


Fig. 1 Chemical structures of penthiopyrad and the two synthetic haptens. Global minimum energy conformer of penthiopyrad calculated by CAChe/CONFLEX/MM3.

and sterol biosynthesis inhibitors.^{3,4} Penthiopyrad was first commercially registered in the United States in 2012 and it has been recently approved by the European Commission.⁵

Nowadays, antibody-based technologies constitute valuable complementary and/or alternative methods for chemical residue and contaminant analysis.⁶ In order to achieve a good acceptance and a widespread implementation of immunosensors in quality control laboratories, appropriate reagents are required and alternative analytical approaches are demanded. Therefore, research projects are needed aiming to increase the scientific knowledge about the processes and factors influencing the analytical properties of antibodies and the development of sensitive procedures.

For small chemicals, such as penthiopyrad, functionalized derivatives of the target molecule containing a linker arm and an activatable chemical group for conjugation to the carrier are usually essential.⁷⁻⁹ The main determinant groups of the target compound must be displayed unmodified in the immunizing conjugate. This rule can generally be achieved by introducing a carbohydrate aliphatic linear spacer at the proper site of the analyte.¹⁰ Moreover, optimum hapten design will benefit not only antibody selectivity and affinity but also assay performance.¹¹ The aim of this study was to investigate different derivatives of penthiopyrad with alternative linker tethering sites and to generate protein conjugates as well as selective and high-affinity antibodies in order to develop sensitive immunoassays for the rapid analysis of this fungicide.

Following those principles, two penthiopyrad haptens – hapten PPa and hapten PPb (Fig. 1) – were designed mimicking as much as possible the structural and electronic properties of the parent analyte. In each of them, the spacer arm was located at opposite positions of the penthiopyrad framework in order to favour alternative display orientations to the immune system. Hapten PPa came from the formal replacement of the pyrazole *N*-methyl group by a C5 carboxylated aliphatic linear chain, which presumably only implied a minimum modification of the electronic and steric properties of the parent compound, whereas in hapten PPb a similar linear spacer arm replaced the branched 4-methylpentan-2-yl substituent at the 2-position of the thiophene ring. In the latter case, there was a greater alteration of the penthiopyrad framework, especially from the steric point of view, but it only affected the proximal part of the hapten in the immunogenic conjugate.



Scheme 1 Schematic representation of the synthesis of hapten PPa (A) and hapten PPb (B).

Since none of the two designed haptens can be prepared by chemical transformations from penthiopyrad or other structurally related accessible compounds, it was necessary to develop a total synthesis of both molecules from commercially available materials. The synthetic sequences developed for the preparation of haptens PPa and PPb are depicted in Schemes 1A and 1B, respectively. Every intermediate of the synthesis as well as the final haptens were purified and fully characterized by spectroscopic methods (for details see the ESI[†], Schemes S1 to S3). A brief discussion of each of the synthetic procedures is also given in the ESI[†]. The synthesis of hapten PPa was accomplished via a six steps sequence in a 28% overall yield, while the synthesis of hapten PPb required seven reaction steps with a calculated overall yield of ca. 30%. In both cases, the overall yields can be considered very satisfactory due to the number and type of reactions that were involved, showing the effectiveness of the developed synthetic sequences.

Carboxyl-functionalized haptens were activated using *N,N'*-disuccinimidyl carbonate in acetonitrile following an adapted strategy that was optimized in previous studies for other compounds,¹² which allowed a straightforward purification and characterization of the resulting active *N*-hydroxysuccinimidyl esters. Immunizing conjugates with BSA, and assay conjugates with OVA and HRP were readily prepared by adding the corresponding pure active ester of the hapten over buffered protein solutions. Data

for hapten activation and conjugation are given in the ESI[†]. Finally, covalent hapten-to-protein complexes were purified by size exclusion chromatography. This step-by-step approach avoided the formation of undesired secondary products during protein coupling, and therefore the same strategy could be employed both for immunizing and assay conjugate preparation, even for polyclonal antibody-based immunoassays. Moreover, since the conjugation reaction was more efficient, lower amounts of hapten were required. The number of conjugated hapten molecules was determined by Matrix-Assisted Laser Desorption Ionization Time-of-Flight Mass Spectrometry (MALDI-TOF/MS). The hapten-to-protein molar ratios (MR) obtained for the immunogens were 19.2 and 15.1 for conjugates BSA-PPa and BSA-PPb, respectively (Fig. S1 in the ESI[†]). The MRs for assay conjugates were 2.7 and 3.2 for conjugates OVA-PPa and OVA-PPb, respectively, and 2.9 and 2.2 for conjugates HRP-PPa and HRP-PPb, respectively (see Figs. S2 and S3 in the ESI[†]).

Animal manipulation was performed in compliance with the laws and guidelines of the Spanish Ministry of Agriculture, Food, and Environment. Antibodies were generated in rabbits by immunization with 0.3 mg of BSA-hapten conjugate using Freund's adjuvants. Partial antibody purification was performed by salting out twice. Further details are provided in the ESI[†]. Antibody-coated direct cELISAs and conjugate-coated indirect cELISAs were

Table 1 Checkerboard assays in two cELISA formats (n = 3)

Direct cELISA								
Tracer conjugate								
HRP-PPa					HRP-PPb			
Ab	[Ab] ^a	[T] ^b	A _{max}	IC ₅₀ (nM)	[Ab]	[T]	A _{max}	IC ₅₀ (nM)
PPa#1	10	10	1.22 ± 0.16	4.39 ± 0.21	3	300	ns ^c	---
PPa#2	10	10	1.24 ± 0.25	4.73 ± 0.62	3	300	0.99 ± 0.14	1.22 ± 0.16
PPb#1	3	100	0.83 ± 0.14	13.79 ± 4.71	3	100	1.09 ± 0.23	7.05 ± 3.07
PPb#2	3	100	0.80 ± 0.14	6.47 ± 1.08	10	30	0.88 ± 0.24	3.14 ± 0.99
Indirect cELISA								
Coating conjugate								
OVA-PPa					OVA-PPb			
Ab	[Ab]	[C] ^d	A _{max}	IC ₅₀ (nM)	[Ab]	[C]	A _{max}	IC ₅₀ (nM)
PPa#1	100	100	0.94 ± 0.12	3.25 ± 0.90	30	100	1.24 ± 0.09	3.29 ± 0.96
PPa#2	100	100	0.95 ± 0.13	2.23 ± 0.36	30	100	1.47 ± 0.45	1.83 ± 0.14
PPb#1	3	100	1.30 ± 0.03	6.06 ± 1.06	30	100	0.90 ± 0.14	22.37 ± 1.67
PPb#2	10	100	1.27 ± 0.10	9.27 ± 3.97	30	100	1.88 ± 0.32	34.84 ± 4.10

^a Antibody dilution (×10³). ^b Tracer concentration in ng mL⁻¹. ^c Signal was lower than 0.8. ^d Conjugate concentration in ng mL⁻¹.

performed using homologous and heterologous conjugates as described in the ESI.† Mean absorbance values were plotted versus the logarithm of analyte concentration, and assay sensitivity to penthiopyrad was estimated as the analyte concentration reducing 50% (IC₅₀) the maximum absorbance (A_{max}).

From each penthiopyrad hapten, two polyclonal antibodies were obtained; namely, antibodies PPa#1 and PPa#2 from hapten PPa, and antibodies PPb#1 and PPb#2 from hapten PPb. Those antibodies were evaluated by checkerboard cELISA with the two hapten conjugates. Eight-point standard curves were prepared by a 10-fold serial dilution, including a blank point. A 10 mM penthiopyrad stock solution in *N,N*-dimethylformamide was used to prepare the first point of the standard curve. One penthiopyrad standard curve and a different antibody or conjugate concentration was assayed in each microplate column. For the direct format, three different antibody dilutions (1/3000, 1/10000, and 1/30000) and four enzyme tracer concentrations (3, 10, 30, and 100 ng mL⁻¹) were assayed in the same plate. For indirect competitive assays, plates were coated with 100 or 1000 ng mL⁻¹ OVA conjugate, and diverse antibody dilutions were assayed from 1/3000 to 1/300000 in one plate. Thus, a series of twelve inhibition curves was retrieved from every plate; each curve corresponding to a particular antibody and assay conjugate combination at a specific concentration.

Sufficient assay signal (over 0.8) at zero dose of analyte was achieved with every antibody/conjugate combination in both assay formats, with the only exception of antibody PPa#1 which did not bind the heterologous enzyme tracer HRP-PPb (Table 1). All assays showed negligible background signals, in spite of using the same coupling procedure for immunizing and assay conjugate preparation. The observed IC₅₀ values were mostly in the low nanomolar range demonstrating the suitability of the prepared haptens and immunogens. The affinities for penthiopyrad of antibodies that were generated from hapten PPa were somewhat higher than those of the antibodies obtained from hapten PPb, indicating that hapten PPa was a better mimic of the target analyte as discussed previously for hapten design.

The IC₅₀ values of PPa-type antibodies could be lowered by using the heterologous conjugates with hapten PPb. Such improvement was also observed with PPb-type antibodies in the indirect assay format, but not in the direct format (Table 1). These results show the importance of preparing different functionalized

haptens, not only for high affinity antibody production but also for sensitive assay development. Linker site heterologies are adequate for assay sensitivity improvement; however, for particular antibodies and assay formats, this could be a too drastic modification making it difficult to achieve sufficient signal. In both cELISAs, the best results were obtained with antibody PPa#2 combined with the heterologous conjugate; IC₅₀ values of 1–2 nM with OVA-PPb or HRP-PPb, according to the assay format. The standard curves of the most sensitive immunoreagent combination by direct and indirect cELISA can be seen in Fig. S4 of the ESI.†

Antibody selectivity was studied by direct cELISA. Cross-reactivity (%) with other fungicides was determined from the quotient between the IC₅₀ values for penthiopyrad and for the corresponding assayed compound, both in molar concentration units. As competitors, different pyrazole bioactive principles (pyraclostrobin, fluxapyroxad, fluopyram, and fluopicolide) and other common fungicides (azoxystrobin, boscalid, pyrimethanil, cyprodinil, fenhexamid, fludioxonil, and tebuconazole) were evaluated. For PPa-type antibodies, no inhibition was found at a concentration of 10 μM with any of the studied compounds. Regarding the two PPb-type antibodies, a moderate inhibition was observed only with fluxapyroxad, which contains a similar pyrazole carboxamide group to that of penthiopyrad – the calculated cross-reactivity was below 0.5% for both PPb-type antibodies. This result is in accordance with Landsteiner's principle,¹³ stating that antibody specificity is predominantly directed towards the hapten moieties positioned opposite to the linker tethering site. In fact, the spacer arm location in hapten PPb was distal to the pyrazole ring.

In conclusion, two haptens were designed with equivalent carboxylated linkers at opposite tethering sites of the target molecule. Haptens PPa and PPb were achieved in six and seven steps, respectively, by total synthesis of the functionalized derivative. Purified active esters of the haptens were employed for efficiently preparing both the immunizing and the assay conjugates. High-affinity polyclonal antibodies to penthiopyrad were generated from both haptens. To our knowledge, these are the first reported haptens, bioconjugates, and antibodies to this modern fungicide. Antibodies from hapten PPa displayed a moderately higher affinity and selectivity than those obtained from hapten PPb. Replacement of the branched aliphatic substituent at the thiophene ring (Fig. 1) by the linear

1 hydrocarbon spacer arm (hapten PPb) did not hamper the
2 generation of antibodies with IC₅₀ values in the nanomolar
3 range – probably because such minor alteration was located at a
4 proximal position – though it was shown to be a slight
5 hindrance to antibody affinity. Finally, antibody PPa#2 was
6 identified as a promising candidate for sensitive immunosensor
7 development and sample analysis.

Valencia Proteomics Unit, a member of ISCIII
ProteoRedProteomics Platform.

Notes and references

^a Department of Biotechnology, Institute of Agrochemistry and Food
Technology, Consejo Superior de Investigaciones Científicas (IATA–
CSIC), Agustí Escardino 7, 46980 Paterna, València, Spain.


^b Department of Organic Chemistry, Universitat de València, Doctor
Moliner 50, 46100 Burjassot, València, Spain. Email:
jvmercader@iata.csic.es.

† Electronic Supplementary Information (ESI) available: General
methods and instruments, experimental details of hapten synthesis
(Schemes S1, S2, and S3), spectrometric characterization data, hapten
activation and conjugation details, MALDI spectra of conjugates,
antibody generation, competitive ELISA procedures, standard curves
(Fig. S1), and ¹H NMR spectra of haptens PPa and PPb. See
DOI: 10.1039/c000000x/

- 1 FRAC Code List 2013, www.frac.info, (accessed April 2014).
- 2 K. Culbreath, T. B. Brenneman, R. C. Kemerait Jr. and G. G.
3 Hammes, *Pest. Manag. Sci.*, 2009, **65**, 66.
- 4 K. L. Fairchild, T. D. Miles and P. S. Wharton, *Crop Protect.*,
5 2013, **49**, 31.
- 6 H. F. Avenot and T. J. Michailides, *Crop Protect.*, 2010, **29**,
7 643.
- 8 Commission Regulation (EU) No 1187/2013, 2013. *Off. J.*
9 *Eur. Union*, **L 313/42**, 1.
- 10 J. M. Van Emon, *J. AOAC Int.*, 2010, **93**, 1681.
- 11 M. Sathe, M. Dervini, G. Broadbent, A. Bodlener, K.
12 Charlton, B. Ravi, M. Rohmer, M. R. Sims and D. C. Cullen,
13 *Anal. Chim. Acta* 2011, **708**, 97.
- 14 X. Cai, K. Tsuchikama and K. D. Janda, *J. Am. Chem. Soc.*,
15 2013, **135**, 2971.
- 16 Z. L. Xu, Y. D. Shen, T. M. Sun, K. Campbell, T. X. Tian, S.
17 W. Zhang, H. T. Lei and T. M. Jiang, *Talanta*, 2013, **103**,
18 306.
- 19 J. Parra, J. V. Mercader, C. Agulló, A. Abad-Fuentes and A.
20 Abad-Somovilla, *Tetrahedron*, 2011, **67**, 624.
- 21 J. Parra, J. V. Mercader, C. Agulló, A. Abad-Somovilla and
22 A. Abad-Fuentes, *Anal. Chim. Acta*, 2012, **715**, 105.
- 23 F. A. Esteve-Turrillas, J. Parra, A. Abad-Fuentes, C. Agulló,
24 A. Abad-Somovilla and J. V. Mercader, *Anal. Chim. Acta*,
25 2010, **682**, 93.
- 26 K. Landsteiner, in *The Specificity of Serological Reactions*,
27 ed. Dover Publications, New York, revised edn., 1962.

Acknowledgements

This work was supported by the Spanish Ministerio de Ciencia e
Innovación (AGL2012-39965-C02-01) and cofinanced by
FEDER funds. J.V.M. is hired by the Consejo Superior de
Investigaciones Científicas (CSIC) under a Ramón y Cajal
contract, cofinanced by Ministerio de Ciencia e Innovación and
by the European Social Fund. E.C.-A. is recipient of a
predoctoral fellowship from the “Atracció de Talent, VLC-
CAMPUS” program of the University of Valencia. The
proteomic analysis was carried out in the SCSIE University of



Two penthiopyrad haptens with opposite linker tethering sites affording antibodies with affinities in the nanomolar range have been prepared with analytical purposes.

Graphical and textual abstract