

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/coxx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Heavy metal boryl chemistry: complexes of cadmium, mercury and lead

Andrey V. Protchenko, Deepak Dange, Andrew D. Schwarz, Christina Y. Tang, Nicholas Phillips, Philip Mountford, Cameron Jones and Simon Aldridge

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Abstract: Synthetic routes to the first boryl complexes of cadmium and mercury are reported via transmetalation from boryllithium; the syntheses of related Group 14 systems highlight the additional factors associated with extension to more redox-active post-transition elements.

Transition metal boryl complexes have attracted much recent attention,¹ in part due to their implication in unique C-H functionalization chemistry.^{2,3} More recently, the synthesis of reagents possessing nucleophilic character at boron^{4,5} has enabled the extension of such chemistry to the f- and early d- block elements.^{6,7} In addition, the large steric profile and strong σ -donor properties of the $\{(HCNDipp)_2B\}$ ligand has led to its use as a spoke ancillary substituent in the synthesis of Main Group carbene systems capable of facile E-H bond activation (E = H, C).⁸ While reagents such as $\{(HCNDipp)_2B\}Li(thf)_2$ are known to be strong reducing agents as well as powerful nucleophiles, the modulation of reactivity achieved for related alkyl/aryl systems by employing a less polar M-C bond,⁹ suggests that post transition metal boryl complexes represent synthetically valuable targets. Thus, for example, zinc boryl complexes have been reported and their conjugate addition to α,β unsaturated ketones explored.¹⁰ While related systems have been reported for the B-metal triads Cu, Ag, Au,^{10a} and Ga, In, Tl,¹¹ well-defined group 12 boryl complexes are confined to two zinc species (Fig. 1).^{10a} In the current study we targeted the first examples of cadmium and mercury boryls using a salt metathesis approach, and sought to explore whether this synthetic methodology could be extended to the more redox active metal lead. Moreover, in addition to homoleptic systems, routes to unsymmetrical (linear) complexes of the type (boryl)MX were sought, as these offer a convenient platform on which to study the σ donor properties of boryl ligands, free from complications involving π back bonding. Existing examples of Hg-B single bonds are confined to exopolyhedral linkages associated with borane clusters,¹² simple 2-centre 2-electron Cd-B and Pb-B bonds are hitherto unknown.¹³ The reactions of $\{(HCNDipp)_2B\}Li(thf)_2$ with $HgBr_2$ in non-polar solvents such as benzene provide a versatile access point to a range of donor-free mercury boryl systems (Scheme 1). Thus, reactions in 1:1 and 2:1 stoichiometries generate complexes of the composition $\{(HCNDipp)_2B\}HgBr$ (**1**) and $\{(HCNDipp)_2B\}_2Hg$ (**2**), respectively. Conversion of **1** into **2** can be effected by the addition of a second equivalent of $\{(HCNDipp)_2B\}Li(thf)_2$ to **1**, and the reverse transformation by mixing





Fig. 1 Previously reported group 12 metal boryl complexes.^{10a}



Scheme 1 Syntheses of borylmercury complexes **1-3**. Key reagents/conditions: (i) $\{(HCNDipp)_2B\}Li(thf)_2$ (0.85 equiv.), benzene, RT, 5 min, 36%; (ii) $\{(HCNDipp)_2B\}Li(thf)_2$ (2.00 equiv.), benzene-*d*₆, RT, 5 min, 43%; (iii) $HgBr_2$ (1.0 equiv.), benzene-*d*₆, RT, 5 min, 96%; (iv) $\{(HCNDipp)_2Ga\}K(OEt_2)_2$ (1.0 equiv.), benzene-*d*₆, RT, 5 min, 35 %.

equimolar quantities of **2** and $HgBr_2$. The composition of both mercury boryl complexes is suggested by multinuclear NMR and micro-analytical measurements and is confirmed by X-ray crystallography (Fig. 2). The monomeric structure of **2** features the linear two-coordinate geometry commonly associated with Hg(II) [$\angle(B-Hg-B) = 179.0(1)^\circ$], and comparison of the Hg-B bond lengths [2.150(3), 2.151(3) Å] with the sum of the respective covalent radii (2.16 Å),¹⁴ is consistent with descriptions as 2-centre 2-electron single bonds. Similar bond lengths have been reported by Hawthorne and co-workers for the Hg-B bonds associated with cyclic mercuracarborand systems.^{12c}

In the case of **1**, by contrast, the solid state structure is based around loosely bound centro-symmetric $\{[(HCNDipp)_2B]_2HgBr\}_2$ dimers, featuring a planar Hg_2Br_2 core [$\angle(Br-Hg-Br) = 80.3(1)^\circ$; $\angle(Hg-Br-Hg) = 99.7(1)^\circ$] and (superficially at least) a T-shaped mercury centre. Comparison of the two different Hg-Br distances [$d(Hg(1)-Br(2)) = 2.486(1) \text{ \AA}$; $d(Hg(1)-Br(1)) = 3.295(1) \text{ \AA}$] is consistent with weak association of the (boryl)HgBr units (*cf.* 2.52/3.40 Å for the sum of the respective covalent/van der Waals radii).¹⁴ The B(3)-Hg(1)-Br(2) angle [169.0(1)°] is, however, 10° more acute than the B-Hg-B angle determined for the strictly





Fig. 2 Molecular structures of **1** (upper left), **2** (upper right), one of the components of the asymmetric unit of **3** (lower left) and **4** (lower right). Here and elsewhere: H atoms omitted and 'Pr groups shown in wireframe format for clarity; thermal ellipsoids set at the 40% probability level. Key bond lengths (Å) and angles (°) for **1**: Hg(1)–B(3) 2.116(5), Hg(1)–Br(2) 2.486(1), Hg(1)–Br(2') 3.295(1), B(3)–Hg(1)–Br(2) 169.0(1). Key metrical parameters for **2-4** are listed in Table 1.

two-coordinate **2**, consistent with a small but significant distortion from linearity imposed by the secondary Hg–Br contacts.

The synthetic utility of **1** in giving access to further examples of heteroleptic mercury boryl systems can be demonstrated by its reaction with the anionic gallyl equivalent $\{[(\text{HCNDipp})_2\text{Ga}]\text{K}(\text{OEt}_2)_2\}$,^{15,16} which yields the mixed group 13 donor species $\{[(\text{HCNDipp})_2\text{B}]\text{Hg}\{[\text{Ga}(\text{NDippCH}_2)_2]\}$ (**3**) via salt metathesis chemistry. The synthesis of **3** is significant in that reactions of mercury(II) dihalides with the same gallyl anion lead to the reductive formation of mercury metal, rather than metathesis.¹⁷ In common with **2**, the structure of **3** features a linear co-ordination geometry at mercury [$\angle(\text{B-Hg-Ga}) = 179.1(1)^\circ$], with the shorter Hg–B distance [2.116(5) Å] measured for **3** being consistent with the weaker *trans* influence of the gallyl ligand compared to its boryl counterpart. Such an assertion is consistent with observations made previously for square planar systems, based on analyses of ¹⁹⁵Pt–³¹P coupling constants.¹⁸ The Hg–Ga linkage – the first of its kind involving three-coordinate gallium – features a bond length [2.476(1) Å] reflective of the difference in the covalent radii of gallium and boron ($\Delta d = 0.38$ Å). The only other example of a Hg–Ga bond, found in $\{[\text{HC}(\text{CMeNDipp})_2]\text{Ga}(\text{SC}_6\text{F}_5)_2\}\text{Hg}$, is markedly longer [2.534(1) Å], presumably due to the tetra-coordinate nature of the gallium centres.¹⁹

While Hg–B bonds have some precedent (albeit not for simple mono-boron ligand systems),¹² unsupported Cd–B and Pb–B bonds are hitherto unknown.¹³ As such, we hypothesized that salt metathesis chemistry utilizing $\{[(\text{HCNDipp})_2\text{B}]\text{Li}(\text{thf})_2\}$ and appropriate cadmium or lead electrophiles might allow access to a range of novel bond types. In the case of cadmium, both bromide and iodide precursors prove to be amenable to such chemistry (Scheme 2). Moreover, the presence of an additional neutral donor – such as *tmeda* (*N,N,N',N'*-tetramethylethylenediamine) modulates the observed reactivity, providing control of product stoichiometry. Thus, in the absence of such a donor, the reaction of cadmium dibromide with $\{[(\text{HCNDipp})_2\text{B}]\text{Li}(\text{thf})_2\}$ generates *bis*(boryl)cadmium complex **4**, irrespective of stoichiometry, with the bromo(boryl) system analogous to **1** presumably




Scheme 2 Syntheses of Cd complexes **4-6**. Key reagents/conditions: (i) $\{[(\text{HCNDipp})_2\text{B}]\text{Li}(\text{thf})_2\}$ (2.00 equiv.), benzene-*d*₆, RT, 12 h, 90%; (ii) $\{[(\text{HCNDipp})_2\text{B}]\text{Li}(\text{thf})_2\}$ (1.00 equiv.), thf, -78°C to 0°C , 5 h, 46%; (iii) $\{[(\text{HCNDipp})_2\text{Ga}]\text{K}(\text{tmeda})_2\}$ (0.50 equiv.), thf, -78°C to 20°C , 12 h, 70%.

being more labile to substitution than (sparingly soluble) CdBr₂. In the case of the more soluble precursor (*tmeda*)CdI₂, the use of one equiv. of boryllithium allows for the controlled introduction of a single boryl ligand yielding a heteroleptic system of the type (*tmeda*)Cd(boryl)I (i.e. **5**).¹⁷ Further substitution can then be effected, for example using a gallylpotassium reagent to give **6**.

By contrast, in the case of lead dihalide precursors (and to some extent their tin analogues), redox chemistry provides a competing reaction pathway to straightforward salt metathesis (Scheme 3). Thus, although the primary product of the reaction of SnCl₂ is $\{[(\text{HCNDipp})_2\text{B}]\}_2\text{Sn}$ (as reported previously),^{8a} a small quantity of a pentanuclear tin cluster species (**7**) is also obtained. **7** features four boryl ligands and consequently a mean metal oxidation state of 0.8; three different tin environments are revealed crystallographically featuring either zero [Sn(1) and Sn(3)], one [Sn(2) and Sn(5)] or two metal-bound boryl ligands [Sn(4)] (Fig. 3).²⁰ In the case of lead(II) bromide, by contrast, the only isolable product is the *bis*(boryl)lead complex, **8**, but the propensity for reduction is signalled by the formation of metallic lead and the relatively low yield of **8** (46%).

The structures of the *bis*(boryl) complexes of both cadmium (**4**) and lead (**8**) have been determined crystallographically, confirming the expected linear [$177.5(1)^\circ$] and bent geometries



Scheme 3 Syntheses of Sn/Pb complexes **7** and **8**. Key reagents/conditions: (i) $\{[(\text{HCNDipp})_2\text{B}]\text{Li}(\text{thf})_2\}$ (2.00 equiv.), Et₂O, RT, 1 h, ca. 4% [+51% Sn(boryl)₂];^{8a} (ii) $\{[(\text{HCNDipp})_2\text{B}]\text{Li}(\text{thf})_2\}$ (2.02 equiv.), benzene, RT, 5 min, 46%.




Fig. 3 Molecular structures of **7** (left), **8** (right). Key bond lengths (Å) and angles (°) for **7**: Sn(2)–B 2.290(5), Sn(5)–B 2.278(6), Sn(4)–B 2.595(7)/2.526(7), Sn(1)–Sn(2) 2.796(1), Sn(1)–Sn(4) 2.883(1), Sn(1)–Sn(5) 2.865(1), Sn(3)–Sn(2) 2.852(1), Sn(3)–Sn(4) 2.888(1), Sn(3)–Sn(5) 2.799(1). Key metrical parameters for **8** are listed in Table 1.

[118.3(1)^o] at the respective metal centres (Figs. 2 and 3). Salient structural parameters are included in Table 1 (along with those of related systems) allowing the following general trends to be identified for post-transition metal boryl complexes: (i) the stronger σ -donor properties of the boryl ligand over gallyl and (less surprisingly) bromide counterparts; (ii) M-B bond lengths within individual groups (*i.e.* 12 and 14) which conform to the trends predicted on the basis of covalent radii, as expected for 2-centre 2-electron single bonds; and (iii) the over-arching influence of the steric demands of the boryl substituent. The latter factor leads to shorter M-B bonds for linear vs. bent *bis*(boryl) derivatives, even to the extent of out-weighting the influence of covalent radius (e.g. for Cd, In, Sn and Hg, Tl, Pb), and presumably results from enhanced steric ‘buttressing’ between pendant Dipp groups as the B-M-B angle narrows.

Table 1 M-B bond lengths and B-M-B angles for *bis*(boryl) and related complexes of the heavier group 12, 13 and 14 metals

	$r(\text{M-B}) / \text{\AA}$	$\angle(\text{B-M-X}) / ^\circ$	$r_{\text{cov}}(\text{M})^{15}$
Zn(boryl) ₂ ^{10a}	2.052(3), 2.053(3)	178.5(1)	1.22
Cd(boryl) ₂ (4)	2.205(2), 2.207(2)	177.5(1)	1.44
Hg(boryl) ₂ (2)	2.150(3), 2.151(3)	179.0(1)	1.32
BrHg(boryl) (1)	2.116(5)	169.0(1)	1.32
(gallyl)Hg(boryl) (3)	2.116(5)	179.1(1)	1.32
In(boryl) ₂ ^{11a}	2.242(3), 2.246(3)	145.4(1)	1.42
Tl(boryl) ₂ ^{11a}	2.167(5), 2.173(6)	177.6(2)	1.45
Sn(boryl) ₂ ^{8a}	2.285(8), 2.294(8)	118.8(3)	1.39
Pb(boryl) ₂ (8)	2.363(4), 2.372(4)	118.3(1)	1.46

Steric considerations also appear to be a factor in modulating the reactivity of heavier p-block *bis*(boryl) systems. Thus for example, mercury system **2** is surprisingly unreactive towards a range of substrates, while the more accessible lead centre in **8** is extremely labile, and its tin counterpart shows a wide range of controlled reactivity involving both insertion and addition processes. Studies of this chemistry will be reported shortly.

We acknowledge financial support from the Leverhulme Trust, OUP John Fell Fund, the ARC and the EPSRC.

Notes and references


^a *Inorganic Chemistry Laboratory, Department of Chemistry, University of Oxford, South Parks Rd, Oxford, UK, OX1 3QR. Tel: 44 (0)1865 285201; E-mail: Simon.Aldridge@chem.ox.ac.uk*

§ Electronic Supplementary Information available: synthetic and characterizing data for all new compounds; CIFs for **1-4**, **7** and **8** (CCDC codes 950737, 950743-4 and 968284-6). See DOI: 10.1039/b000000x/

- For reviews of metal boryl chemistry see, for example: (a) G.J. Irvine, M.J.G. Lesley, T.B. Marder, N.C. Norman, C.R. Rice, E.G. Robins, W.R. Roper, G.R. Whittell and L.J. Wright, *Chem. Rev.*, 1998, **98**, 2685; (b) S. Aldridge and D.L. Coombs, *Coord. Chem. Rev.*, 2004, **248**, 535 (c) D.L. Kays and S. Aldridge, *Struct. Bonding (Berlin)*, 2008, **130**, 29; (d) L. Dang, Z. Lin and T.B. Marder, *Chem. Commun.*, 2009, 3987; (e) H. Braunschweig, R.D. Dewhurst and A. Schneider, *Chem. Rev.* 2010, **110**, 3924.
- I.A.I. Mkhaliid, J.H. Barnard, T.B. Marder, J.M. Murphy and John F. Hartwig, *Chem. Rev.* 2010, **110**, 890.
- For reviews of boryl complexes in other hydrocarbon functionalization processes see: (a) H.C. Brown and B. Singaram, *Pure Appl. Chem.*, 1987, **59**, 879; (b) K. Burgess and M.J. Ohlmeyer, *Chem. Rev.*, 1991, **91**, 1179; (c) I. Beletskaya and A. Pelter, *Tetrahedron*, 1997, **53**, 4957; (d) T.B. Marder and N.C. Norman, *Top. Catal.*, 1998, **5**, 63; (e) T. Ishiyama and N. Miyaara, *J. Organomet. Chem.*, 2000, **611**, 392; (f) H.E. Burks and J.P. Morken, *Chem. Commun.*,

- 2007, 4717; (g) P.J. Guiry, *ChemCatChem.*, 2009, **1**, 233.
- (a) Y. Segawa, M. Yamashita and K. Nozaki, *Science*, 2006, **314**, 5796; (b) Y. Segawa, Y. Suzuki, M. Yamashita and K. Nozaki, *J. Am. Chem. Soc.*, 2008, **47**, 16069.
- For a recent review of the chemistry of boryllithium reagents, see: M. Yamashita and K. Nozaki, *J. Synth. Org. Chem. Jpn.*, 2010, **68**, 4.
- T. Terabayashi, T. Kajiwarra, M. Yamashita and K. Nozaki, *J. Am. Chem. Soc.*, 2009, **131**, 14162.
- L.M.A. Saleh, K.H. Birjkumar, A.V. Protchenko, A.D. Schwarz, S. Aldridge, C. Jones, N. Kaltsoyannis and P. Mountford, *J. Am. Chem. Soc.*, 2011, **133**, 3836.
- (a) A.V. Protchenko, K.H. Birjkumar, D. Dange, A.D. Schwarz, D. Vidovic, C. Jones, N. Kaltsoyannis, P. Mountford and S. Aldridge, *J. Am. Chem. Soc.*, 2012, **134**, 6500; (b) A.V. Protchenko, A.D. Schwarz, M.P. Blake, C. Jones, N. Kaltsoyannis, P. Mountford and S. Aldridge, *Angew. Chem., Int. Ed.*, 2013, **52**, 568.
- See, for example C. Elschenbroich in *Organometallics: A Concise Introduction* 2nd edition, Wiley-VCH, Weinheim, 2006, chp 6.
- (a) T. Kajiwarra, T. Terabayashi, M. Yamashita and K. Nozaki, *Angew. Chem., Int. Ed.*, 2008, **47**, 6606; (b) Y. Segawa, M. Yamashita and K. Nozaki, *Angew. Chem., Int. Ed.*, 2007, **46**, 6710. See also: (c) Y. Nagashima, R. Takita, K. Yoshida, K. Hirano and M. Uchiyama, *J. Am. Chem. Soc.*, 2013, **135**, 18730.
- (a) A.V. Protchenko, D. Dange, J.R. Harmer, C.Y. Tang, A.D. Schwarz, M.J. Kelly, N. Phillips, R. Tirfion, K.H. Birjkumar, C. Jones, N. Kaltsoyannis, P. Mountford and S. Aldridge, *Nature Chem.*, in press, DOI: 10.1038/nchem.1870. For a Ga(III) boryl see: (b) N. Dettenrieder, C. Schädle, C. Maichle-Mössmer, P. Sirsch and R. Anwander, *J. Am. Chem. Soc.*, 2014, **136**, 886.
- (a) A.Y. Usiatinsky, O.M. Khitrova, P.V. Petrovskii, F.M. Dolgushin, A.I. Yanovsky, Y.T. Struchkov and V.I. Bregadze, *Mendeleev Commun.*, 1994, 169; (b) K.F. Shaw, B.D. Reid and A.J. Welch, *J. Organomet. Chem.*, 1994, **482**, 207; (c) Z. Zheng, C.B. Knobler, C.E. Curtis and M.F. Hawthorne, *Inorg. Chem.*, 1995, **34**, 432; (d) Z. Zheng, M. Diaz, C.B. Knobler and M.F. Hawthorne, *J. Am. Chem. Soc.*, 1995, **117**, 12338; (e) I.A. Lobanova, V.I. Bregadze, S.V. Timofeev, P.V. Petrovskii, Z.A. Starikova and F.M. Dolgushin, *J. Organomet. Chem.*, 2000, **597**, 48; (f) S.V. Timofeev, I.A. Lobanova, A.R. Kudinov, V.I. Meshcheryakov, O.L. Tok, P.V. Petrovskii, F.M. Dolgushin, Z.A. Starikova and V.I. Bregadze, *Izv. Akad. Nauk SSSR, Ser. Khim. (Russ.)*, 2000, 1609; (g) S.V. Timofeev, I.A. Lobanova, P.V. Petrovskii, Z.A. Starikova and V.I. Bregadze, *Izv. Akad. Nauk SSSR, Ser. Khim. (Russ.)*, 2001, 1604; (h) I.A. Lobanova, S.V. Timofeev, A.R. Kudinov, V.I. Meshcheryakov, P.V. Petrovskii, Z.A. Starikova and V.I. Bregadze, *Izv. Akad. Nauk SSSR, Ser. Khim. (Russ.)*, 2002, 480; (i) V.I. Bregadze, I.A. Lobanova, S.V. Timofeev, A.R. Kudinov, V.I. Meshcheryakov, O.L. Tok, P.V. Petrovskii, Z.A. Starikova; *J. Organomet. Chem.*, 2002, **657**, 171.
- Compounds have been structurally characterized featuring M-H-B interactions (M = Cd or Pb) or incorporating the metal in a delocalized cluster bonding framework, but direct metal-boron interactions are limited to a single plumbylene/B(C₆F₅)₃ complex: H. Arp, J. Baumgartner and C. Marschner, *J. Am. Chem. Soc.*, 2012, **134**, 6409.
- (a) B. Cordero, V. Gómez, A.E. Platero-Prats, M. Revés, J. Echeverría, E. Cremades, F. Barrigán and S. Alvarez, *Dalton Trans.*, 2008, **21**, 2832.
- R.J. Baker, R.D. Farley, C. Jones, M. Kloth and D.M. Murphy, *J. Chem. Soc., Dalton Trans.*, 2002, 3844.
- Attempts to synthesize Hg(I) systems of the type {(HCN-Dipp)₂B}Hg-Hg{B(NDippCH)₂} via reduction of **1** with KC₈ lead instead to the predominant formation of **2** and elemental mercury.
- For related chemistry leading to the formation of a system of the type (tmeda)Cd(gallyl)I see; O. Bonello, C. Jones, A. Stasch and W. Woodul, *Organometallics*, 2010, **29**, 4914.
- C. Jones, D.P. Mills, R.P. Rose and A. Stasch, *Dalton Trans.*, 2008, 4395.
- G. Prabusankar, C. Gemel, M. Winter, R.W. Seidel and R.A. Fischer, *Chem.-Eur. J.*, 2012, **16**, 6041.
- For a recent example of a metal-rich tin cluster featuring bulky terphenyl ligands, see E. Rivard, J. Steiner, J.C. Fettingier, J.R. Giuliani, M.P. Augustine and P.P. Power, *Chem. Commun.*, 2007, 4919.

Table of Contents



Heavy metal boryl complexes: the first boryl complexes of cadmium and mercury are reported via transmetalation from boryllithium; the syntheses of related tin and lead systems highlight the additional factors associated with extension to more redox-active post-transition elements.