

# ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

# Nickel-catalyzed chelation-assisted direct arylation of unactivated C(sp<sup>3</sup>)-H bonds with aryl halides

Mingliang Li, Jiaxing Dong, Xiaolei Huang, Kaizhi Li, Qian Wu, Feijie Song and Jingsong You\*

Received (in XXX, XXX) XthXXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXXXX 20XX


DOI: 10.1039/b000000x

In this work, we have disclosed the nickel-catalyzed unactivated  $\beta$ -C(sp<sup>3</sup>)-H bond arylation of aliphatic acid derivatives with aryl iodides/bromides via a bidentate chelation-assistance of an 8-aminoquinoline moiety. These preliminary results indicate the intrinsic catalytic potential of nickel metal for unactivated C(sp<sup>3</sup>)-H bond arylation.

The C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bonds extensively exist in synthetic products, naturally occurring molecules, pharmaceuticals, and bioactive compounds. The construction of C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bonds has attracted the great interest of organic chemists for a long time. Transition metal-catalyzed direct C-H bond functionalization has been one of the most promising methods for the C-C bond formation.<sup>1</sup> The C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bond formation via catalytic direct C(sp<sup>3</sup>)-H bond arylation relies heavily on palladium(II) catalysis.<sup>2-4</sup> In recent years, the nickel-catalyzed direct C-H bond functionalization has received more and more attentions due to the abundance, low cost and particular electronic properties of nickel metal.<sup>5</sup> In spite of significant efforts, the Ni-catalyzed direct C(sp<sup>3</sup>)-H arylation remains rare. In 2011, Hartwig and co-workers disclosed the nickel-catalyzed  $\alpha$ -arylation and heteroarylation of ketones with aryl halides.<sup>6</sup> Very recently, Lei and co-workers reported the nickel-catalyzed oxidative C(sp<sup>3</sup>)-H bond arylation of tetrahydrofuran and 1,4-dioxane with arylboronic acids.<sup>7</sup> However, these direct arylation reactions occurred at the activated  $\alpha$ -C(sp<sup>3</sup>)-H bonds adjacent to oxygen atoms (ether derivatives) and carbonyl group (ketones), and the nickel-catalyzed arylation of unactivated C(sp<sup>3</sup>)-H bonds is still a highly challenging subject.

The chelation-assistance strategy has become an efficient means for the functionalization of unactivated C-H bonds.<sup>8</sup> In particular, the 8-aminoquinoline group has recently attracted wider and wider attention since the seminal work of Daugulis in 2005.<sup>3a,3d,3j</sup> Various metal catalysts (e.g., Pd, Rh, and Ru) have been applied extensively, whereas the chelation-assisted examples involving nickel catalysts are rare. In 2011, Chatani and co-workers uncovered the first nickel-catalyzed *ortho*-C(sp<sup>2</sup>)-H bond activation to achieve regioselective oxidative cycloaddition of aromatic amides to alkynes with the bidentate auxiliary strategy.<sup>5c</sup> Lately, the same group developed the nickel-catalyzed chelation-assisted *ortho*-C(sp<sup>2</sup>)-H alkylation of aromatic amides.<sup>5d</sup> Inspired by these pioneering works, it is reasonable to assume that the activation of unactivated C(sp<sup>3</sup>)-H bonds would

be achieved through nickel catalysis with the assistance of bidentate directing group.<sup>9</sup> Herein, we wish to report the nickel-catalyzed unactivated primary  $\beta$ -C(sp<sup>3</sup>)-H bond arylation of aliphatic acid derivatives with aryl iodides/bromides (eqn (1)). During the submission of this manuscript, Chatani et al. reported the direct arylation of C(sp<sup>3</sup>)-H bonds of aliphatic amides by using the same directing group.<sup>10</sup> However, only aryl iodides were employed as the coupling partner in their work.




Considering 8-aminoquinoline has been proven to be a powerful bidentate auxiliary in transition metal-catalyzed direct C-H bond activation, a variety of  $\alpha,\alpha$ -disubstituted propionic acids were first decorated with 8-aminoquinoline to *N*-(quinolin-8-yl)propanamide derivatives. In our initial exploration, it was found that the reaction of 2-benzyl-3-(4-methoxyphenyl)-2-methyl-*N*-(quinolin-8-yl)propanamide **1a** with 1-iodo-4-methoxybenzene **2a** gave the desired product **3a** in 24% yield in the presence of Ni(OTf)<sub>2</sub>, PPh<sub>3</sub> and K<sub>2</sub>CO<sub>3</sub> in 1,4-dioxane at 140 °C for 24 h (ESI†, Table S1, entry 1). After screening various bases (e.g., K<sub>2</sub>CO<sub>3</sub>, Na<sub>2</sub>CO<sub>3</sub>, K<sub>3</sub>PO<sub>4</sub> and Li<sub>2</sub>CO<sub>3</sub>), Na<sub>2</sub>CO<sub>3</sub> gave the best yield of 35% (ESI†, Table S1, entries 1-4). With addition of DMSO and PivOH, the product yield was improved greatly to 83% at 160 °C (ESI†, Table S1, entry 9). Other nickel(II) catalysts such as Ni(OAc)<sub>2</sub> and NiCl<sub>2</sub> provided **3a** in moderate yields under otherwise identical conditions (ESI†, Table S1, entries 13 and 14). Interestingly, nickel(0) catalyst such as Ni(cod)<sub>2</sub> could also deliver **3a** in 62% yield (ESI†, Table S1, entry 15). The yield decreased to 54% when the reaction was performed in the absence of PPh<sub>3</sub> (ESI†, Table S1, entry 10). In addition, other frequently used directing groups such as pyridin-2-ylmethanamine and 2-(methylthio)aniline could not promote the occurrence of arylation. Thus, the best result was obtained by using Ni(OTf)<sub>2</sub> (10 mol%) as the catalyst, PPh<sub>3</sub> (20 mol%) as the ligand, PivOH (0.2 equiv) and DMSO (3.5 equiv) as the additives in dry 1,4-dioxane at 160 °C.

With the optimized reaction conditions in hand, we investigated the generality of the reaction of **1a** with aryl iodides as shown in Table 1. We found that the reactions of aryl iodides with electron-donating groups (except the *ortho*-substituted

methoxy group) could deliver the cross-coupled products in satisfactory yields (Table 1, **3a-3e**). Aryl iodides bearing electron-withdrawing groups such as *p*-CH<sub>3</sub>CO-, *p*-CO<sub>2</sub>Me, and *p*-CONEt<sub>2</sub> also proceeded smoothly to afford the desired products **5f**, **3g** and **3h** in 55%, 60% and 62% yields, respectively. Notably, a variety of functional groups such as alkyloxy, ketone, ester, amide, acetamido, and even free amino groups were tolerated under the optimized reaction conditions, which is useful in further synthetic transformations. We were pleased to find that aryl bromides could also undergo the β-C(sp<sup>3</sup>)-H bond arylation in synthetically useful yields (Table 1). It is worth to note that transition metal-catalyzed direct C(sp<sup>3</sup>)-H bond arylation with aryl bromides is still a difficult task in the synthetic organic community and is underrepresented to date. To the best of our knowledge, this is the first example to achieve transition metal-catalyzed chelation-assisted direct arylation of unactivated C(sp<sup>3</sup>)-H bonds with aryl bromides. These preliminary results indicated the intrinsic potential of nickel catalysis in the C(sp<sup>3</sup>)-H bond arylation with aryl bromides.

**Table 1** Nickel-catalyzed β-C(sp<sup>3</sup>)-H bond arylation of aliphatic amide **1a** with aryl halides<sup>a</sup>




<sup>a</sup> Conditions: Ni(OTf)<sub>2</sub> (10 mol%), PPh<sub>3</sub> (20 mol%), Na<sub>2</sub>CO<sub>3</sub> (2.0 equiv), PivOH (0.2 equiv), DMSO (3.5 equiv), amide **1a** (0.2 mmol), and aryl iodide (3.0 equiv) or aryl bromide (4.0 equiv) in dry 1,4-dioxane (1 mL) at 160 °C for 36 h. Isolated yields. <sup>b</sup> **2a** (2.0 equiv) for 24 h. <sup>c</sup> 22% of **1a** was recovered.

To explore the effect of the R substituent of the phenyl ring of aliphatic acid derivatives **1** on the primary β-C(sp<sup>3</sup>)-H bond arylation, an array of amides **1** were investigated accordingly. To our delight, α,α-dibenzyl substituted propionic acid derivatives bearing either electron-donating or electron-withdrawing substituent on the phenyl ring afforded the desired products in synthetically useful yields (Table 1, **3a**; Table 2, **3f-3l**). Notably, the current methodology was compatible with some important functional groups such as Ac, CO<sub>2</sub>Me, CONEt<sub>2</sub>, CN and CHO. The coupling reaction of 2-methyl-2-(naphthalen-2-ylmethyl)-3-phenyl-N-(quinolin-8-yl)propanamide with **2a** could also generate the β-arylated product **3m** in 79% yield.

To examine the synthetic usefulness of our protocol, we turned our attention to other aliphatic acid derivatives, while 2,2-dibenzyl-substituted propanamides could smoothly afford the arylated products **3**. As shown in Table 3, the nickel-catalyzed


C(sp<sup>3</sup>)-H bond arylation exhibited a relatively broad substrate scope for aliphatic acid derivatives. 2-Methyl-3-phenyl-acrylamide reacted with 1-iodo-4-methoxybenzene **2a** to afford the coupled product **4** in 58% yield. Isobutyramides could couple with **2a** to afford the single arylated products. For example, the reaction of 2,2-dimethyl-3-phenylpropanamide with **2a** delivered the desired product **5** in 60% yield. Isobutyramide with an α-ethyl substituent proceeded well to yield the arylated product **6** in 64% yield. Interestingly, the diarylated products could be obtained by increasing the amount of 1-iodo-4-methoxybenzene **2a** (Table 3, **7** and **8**). It should be noted that 2-methyl-2-phenylpropanamide selectively went through C(sp<sup>3</sup>)-H bond activation of the methyl group instead of C(sp<sup>2</sup>)-H bond of the phenyl ring to generate the arylated product **8**. In addition to the methyl group of quaternary carbon centers, the methyl group of tertiary carbon centers could also undergo direct C(sp<sup>3</sup>)-H bond arylation with aryl iodide. The reaction of *N*-(quinolin-8-yl)isobutyramide with **2a** provided the desired product **9** in 50% yield.

**Table 2** Nickel-catalyzed β-C(sp<sup>3</sup>)-H bond arylation of various aliphatic amides with 1-iodo-4-methoxybenzene **2a**<sup>a</sup>



<sup>a</sup> Conditions: Ni(OTf)<sub>2</sub> (10 mol%), PPh<sub>3</sub> (20 mol%), Na<sub>2</sub>CO<sub>3</sub> (2.0 equiv), PivOH (0.2 equiv), DMSO (3.5 equiv), amide **1** (0.2 mmol) and **2a** (3.0 equiv) in dry 1,4-dioxane (1 mL) at 160 °C for 36 h. Isolated yields.

**Table 3** Scope of aliphatic amides in Ni-catalyzed β-C(sp<sup>3</sup>)-H bond arylation<sup>a</sup>



<sup>a</sup> Conditions: Ni(OTf)<sub>2</sub> (10 mol%), PPh<sub>3</sub> (20 mol%), Na<sub>2</sub>CO<sub>3</sub> (2.0 equiv), PivOH (0.2 equiv), DMSO (3.5 equiv), amide **1** (0.2 mmol), and 1-iodo-4-methoxybenzene **2a** (3.0 equiv) in dry 1,4-dioxane (1 mL) at 160 °C for 36 h. Isolated yields. <sup>b</sup> **2a** (2.0 equiv) for 24 h. <sup>c</sup> 2-Nitrobenzoic acid instead of PivOH was used. <sup>d</sup> At 150 °C. <sup>e</sup> **2a** (4.0 equiv).

In summary, we have developed the nickel-catalyzed unactivated  $\beta$ -C(sp<sup>3</sup>)-H bond arylation of aliphatic acid derivatives with aryl iodides via a bidentate chelation-assistance of an 8-aminoquinoline moiety. Besides aryl iodides, aryl bromides are also capable of undergoing the cross-coupling reactions to a certain extent. Further studies to extend the scopes and clarify the detailed mechanism<sup>11</sup> are ongoing in our laboratory.

We thank the financial support from the National Basic Research Program of China (973 Program, 2011CB808601), and the National NSF of China (Nos 21272160, 21025205, and 21321061).

## Notes and references

Key Laboratory of Green Chemistry and Technology of Ministry of Education, College of Chemistry, and State Key Laboratory of Biotherapy, West China Medical School, Sichuan University, 29 Wangjiang Road, Chengdu 610064, PR China. Fax: (+86) 28-85412203; E-mail: jsyou@scu.edu.cn

† Electronic Supplementary Information (ESI) available: detailed experimental procedures and analytical data. See DOI:10.1039/b000000x/

- 1 For selected reviews on transition metal-catalyzed C-C bond formation via C-H bond cleavages, see: (a) X. Chen, K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2009, **48**, 5094; (b) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; (c) C. C. C. Johansson and T. J. Colacot, *Angew. Chem., Int. Ed.*, 2010, **49**, 676; (d) L. Ackermann, *Chem. Commun.*, 2010, **46**, 4866; (e) C. S. Yeung and V. M. Dong, *Chem. Rev.*, 2011, **111**, 1215; (f) L. Ackermann, *Chem. Rev.*, 2011, **111**, 1315; (g) C. Liu, H. Zhang, W. Shi and A. Lei, *Chem. Rev.*, 2011, **111**, 1780; (h) S.-Y. Zhang, F.-M. Zhang and Y.-Q. Tu, *Chem. Soc. Rev.*, 2011, **40**, 1937; (i) P. B. Arockiam, C. Bruneau and P. H. Dixneuf, *Chem. Rev.*, 2012, **112**, 5879; (j) D. Y.-K. Chen and S. W. Youn, *Chem.-Eur. J.*, 2012, **18**, 9452; (k) J. Yamaguchi, A. D. Yamaguchi and K. Itami, *Angew. Chem., Int. Ed.*, 2012, **51**, 8960.
- 2 For selected reviews, see: (a) R. Jazzar, J. Hitce, A. Renaudat, J. Sofack-Kreutzer and O. Baudoin, *Chem.-Eur. J.*, 2010, **16**, 2654; (b) O. Baudoin, *Chem. Soc. Rev.*, 2011, **40**, 4902; (c) H. Li, B.-J. Li and Z.-J. Shi, *Catal. Sci. Technol.*, 2011, **1**, 191.
- 3 For selected examples of palladium-catalyzed C(sp<sup>3</sup>)-H bond activation, see: (a) V. G. Zaitsev, D. Shabashov and O. Daugulis, *J. Am. Chem. Soc.*, 2005, **127**, 13154; (b) M. Lafrance, S. I. Gorelsky and K. Fagnou, *J. Am. Chem. Soc.*, 2007, **129**, 14570; (c) L.-C. Campeau, D. J. Schipper and K. Fagnou, *J. Am. Chem. Soc.*, 2008, **130**, 3266; (d) D. Shabashov and O. Daugulis, *J. Am. Chem. Soc.*, 2010, **132**, 3965; (e) W. R. Gutekunst and P. S. Baran, *J. Am. Chem. Soc.*, 2011, **133**, 19076; (f) M. Wasa, K. M. Engle, D. W. Lin, E. J. Yoo and J.-Q. Yu, *J. Am. Chem. Soc.*, 2011, **133**, 19598; (g) G. He and G. Chen, *Angew. Chem., Int. Ed.*, 2011, **50**, 5192; (h) M. Nakanishi, D. Katayev, C. Besnard and E. P. Kündig, *Angew. Chem., Int. Ed.*, 2011, **50**, 7438; (i) M. Wasa, K. S. L. Chan, X.-G. Zhang, J. He, M. Miura and J.-Q. Yu, *J. Am. Chem. Soc.*, 2012, **134**, 18570; (j) L. D. Tran and O. Daugulis, *Angew. Chem., Int. Ed.*, 2012, **51**, 5188; (k) S. Aspin, A.-S. Goutierre, P. Larini, R. Jazzar and O. Baudoin, *Angew. Chem., Int. Ed.*, 2012, **51**, 10808; (l) M. Fan and D. Ma, *Angew. Chem., Int. Ed.*, 2013, **52**, 12152; (m) N. Rodríguez, J. A. Romero-Revilla, M. Á. Fernández-Ibáñez and J. C. Carretero, *Chem. Sci.*, 2013, **4**, 175; (n) A. Millet, P. Larini, E. Clot and O. Baudoin, *Chem. Sci.*, 2013, **4**, 2241.
- 4 For selected examples of other metal-catalyzed C(sp<sup>3</sup>)-H bond arylation, see: (a) S. J. Pastine, D. V. Gribkov and D. Sames, *J. Am. Chem. Soc.*, 2006, **128**, 14220; (b) G. Deng, L. Zhao and C.-J. Li, *Angew. Chem., Int. Ed.*, 2008, **47**, 6278; (c) Y.-Z. Li, B.-J. Li, X.-Y. Lu, S. Lin and Z.-J. Shi, *Angew. Chem., Int. Ed.*, 2009, **48**, 3817; (d) N. Yoshikai, A. Mieczkowski, A. Matsumoto, L. Ilies and E. Nakamura, *J. Am. Chem. Soc.*, 2010, **132**, 5568; (e) N. Dastbaravardeh, M. Schnürch and M. D. Mihovilovic, *Org. Lett.*, 2012, **14**, 3792; (f) R. Shang, L. Ilies, A. Matsumoto and E. Nakamura, *J. Am. Chem. Soc.*, 2013, **135**, 6030.


- 5 (a) J. Yamaguchi, K. Muto and K. Itami, *Eur. J. Org. Chem.*, 2013, **19**; (b) Y. Nakao, E. Morita, H. Idei and T. Hiyama, *J. Am. Chem. Soc.*, 2011, **133**, 3264; (c) H. Shiota, Y. Ano, Y. Aihara, Y. Fukumoto and N. Chatani, *J. Am. Chem. Soc.*, 2011, **133**, 14952; (d) Y. Aihara and N. Chatani, *J. Am. Chem. Soc.*, 2013, **135**, 5308; (e) X. Wu, Y. Zhao and H. Ge, *J. Am. Chem. Soc.*, 2014, **136**, 1789.
- 6 S. Ge and J. F. Hartwig, *J. Am. Chem. Soc.*, 2011, **133**, 16330.
- 7 D. Liu, C. Liu, H. Li and A. Lei, *Angew. Chem., Int. Ed.*, 2013, **52**, 4453.
- 8 For selected reviews and highlights, see: (a) G. Rouquet and N. Chatani, *Angew. Chem., Int. Ed.*, 2013, **52**, 11726; (b) M. Corbet and F. D. Campo, *Angew. Chem., Int. Ed.*, 2013, **52**, 9896.
- 9 K. Li, G. Tan, J. Huang, F. Song and J. You, *Angew. Chem., Int. Ed.*, 2013, **52**, 12942.
- 10 Y. Aihara and N. Chatani, *J. Am. Chem. Soc.*, 2014, **136**, 898.
- 11 Although the detailed mechanism is not clear at this stage, a plausible catalytic cycle is proposed in ESI† (Scheme S1).

## Graphic Abstract

---

## Nickel-catalyzed chelation-assisted direct arylation of unactivated C(sp<sup>3</sup>)–H bonds with aryl halides

Mingliang Li, Jiaying Dong, Xiaolei Huang, Kaizhi Li, Qian Wu, Feijie Song and Jingsong You\*



The nickel-catalyzed chelation-assisted unactivated  $\beta\text{-C}(\text{sp}^3)\text{-H}$  bond arylation of aliphatic acid derivatives with aryl iodides/bromides has been achieved.