


ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE


Lewis Base Mediated Halogenation/Semipinacol Rearrangement of Diazo Compounds: A New Access to α -Halo-Quaternary Ketones†Haibin Mao,^a Zhongkai Tang,^a Hongwen Hu,^a Yixiang Cheng,^a Wen-Hua Zheng^a and Chengjian Zhu^{*a,b}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x



A novel halogenation/semipinacol rearrangement of α -diazo alcohol catalyzed by Lewis base has been developed through a carbene-free mechanism. This semipinacol transposition, initiated by an electrophilic halogenation ($X = \text{Cl}^+$, Br^+ , I^+) of diazo carbon event, furnished a convenient synthetic route for the efficient synthesis of α -halo-quaternary ketones under mild conditions.

As the metal carbene precursors, diazo compounds have been widely applied in organic synthesis which participate in metal-catalyzed rearrangements, cycloadditions, X-H ($X = \text{C}, \text{N}, \text{O}, \text{Si}, \text{S}$, etc.) bond insertions, and ylide-forming reactions (Scheme 1A).¹ On the other hand, diazo group transformations can also be realized through the carbene-free mechanism since diazo carbon is electron-rich in nature² from the resonance structure $\square$ (Scheme 1B). Diazo carbon can react with the electron-deficient species ($E^+ = \text{H}^+$,³ organoboron,⁴ carbonyl activated by Lewis acids,⁵ imine activated by Brønsted acids,⁶ etc.) to form a procarbenium ion intermediate $\square$ which is quenched by the electron-rich species subsequently with concomitant expulsion of N_2 (Scheme 1B). It is observed that the long-standing interest in the diazo chemistry was focused on the reactions based on the process of carbene, particularly in the transition-metal catalysis. While in consideration of environmental friendliness, convenient operation and diversity of diazo transformation, the development of efficient and elegant functionalization of diazo compounds processing carbene-free mechanism is really of great significance for further application of diazo compounds in organic synthesis.




Scheme 1 Activation patterns of the diazo carbon

A) Halogenation/Semipinacol Rearrangement of Allylic Alcohols:

B) Halogenation/Semipinacol Rearrangement of α -Diazo Alcohols:

Scheme 2 Semipinacol rearrangement initiated by an electrophilic halogenation

Very recently, chiral Lewis bases (LB) were used to catalyze the asymmetric halogenation of alkenes.^{7,9} Especially, Tu developed a novel asymmetric halogenation/semipinacol rearrangement⁸ reaction of allylic alcohols catalyzed by chiral Lewis bases for the synthesis of chiral α -quaternary β -haloketones (Scheme 2A).⁹ Furthermore, Murphy realized the *gem*-dihalogenation of diazoacetate derivatives employed Lewis bases as catalysts.¹⁰ Inspired by this previous work, and also in conjunction with our ongoing interest in the synthesis and applications of diazo compounds without metal catalyst,¹¹ we describe herein our results on Lewis base catalyzed halogenation/semipinacol rearrangement of diazo compounds (Scheme 2B). This strategy opens a new access to α -halo-quaternary ketones via the carbon-chain extension or ring expansion of ketones. These organohalo compounds are a very important class of synthetic intermediates that are useful for construction of α -quaternary ketones.¹² Moreover, we also make a preliminary effort to the catalytic, asymmetric version of this reaction, which led to moderate enantioselectivity.


Table 1 Optimization of the reaction conditions.^a


entry	cat	solvent	<i>t</i>	yield ^b (%)
1	none	THF	2 h	72
2	Py	THF	40 min	72
3	Et ₃ N	THF	12 h	71
4	DABCO	THF	20 min	78
5	DBU	THF	1 h	68
6	DABCO	Toluene	1 h	76
7	DABCO	DCM	10 min	72
8	DABCO	EA	10 min	68
9	DABCO	MeCN	1 min	67

^a Unless otherwise noted, the reaction was carried out with 0.20 mmol of **2a**, 0.25 mmol of **3a**, 10 mol % of catalyst (0.02 mmol). ^b Yield of isolated product.

We began by testing the halogenation/semipinacol rearrangement of α-diazo alcohols **2a**, which could be easily prepared from the commercially available benzophenone and ethyl diazoacetate.¹³ We observed that combination of **2a** and 1,3-dichloro-5,5-dimethylhydantoin (DCDMH, **3a**) in THF at room temperature afforded the corresponding product in 72% yield (Table 1, entry 1). To further improve reaction efficiency, we examined a number of Lewis bases (Table 1, entries 2-5). The addition of 10 mol% triethylamine prolonged reaction time obviously compared with the background reaction (Table 1, entry 3). Surprisingly, when 10 mol% of DABCO was used, the desired α-chloro-quaternary ketone **4a** was obtained in 78% isolated yield in 20 min (Table 1, entry 4). Next, a variety of common organic solvents were screened (Table 1, entries 4, 6-9). The experimental results revealed that increased solvent polarity led to the increase of reaction efficiency. However, this was associated with some by-product resulted in significantly reduced yields.


With the optimal reaction conditions in hand, we investigated the substrate scope of this halogenation/semipinacol rearrangement (Table 2). Various α-diazo alcohols **2** derived from symmetrical benzophenone derivatives with halogen or electron-donating group substituents were suitable substrates for the chlorination (Table 2, entries 1-5) or bromination reaction (Table 2, entries 11-14) leading to the desired products in excellent yields (up to 93%). Generally, chlorination reaction gave lower yields and lower reaction efficiency, while bromination reaction gave higher yields and higher reaction efficiency. Besides the carbon-chain extension, this method could effectively achieve the ring expansion of cyclic ketones (Table 2, entries 6, 15). Interestingly, the chlorination reaction was performed well over substrates derived from dissymmetrical alkyl aryl ketones, giving

Table 2 Substrate scope studies.^a


entry	substrate	X	product	yield ^b (%)
1	2a : R ¹ =R ² = Ph	3a :Cl	4a	78
2	2b : R ¹ =R ² = 4-F-C ₆ H ₄	Cl	4b	80
3	2c : R ¹ =R ² = 4-Cl-C ₆ H ₄	Cl	4c	71
4	2d : R ¹ =R ² = 4-Me-C ₆ H ₄	Cl	4d	81
5	2e : R ¹ =R ² = 4-OMe-C ₆ H ₄	Cl	4e	85
6	2f	Cl	4f	91
7	2g : R ¹ =Ph, R ² =Me	Cl	4g	76
8	2h : R ¹ =Ph, R ² = Et	Cl	4h	73
9	2i : R ¹ =Ph, R ² = ⁱ Pr	Cl	4i	78
10	2j	Cl	4j	61
11	2a : R ¹ =R ² = Ph	3b :Br	4k	92
12	2b : R ¹ =R ² = 4-F-C ₆ H ₄	Br	4l	93
13	2c : R ¹ =R ² = 4-Cl-C ₆ H ₄	Br	4m	82
14	2d : R ¹ =R ² = 4-Me-C ₆ H ₄	Br	4n	91
15	2f	Br	4o	95
16 ^c	2a : R ¹ =R ² = Ph	NIS	4p	78

^a The reaction was carried out with 0.20 mmol of **2**, 0.25 mmol of **3**, 10 mol % of DABCO (0.02 mmol) in THF (2.0 mL) at room temperature. ^b Yield of isolated product. ^c The reaction was carried out with 0.20 mmol of **2a**, 0.25 mmol of NIS in THF (0.2 mL) under argon atmosphere at room temperature for 20 hours.

the products chemoselectively in good yields. When employing chain alkyl aryl ketones as substrates, electron-rich phenyl ring migration had priority in the selective formal C–C insertion process (Table 2, entries 7-9). While when the diazo compound **2j** derived from α-tetralone was used, alkyl migration occupied a dominant (Table 2, entry 10) (see the Supporting Information for possible reasons). In addition, under the dry and oxygen-free argon atmosphere, α-iodo-quaternary ketone could be synthesized efficiently processing iodination/semipinacol rearrangement (Table 2, entry 16). It is noteworthy that DABCO would seriously reduce the efficiency of iodination reaction.



Scheme 3 Catalytic Asymmetric Variant of this Process

We also attempted to develop a catalytic asymmetric version of this halogenation/semipinacol rearrangement for access to enantioenriched α -halo-quaternary ketones. After a series of efforts (see the Supporting Information for further details), we discovered that product **4o** could be obtained in 86% yield with 72% ee when employing 20 mol % cinchona alkaloid $(\text{DHQD})_2\text{PHAL}$ as catalyst (Scheme 3).

In conclusion, we have developed a novel halogenation/semipinacol rearrangement of α -diazo alcohol catalyzed by Lewis base. This methodology greatly expands the synthetic applications of diazo compounds through the carbene-free mechanism. The protocol provides a straightforward way to generate the α -halo-quaternary ketones under mild conditions. We believe that the resulting densely functionalized ketones bearing α -halo quaternary carbon could be highly valuable and versatile intermediates for further transformations. To further realize asymmetric version of this reaction, we examined a number of chiral tertiary amines and varied reaction parameters, which led to the moderate enantioselectivity. Efforts to realize the highly enantioselective version of this reaction and further exploration of the asymmetric reaction scope are ongoing in our laboratory.

We gratefully acknowledge the National Natural Science Foundation of China (21372114, 21172106, 21074054), National Science Fund for Talent Training in Basic Science (no. J1103310), the National Basic Research Program of China (2010CB923303) and the Research Fund for the Doctoral Program of Higher Education of China (20120091110010) for their financial support.

Notes and references

^a School of Chemistry and Chemical Engineering, State Key Laboratory of Coordination Chemistry, Nanjing University, Nanjing, 210093 P. R. China. Fax: (+86)25-83594886, E-mail: cjzhu@nju.edu.cn

^b State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Shanghai, 200032, P. R. China

† Electronic Supplementary Information (ESI) available: complete experimental procedures, characterization data and copies of NMR spectra for all the new compounds. See DOI: 10.1039/b000000x/

Notes and references

- (a) M. Regitz and G. Maas, *Diazo Compounds: Properties and Synthesis*, Academic Press, London, 1986, 65-198; (b) A. Padwa and M. D. Weingarten, *Chem. Rev.* 1996, **96**, 223; (c) M. P. Doyle, M. A. McKerver and T. Ye, *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*, Wiley, New York, 1998; (d) H. M. L. Davies and R. E. J. Beckwith, *Chem. Rev.* 2003, **103**, 2861; (e) Z. Zhang and J. Wang, *Tetrahedron* 2008, **64**, 6577; (f) M. P. Doyle, R. Duffy, M. Ratnikov and L. Zhou, *Chem. Rev.* 2009, **110**, 704; (g) H. Lu and X. P. Zhang, *Chem. Soc. Rev.* 2011, **40**, 1899; (h) H. M. L. Davies and Y. Lian, *Acc. Chem. Res.* 2012, **45**, 923; (i) S.-F. Zhu and Q.-L. Zhou, *Acc. Chem. Res.* 2012, **45**, 1365.
- For reviews, see: (a) Y. Zhang and J. Wang, *Chem. Commun.* **2009**, 5350; (b) Y. Zhang and J. Wang, *Synlett* **2005**, 2886.

- (a) D. Hébrault, D. Uguen, A. De Cian and J. Fischer, *Tetrahedron Lett.* 1998, **39**, 6703; (b) T. Hashimoto, S. Isobe, C. K. A. Callens and K. Maruoka, *Tetrahedron* 2012, **68**, 7630.
- For selected examples, see: (a) C. Peng, W. Zhang, G. Yan and J. Wang, *Org. Lett.* 2009, **11**, 1667; (b) H. Li, L. Wang, Y. Zhang and J. Wang, *Angew. Chem., Int. Ed.* 2012, **51**, 2943; (c) H. Li, X. Shanguan, Z. Zhang, S. Huang, Y. Zhang and J. Wang, *Org. Lett.* 2013, **16**, 448. For review, see: (d) H. Li, Y. Zhang and J. Wang *Synthesis* 2013, **45**, 3090.
- (a) W.-J. Liu, B.-D. Lv and L.-Z. Gong, *Angew. Chem., Int. Ed.* 2009, **48**, 6503; (b) W. Li, J. Wang, X. Hu, K. Shen, W. Wang, Y. Chu, L. Lin, X. Liu and X. Feng, *J. Am. Chem. Soc.* 2010, **132**, 8532; (c) W. Li, X. Liu, X. Hao, Y. Cai, L. Lin and X. Feng, *Angew. Chem., Int. Ed.* 2012, **51**, 8644; (d) W. Li, X. Liu, F. Tan, X. Hao, J. Zheng, L. Lin and X. Feng, *Angew. Chem., Int. Ed.* 2013, **52**, 10883; (e) L. Gao, B. C. Kang, G.-S. Hwang and D. H. Ryu, *Angew. Chem., Int. Ed.* 2012, **51**, 8322; (f) L. Gao, B. C. Kang and D. H. Ryu, *J. Am. Chem. Soc.* 2013, **135**, 14556.
- (a) J. C. Antilla and W. D. Wulff, *J. Am. Chem. Soc.* 1999, **121**, 5099; (b) J. C. Antilla and W. D. Wulff, *Angew. Chem., Int. Ed.* 2000, **39**, 4518; (c) A. P. Patwardhan, V. R. Pulgam, Y. Zhang and W. D. Wulff, *Angew. Chem., Int. Ed.* 2005, **44**, 6169; (d) A. A. Desai and W. D. Wulff, *J. Am. Chem. Soc.* 2010, **132**, 13100; (e) M. J. Vetticatt, A. A. Desai and W. D. Wulff, *J. Am. Chem. Soc.* 2010, **132**, 13104; (f) G. Hu, A. K. Gupta, R. H. Huang, M. Mukherjee and W. D. Wulff, *J. Am. Chem. Soc.* 2010, **132**, 14669; (g) L. Huang and W. D. Wulff, *J. Am. Chem. Soc.* 2011, **133**, 8892; (h) T. Hashimoto, N. Uchiyama and K. Maruoka, *J. Am. Chem. Soc.* 2008, **130**, 14380; (i) T. Akiyama, T. Suzuki and K. Mori, *Org. Lett.* 2009, **11**, 2445.
- Asymmetric halocyclizations catalyzed by chiral Lewis bases. For selected review, see: (a) U. Hennecke, *Chem. Asian J.* 2012, **7**, 456. For selected examples, see: (b) T. Ishimaru, N. Shibata, T. Horikawa, N. Yasuda, S. Nakamura, T. Toru and M. Shiro, *Angew. Chem., Int. Ed.* 2008, **47**, 4157; (c) D. C. Whitehead, R. Yousefi, A. Jaganathan and B. Borhan, *J. Am. Chem. Soc.* 2010, **132**, 3298; (d) R. Yousefi, D. C. Whitehead, J. M. Mueller, R. J. Staples and B. Borhan, *Org. Lett.* 2011, **13**, 608; (e) A. Jaganathan, A. Garzan, D. C. Whitehead, R. J. Staples and B. Borhan, *Angew. Chem., Int. Ed.* 2011, **50**, 2593; (f) L. Zhou, C. K. Tan, X. Jiang, F. Chen and Y.-Y. Yeung, *J. Am. Chem. Soc.* 2010, **132**, 15474; (g) C. K. Tan, L. Zhou and Y.-Y. Yeung, *Org. Lett.* 2011, **13**, 2738; (h) L. Zhou, J. Chen, C. K. Tan and Y.-Y. Yeung, *J. Am. Chem. Soc.* 2011, **133**, 9164; (i) X. Jiang, C. K. Tan, L. Zhou and Y.-Y. Yeung, *Angew. Chem., Int. Ed.* 2012, **51**, 7771; (j) W. Zhang, S. Zheng, N. Liu, J. B. Werness, I. A. Guzei and W. Tang, *J. Am. Chem. Soc.* 2010, **132**, 3664; (k) G. E. Veitch and E. N. Jacobsen, *Angew. Chem., Int. Ed.* 2010, **49**, 7332.
- For selected review of semipinacol rearrangement, see: (a) B. Wang and Y. Q. Tu, *Acc. Chem. Res.* 2011, **44**, 1207; (b) Z.-L. Song, C.-A. Fan and Y.-Q. Tu, *Chem. Rev.* 2011, **111**, 7523.
- Asymmetric halogenation/semipinacol rearrangement catalyzed by chiral Lewis bases. For selected review, see: (a) S.-H. Wang, B.-S. Li and Y.-Q. Tu, *Chem. Commun.* 2014, **50**, 2393. For selected examples, see: (b) M. Wang, B. M. Wang, L. Shi, Y. Q. Tu, C.-A. Fan, S. H. Wang, X. D. Hu and S. Y. Zhang, *Chem. Commun.* **2005**, 5580; (c) H. Li, F.-M. Zhang, Y.-Q. Tu, Q.-W. Zhang, Z.-M. Chen, Z.-H. Chen and J. Li, *Chem. Sci.* 2011, **2**, 1839; (d) Z.-M. Chen, Q.-W. Zhang, Z.-H. Chen, H. Li, Y.-Q. Tu, F.-M. Zhang and J.-M. Tian, *J. Am. Chem. Soc.* 2011, **133**, 8818; (e) Z.-M. Chen, B.-M. Yang, Z.-H. Chen, Q.-W. Zhang, M. Wang and Y.-Q. Tu, *Chem. Eur. J.* 2012, **18**, 12950.
- J. Tao, R. Tran and G. K. Murphy, *J. Am. Chem. Soc.* 2013, **135**, 16312.
- (a) H. Mao, A. Lin, Y. Shi, Z. Mao, X. Zhu, W. Li, H. Hu, Y. Cheng and C. Zhu, *Angew. Chem., Int. Ed.* 2013, **52**, 6288; (b) H. Mao, A. Lin, Z. Tang, H. Hu, C. Zhu and Y. Cheng, *Chem. Eur. J.* 2014, **20**, 2454.
- K. Shibatomi, Y. Soga, A. Narayama, I. Fujisawa and S. Iwasa, *J. Am. Chem. Soc.* 2012, **134**, 9836.
- A. Gioiello, F. Venturoni, B. Natalini and R. Pellicciari, *J. Org. Chem.* 2009, **74**, 3520.