

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Isomerisation of 1,4-dichlorobenzene using highly acidic alkali chloroaluminate melts

Cite this: DOI: 10.1039/x0xx00000x

J. Messner^a, P. S. Schulz^a, N. Taccardi^a, Sven Kuhlmann,^b and P. Wasserscheid^{a*}

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The isomerisation reaction of 1,4-dichlorobenzene to the thermodynamically favoured and technically desired 1,3-dichlorobenzene has been studied comparing highly acidic chloroaluminate melts with organic imidazolium and alkali metal ions. Interestingly, the inorganic melts show much higher reactivity and full recyclability if small AlCl₃ losses are compensated and the reaction is carried out under slight HCl pressure.

Traditionally, Lewis acid catalysed reactions suffer from the need of hydrolysis for product isolation, causing the complete loss of the applied catalyst (e.g. AlCl₃). Hence, liquid-liquid biphasic reaction systems with acidic ionic liquids are of great interest for more sustainable acid catalysis. Over the recent years, it has been demonstrated in a large number of publications¹ that acidic ionic liquids allow a straightforward catalyst recycling by liquid phase separation, if none of the reactants form stable complexes with the acidic anions of the melt. Friedel-Crafts alkylations of aromatic compounds^{2, 3} or alkane isomerisation reactions,⁴ are among the prominent examples of this very successful strategy. Moreover, the physico-chemical properties of chloroaluminate ionic liquids, such as their melting points,⁵ viscosities⁶ and acidities⁷ are well understood. In contrast, acid catalysis with acidic alkali chloroaluminate melts is much less developed. This is mainly due to the marked difference in physical properties, such as e. g. melting points. While imidazolium chloroaluminates, such as 1-ethyl-3-methylimidazolium chloride ([EMIM]Cl)/AlCl₃ form liquid systems over a wide range of composition at room temperature,⁶ the respective LiCl/AlCl₃ displays, a melting point of 170 °C for $\chi(\text{AlCl}_3) = 0.67$.⁸ Despite these differences, anion speciation in both organic and inorganic chloroaluminates leads to quite a similar picture. Raman spectroscopy,^{9, 10} mass spectroscopy (MS)¹¹ and ²⁷Al NMR¹² have been applied to identify Cl⁻, [AlCl₄]⁻, [Al₂Cl₇]⁻ and [Al₃Cl₁₀]⁻ in organic and inorganic chloroaluminate melts. For systems with a molar fraction $\chi(\text{AlCl}_3) < 0.5$ only Cl⁻ and [AlCl₄]⁻ were found. Whereas [Al₂Cl₇]⁻ and [Al₃Cl₁₀]⁻ are the dominating species in melts with $\chi(\text{AlCl}_3) > 0.5$.^{13, 14} However, in contrast to organic chloroaluminates (where only [AlCl₄]⁻ is found for $\chi(\text{AlCl}_3) = 0.50$),¹⁴ inorganic chloroaluminate melts have been

shown to contain acidic [Al₂Cl₇]⁻ species in equimolar chloride salt/AlCl₃ mixtures.¹⁵ Simulation studies by Salanne *et al.*¹⁶ have confirmed this finding and claimed an increasing content of [Al₂Cl₇]⁻ in inorganic chloroaluminate melts of equimolar composition by going from the potassium over the sodium to the lithium system. IR measurements resulted in additional evidence for this important difference between organic and inorganic chloroaluminate melts.¹⁷ However, although spectroscopically confirmed, this property was so far poorly exploited for superior performance in technical relevant catalytic transformations. For both, organic and inorganic chloroaluminate systems, a further increase of the acidity is possible by adding hydrogen chloride gas (HCl). As such addition leads to the formation of super acids, the individual acidity of the melt system is a direct function of the chloride salt/AlCl₃ ratio and the partial pressure of HCl gas.^{18, 19}

In this contribution we describe the use of acidic chloroaluminate melts, both with organic and inorganic cations, in the isomerisation reaction of 1,4-dichlorobenzene (1,4-DCB). The technical interest in this reaction originates from the fact that the chlorination of benzene with gaseous chlorine and iron chloride as catalyst leads typically to a kinetic product mixture of DCBs that contains much less 1,3-DCB than industrially needed. Due to the ortho-/ para-directing effect of the primarily attached chloro substituent, the kinetic mixture contains mainly the 1,4- and 1,2-isomers and only small amounts of 1,3-DCB. However, 1,3-DCB is of high commercial interest as feedstock for herbicides, insecticides, dyes and pharmaceuticals synthesis. Direct and selective synthesis routes to 1,3-DCB exist and are based on chlorination of 1,3-dinitrobenzene or Sandmeyer reaction of 3-chloroaniline. Both routes are much more expensive and much less atom efficient than direct benzene chlorination. Thus, the isomerisation of a kinetic DCB mixture to the 1,3-DCB-rich thermodynamic mixture is of significant technical interest. The thermodynamic equilibrium composition of DCB isomers at 150 °C consists of 60 % 1,3-DCB, 32 % 1,4-DCB and 8 % 1,2-DCB.

The isomerisation of 1,4-DCB is not a new transformation. Olah *et al.*²⁰ investigated this reaction already in the 60's using water promoted aluminium chloride as catalyst. The isomerisation with AlCl₃/water is highly selective to form only the three DCB isomers together with small amounts of polychlorinated biphenyls. However, in the homogeneous Olah protocol, the AlCl₃ catalyst has to be

hydrolysed for product isolation, leading to complete loss of the catalyst.

In this contribution we report on alternative liquid-liquid biphasic reaction systems for the isomerisation of DCB that allows easy product isolation and complete recycling of the highly acidic catalyst phase. For this purpose, we applied organic and inorganic, acidic chloroaluminate melts. In detail, we investigated the isomerisation of 1,4-DCB comparing AlCl_3 -[EMIM]Cl and AlCl_3 -MCl (MCl= LiCl or LiCl-NaCl) as reaction phases. As the DCB isomerisation requires strongly acidic conditions, all applied melts contained a molar ratio of $\chi(\text{AlCl}_3) = 0.67$. This was the maximal molar fraction of AlCl_3 that allowed a reliable performance comparison between the different systems. In fact, the melting points of the AlCl_3 -LiCl system⁸ and the imidazolium system⁶ increase drastically above this composition. In order to achieve comparable starting points for the reaction, all melts were prepared and heated to 170 °C prior to adding the 1,4-DCB substrate. In case of the reference system, 1,4-DCB was placed in a preheated flask at 170 °C and a mixture of AlCl_3 and $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ as water donor was added directly to the organic component (for details of the experimental procedures see electronic supporting information –ESI). For the experiments shown in Figure 1, samples were taken every hour, dissolved in dichloromethane, extracted with water and analysed by GC.




Figure 1 Isomerisation of 1,4-dichlorobenzene with four different catalytic systems: (□) reference system 20 mol % AlCl_3 and 10 mol % H_2O in form of $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$; (△) 20 mol % AlCl_3 and 10 mol % LiCl; (▽) 20 mol % AlCl_3 , 4 mol % LiCl and 6 mol % NaCl; (◇) 20 mol % AlCl_3 and 10 mol % [EMIM]Cl, at 170 °C.

Most interestingly, there is a drastic difference in reactivity between the inorganic chloroaluminate melts and the imidazolium system. While the latter hardly shows any isomerisation activity under the applied conditions, the inorganic melts show significant activity and after 22 h reaction time exceed the level of conversion that was realized with the homogeneous Olah system²⁰ (25 % vs. 23 % conversion). Remarkably, the Olah system shows very high reactivity in the first hour (19 % conversion) while further reaction is very sluggish and only little extra conversion is observed in the following 23 hours. Comparing all chloroaluminate melts with the Olah reference system, it has to be kept in mind that all molten salt systems operate under liquid-liquid biphasic conditions while the AlCl_3 is homogeneous in nature. In addition, it has to be stated that the Olah system was not easy to handle at temperature of 170 °C due to AlCl_3 sublimation and huge amounts of HCl gas evolving from the system. These problems were effectively minimized in all molten salt systems. In further experiments, we addressed the optimisation of the reaction conditions for the 1,4-DCB isomerisation using AlCl_3 -LiCl. In particular, we were interested in two aspects: a) the

influence of HCl on the isomerisation reaction (formed from traces of humidity in the LiCl); b) the question whether recycling of the acidic melt is possible (saving large amounts of AlCl_3 and avoid waste production from the AlCl_3 hydrolysis).

In order to reveal the influence of dissolved HCl in the acidic molten salt, we compared the 1,4-DCB isomerisation reaction in an open flask with an otherwise identical reaction (4h reaction time, 170 °C) in a glass autoclave (see ESI for experimental details) where all the formed HCl gas was kept in the system (Figure 2).




Figure 2 Isomerisation of 1,4-DCB with AlCl_3 -LiCl in an open flask compared to the reaction in a closed autoclave; 170 °C, 4 h, 20 mol % AlCl_3 and 10 mol % LiCl.

Very clearly, the isomerisation of 1,4-DCB was observed to proceed much more effectively in the autoclave experiment, with 1,3-DCB yields exceeding 45 %. Crossing this conversion limit was found to be of practical relevance, as the DCB isomeric mixtures are liquids at room temperature with a molar fraction of the 1,3-isomer above 45 %. Thus, after the reaction the liquid organic layer can be easily decanted from the now solid molten salt at room temperature.

For recycling, the resulting solid molten salt was re-molten and fresh 1,4-DCB was added. Two special precautions proved to be necessary to allow recycling with constant yield in 1,3-DCB.




Figure 3 Isomerisation of 1,4-DCB with AlCl_3 - LiCl (BR) ($X_{\text{AlCl}_3} = 0.67$) and melt recycling (Re) under various conditions: A) substitution of leached AlCl_3 , no HCl-gas added, B) recycling with HCl-gas added, no addition of AlCl_3 C) recycling with added AlCl_3 and HCl-gas [170 °C, 4 h, 250 g (1.7 mol) 1,4-DCB, 20 mol % (45.4 g) AlCl_3 , 10 mol % (7.21 mol) LiCl, recycling with 1 bar HCl and / or the addition of 4.54 g AlCl_3].

First, the AlCl_3 content of the organic product phase was analysed by ICP to determine the level of Al-leaching. This level was found to be about 3 % of the previously applied AlCl_3 . Replacing this loss was found to be beneficial for the catalyst recycling. Secondly, it turned out that recycling is much more effective if after the product removal – a process that requires the opening of the reactor under inert gas – 1 bar of HCl-gas was added to the reaction system. Note that in an industrial scenario HCl is always omnipresent around an arene chlorination plant. With these two measures, the recycling of the acid catalyst phase worked indeed in an excellent manner with no loss of activity over at least five recycling runs (Figure 3, C). Thus, in comparison to the Olah system²¹, the here described acidic lithium chloroaluminate system has three distinct advantages for DCB isomerisation: a) drastic reduction of AlCl_3 sublimation; b) straight forward product isolation; and c) re-use of more than 95 mol-% of the applied AlCl_3 in subsequent isomerisation reactions.

In order to evaluate the scope of the new catalytic system for the isomerisation of other dihaloarene compounds, we also studied the isomerisation of 1-bromo-4-chlorobenzene (BCB) (see ESI for details). It was found that the reaction of the BCB substrate required less harsh reaction conditions compared to the fully chlorinated substrate as expected from literature reports.²¹




Figure 4 Isomerisation reaction of 1,4-BCB with AlCl_3 – LiCl vs. AlCl_3 -[EMIM]Cl (both X_{AlCl_3} = 0.67); 170 °C, 4 h, 20 mol % AlCl_3 .

Interestingly, the acidic $[\text{EMIM}]^+$ chloroaluminate melt exhibits reasonable catalytic activity towards the isomerisation of BCB still being much less reactive than its inorganic counterpart. While the AlCl_3 -LiCl system reached 40 % of the 1,3 isomer after less than 1 hour of reaction time, the same final yield took 22 h with the $[\text{EMIM}]^+$ melt of the same Al-content. A reasonable explanation for this marked difference in reactivity can be derived from the fact that the inorganic melt systems contain a higher concentration of acidic anions compared to the organic one.

In conclusion we could demonstrate that alkali chloroaluminate melts - and in particular lithium chloroaluminate melts – form highly active and recyclable catalyst phases for the technical relevant reaction of dihaloarene isomerisation.²² Remarkable differences in reactivity have been found between inorganic chloroaluminate systems and their imidazolium analogues at the same molar ratio of AlCl_3 . It has been further demonstrated that HCl partial pressure plays an important role for the catalyst recycling and thus obviously for the reactivity of the isomerisation system. Replenishing AlCl_3 losses and applying 1 bar HCl during reaction, resulted in a fully recyclable catalyst system for at least five dichlorobenzene isomerisation cycles.

Notes and references

The authors thank LANXESS Deutschland GmbH for financial support and for permission to publish this research.

^a Lehrstuhl für Chemische Reaktionstechnik der Universität Erlangen-Nürnberg (FAU), Egerlandstrasse 3, D-91058 Erlangen, Germany.

E-mail: peter.wasserscheid@fau.de; Fax: +49 9131 85 27421; Tel: +49 9131 85 27420

^b LANXESS Chemical (China) Co. Ltd.; 5 Corporate Avenue, No 150 Hubin Road, HuangPu District, Shanghai, 200021, P.R. of China.

Electronic Supplementary Information (ESI) available: [Experimental Procedures, ICP, GC, GC-MS data]. See DOI: 10.1039/c000000x/

1. M. Earle, in *Ionic Liquids in Synthesis*, eds. T. Welton and P. Wasserscheid, WILEY-VCH, 2008, Weinheim.
2. S. J. Nara, J. R. Harjani and M. M. Salunkhe, *J. Org. Chem.*, 2001, **66**, 8616-8620.
3. J. A. Boon, J. A. Levisky, J. L. Pflug and J. S. Wilkes, *J. Org. Chem.*, 1986, **51**, 480-483.
4. C. Meyer and P. Wasserscheid, *Chem. Commun.*, 2010, **46**, 7625-7627.
5. J. Estager, J. D. Holbrey and M. Swadzba-Kwasny, *Chem. Soc. Rev.*, 2014, **43**, 847-886.
6. A. A. Fannin, Jr., D. A. Floreani, L. A. King, J. S. Landers, B. J. Piersma, D. J. Stech, R. L. Vaughn, J. S. Wilkes and J. L. Williams, *J. Phys. Chem.*, 1984, **88**, 2614-2621.
7. J. Estager, A. A. Oliferenko, K. R. Seddon and M. Swadzba-Kwasny, *Dalton Trans.*, 2010, **39**, 11375-11382.
8. R. A. Carpio, A. A. Fannin, Jr., F. C. Kibler, Jr., L. A. King and H. A. Oye, *J. Chem. Eng. Data*, 1983, **28**, 34-36.
9. R. J. Gale, B. Gilbert and R. A. Osteryoung, *Inorg. Chem.*, 1978, **17**, 2728-2729.
10. S. J. Cyvin, P. Klaboe, E. Rytter and H. A. Øye, *J. Chem Phys*, 1970, **52**, 2776-&.
11. G. Franzen, B. P. Gilbert, G. Pelzer and E. DePauw, *Org. Mass Spectrom.*, 1986, **21**, 443-444.
12. F. Taulelle and A. I. Popov, *J. Solution Chem.*, 1986, **15**, 463-471.
13. E. Rytter, H. A. Øye, S. J. Cyvin, B. N. Cyvin and P. Klaboe, *J. Inorg. Nucl. Chem.*, 1973, **35**, 1185-1198.
14. H. A. Oye, M. Jagtoyen, T. Oksefjell and J. S. Wilkes, *Mater. Sci. Forum*, 1991, **73-75**, 183-189.
15. R. S. Juvet, Jr., V. R. Shaw and M. A. Khan, *J. Amer. Chem. Soc.*, 1969, **91**, 3788-3792.
16. M. Salanne, L. J. A. Siqueira, A. P. Seitsonen, P. A. Madden and B. Kirchner, *Condens. Matter*, 2013, 1-12,.
17. J. Hvistendahl, P. Klaboe, E. Rytter and H. A. Oeye, *Inorg. Chem.*, 1984, **23**, 706-715.
18. G. P. Smith, A. S. Dworkin, R. M. Pagni and S. P. Zingg, *J. Am. Chem. Soc.*, 1989, **111**, 5075-5077.
19. G. P. Smith, A. S. Dworkin, R. M. Pagni and S. P. Zingg, *J. Am. Chem. Soc.*, 1989, **111**, 525-530.
20. G. A. Olah, W. S. Tolgyesi and R. E. A. Dear, *J. Org. Chem.*, 1962, **27**, 3449-3455.
21. G. A. Olah, W. S. Tolgyesi and R. E. A. Dear, *J. Org. Chem.*, 1962, **27**, 3455-3464.
22. S. Kuhlmann, U. Böger, P. Wasserscheid, S. Schlenk, J. Messner and H.-M. Weber, WO 2013/189848 to Lanxess Deutschland GmbH [Chem. Abstr. 2013, 993608].