

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Oxidative Coupling of Alkenes with Amides Using Peroxides: Selective Amide C(sp³)-H versus C(sp²)-H Functionalization

Xu-Heng Yang,^a Wen-Ting Wei,^a Hai-Bing Li,^a Ren-Jie Song,^a and Jin-Heng Li^{*ab}

Received (in XXX, XXX) Xth XXXXXXXXXX 200X, Accepted Xth XXXXXXXXXX 200X

First published on the web Xth XXXXXXXXXX 200X


DOI: 10.1039/b000000x

A new oxidative coupling of unactivated terminal alkenes with amides using peroxides is described, in which mono- and difunctionalization of alkenes are selectively achieved. Within this reaction with amides, the chemoselectivity toward the functionalization of the C(sp³)-H bonds adjacent to the nitrogen atom or the functionalization of the carbonyl C(sp²)-H bonds across alkenes relies on the reaction conditions.

Alkenes are abundant organic molecules and simple chemical feedstock, and methods for their direct, selective functionalization are attractive as an important means to assemble more complex molecular entities. The transition-metal-catalyzed Heck couplings of alkenes with organic electrophiles have become one of the most efficient and common methods for the alkene functionalization due to their operational simpleness, the use of commercially available starting materials and the versatility of the resulting alkenes in synthesis.¹ Despite their widespread applications, the Heck coupling routes are typically protracted by the need to pre-functionalize the C-H bond toward the C-X bond (X = halides and pseudohalides). Therefore, an attractive alternative would involve a C-H functionalization that directly couples unactivated alkenes with broad diversification of functional groups and excellent selectivity control.

The oxidative coupling methodology involving the C-H functionalization has attracted much attention for replacing the conventional cross-coupling procedures, in part due to its step-economy by the avoidance of the pre-functionalization process.²⁻⁵ Despite remarkable advances in the field, approaches for the coupling of alkenes, particularly unactivated alkenes, with the carbonyl C(sp²)-H bonds are rare and limited.²⁻⁵ Generally, these oxidative transformations focus on the coupling with the aldehyde C(sp²)-H bonds (Scheme 1a), including 1) the coupling of electron-deficient alkenes with aldehydes (often alkyl aldehydes) initiated by air or *N*-hydroxyphthalimide (NHPI) combined with dibenzoyl peroxide (BPO) leading to saturated ketones,³ 2) the Heck-type coupling of unactivated alkenes with aryl aldehydes by using the [(Cp*RhCl₂)₂]/C₅H₂Ph₄/Cu(OAc)₂ or CuCl₂/*tert*-Butyl hydroperoxide (TBHP) system for synthesizing α,β -unsaturated ketones,⁴ and 3) the difunctionalization of arylalkenes with aldehydes and TBHP catalyzed by FeCl₂ accessing β -peroxy ketones.⁵ However, methods for the oxidative coupling of alkenes with the formyl C(sp²)-H bonds of formamides have not been reported.⁶ Herein, we report a new oxidative coupling of terminal alkenes with amides using

peroxides and a catalytic amount of bases (Scheme 1b); the chemoselectivity of this method can be controlled by changing the reaction conditions: While a catalytic amount of ^tBuOK combined with TBHP delivers β -amino ketones⁵ from alkenes and the C(sp³)-H bonds adjacent to the nitrogen atom of amides, the FeCl₃/DABCO/*di-tert*-butyl peroxide (DTBP) system assembles α,β -unsaturated amides by using alkenes to react the carbonyl C(sp²)-H bonds of formamides. Notably, the oxygen atoms in the new formed carbonyl groups are from hydroperoxides.




Scheme 1 Oxidative Coupling of Alkenes with Carbonyl Compounds.

We commenced our investigations by optimizing the reaction conditions for the oxidative coupling of 4-vinyltoluene (**1a**) with *N,N*-dimethylformamide (DMF; **2a**) (Table 1).⁷ Initially, alkene **1a** was treated with DMF **2a** and TBHP at 110 °C for 24 h, but no coupled products were observed (entry 1). Gratifyingly, we found that a catalytic amount of ^tBuOK could trigger the reaction, and 10 mol% ^tBuOK was preferred to form β -amino ketone **3aa** in 70% yield (entries 2-4). Other bases, including K₂CO₃, DBU and DABCO, could effect the reaction, but they were less efficient than ^tBuOK (entries 4-7). Screening of peroxides revealed that only hydroperoxides, TBHP, TAHP (*tert*-amyl hydroperoxide) and CHP (cumene hydroperoxide), could initiate the coupling (entries 2 and 8-10), and other peroxides without hydroxyl group, DTBP and BPO, have no activities at all (entries 11 and 12). However, no product **3aa** was observed without oxidants (entry 14). After varying the temperatures, the reaction at 110 °C provided the best results (entry 2 vs. entries 14-15). Notably, the reaction was successfully performed in MeCN, delivering **3aa** in 70% yield (entry 16).

Interestingly, addition of FeCl₃ could shift the chemoselectivity: β -amino ketone **3aa** and (*E*)-*N,N*-dimethyl-3-*p*-tolylacrylamide (**4aa**)⁵ were obtained in 5 and 8% yield, respectively (entry 17). Evaluation of various reaction

parameters revealed that 30 mol% FeCl₃, 30 mol% DABCO and 5 equiv DTBP at 110 °C for 48 h gave the best results: product **4aa** was obtained exclusively in 85% yield (entries 17-19 and 21 vs. entry 20). It should be noted that FeCl₂ can effect the reaction, albeit lowering the yield to 53% (entry 21).

Table 1 Screening Optimal Conditions^a




Entry	[O] (equiv)	Base (mol%)	Solvent	T (°C)	Isolated yield (%)	
					3aa	4aa
1	TBHP (2)	—	DMF	110	0	0
2	TBHP (2)	^t BuOK (5)	DMF	110	26	0
3	TBHP (2)	^t BuOK (10)	DMF	110	70	0
4	TBHP (2)	^t BuOK (20)	DMF	110	69	0
5	TBHP (2)	K ₂ CO ₃ (10)	DMF	110	65	0
6	TBHP (2)	DBU (10)	DMF	110	46	0
7	TBHP (2)	DABCO (10)	DMF	110	19	0
8 ^b	TBHP (2)	^t BuOK (10)	DMF	110	66	0
9	TAHP (2)	^t BuOK (10)	DMF	110	61	0
10	CHP (2)	^t BuOK (10)	DMF	110	28	0
11	DTBP (2)	^t BuOK (10)	DMF	110	trace	0
12	BPO (2)	^t BuOK (10)	DMF	110	trace	0
13	—	^t BuOK (10)	DMF	110	0	0
14	TBHP (2)	^t BuOK (10)	DMF	80	10	0
15	TBHP (2)	^t BuOK (10)	DMF	120	69	0
16 ^c	TBHP (2)	^t BuOK (10)	MeCN	110	70	0
17 ^d	TBHP (2)	^t BuOK (10)	DMF	110	8	5
18 ^d	TBHP (2)	DABCO (10)	DMF	110	trace	9
19 ^d	DTBP (2)	DABCO (10)	DMF	110	0	43
20 ^d	DTBP (5)	DABCO (30)	DMF	110	0	85
21 ^e	DTBP (5)	DABCO (30)	DMF	110	0	53

^a Reaction conditions: **1a** (0.4 mmol), **2a**, [O] (2 equiv), base and solvent (2 mL) for 24 h. TBHP (5 M in decane). ^b TBHP (70% in water). ^c DMF (4 mmol). ^d FeCl₃ (30 mol%) was added for 48 h. ^e FeCl₂ (30 mol%) was added for 48 h.

With the optimal reaction conditions in hand, we first sought to coupling of various terminal alkenes **1** with amides **2** to synthesize diverse β -amino ketones **3** (Table 2). In the presence of TBHP and ^tBuOK, β -amino ketone **3ba** was formed in 56% yield from styrene (**1b**) and DMF **2a**. Interestingly, a variety of substituted arylalkenes were also compatible with the optimal conditions, and substituents, including Cl, Br, MeO and CN, on the aromatic ring were well-tolerated (**3ca-ea** and **3ge**). Treatment of Cl- or Br-substituted alkenes **1c** and **1d** with DMF **2a**, TBHP and ^tBuOK successfully afforded **3ca** and **3da** in 62 and 61% yield, respectively. *ortho*-MeO-substituted alkene **1d** also delivered **3ea** in moderate yield. Unfortunately, oct-1-ene (**1f**), an alkylalkene, was not a suitable substrate (**3fa**). The optimal conditions were applicable to various amides **2b-2e** (**3ab-ae**), but failed to an amine **2f** (**3af**). Two cycloalkylamines, piperidine-1-carbaldehyde (**2b**) and morpholine-4-carbaldehyde (**2c**), could couple with alkene **1a**, TBHP and ^tBuOK, providing **3ab** and **3ac** in moderate yields. 1-

Methylpyrrolidin-2-one (**2d**) was also a viable substrate, generating **3ad** in 63% yield, in which alkene **1a** regioselectively attacked on the 5 position (the methylene group), not the 1-methyl group. Moreover, the reaction of *N,N*-dimethylacetamide (**2e**) with various arylalkenes **1a-c** or **1g**, TBHP and ^tBuOK was successfully performed, furnishing **3ae**, **3ce** and **3ge** in good yields.


Table 2 Coupling of Alkenes (**1**) with Amides (**2**), TBHP and ^tBuOK^a



^a Reaction conditions: **1** (0.4 mmol), **2** (4 mmol), ^tBuOK (10 mol%), TBHP (2 equiv; 5 M in decane) and MeCN (2 mL) at 110 °C for 24 h.

As shown in Table 3, we turned our attention to assemble α,β -unsaturated amides **4** from alkenes **1** and formamides **2** in the presence of FeCl₃, DTBP and DABCO. Arylalkenes **1b-c** and **1h-i** even with Br, NO₂ and Cl groups on the aryl ring successfully underwent the coupling with DMF **2a**, giving **4ba-ca** and **4ha-ia** in 53-77% yield, and the order of their reactivity is as follow: electron-rich > electron-deficient. However, alkylalkene **1f** had a relatively lower reactivity (**4fa**). Gratifyingly, 1,1-diphenylethylene (**1j**) and (*E*)-phenylbuta-1,3-diene (**1k**) delivered **4ja** and **4ka** smoothly in 55% and 50% yields, respectively. It is important to note that this coupling generally proceeds in moderate to good yields with an extensive range of other formamides, including *N,N*-dialkylformamides **1b-c** and **1g**, *N*-methyl-*N*-phenylformamide (**1h**), *N,N*-diphenylformamide (**1i**) and *N*-phenylformamide (**1j**) (**4bb-bc** and **4bg-bj**).


The results of entries 3 and 9-13 in Table 1 showed that only hydroperoxides could trigger the reaction, suggesting that the oxygen atoms in the new formed carbonyl group of products **3** may be from hydroperoxides. To verify these, a control experiment between alkene **1a** and DMF **2a** using 4 equiv H₂¹⁸O was performed: product **3aa** did not include the ¹⁸O atom, ruling out the oxygen atom from H₂O. Moreover, the presence of radical inhibitors, TEMPO, hydroquinone and BHT, resulted in no detectable **3aa**, implying that the current reaction includes a radical process.

Table 3 Fe-Catalyzed Coupling of Alkenes (**1**) with Formamides (**2**)^a

^a Reaction conditions: **1** (0.4 mmol), **2** (2 mL), FeCl₃ (30 mol%), DBACO (30 mol%) and DTBP (5 equiv) at 110 °C for 48 h.

Possible mechanisms outlined in Scheme 2 were proposed for the oxidative coupling.^{2-5,8} Initially, alkyl radical **A** is formed by abstracting the α -H-atom alpha to the N-atom of DMF **1a** with ^tBuOO[•], which is generated from TBHP under heating with the aid of bases.^{2,8} Addition of alkyl radical **A** to alkene **1** affords the other alkyl radical **B**. The reaction of alkyl radical **B** and ^tBuOO[•] occurs to afford intermediate **C**. Finally, the O-O bond of intermediate **C** is cleaved by ^tBuOK, followed by oxidation with TBHP produces **3** and ^tBuOH.

In the presence of Fe²⁺ species, DTBP is split into Fe³⁺(O^tBu) and ^tBuO[•] radical.²⁻⁵ Hydrogen-abstraction of DMF **2a** by ^tBuO[•] radical delivers carbonyl radical **D**, and then addition across alkene **1** affords alkyl radical **E**. Single-electron-transfer between alkyl radical **E** and Fe³⁺(O^tBu) gives alkyl cation **F**, Fe²⁺ species and ^tBuO⁻. Finally, β -H elimination of alkyl cation **F** with ^tBuO⁻ by using DABCO base gives **4** and ^tBuOH.

**Scheme 2** Possible Mechanisms.

The presence of FeCl₃ lowered the reactivity of alkenes **1** for reacting with the C(sp³)-H bonds of amines is because [•]OH radical is consumed to form the Fe^{III}(OH) species, compressing the generation of ^tBuOO[•] radical.

In summary, we have developed the first oxidative coupling of unactivated terminal alkenes with amides using peroxides for selective synthesis of β -amino ketones and α,β -

unsaturated amides. This method proceeds via a tandem C-H/alkene functionalization step that occurs through an oxidative radical pathway with a broad substrate scope and excellent selectivity control.

This research was supported by the NSFC (No. 21172060) and Hunan Provincial Natural Science Foundation of China (No. 13JJ2018).

Notes and references

- ^a State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Chemistry and Chemical Engineering, Hunan University, Changsha 410082, China. Fax: 0086731 8871 3642; Tel: 0086731 8882 2286; E-mail: jhli@hnu.edu.cn
- ^b State Key Laboratory of Applied Organic Chemistry Lanzhou University, Lanzhou 730000, China.
- † Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/
- ‡ Footnotes should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.
- (a) R. F. Heck and J. P. Nolley, *J. Org. Chem.*, 1972, **37**, 2320; (b) T. Mizoroki, K. Mori and A. Ozaki, *Bull. Chem. Soc. Jpn.*, 1971, **44**, 581; for selected reviews: (c) A. de Meijere and F. E. Meyer, Jr. *Angew. Chem. Int. Ed. Engl.*, 1994, **33**, 2379; (d) M. Oestreich, *The Mizoroki-Heck reaction*, Wiley, Hoboken, NJ, 2008; (e) I. P. Beletskaya and A. V. Cheprakov, *Chem. Rev.*, 2000, **100**, 3009; (f) D. Mc Cartney and P. J. Guiry, *Chem. Soc. Rev.*, 2011, **40**, 5122.
 - (a) C.-J. Li, *Acc. Chem. Res.*, 2009, **42**, 335; (b) G. P. Mcglacken, L. M. Bateman and L. Bateman, *Chem. Soc. Rev.*, 2009, **38**, 2447; (c) D. Shabashov, *Acc. Chem. Res.*, 2009, **42**, 1074; (d) J. A. Ashenurst, *Chem. Soc. Rev.*, 2010, **39**, 540; (e) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, *Chem. Soc. Rev.*, 2011, **40**, 5068; (f) J. Le Bras and J. Muzart, *Chem. Rev.*, 2011, **111**, 1170; (g) C. Liu, H. Zhang, W. Shi and A. Lei, *Chem. Rev.*, 2011, **111**, 1780; (h) Y.-X. Xie, R.-J. Song, J.-N. Xiang and J.-H. Li, *Chin. J. Org. Chem.*, 2012, **32**, 1555; (i) C. Zhang, C. Tang and N. Jiao, *Chem. Soc. Rev.*, 2012, **41**, 3464; (j) F. Guo, M. D. Clift and R. J. Thomson, *Eur. J. Org. Chem.*, 2012, 4881; (k) M. Grzybowski, K. Skonieczny, H. Butenschön and D. T. Gryko, *Angew. Chem. Int. Ed.*, 2013, **52**, 9900; (l) S. A. Girard, T. Knauber and C.-J. Li, *Angew. Chem. Int. Ed.*, 2014, **53**, 74.
 - (a) V. Chudasama, R. J. Fitzmaurice and S. Caddick, *Nat. Chem.*, 2010, **2**, 592; (b) S. Tsujimoto, T. Iwahama, S. Sakaguchi and Y. Ishii, *Chem. Commun.*, 2001, 2352; (c) S. Tsujimoto, S. Sakaguchi and Y. Ishii, *Tetrahedron Lett.*, 2003, **44**, 5601.
 - (a) Z. Shi, N. Schröder and F. Glorius, *Angew. Chem. Int. Ed.*, 2012, **51**, 8092. (b) J. Wang, C. Liu, J. Yuan and A. Lei, *Angew. Chem. Int. Ed.*, 2013, **52**, 2256.
 - (a) W. Liu, Y. Li, K. Liu and Z. Li, *J. Am. Chem. Soc.*, 2011, **133**, 10756; (b) L. Lv, B. Shen and Z. Li, *Angew. Chem. Int. Ed.*, 2014, **53**, 4164.
 - Although there are few papers on the transition metal-catalyzed reactions between alkenes and formamides, only the corresponding hydroamidation products were obtained. Moreover, these transition-metal-catalyzed methods are restricted to expensive noble metal catalysts (Ru, Ni or Pd) and harsh reaction conditions (high reaction temperatures and/or labile additives): (a) Y. Tsuji, S. Yoshii, T. Ohsumi, T. Kondo and Y. Watanabe, *J. Organomet. Chem.*, 1987, **331**, 379; (b) T. Kondo, T. Okada and T. Mitsudo, *Organometallics*, 1999, **18**, 4123; (c) S. Ko, H. Han and S. Chang, *Org. Lett.*, 2003, **5**, 2687; (d) D. C. D. Nath, C. M. Fellows, T. Kobayashi and T. Hayashi, *Aust. J. Chem.*, 2006, **59**, 218; (e) Y. Nakao, H. Idei, K. S. Kanyiva and T. Hiyama, *J. Am. Chem. Soc.*, 2009, **131**, 5070; (f) T. Fujihara, Y. Katafuchi, T. Iwai, J. Terao and Y. Tsuji, *J. Am. Chem. Soc.*, 2010, **132**, 2094.
 - The detail data of screening optimal conditions are summarized in Table S1 of Supplementary Information.
 - (a) A. Studer, *Chem. Eur. J.*, 2001, **7**, 1159; (b) A. Studer, *Chem. Soc. Rev.*, 2004, **33**, 267; (c) A. Studer and T. Schulte, *Chem. Rec.*, 2005, **5**, 27; (d) X.-L. Lian, H. Lei, X.-J. Quan, Z.-H. Ren, Y.-Y. Wang and Z.-H. Guan, *Chem. Commun.*, 2013, **49**, 8196; (e) M.-N. Zhang, R.-R. Hui, Z.-H. Ren, Y.-Y. Wang and Z.-H. Guan, *Org. Lett.*, 2014, **16**, 3082.