

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Total Syntheses of (\pm)-Musellarins A-C

Cite this: DOI: 10.1039/x0xx00000x

Zhilong Li, Tsz-Fai Leung and Rongbiao Tong*

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The first, diastereoselective total syntheses of musellarins A-C were achieved concisely with 7.8%-9.8% yields in 15-16 steps. The key synthetic features include (i) Achmatowicz rearrangement, Kishi reduction, and Friedel-Crafts cyclization to construct the tricyclic framework and (ii) Heck coupling of aryldiazonium salt to introduce the aryl on the dihydropyran in a 2,6-*trans* fashion at the final stage of synthesis.

Diarylheptanoid natural products are a family of secondary plant metabolites isolated from various sources, featuring the structural motif of a seven-carbon linkage of two aromatic rings.¹ Many diarylheptanoids displayed potent biological activities such as antioxidant, anticancer, antibacterial, antifungal, antiosteoporosis, antihepatotoxicity, and melanogenesis inhibition, and therefore have been increasingly recognized as the potential therapeutic agents.¹ Among over 400 known diarylheptanoid natural products, musellarins A-E (Figure 1) represent an unusual structural class





Figure 1. Musellarins and other types of diarylheptanoids

for the presence of the rare bicyclic tetrahydropyran motif. Musellarin A was reported in 2002 by Kinghorn² and co-workers from hybrid plant fruits of *musa x paradisiaca* in Peru as the first example of diarylheptanoids containing such bicyclic functionality, which had been established by extensive NMR studies and confirmed by single crystal X-ray diffraction analysis. In 2011, musellarin A was isolated again by Zhao³ and coworkers from monotypic plant *Musella lasiocarpa* in Yunan, China, in 0.0006% yield (18 mg from 3 kg of dry plant

materials). Zhao *et al.* also obtained several new congeners of musellarin A: musellarins B-E containing this rare bicyclic skeleton. Although the biological activities of musellarins C-E have not been reported, musellarin A significantly induced quinone reductase activity against Hepalcl7 cells,² and musellarin B has shown moderate cytotoxicity against several cancer cell lines including HL-60 (IC₅₀ 21.3 μ M), SMMC-7721 (IC₅₀ 26.7 μ M) and A-549 (IC₅₀ 25.1 μ M).³ The potential bioactivity and unusual structures of musellarins within the family of diarylheptanoids prompted us to undertake their synthetic studies. Herein, we report the first total syntheses of musellarins A-C featuring a late stage stereoselective installation of the second aryl via Heck coupling of aryl diazonium salts.


Retrosynthetically (Scheme 1), we planned to introduce the aryl group with different substituents via Heck coupling of enol ether **2** and aryl diazonium salts,⁴ which was expected to simultaneously deliver the desired stereochemistry at C2 and the appropriate double bond at C3-C4. It was noted that the



Scheme 1. Retrosynthetic Analysis of Musellarins A-C

similar Heck couplings yielding the 2,6-*trans* disubstituted tetrahydropyran with excellent diastereoselectivity has been elegantly exploited by Schmidt in the total synthesis of centrolobine and various diarylheptanoid analogues (Type IV, Figure 1).⁵ The enol ether **2** could be prepared from ketone **3** via Pd-catalyzed reduction of its corresponding enol triflate. The ketone **3** was envisioned to be derived from intramolecular Friedel-Crafts cyclization⁶ of γ -aryl enone **4**, which could be readily synthesized from furfuryl alcohol **6** through Achmatowicz rearrangement⁷ and Kishi reduction.⁸ Wittig olefination of **7** and **8** would deliver the requisite alcohol **6** after hydrogenation.

Our synthesis (Scheme 2) began with preparation of alcohol **10**, which was obtained in excellent yield from commercially available isovanillin **9** by protection of the alcohol as triisopropylsilyl ether and reduction of the resulting aldehyde with NaBH₄. The alcohol **10** was then converted into the phosphonium salt **7** in two straightforward steps. Wittig olefination of aldehyde **8** and phosphorus ylide generated *in situ* from **7** produced a 2:3 *E/Z* mixture of alkene **11** in 71%



Scheme 2. Synthesis of the Tricyclic Compound **3**

yield. The subsequent hydrogenation of **11** with Pd/C in methanol (or ethyl acetate), however, resulted in a rather complex mixture, partially due to over-reduction. Gratifyingly, we found that magnesium⁹ (turnings) in methanol at room temperature could effectively saturate the double bond with concomitant deacetylation¹⁰ to provide the furfuryl alcohol **6** in 68% yield. Achmatowicz rearrangement of **6** was smoothly promoted by *m*-CPBA at 0 °C to provide the dihydropyranone hemiacetal **5** in 85% yield. Kishi reduction of **5** using a classical combination of trifluoroacetic acid and triethylsilane generated a 2:1 mixture of dihydropyranone **4** and a surprising Friedel-Crafts cyclization adduct **3** in 78% combined yield.




Figure 2. ORTEP Diagram of Compound **3**

As we planned in our design stage, the intramolecular Friedel-Crafts cyclization of dihydropyranone **4** occurred in the presence of BF₃-Et₂O at -78 °C to provide quantitatively the tricyclic compound **3**, which structure was unambiguously confirmed by X-ray diffraction analysis (Figure 2).

Encouraged by the unexpected result that Kishi reduction/Friedel-Crafts cyclization could occur in a one-pot manner, we attempted to improve the yield of **3** in the one-pot process (Table 1). Under classical Kishi reduction conditions

Table 1. Attempted Conditions for One-pot Kishi Reduction and Intramolecular Friedel-Crafts Cyclization^a


entry	acid (equiv.)	R ₃ SiH (equiv)	Temp (°C) /Time (h)	ratio ^b (4:3)	yield ^c (%)
1	TFA (10)	Et ₃ SiH	-40/1	2:1	78
2	TFA (10)	<i>i</i> -Pr ₃ SiH	-40/1	1:1	41
3	TFA (10)	Et ₃ SiH	-40/12	complex	N/A
4	BF ₃ -Et ₂ O (2)	Et ₃ SiH	-78/1	14:1	74
5	BF ₃ -Et ₂ O (5)	Et ₃ SiH	-78/5	complex	N/A
6	BF ₃ -Et ₂ O (0.5)	Et ₃ SiH	-78→rt/24	5:1	45
7	TMSOTf (1)	Et ₃ SiH	-78/1	complex	N/A
8	TESOTf (0.3)	Et ₃ SiH	-78/3	complex	N/A
9	TBSOTf (0.3)	Et ₃ SiH	-78/3	complex	N/A
10	TiCl ₄ (1.2)	Et ₃ SiH	-78/1	0:1	30
11	SnCl ₄ (1)	Et ₃ SiH	-78/1	complex	N/A

Notes: a: the reaction was run with 20 mg of compound **5**. b: ratio was determined by NMR analysis of the crude reaction mixture. c: combined yield after flash column chromatography on silica gel. TFA: trifluoroacetic acid. TMSOTf: trimethylsilyl trifluoromethanesulfonate. TESOTf: triethylsilyl trifluoromethanesulfonate. TBSOTf: *tert*-butyldimethylsilyl trifluoromethanesulfonate.

reported previously (entries 1 and 4), TFA (or BF₃-Et₂O) and triethylsilane gave the dihydropyranone **4** as the major product in a good yield if the reaction was quenched within 1 hr. Extending the reaction time (entry 3) or excess equivalent of BF₃-Et₂O (entry 5) did not lead to any further conversion of **4** to **3**. Instead, significant decomposition was observed. When the bulkier triisopropylsilane (entry 2) was used as the hydride donor, intramolecular Friedel-Crafts cyclization produced a 1:1 mixture of **4** and **3** in a low combined yield. Since other conditions employing triisopropylsilane and phenyldimethylsilane did not give a better yield of **3**, triethylsilane would be used for further attempts. Reducing the equivalent of BF₃-Et₂O (entry 6) slowed considerably the reduction and cyclization, which required elevated temperature and longer reaction time for complete consumption of **5** but gave a lower yield as compared to entry 4. Other strong Lewis acids (entries 7, 8, 9, and 11) resulted in decomposition with trace amount of desired compounds **3** or **4**. Interestingly, TiCl₄-Et₃SiH gave exclusively the tricyclic compound **3** in 30% yield. Unfortunately, the overall yield was synthetically too low and could not be improved under various conditions. It appeared to us that two-step operations would be the choice: Kishi reduction with TFA/Et₃SiH or BF₃-Et₂O/Et₃SiH (entries 1 and 4) and independent Friedel-Crafts cyclization with BF₃-Et₂O as shown in Scheme 2.

Next, we directed our attention to synthesize the tricyclic enol ether **2** via a two-step deoxygenation¹¹ of ketone **3** (Scheme 3). To this end, ketone **3** needed to be regioselectively deprotonated with a base to generate an enol anion, which would be triflated with PhNTf₂. However, treatment of ketone **3** with PhNTf₂ and a bulky base such as KHMDS or LDA at -78


°C led to exclusive formation of vinyl triflate **12a** in good yield.¹² When the more steric demanding base LiTMP was used, a regiomer mixture of **12a** and **12b** could be obtained



Scheme 3. Synthesis of the Key Intermediate Enol Ether **2**

in lower yield (40-50%) with a 1:1 ratio of diastereomers, which could not be easily separated by column chromatography. In view of difficulty in efficient preparation of the enol triflate **12b** by this straightforward method, we set out to explore the possibility of olefin isomerization¹³ for the synthesis of enol ether **2** from **12a**. First, vinyl triflate **12a** was transformed into alkene **13** in excellent yield by palladium-catalyzed reduction.¹¹ To our delight, the isomerization of **13** with Wilkinson catalyst¹⁴ in the presence of DBU proceeded smoothly at reflux to provide enol ether **2** in 63% yield, which served as the common key intermediate for the synthesis of musellarins A-C via Heck coupling with aryl diazonium salts.

Three aryl diazonium salts (**15a-c**), corresponding to musellarins A-C, were prepared (Scheme 4) by using similar protocols in the literature.⁵ Specifically, acetaminophen (**14**), a




Scheme 4. Synthesis of the Aryl Diazonium Salts

widely used over-the-counter analgesic, could be successfully transformed into the aryl diazonium salt **15a**^{5e} with 58% yield in two steps. The preparation of aryl diazonium salts **15b** and **15c** could be achieved in excellent yields from **16** and **18**, respectively, via three-step sequence: acetylation, palladium-catalyzed reduction of nitro group to amine, and diazotization. It was noteworthy that acetyl protection of the phenol was essential to the success since 4-amino-phenols derived from direct reduction of nitrophenols (**16** and **18**) were unstable and could not be converted to the corresponding diazonium salts.

With both coupling partners in hand, we then explored the Heck coupling using the conditions developed by Schmidt⁵

(Scheme 5). Gratifyingly, Heck reaction of the enol ether **2** with aryl diazonium salts **15a-c** using Pd(OAc)₂ as a precatalyst



Scheme 5. Completion of Total Syntheses of Musellarins A-C

occurred smoothly to provide the corresponding coupling products, which immediately were subjected to desilylation with TBAF and/or deacetylation with K₂CO₃/MeOH to furnish musellarins A-C, respectively, in excellent yields with good diastereoselectivity (dr 7:1~9:1). Separation of the musellarins A-C from their minor diastereomers by preparative TLC (silica gel) led to analytically pure musellarins A-C. All spectroscopic data of our synthetic samples were identical to those reported for natural musellarins A-C.¹⁵

In summary, we have accomplished the first, concise total syntheses of musellarins A-C in a longest linear sequence of 15-16 steps from commercially available materials. The synthetic strategy was enabled by (i) exploitation of Achmatowicz rearrangement, Kishi reduction and Friedel-Crafts alkylation to construct the tricyclic framework **3** with a *cis* ring conjunction and (ii) high-yielding Heck coupling of aryl diazonium salts in good *trans*-diastereoselectivity at the final stage of synthesis. Importantly, the flexibility and convergence of the synthetic route developed here would permit an expedient entry to musellarins and their analogues for further biological studies.

This research was financially supported by HKUST and Research Grant Council of Hong Kong (ECS 605912, GRF 605113, and GRF 16305314). The authors are also grateful to Prof. Dr. Ian Williams and Dr. Herman Sung (HKUST) for the single crystal X-ray diffraction analysis.

Notes and references

Department of Chemistry, The Hong Kong University of Science and Technology, Clearwater Bay, Kowloon, Hong Kong, China.

Email: rtong@ust.hk

† Electronic Supplementary Information (ESI) available: Detailed experimental procedures, characterizations and copies of ¹H- and ¹³C-NMR spectra of new compounds, and X-ray crystallographic data (CCDC 1012651). See DOI: 10.1039/c000000x/

- For recent reviews, see: (a) H. Lv and G. She, *Rec. Nat. Prod.*, 2012, **6**, 321; (b) J. Zhu, G. Islas-Gonzalez and M. Bois-Choussy, *Org. Prep. Proc. Int.*, 2000, **32**, 505; (c) G. M. Keserü and M. Nógrádi, *Stud. Nat. Prod. Chem.*, 1995, **17**, 357.
- D. S. Jang, E. J. Park, M. E. Hawthorne, J. S. Vigo, J. G. Graham, F. Cabieses, B. D. Santarsiero, A. D. Mesecar, H. H. S. Fong, R. G. Mehta, J. M. Pezzuto and A. D. Kinghorn, *J. Agric. Food Chem.*, 2002, **50**, 6330.
- L.-B. Dong, J. He, X.-Y. Li, X.-D. Wu, X. Deng, G. Xu, L.-Y. Peng, Y. Zhao, Y. Li, X. Gong and Q.-S. Zhao, *Nat. Prod. Bioprospect.*, 2011, **1**, 41.
- For reviews on Heck coupling of aryl diazonium salts, see: (a) A. Roglans, A. Pla-Quintana and M. Moreno-Mañas, *Chem. Rev.*, 2006, **106**, 4622; (b) J. G. Taylor, A. V. Moro and C. R. D. Correia, *Eur. J. Org. Chem.*, 2011, 1403; (c) F.-X. Felpin, L. Nassar-Hardy, F. Le Callonnec and E. Fouquet, *Tetrahedron*, 2011, **67**, 2815. For selected examples employing Heck coupling of aryl diazonium salts, see: (d) K. Kikukawa and T. Matsuda, *Chem. Lett.*, 1977, 159; (e) E. A. Severino, E. R. Costenaro, A. L. L. Garcia and C. R. D. Correia, *Org. Lett.*, 2003, **5**, 305; (f) A. L. L. Garcia, M. J. S. Carpes, A. C. B. M. de Oca, M. A. G. dos Santos, C. C. Santana and C. R. D. Correia, *J. Org. Chem.*, 2005, **70**, 1050; (g) J. C. Pastre and C. R. D. Correia, *Org. Lett.*, 2006, **8**, 1657; (h) F.-X. Felpin, E. Fouquet and C. Zakri, *Adv. Synth. Catal.*, 2008, **350**, 2559; (i) F.-X. Felpin, O. Iburguren, L. Nassar-Hardy and E. Fouquet, *J. Org. Chem.*, 2009, **74**, 1349; (j) F.-X. Felpin, O. Iburguren, L. Nassar-Hardy and E. Fouquet, *J. Org. Chem.*, 2009, **74**, 1349; (k) F.-X. Felpin, K. Miqueu, J.-M. Sotiropoulos, E. Fouquet, O. Iburguren and J. Laudien, *Chem. Eur. J.*, 2010, **16**, 5191; (l) J. Laudien, E. Fouquet, C. Zakri and F.-X. Felpin, *Synlett*, 2010, 1539.
- (a) B. Schmidt and F. Höfter, *Chem. Eur. J.*, 2009, **15**, 11948; (b) B. Schmidt, R. Berger and F. Höfter, *Org. Biomol. Chem.*, 2010, **8**, 1406; (c) B. Schmidt, *Chem. Commun.*, 2003, 1656; (d) B. Schmidt, F. Höfter, A. Kelling and U. Schilde, *J. Org. Chem.*, 2011, **76**, 3357; (e) B. Schmidt, F. Höfter, R. Berger and S. Jessel, *Adv. Synth. Catal.*, 2010, **352**, 2463.
- For selected recent examples, see: (a) M. A. Grundl and D. Trauner, *Org. Lett.*, 2006, **8**, 23; (b) M. A. Grundl, A. Kaster, E. D. Beaulieu and D. Trauner, *Org. Lett.*, 2006, **8**, 5429; (c) S. Tang, Y. Xu, J. He, Y. He, J. Zheng, X. Pan and X. She, *Org. Lett.*, 2008, **10**, 1855; (d) T. Lomberget, E. Bentz, D. Bouyssi and G. Balme, *Org. Lett.*, 2003, **5**, 2055; (e) R. Hayashi and G. R. Cook, *Org. Lett.*, 2007, **9**, 1311; (f) E. Fillion and D. Fishlock, *J. Am. Chem. Soc.*, 2005, **127**, 13144.
- (a) O., Jr. Achmatowicz, P. Bukowski, B. Szechner, Z. Zwierzchowska and A. Zamojski, *Tetrahedron*, 1971, **27**, 1973; (b) L. Zhu, L. Song and R. Tong, *Org. Lett.*, 2012, **14**, 5892; (c) J. Ren, Y. Liu, L. Song and R. Tong, *Org. Lett.*, 2014, **16**, 2986. For leading reviews on Achmatowicz rearrangement, see: (d) B. H. Lipshutz, *Chem. Rev.*, 1986, **86**, 795; (e) J. M. Harris, M. Li, J. G. Scott and G. A. O'Doherty, Achmatowicz Approach to 5,6-Dihydro-2H-pyran-2-one Containing Natural Products. In *Strategy and Tactics in Organic Synthesis*, Harmata, M., Ed.; Elsevier, London, 2004, Vol. **5**, pp 221-253. For seminal contributions from O'Doherty group, see: (f) Q. Chen, Y. Zhong and G. A. O'Doherty, *Chem. Commun.*, 2013, **49**, 6806, and references cited therein; (g) R. S. Babu and G. A. O'Doherty, *J. Am. Chem. Soc.*, 2003, **125**, 12406; (h) R. S. Babu, M. Zhou and G. A. O'Doherty, *J. Am. Chem. Soc.*, 2004, **126**, 3428; (i) H. Guo and G. A. O'Doherty, *Org. Lett.*, 2006, **8**, 1609.
- (a) K. C. Nicolaou, T. M. Baker and T. Nakamura, *J. Am. Chem. Soc.*, 2011, **133**, 220; (b) K. L. Jackson, J. A. Henderson, H. Motoyoshi and A. J. Phillips, *Angew. Chem., Int. Ed.*, 2009, **48**, 2346; (c) J. A. Henderson, K. L. Jackson and A. J. Phillips, *Org. Lett.*, 2007, **9**, 5299; (d) J. M. Um, K. N. Houk and A. J. Phillips, *Org. Lett.*, 2008, **10**, 3769; (e) M. D. Lewis, J. K. Cha and Y. Kishi, *J. Am. Chem. Soc.*, 1982, **104**, 4976; (f) For the initial use of TFA in Kishi reductions, see: G. A. Kraus, M. T. Molina and J. A. Walling, *J. Chem. Soc., Chem. Commun.*, 1986, **21**, 1568.
- Magnesium in methanol has been widely used in reduction of α,β -unsaturated esters or nitriles, see: (a) J. A. Profitt, D. S. Watt and E. J. Corey, *J. Org. Chem.*, 1975, **40**, 127; (b) I. K. Youn, G. H. Yon and C. S. Pak, *Tetrahedron Lett.*, 1986, **27**, 2409; (c) W.-D. Z. Li and K. Wei, *Org. Lett.*, 2004, **6**, 1333; (d) J. Cabares and L. Mavoungou-Gomes, *Bull. Soc. Chim. Fr.*, 1986, 401. For a review, see: (e) G. H. Lee, I. K. Youn, E. B. Choi, H. K. Lee, G. H. Yon, H. C. Yang and C. S. Pak, *Curr. Org. Chem.*, 2004, **8**, 1263.
- Y.-C. Xu, E. Lebeau and C. Walker, *Tetrahedron Lett.*, 1994, **35**, 6207.
- (a) S. Cacchi, E. Morera and G. Ortar, *Tetrahedron Lett.*, 1984, **25**, 4821; (b) S. Cacchi, P. G. Ciattini, E. Morera and G. Ortar, *Tetrahedron Lett.*, 1986, **27**, 5541; (c) S. Cacchi, E. Morera and G. Ortar, *Org. Synth.*, 1990, **68**, 138; (d) S. Cacchi, E. Morera and G. Ortar, *Org. Synth.*, 2011, **88**, 260.
- For triflation of dihydrofuranone, enol triflate was obtained, see: (a) T. O. Painter, J. R. Bunn, F. J. Schoenen, J. T. Douglas, V. W. Day and C. Santini, *J. Org. Chem.*, 2013, **78**, 3720; (b) F. Tang and K. D. Moeller, *Tetrahedron*, 2009, **65**, 10863.
- For selected examples involving transition metal-catalyzed isomerization of alkenes to enol ethers, see: (a) E. J. Corey and J. W. Suggs, *J. Org. Chem.*, 1973, **38**, 3224; (b) M. A. Leeuwenburgh, H. S. Overkleeft, G. A. van der Marel and J. H. van Boom, *Synlett*, 1997, 1263; (c) H. Oguri, S. Tanaka, T. Oishi and M. Hirama, *Tetrahedron Lett.*, 2000, **41**, 975; (d) P. A. V. van Hooft, F. E. Oualid, H. S. Overkleeft, G. A. van der Marel, J. H. van Boom and M. A. Leeuwenburgh, *Org. Biomol. Chem.*, 2004, **2**, 1395; (e) B. Schmidt, *J. Org. Chem.*, 2004, **69**, 7672.
- R. Young and G. Wilkinson, *Inorg. Synth.*, 1977, **17**, 75.
- See Electronic Supplementary Information†.