

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Visible Light-Promoted Metal-Free sp^3 -C–H Fluorination

Cite this: DOI: 10.1039/x0xx00000x

Ji-Bao Xia, Chen Zhu and Chuo Chen*

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Photoexcited acetophenone can catalyze the fluorination of unactivated $C(sp^3)$ -H groups. While acetophenone, a colorless oil, only has a trace amount of absorption in the visible light region, its photoexcitation can be achieved by irradiation with light generated by a household compact fluorescent lamp (CFL). This operational simple method provides improved substrate scope for the direct incorporation of a fluorine atom into simple organic molecules. CFL-irradiation can also be used to promote certain classic UV-promoted photoreactions of colorless monoarylketones and enones/enals.

Direct C–H functionalization is a powerful new method for small-molecule synthesis.^{1,2} Recently, we have reported that visible light can activate diarylketones to catalyze $C(sp^3)$ -H fluorination at benzylic positions.^{3a} 9-Fluorenone catalyzes benzylic C–H monofluorination while xanthone catalyzes benzylic C–H difluorination (Fig. 1a). We now show that the violet light (375–400 nm) generated by a household compact fluorescent lamp (CFL) can activate acetophenone, a monoarylketone, to catalyze the fluorination of unactivated $C(sp^3)$ -H groups (Fig. 1b). Acetophenone is a colorless oil that has only a trace amount of absorption above 375 nm ($n \rightarrow \pi^*$ transition $\lambda_{max} \sim 325$ nm). Herein, we demonstrate that CFL-irradiation (>375 nm) can effectively promote its photoexcitation. We further show that certain classic UV-promoted photoreactions of monoarylketones and enones/enals can also be induced by CFL-irradiation.

Several catalytic C–H fluorination reactions have been developed to introduce fluorine atoms directly without prior functionalization.^{4–13} For example, the Sanford,⁴ Yu,⁵ and Doyle⁶ groups have each reported an efficient palladium catalyst system while the Groves⁷ and the Lectka⁸ groups have disclosed a manganese and a copper catalyst system, respectively. The Inoue,⁹ Tang,¹⁰ and Hartwig¹¹ groups have also shown that *N*-oxyl radical and silver salts can catalyze or promote C–H fluorination. More recently, photolytic activation methods have also been reported by Britton,¹² Lectka,^{8c} and Tan¹³ using decatungstate, 1,2,4,5-tetracyanobenzene (TCB), and anthraquinone (AQN) as the catalyst, respectively.




Fig. 1 Photolytic C–H fluorination reactions catalyzed by photoexcited arylketones. The reactivity and selectivity of the catalysts can be tuned by the ketone substituent groups.

We have been interested in developing vanadium and ketone catalysts for C–H fluorination reactions.³ Previously, we showed that CFL-irradiation¹⁴ could activate 9-fluorenone, a yellow solid, to catalyze the benzylic fluorination of ethylbenzene (**1**) with Selectfluor to give **3** (Table 1, entries 1 and 2).^{3a} Intriguingly, we recently found that the photolytic fluorination of 4-ethylacetophenone (**2**) did not require 9-fluorenone (entries 3 and 4). We suspected that CFL-irradiation led to the photoexcitation of **2**. The resulting diradical-like **2*** mediated this C–H fluorination reaction by abstracting a benzylic hydrogen atom from another molecule of **2**. Subsequent fluorine atom transfer from Selectfluor gave **4**.

4-Ethylacetophenone (**2**) is a monoarylketone that has no apparent absorption above 375 nm ($n \rightarrow \pi^*$ transition $\lambda_{max} \sim 325$ nm). Its photoexcitation has been traditionally carried out by irradiation with <360 nm UV light. Indeed, UV-irradiation improved the

quantum yield significantly but also promoted considerable decomposition (entries 5 and 6). In contrast, irradiation with violet light (375–465 nm, λ_{max} 419 nm) resulted in a clean transformation (entry 7), suggestive of productive absorption at >375 nm.

Table 1 Effects of the light source and the ketone catalyst on the benzylic fluorination of **1** and **2**

Entry	Substrate	Catalyst	Light source	Time	Yield ^[j]
1	1	9-fluorenone	1 × CFL ^[a]	24 h	89% (85%) ^[j]
2	1	–	1 × CFL ^[a]	24 h	0%
3	2	9-fluorenone	1 × CFL ^[a]	24 h	80% (76%) ^[j]
4	2	–	1 × CFL ^[a]	30 h	89%
5	2	–	16 × RPR-3000A ^[b]	30 min	46%
6	2	–	16 × RPR-3500A ^[c]	1 h	73%
7	2	–	16 × RPR-4190A ^[d]	6 h	96%
8	1	acetophenone	16 × RPR-4190A ^[d]	6 h	88%
9	1	acetophenone	16 × RPR-4190A ^[d,e]	20 h	70%
10	1	acetophenone	16 × RPR-4190A ^[d,f]	20 h	0%
11	1	acetophenone	1 × violet LED ^[g]	3 h	85%
12	1	acetophenone	1 × CFL ^[a]	20 h	61%
13	1	acetophenone	1 × CFL ^[a,e]	20 h	50%
14	1	acetophenone	1 × CFL ^[a,i]	20 h	0%
15	1	acetone	1 × CFL ^[a]	20 h	0%
16	1	acetophenone	– ^[h]	24 h	0%

^[a] 19 W, household lamp. ^[b] 21 W, 250–375 nm. ^[c] 24 W, 300–420 nm. ^[d] 375–465 nm. ^[e] With a 375 nm longpass filter. ^[f] With a 400 nm longpass filter. ^[g] 9 W, 370–405 nm. ^[h] 50 °C. ^[i] Determined by ¹⁹F NMR using C₆H₅F as an external standard. ^[j] Isolated yield.

Because acetophenone and 4-ethylacetophenone (**2**) have very similar UV absorption spectra, we hypothesized that violet light could also activate acetophenone to catalyze the benzylic fluorination of ethylbenzene (**1**). Indeed, a clean fluorination occurred to give **3** in 88% yield when irradiating a solution of **1** with violet light (375–465 nm) in the presence of 5 mol % acetophenone and 2 equiv Selectfluor (entry 8).

Similar to **2**, acetophenone is a colorless oil that has no apparent absorption above 375 nm. To exclude the possibility of UV-contamination,¹⁵ we repeated the fluorination of **1** catalyzed by acetophenone (reagent grade, >98%; analytical grade, ≥99.5%; or freshly redistilled from calcium hydride) in the presence of a 375 nm longpass filter. The reaction proceeded slower but smoothly (entry 9). Consistently, using a 370–405 nm LED flashlight as the light source led to the formation of **3** in 85% yield in only 3 h (entry 11). No reaction occurred in the presence of a 400 nm longpass filter (entry 10). CFL-irradiation also promoted the acetophenone-catalyzed benzylic fluorination of **1** in the presence of a 375 nm filter but not a 400 nm filter (entries 12–14). No reaction occurred when using acetone ($n \rightarrow \pi^*$ transition $\lambda_{\text{max}} \sim 280$ nm) as the catalyst (entry 15) or heating the solution at 50 °C in dark (entry 16). Therefore, we believe that short violet light (375–400 nm)¹⁶ could effectively promote the photoexcitation of acetophenone under the regular reaction conditions.

We envisioned that, with a larger n/π^* energy gap, acetophenone would be more reactive than diarylketones toward unactivated C–H

groups. To test this hypothesis, we examined the fluorination of cyclohexane (**5**) catalyzed by a series of different ketones (Table 2). Indeed, acetophenone was found to be the most effective catalyst (entry 1). Introducing a methoxy group to acetophenone led to a slightly reduced catalyst activity (entry 2) while adding a nitro group resulted in a low yield of **6** (entry 3). Trifluoroacetophenone and benzaldehyde are also less effective catalysts (entries 4 and 5). Consistent with our previous observations, benzaldehyde reacted with Selectfluor to give benzoyl fluoride under the reaction conditions.^{3a,17} Benzophenones and 9-fluorenone were less effective despite significantly better absorption (entry 6–9). Xanthone was nearly as good as acetophenone when a large excess of **5** was used (entry 10). However, acetophenone showed better catalyst activity when the amount of **5** was reduced to 5 equiv (entries 11 and 12). The substrate to Selectfluor ratio could be further reduced to ~1.5:1 for acetophenone (entries 13 and 14).

Table 2 Effects of the catalyst on the fluorination of **5**^[a]

Entry	5 (equiv)	ketone catalyst	Yield ^[b]
1	23	R = CH ₃ , X = H	90%
2	23	R = CH ₃ , X = OMe	82%
3	23	R = CH ₃ , X = NO ₂	18%
4	23	R = CF ₃ , X = H	63%
5	23	R = H, X = H	59%
6	23	X = H	78%
7	23	X = OMe	76%
8	23	X = NMe ₂	17%
9	23	X = –	75%
10	23	X = O	87%
11	5	X = O	81%
12	5		88%
13	1.5		76%
14	1.0		59%

^[a] Reaction conditions: 0.01 mmol catalyst, 0.2 mmol Selectfluor, 2 mL CH₃CN. ^[b] Determined by ¹⁹F NMR using C₆H₅F as an external standard.

The scope of this acetophenone-catalyzed photolytic C–H fluorination reaction is shown in Fig. 2. The fluorination of cycloalkanes proceeded smoothly. Monofluorides **7–10** were obtained in good yields with only a small (~5%) amount of difluorination products as judged by the ¹⁹F NMR spectra of the crude reaction mixtures. The site selectivity of this reaction follows the general trend of innate C–H oxidation reactions (3° > 2° > 1°).¹⁸ For example, the fluorination of norbornane provided **11** with good regio- and stereoselectivity (*exo/endo*=15/1) because of hyperconjugation effects.^{18a} The fluorination of propionic acid, butyric acid, and isovaleric acid gave **12**, **13**, and **14** in <5%, 55%, and 81% yield, respectively. The β - and γ -positions of cycloheptanone were fluorinated at comparable rates (β : γ =1:1). While attempts to isolate β -fluorocycloheptanone (**15- β**) resulted in elimination of the fluoride, γ -fluorocycloheptanone (**15- γ**) could be isolated in 35% yield. Fluorination of *N*-phthaloyl-L-valine methyl ester gave **16** in 85% yield. Monoterpenes 1,4-cineole and L-menthone reacted selectively at the tertiary positions to give **17** and **18** in 73% and 71% yield, respectively. We have also re-examined

the fluorination of *N*-phthaloyl-L-valine and L-menthone under UV-irradiation (250–375 nm) conditions. The reactions proceeded significantly faster but the products degraded under the reaction conditions, leading to low yields (<40%) of **16** and **18**.




Fig. 2 Scope of the acetophenone-catalyzed photolytic C–H fluorination reaction. ^[a] *Endo*-2-fluoronorborene was also formed in 4% NMR yield. ^[b] Isolated as the corresponding benzyl ester. ^[c] Substrate (1.0 equiv), Selectfluor (1.5 equiv). ^[d] Acetophenone (20 mol %).




Fig. 3 Site selectivity of the acetophenone-catalyzed photolytic C–H fluorination reaction. Reaction conditions: CFL-irradiation, **19** (5.0 equiv), **20** and **21** (1.5 equiv), acetophenone (5 mol %), Selectfluor (1.0 equiv).

We have used hexane (**19**), 2-hexanone (**20**), and sclareolide (**21**) to further study the site selectivity of this acetophenone-catalyzed photolytic C–H fluorination reaction (Fig. 3). The fluorination of **19** occurred primarily on the methylene groups (C1:C2:C3=1:17:7). Introducing a carbonyl group deactivated the C3 methylene group, rendering the C4 and C5 methylenes the only detectable reaction sites of **20** (C4:C5=1:1.3). Similar to other innate radical C–H oxidation reactions,^{7a,18} sesquiterpenoid **21** reacted predominately at the C2 position (C2/C3 = 3.6/1).

Mechanistically, it is rather intriguing that CFL-irradiation could promote the photoexcitation of acetophenone that is a colorless oil. Recently, the Melchiorre and Bach groups have demonstrated that Lewis base and Lewis acid can react with carbonyl groups to induce a bathochromic shift.¹⁹ However, we did not observe significant bathochromic shift when mixing acetophenone, benzophenone, or 9-fluorenone with Selectfluor and **1** in acetonitrile (Fig. 4 and ESI), indicating that the photoexcitation of acetophenone was not facilitated by Selectfluor through a fluoronium transfer. In contrast, we detected a trace amount of absorption between 375 and 400 nm

with acetophenone at high concentrations. We thus believe that this weak absorption is responsible for driving this C–H fluorination reaction.²⁰




Fig. 4. The *R* band (*n*→ π^* transition) of acetophenone extends to ~400 nm at high concentrations.




Fig. 5. CFL-irradiation promoted Norrish type II cleavage and Norrish–Yang cyclization of **22**.

We believe that, similar to our previous fluorenone-catalyzed fluorination reactions,^{3a} acetophenone functioned as a C–H abstraction catalyst instead of an electron- or energy-transfer catalyst.^{8c,13} To test this hypothesis and to further exclude the possibility that Selectfluor reacted with acetophenone to form a transient reactive species not detectable by UV, we examined if CFL-irradiation could also promote unimolecular C–H abstraction of monoarylketones. Indeed, CFL-irradiation induced the Norrish type II cleavage and Norrish–Yang cyclization of valerophenone (**22**) (*n*→ π^* transition λ_{max} ~325 nm) to give **23**–**26** (Fig. 5). Addition of trifluoroacetic acid did not accelerate these reactions or cause a bathochromic shift of the *R* band of **22**, indicating that the photoactivation of **22** was not promoted by adventitious acid protonating its carbonyl group. Furthermore, results of the light-dark cycle experiments suggest that acetophenone-catalyzed fluorination of cyclooctane is not a free radical chain reaction (ESI).²¹ The unimolecular photoreaction of **22** further suggest that acetophenone can function as a C–H abstraction catalyst instead of a photosensitizer in activating Selectfluor toward C–H abstraction.^{8d,13}

Further support for the hypothesis that CFL-irradiation can promote the photoexcitation of simple monoarylketones follows the observation that CFL-irradiation could also promote photoreactions of colorless enones/enals (*n*→ π^* transition λ_{max} ~320 nm). For example, photolysis of cyclopentenone (**27**) in 2-propanol (**28**) gave **29** through an etheral C–H abstraction by **27*** (Fig. 6). Results of the light-dark cycle experiments also suggest this C–H abstraction/conjugate addition reaction is not a free radical chain reaction (ESI).²¹ Additionally, we found that [2+2] cycloaddition of

(*Z*)-enal **30** could also be induced by CFL-irradiation to give **31** together with the *E/Z* isomerization product **32** that also cyclized to **31** very slowly.




Fig. 6. CFL-irradiation promoted photoreactions of **27** and **30**.

In conclusion, we have shown that, despite low quantum yields, short violet light (375–400 nm) generated by a low-energy household CFL can promote photoreactions of monoarylketones and enones/enals that have an *R* band $\lambda_{\text{max}} \sim 320$ nm. By avoiding the harmful high-energy UV light, these photoreactions can be performed without using specialized photochemical equipment and give fewer side-reactions. Using this mild photolysis method, photoexcited acetophenone can be readily generated to catalyze the fluorination of unactivated C(sp³)–H groups. This new fluorination reaction is operationally simple and utilizes cheap, readily available catalyst. Further investigation of the utility of this photolytic fluorination method is underway.

Acknowledgements

Financial Support was provided by NIH (NIGMS R01-GM079554), the Welch Foundation (I-1596) and UT Southwestern. We thank Mr. Wenhan Zhang of Prof. Joseph Ready's lab (UT Southwestern) for repeating and confirming the reactions in Figures 4 and 5.

Notes and references

Department of Biochemistry, UT Southwestern Medical Center, Dallas, Texas, 75390, USA. E-mail: Chuo.Chen@UTSouthwestern.edu; Fax: +1 214 6480320; Tel: +1 213 6485048

- (a) Y. Ishihara and P. S. Baran, *Synlett*, 2010, 1733; (b) J. L. Roizen, M. E. Harvey and J. Du Bois, *Acc. Chem. Res.*, 2011, **45**, 911; (c) K. M. Engle, T.-S. Mei, M. Wasa and J.-Q. Yu, *Acc. Chem. Res.*, 2012, **45**, 788; (d) S. R. Neufeldt and M. S. Sanford, *Acc. Chem. Res.*, 2011, **45**, 936.
- (a) J. Lu, X. Tan and C. Chen, *J. Am. Chem. Soc.*, 2007, **129**, 7768; (b) J.-B. Xia, K. W. Cormier and C. Chen, *Chem. Sci.*, 2012, **3**, 2240; (c) C. Zhu, J.-B. Xia and C. Chen, *Tetrahedron Lett.*, 2014, **55**, 232.
- (a) J.-B. Xia, C. Zhu and C. Chen, *J. Am. Chem. Soc.*, 2013, **135**, 17494; (b) J.-B. Xia, Y. Ma and C. Chen, *Org. Chem. Front.*, 2014, **1**, 468.
- (a) K. L. Hull, W. Q. Anani and M. S. Sanford, *J. Am. Chem. Soc.*, 2006, **128**, 7134; (b) K. B. McMurtrey, J. M. Racowski and M. S. Sanford, *Org. Lett.*, 2012, **14**, 4094; (c) J. M. Racowski, J. B. Gary and M. S. Sanford, *Angew. Chem. Int. Ed.*, 2012, **51**, 3414.
- (a) X. Wang, T.-S. Mei and J.-Q. Yu, *J. Am. Chem. Soc.*, 2009, **131**, 7520; (b) K. S. L. Chan, M. Wasa, X. Wang and J.-Q. Yu, *Angew. Chem. Int. Ed.*, 2011, **50**, 9081.
- M.-G. Braun and A. G. Doyle, *J. Am. Chem. Soc.*, 2013, **135**, 12990.
- (a) W. Liu, X. Huang, M.-J. Cheng, R. J. Nielsen, W. A. Goddard III and J. T. Groves, *Science*, 2012, **337**, 1322; (b) W. Liu and J. T. Groves, *Angew. Chem. Int. Ed.*, 2013, **52**, 6024; (c) X. Huang, W. Liu, H. Ren, R.

- Neelamegam, J. M. Hooker and J. T. Groves, *J. Am. Chem. Soc.*, 2014, **136**, 6842.
- (a) S. Bloom, C. R. Pitts, D. C. Miller, N. Haselton, M. G. Holl, E. Urheim and T. Lectka, *Angew. Chem. Int. Ed.*, 2012, **51**, 10580; (b) S. Bloom, C. R. Pitts, R. Woltornist, A. Griswold, M. G. Holl and T. Lectka, *Org. Lett.*, 2013, **15**, 1722; (c) S. Bloom, J. L. Knippela and T. Lectka, *Chem. Sci.*, 2014, **5**, 1175. (d) C. R. Pitts, S. Bloom, R. Woltornist, D. J. Auvenshine, L. R. Ryzhkov, M. A. Siegler and T. Lectka, *J. Am. Chem. Soc.*, 2014, **136**, 9780. (e) S. Bloom, J. L. Knippel, M. G. Holl, R. Barber and T. Lectka, *Tetrahedron Lett.*, 2014, **55**, 4576.
- Y. Amaoka, M. Nagatomo and M. Inoue, *Org. Lett.*, 2013, **15**, 2160.
- P. Xu, S. Guo, L. Wang and P. Tang, *Angew. Chem. Int. Ed.*, 2014, **53**, 5955.
- P. S. Fier and J. F. Hartwig, *Science*, 2013, **342**, 956.
- S. D. Halperin, H. Fan, S. Chang, R. E. Martin and R. Britton, *Angew. Chem. Int. Ed.*, 2014, **53**, 4690.
- C. W. Kee, K. F. Chin, M. W. Wong and C.-H. Tan, *Chem. Commun.*, 2014, **50**, 8211.
- For examples of CFL/LED-promoted photoreactions, see: (a) T. P. Yoon, *ACS Catal.*, 2013, **3**, 895; (b) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322; (c) J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102; (d) J. Xun and W.-J. Xiao, *Angew. Chem. Int. Ed.*, 2012, **51**, 6828.
- (a) M. Khazova and J. B. O'Hagan, *Radiat. Prot. Dosimetry*, 2008, **131**, 521; (b) T. Mironava, M. Hadjiargyrou, M. Simon and M. H. Rafailovich, *Photochem. Photobiol.*, 2012, **88**, 1497.
- Light in the range of 375–400 nm has been referred to as both UV and visible light.
- R. E. Banks, N. J. Lawrence and A. L. Popplewell, *Synlett*, 1994, 831.
- (a) T. Newhouse and P. S. Baran, *Angew. Chem. Int. Ed.*, 2011, **50**, 3362; (b) M. S. Chen and M. C. White, *Science*, 2010, **327**, 566.
- (a) E. Arceo, I. D. Jurberg, A. Álvarez-Fernández and P. Melchiorre, *Nat. Chem.*, 2013, **5**, 750; (b) R. Brimioulle and T. Bach, *Science*, 2013, **342**, 840.
- (a) D. L. Walker and B. Fraser-Reid, *J. Am. Chem. Soc.*, 1975, **97**, 6251; (b) Z. Benko and B. Fraser-Reid, *J. Org. Chem.*, 1988, **53**, 2066; (c) M. Fagnoni, D. Dondi, D. Ravelli and A. Albini, *Chem. Rev.*, 2007, **107**, 2725.
- These experiments do not exclude that possibility that a radical chain reaction with non-negligible rates of chain termination is operating.