

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Cytotoxic and Cytostatic Gold(III) Corrole

Cite this: DOI: 10.1039/x0xx00000x


Darius R. Teo,^{a,b} Harry B. Gray,^{a,*} Punrajit Lim,^b John Termini,^{b,*} Elena Domeshek,^c and Zeev Gross^{c,*}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

We have synthesized and characterized a water-soluble gold(III) corrole (**1-Au**) that is highly toxic to cisplatin-resistant cancer cells. Relative to its 1-Ga analogue, axial ligands bind only weakly to **1-Au**, which likely accounts for its lower affinity for human serum albumin (HSA). We suggest that the cytotoxicity of **1-Au** may be related to this lower HSA affinity.

The seminal report on the scalable synthesis of the first corrole that is stable in its free-base form, 5,10,15-tris(pentafluorophenyl)corrole,¹ was followed one year later by the introduction of a positively charged derivative as an anticancer agent.² More recent health-related research has focused on the iron(III), manganese(III), aluminium(III), and gallium(III) complexes of 2,17-bis-sulfonato-5,10,15-tris(pentafluorophenyl)corrole (Fig. 1: **1-Fe**, **1-Mn**, **1-Al**, and **1-Ga**).^{3,4,5}



Scheme 1 Synthesis of **1-Au** and previously reported metal complexes of **1-H₃**.

^a Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, United States

^b Department of Molecular Medicine, Beckman Research Institute of the City of Hope, 1500 E. Duarte Road, Duarte, California 91010, United States

^c Schulich Faculty of Chemistry, Technion—Israel Institute of Technology, Haifa 32000, Israel

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/c000000x/

These well characterized amphipolar complexes form tight conjugates with a variety of proteins that have been used as trafficking and cell penetrating vehicles in physiologically relevant media.⁶ Closed-shell metallocorroles have been employed for optical imaging, whereas certain others exhibit pro-oxidant, anti-oxidant, and cell-killing properties, as follows: **1-Mn** catalyzes the peroxidase-like enantioselective oxidation of sulfides by hydrogen peroxide,⁷ **1-Fe** is the best synthetic catalyst reported to date for the catalase-like decomposition of hydrogen peroxide to water and molecular oxygen,⁸ **1-Al** is a versatile optical imaging agent,⁹ and **1-Ga** is cytotoxic to breast cancer cells.¹⁰ What distinguishes corroles from most other ligands is that none of its metal-ion derivatives is demetallated under physiologically relevant conditions.¹¹

During the last decade, several gold(I) and gold(III) compounds were reported to manifest impressive antitumor properties, making up a promising class of experimental anticancer agents.¹² Among these agents, a cytotoxic gold(III) porphyrin complex, [Au(TPP)]Cl, was found to be particularly active, as it blocked the self-renewal ability of cancer stem-like cells.¹³ Most notably, [Au(TPP)]Cl was neither demetallated nor reduced during circulation in biological media. This key finding stimulated our interest in gold(III) corroles, which are redox-stable, low-spin square planar complexes, similar to gold(III) porphyrins and platinum(II) anticancer drugs such as cisplatin.¹⁴

Here we report a water-soluble gold(III) complex of 2,17-bis-sulfonato-5,10,15-tris(pentafluorophenyl)corrole (**1-Au**): we have studied its association with human serum albumin (HSA), the most abundant (and potential drug transporting) protein present in plasma.¹⁵ In the course of our investigations, we found **1-Au** to be cytotoxic and cytostatic to cisplatin-resistant cancer cell lines.

Two methods for the preparation of gold(III) corroles have been reported: [ClAu(tht)] (tht = tetrahydrothiophene) was used

for insertion into a β -brominated corrole in 85% yield,¹⁶ whereas Au(OAc)₃ was utilized for the synthesis of several β -unsubstituted gold(III) triarylcorroles in 25-35% yield.¹⁷ The above reagents, as well as chloroauric acid and its salts were tested in an attempted synthesis of **1-Au**, but only gold(III) acetate led to the desired complex. The treatment of 2,17-bis-sulfonato-5,10,15-tris(pentafluorophenyl)corrole (**1-H₃**) with excess Au(OAc)₃ in pyridine at room temperature allowed isolation of **1-Au** in 25% yield (Scheme 1).²¹




Fig. 1 a) Absorption spectrum of **1-Au**. b) Changes in the CD spectra of **1-Au**, with increasing amounts of HSA. [**1-Au**] = 5.0×10^{-5} M, [HSA] = $0 - 5 \times 10^{-5}$ M.




Fig. 2 ¹⁹F NMR spectrum of **1-Au**, in CD₃OD.

The electronic spectrum of **1-Au** is similar to those of other metal complexes of the same corrole (Fig. 1a); and, as expected, the ¹H NMR spectrum displays two singlets (at 10.19 and 9.16 ppm), assigned to C3-H and C18-H protons adjacent to the sulfonic acid moieties. Analysis of the ¹⁹F-NMR spectrum (Fig. 2) reveals that the F atoms above and below the macrocycle plane experience identical magnetic environments: each of three different C₆F₅ groups contributes only one chemical shift for the *ortho*- and *meta*-F's that it contains. The apparent relatively high symmetry and the diamagnetism of **1-Au** are fully consistent with expectation for a square planar low-spin 5d⁸ gold(III) complex. MS spectroscopy further confirmed the structure and the low affinity for axial ligands, as indicated by the molecular mass without any additives from coordinating solvents (pyridine, methanol, ethyl acetate) that were used in the synthesis and for chromatographic purification.

Examination of **1-Au** in aqueous PBS and PBS/glutathione solutions revealed no changes in the visible electronic spectrum with time, which is one indication that it has the stability required for consideration as a drug candidate. Four NCI60 cell lines, representative of breast, prostate, skin, and melanoma

cancers, were selected for investigations of **1-Au** treatments on their survival (by the MTA assay) and proliferation (by the BrdU assay).²² Interestingly, **1-Au** was found to possess 3.3 to 10.0-fold greater cytotoxicity than that of **1-Ga** (Fig. 3a-b) across all four cancer cell lines. The cytotoxicity of **1-Au** also was greater than that of cisplatin in the prostate, skin, and melanoma cancer cell lines (Table 1). Cytostatic activity also was examined for the two corroles: that of **1-Au** was 4.3 to more than 13.8-fold larger than that of **1-Ga** across all four cancer cell lines (Fig. 3c-d and Table 2), clearly showing that **1-Au** inhibits DNA replication far more than **1-Ga**. The striking similarity in the MTS and BrdU IC₅₀ values for **1-Au** further suggests that the dominant mechanism of cell-killing by **1-Au** is through cell cycle arrest, a result of DNA replication inhibition. Our proposed mechanism for **1-Au** is well-supported by our previous work that showed late M cell cycle arrest across several different cell lines by **1-Ga**, **1-AI** and **1-Mn**.²³




Fig. 3 Dose-response cytotoxicity and cytostaticity curves for **1-Au** (a,c) and **1-Ga** (b,d), presented as percentages relative to cells treated in a corrole-free medium with the same concentration of DMSO (1%) used for dissolving the corroles.

Table 1 Comparison of IC₅₀ cytotoxicity values of **1-Au**, **1-Ga** and

| Cell line | DU145 | SK-MEL-28 | MDA-MB-231 | OVCAR-3 | |
|-----------------------------|------------------|-----------|------------|---------|-------|
| Tumor type | prostate | melanoma | breast | ovarian | |
| IC₅₀ (μM) | 1-Au | 47.9 | 26.8 | 19.7 | 27.5 |
| | 1-Ga | 158.9 | 131.4 | 129.2 | 274.2 |
| | cisplatin | 81.0 | 36.8 | 44.8 | 22.8 |

cisplatin in human cancer cell lines.

Table 2 Comparison of IC₅₀ cytostaticity values for **1-Au** and **1-Ga**.

| Cell line | | DU145 | SK-MEL-28 | MDA-MB-231 | OVCAR-3 |
|------------------|-------------|--------|-----------|------------|---------|
| IC ₅₀ | 1-Au | 68.5 | 35.6 | 75.1 | 25.4 |
| (μ M) | 1-Ga | >350.0 | 201.9 | 322.5 | >350.0 |

Extensive work on **1-Ga** demonstrated the importance of its spontaneous noncovalent conjugation with proteins (owing to the amphipolar nature of the macrocycle), which serves for both its circulation and assisted cell membrane penetration. In order to gain insight into why the gold(III) corrole performs better than its gallium analogue, we examined its interaction with human serum albumin. First we employed circular dichroism (CD), since a signal in the visible range will only be obtained if **1-Au** conjugates with HSA and thus experiences a chiral environment. A strong signal was obtained upon titration of **1-Au** with HSA, with a positive Cotton effect and ellipticity inversion at 420 nm (Fig. 1b).²⁴ No change in the shape of the signal accompanied the titration, quite unlike our observations in related CD experiments where we found that 8-10 equivalents of **1-Ga** bind to HSA.¹⁵

Additional evidence for marked differences in binding affinities of the two corroles was obtained from quenching of the fluorescence of the only tryptophan residue present in HSA by **1-Au** and **1-Ga** (Fig. 4a and 4b, respectively).




Fig 4 Quenching of HSA Trp²¹⁴ emission ($\lambda_{exc} = 340$ nm) upon titration with a) **1-Ga** or b) **1-Au**, with [corrole]:[HSA] ratios of 0, 0.6, 0.7, 1, 1.4, 2, 2.8, 4.1, and 6.8.

The comparison clearly demonstrates that the affinity of **1-Au** for HSA (or more precisely to binding sites that are close to Trp²¹⁴) is much lower than that of **1-Ga**. At the highest corrole/HSA ratio, the Trp²¹⁴ emission was completely quenched for **1-Ga** (the residual emission is due to tyrosines), but not for **1-Au**.




Fig. 5 ESI-TOF mass spectra obtained from aqueous solutions of 5 μ M HSA that were incubated with 30 μ M of a) **1-Au** and b) **1-Ga**. HSA represents the free protein, while HSA + L and HSA + 2L represent HSA conjugated with the respective metallocorroles in a 1:1 and 1:2

ratios, respectively. The m/z values for HSA in each of the traces represent different ionization states (z), from +14 to +16.

Another qualitative indication of the weaker binding of **1-Au** was obtained by mass spectroscopy.²⁵ Analysis of the ESI-MS spectra of 1:6 solutions of HSA and corrole revealed that molecular clusters that correspond to HSA-bound conjugates were much more pronounced for **1-Ga** than **1-Au** (Fig. 5). The relatively weak binding of **1-Au** to albumin (present in the *in vitro* assays used for evaluating its cytotoxicity and cytostaticity) could be one of the factors contributing to its higher activity relative to **1-Ga**. The low affinity of **1-Au** for basic amino acid residues means that more uncomplexed agent likely will be available for intervention in intracellular processes. In support of this proposal, several investigators have shown that the interactions of many metallodrugs (including cisplatin and gold(III) pyridyls) with albumin are major influences on their biological activities.^{26,27,28}

We have discovered that **1-Au** is a more potent anticancer agent than **1-Ga**, with three to ten-fold greater cytotoxicity and four to more than ten-fold greater cytostaticity in four cancer cell lines. In our hands, **1-Au** also exhibited greater cytotoxicity than cisplatin in three out of the four cell lines examined. We suggest that the diminished binding of **1-Au** to HSA relative to **1-Ga** is responsible for its greater cell killing ability. The correlation between cytotoxicity and cytostaticity further suggests that the main mechanism of action is via mitotic arrest, as established for other metal complexes of the same corrole.²³ Although the anticancer potency of **1-Au** is not as great as that of [Au(PPP)]Cl, it is substantially more cytotoxic than water-soluble gold(III) porphyrins.²⁹ The take-home lesson is clear: we must place much more emphasis on investigations of less hydrophilic corroles in our search for better anticancer agents. More in depth investigations of the mechanism of action of the most active gold corroles, as well as exploration of their potential for photodynamic therapy,³⁰ will be pursued on these derivatives.

Acknowledgements

We thank Roger E. Moore of the Mass Spectrometry and Proteomics Core (City of Hope) for technical assistance. Our work is supported by the Caltech-City of Hope Biomedical Research Initiative (H.B.G., J.T., Z.G.) and the Caltech Summer Undergraduate Research Fellowship program (D.R.T.).

Notes and references

- Z. Gross, N. Galili and I. Saltsman, *Angew. Chem., Int. Ed.*, 1999, **38**, 1427.
- D. Aviezer, S. Cotton, M. David, A. Segev, N. Khaselev, N. Galili, Z. Gross and A. Yayon, *Cancer Res.*, 2000, **60**, 2973.
- A. Haber, A. Abu-Younis Ali, M. Aviram and Z. Gross, *Chem. Commun.*, 2013, **49**, 10917.
- H. Agadjanian, J. Ma, A. Rentsendorj, V. Valluripalli, J. Y. Hwang, A. Mahammed, D. L. Farkas, H. B. Gray, Z. Gross and L. Medina-Kauwe, *Proc. Natl. Acad. Sci.*, 2009, **106**, 6105.

- 5 C. M. Lemon and P. J. Brothers, *J. Porphyrins Phthalocyanines*, 2011, **15**, 809.
- 6 Z. Okun, L. Kuperschmidt, M. B. Youdim and Z. Gross, *Anti-cancer Agents Med. Chem.*, 2011, **11**, 380.
- 7 I. Nigel-Etinger, A. Mahammed and Z. Gross, *Catal. Sci. Tech.*, 2011, **1**, 578.
- 8 A. Mahammed and Z. Gross, *Chem. Commun.*, 2010, **46**, 7040.
- 9 C. M. Blumenfeld, R. H. Grubbs, R. A. Moats, H. B. Gray, K. Sorasaene, *Inorg. Chem.*, 2013, **52**, 4774.
- 10 H. Agadjanian, J. Weaver, A. Mahammed, A. Rentsendorj, S. Bass, J. Kim, I. Dmochowski, R. Margalit, H. Gray, Z. Gross and L. Medina-Kauwe, *Pharm. Res.*, 2006, **23**, 367.
- 11 L. Yun, H. Vazquez-Lima, H. Fang, Z. Yao, G. Geisberger, C. Dietl, A. Ghosh, P. J. Brothers and X. Fu, *Inorg. Chem.*, 2014, **53**, 7047.
- 12 A. Bindoli, M. P. Rigobello, G. Scutari, C. Gabbiani, A. Casini and L. Messori, *Coord. Chem. Rev.*, 2009, **253**, 1692.
- 13 C. T. Lum, A. S.-T. Wong, M. C. M. Lin, C.-M. C.-M. Che and R. W.-Y. Sun, *Chem. Commun.*, 2013, **49**, 4364.
- 14 S. E. Sherman, D. Gibson, A. H. Wang and S. J. Lippard, *Science*, 1985, **230**, 412.
- 15 A. Mahammed, H. B. Gray, J. J. Weaver, K. Sorasaene and Z. Gross, *Bioconjugate Chem.*, 2004, **15**, 738.
- 16 E. Rabinovich, I. Goldberg and Z. Gross, *Chem.–Eur. J.*, 2011, **17**, 12294.
- 17 K. E. Thomas, A. B. Alemayehu, J. Conradie, C. Beavers and A. Ghosh, *Inorg. Chem.*, 2011, **50**, 12844.
- 18 A. Mahammed and Z. Gross, *J. Chem. Soc.*, 2005, **127**, 2883.
- 19 I. Saltsman, A. Mahammed, I. Goldberg, E. Tkachenko, M. Botoshansky and Z. Gross, *J. Chem. Soc.*, 2002, **124**, 7411.
- 20 A. Mahammed, Z. Gross, *J. Inorg. Biochem.*, 2002, **88**, 305.
- 21 Synthesis of **1-Au**. **1-H₃** (37 mg, 0.046 mmol) was added to a solution of gold acetate (52 mg, 0.14 mmol) in pyridine under N_{2(g)} atmosphere. The reaction mixture was stirred overnight, after which solvent was evaporated under reduced pressure without heating. The product was purified by silica-gel chromatography (silica gel, MeOH:EA 1:15 as eluent), affording **1-Au** as a violet solid in 25% yield. **1-Au**: UV/Vis (CH₃CN) λ_{max} (ε × 10⁴): 418 nm (1.9), 531 nm (0.22), 568 nm (0.67). ¹H-NMR (CD₃OD, 400 MHz): δ = 10.19 (s, 1H, pyrrole), 9.16 (s, 1H, pyrrole), 9.00 (d, J=5.1 Hz, 2H, pyrrole), 8.97 (d, J₁=5.1 Hz, 2H, pyrrole). ¹⁹F-NMR (376.75 MHz, CD₃OD): δ = -138.41 (dd, J₁=6.5 Hz, J₂=23.0 Hz, 2 *ortho*-F), -140.15 (m, 4 *ortho*-F), -155.35 (t, J=20.8 Hz, 1 *para*-F), -155.14 (t, J₁=19.1 Hz, 1 *para*-F), -158.13 (t, J=19.1 Hz, 1 *para*-F), -164.69 (dt, J₁=7.15 Hz, J₂=22.1 Hz, 2 *meta*-F), -164.87 (dt, J₁=7.3 Hz, J₂=22.1 Hz, 2 *meta*-F), -168.03 (dt, J₁=6.7 Hz, J₂=22.77 Hz, 2 *meta*-F). MS-ES⁺ (TOF ES⁺): m/z: 1148.77 (calcd for [C₃₇AuH₇F₁₅N₄O₆S₂]⁺: 1149.10). Purity was further confirmed by analytical HPLC, performed on a 150 x 4.6 mm Kromasil C18 reverse phase column and H₂O/CH₃CN as the eluent.
- 22 Cells were grown in RPMI 1640 medium (Mediatech, Inc.) supplemented with 10% fetal bovine serum (Omega Scientific, Inc., Tarzana, CA), and maintained at 37°C under 5 % CO₂ in a humidified incubator. Cytotoxicity was determined by cell viability measurements using the CellTiter 96 Aqueous One Cell Proliferation Assay (MTS) from Promega (Madison, WI). Cytostatic activity was measured by the BrdU incorporation assay using the Cell Proliferation ELISA, BrdU (colorimetric assay) from Calbiochem (San Diego, CA). Cells were plated in seven 96-well dishes (5.6 × 10⁴ cells per well; 90 μL per well) 24 h prior to treatment. Drug treatment was initiated by adding 10 μL of fresh media containing the respective concentrations of metalocorrole or cisplatin in DMSO to make up 0.1 mL per well, followed by incubation for 72 h at 37 °C, 5% CO₂. For control wells, the same volume (10 μL) of fresh media containing 10% DMSO was added to make up 0.1 mL per well. The final amount of DMSO per well was kept at 1%. The MTS and BrdU assays were performed according to the manufacturers' directions. All drug treatments were performed in the dark to minimize the effects of metalocorrole photochemistry. Experiments were performed in triplicate. The means from triplicate spectrophotometric data were analyzed by nonlinear regression analysis using GraphPad Prism 5 (GraphPad Software, Inc., La Jolla, CA) to obtain a growth equation for each treatment. The rate of cell proliferation (BrdU) or growth (MTS) of corrole-treated cells relative to control cells is reported as percent of untreated cells.
- 23 P. Lim, A. Mahammed, Z. Okun, I. Saltsman, Z. Gross, H. B. Gray and J. Termini, *Chem. Res. Toxicol.*, 2011, **25**, 400.
- 24 CD spectra were recorded on a JASCO J-815 Circular Dichroism Spectrometer using 5.0 × 10⁻⁵ M corrole solution in 0.1 M sodium phosphate buffer at and adding microliter portions of 0.5 mM HSA solution.
- 25 A 10 mM NH₄OAc buffer was filtered through a 47 mm MF-Millipore membrane filter with 0.22 μm pore size (EMD Millipore, US) and used for sample preparation. The concentration of metalocorrole and HSA were kept at 30 μM and 5 μM, respectively. **1-Au** and **1-Ga** were allowed to incubate with HSA for 60-90 min at 25 °C. A Waters (Manchester, UK) Synapt G2 HDMS high definition mass spectrometer with ion mobility MS equipped with a Waters nanoAcquity 2D UHPLC system was used for measurements.
- 26 C. D. Sanghvi, P. M. Olsen, C. Elix, S. Peng, D. Wang, Z. Chen, D. M. Shin, K. I. Hardcastle, C. E. MacBeth and J. F. Eichler, *J. Inorg. Biochem.*, 2013, **128**, 68.
- 27 W. Hu, Q. Luo, K. Wu, X. Li, F. Wang, Y. Chen, X. Ma, J. Wang, J. Liu, S. Xiong and P. J. Sadler, *Chem. Commun.*, 2011, **47**, 6006.
- 28 D. Gibson, *Dalton Trans.*, 2009, 10681.
- 29 R. W.-Y. Sun and C.-M. Che, *Coord. Chem. Rev.*, 2009, **253**, 1682.
- 30 There is increased cytotoxicity of **1-Ga** when used in combination with light: J. Y. Hwang, J. Lubow, D. Chu, J. Sims, F. Alonso-Valenteen, H. B. Gray, Z. Gross, D. L. Farkas, and L. K. Medina-Kauwe *J. Controlled Release* **2012**, **163**, 368.

Table of Contents

The bis-sulfonated Au(III) corrole (**1-Au**) was found to be much more cytotoxic and cytostatic than the Ga(III) analog **1-Ga**, which might be attributed to the lower affinity of **1-Au** to serum albumin.

