

# ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## COMMUNICATION

# Morphology-driven absorption and emission colour changes in liquid-crystalline, cyclometallated platinum(II) complexes.

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,

Accepted 00th January 2012


DOI: 10.1039/x0xx00000x

www.rsc.org/

Valery N. Kozhevnikov,<sup>\*a</sup> Bertrand Donnio<sup>b,c</sup>, Benoît Heinrich<sup>b</sup> and Duncan W. Bruce<sup>\*d</sup>

**Platinum(II) complexes of 1,3-bis(2-pyridyl)benzene containing two alkyl chains are unexpectedly mesomorphic and capable of changing absorption and emission colour depending on the phase obtained after thermal treatment.**

Optical properties of square-planar Pt<sup>II</sup> complexes are often determined by the organisation of individual molecules in the bulk.<sup>1,2</sup> While in dilute solution absorption and emission are normally determined by individual molecules, whereas in the solid state the way complexes interact can play a crucial role. Molecular aggregates with strong Pt···Pt interactions often show drastically different absorption and emission characteristics from other aggregates in which the metal-metal interaction is weak or absent. Such behaviour is explained by the fact that short Pt-Pt distance allows formation of metal-metal-to-ligand-charge-transfer (MMLCT) states which leads to significant red-shift in absorption and emission. Optical responses of this kind have attracted considerable recent interest for the design of sensors and stimulus-responsive materials. Many such materials are based on crystalline samples, but 'soft' systems such as gels<sup>3</sup>, fibres<sup>4</sup> and polymers<sup>5</sup> have also been identified. Liquid-crystals are also interesting in this regard. Molecules in the liquid-crystalline state display much higher degrees of (anisotropic) order than classical liquids, but the system is much less ordered than the crystalline state. In addition, the occasional presence of monotropic phases means that kinetically stabilised states can be obtained, sometimes by fast cooling, allowing access to a wider range of self-organised structures. Furthermore, these metastable states can, on occasion, be rather long-lived.<sup>6</sup> Liquid-crystalline, cyclometallated complexes of Pt<sup>II</sup><sup>7,8,9,10</sup> are often highly luminescent and allow observation of morphology-dependent effects in both absorption and emission modes.<sup>11</sup> Here we report simple liquid-crystalline Pt<sup>II</sup> complexes of 1,3-bis(2-pyridyl)benzene with just two aliphatic chains, whose




**Fig. 1** Synthesis of the dialkyl-functionalised Pt complexes: (i) 5-bromo-2-iodopyridine,  $[Pd(PPh_3)_4]$ ,  $K_3PO_4$ , DMF, 100 °C, 14 h, 69%; (ii) 1-alkyne, CuI, TEA,  $[Pd(PPh_3)_4]$ , DMF, reflux, 12 h, 45-63%; (iii)  $H_2$ , Pd/C, RT, 2 d, 89-92%; (iv)  $K_2[PtCl_4]$ , AcOH, reflux, 24 h, 59-64%.

thermal behaviour is accompanied by drastic changes both in absorption and emission colour.

The synthesis of the ligand and the Pt<sup>II</sup> complex are depicted in Figure 1. A Suzuki-Miyaura coupling of the diboronate derivative **1** with 2-iodo-5-bromopyridine gives the dibromo derivative **2**. This was then reacted with terminal alkynes under Sonogashira conditions to give the alkynyl derivatives **3**, which were then reduced to the alkyl derivatives **4**. The target platinum complexes **5**, were finally obtained by heating a 1:1 mixture of the

ligand and potassium tetrachloroplatinate(II) in acetic acid under reflux.

Complexes **5** show interesting thermochromic behaviour. Thus, the yellow solid **5**(12), obtained from solution, melts at 119 °C to form a yellow, isotropic liquid. Cooling from the isotropic phase leads to the formation of a viscous, birefringent, monotropic mesophase at 82 °C (Fig. 1a). Interestingly, the formation of the mesophase is accompanied by the change in colour of the sample from yellow to red. The mesophase is fluid immediately after formation but on further cooling progressively becomes more viscous, forming a glassy state at approximately 75 °C without visible changes in appearance. This film stayed red for a period of several days, trapped in the metastable, glassy mesophase. Crystallisation did, however, occur slowly, reforming the crystalline phase and as can be seen in Fig 1b, this process of reforming the crystalline phase is accompanied by the change in colour from red to yellow. Crystallisation was found to be much faster at temperatures between 75-80 °C, no doubt on account of the greater mobility in the molten phase as the molecules explored different spatial organisation to lead to the crystalline state.




**Fig.2.** Complex **5**(12): a) the optical texture observed under polarised microscope; b) visual appearance of the sample at the transition between yellow isotropic phase and red mesophase; c) SAXS diffraction patterns of **5**(12) at different temperatures; d) emission and excitation spectra for the yellow (blue line) and red (black line) phases

Re-heating the red mesophase then leads to clearing at 93 °C, which is the true thermodynamic transition temperature, and is accompanied by a colour change from red back to yellow. Evidently there is significant supercooling of this transition when the temperature is lowered from the clearing point.

In order to gain a better understanding of the nature of the red and yellow phases, small-angle X-ray scattering (SAXS) experiments were carried out (Fig. 2a). Characteristic of the original yellow crystalline solid obtained from solution showed many sharp reflections with a primary  $d$ -spacing = 24.55 Å. While crystallisation prevented collection of data in the fluid red mesophase, a diffraction pattern was obtained in the glassy state and showed a  $d$ -spacing = 35.4 Å and a broad reflection close to  $2\theta = 20^\circ$ , expected where molten terminal chains are present. There was also evidence for other, broad reflections as well as a sharper reflection corresponding to a  $d$ -spacing of about 10 Å. The data are insufficient to assign the phase symmetry, but the combination of the sharper, low-angle reflection and the broad wide-angle reflection supports assignment as a liquid-crystalline phase. This is consistent with the presence of a birefringent texture seen using polarised microscopy (Fig. 2a), although it was not possible to obtain a well-developed texture that would allow mesophase characterisation. If the sample was left to crystallise at room temperature for two weeks, SAXS showed a pattern close to that of the pristine yellow solid with a primary  $d$ -spacing = 24.55 Å. The presence of additional reflections suggests the possible co-existence of a second crystalline modification, which may be metastable, being formed during crystallisation.

Similar behaviour was found for complex **5**(15) but at slightly different temperatures. Thus, the yellow crystalline solid melts at 120 °C and on cooling, the red, monotropic mesophase begins to form at 90 °C, forming a glass at approximately 80-70 °C on further cooling. The mesophase-isotropic liquid transition takes place at 102 °C on re-heating the red mesophase. Again, crystallisation prevented collection of data in the fluid mesophase.

In order to suppress crystallisation, a mixture of **5**(12) and **5**(15) (1:1 by mass) was prepared by dissolving the mixture of compounds in  $\text{CH}_2\text{Cl}_2$  followed by evaporation. In this way, crystallisation was suppressed significantly and it was possible to obtain an X-ray pattern for the fluid liquid-crystalline phase (83 °C, Fig. 3). The data can be assigned to a lamellar phase with a periodicity of about 38.5 Å, which distance is reasonable given that the data obtained for pure **5**-**12** suggested a lamellar phase with  $d = 35.4$  Å. The observation of a mesophase in complexes **5** is perhaps unexpected for such a simple system. This is an interesting phenomena, which deserves further investigation.



**Fig. 3** SAXS diffractogram of a mixture of **5(12)** and **5(15)** at 83 °C

In term of photophysical properties, both absorption and emission characteristics are affected by the state of the sample. The absorption maximum for the yellow crystalline phase is at 410 nm, while for the red phase it is at 520 nm. The luminescence profile also changes from an orange-red colour for the yellow phase ( $\lambda_{\text{max}} \approx 486, 658$  nm) to the near infrared region ( $\lambda_{\text{max}} \approx 730$  nm) for the red phase. These data correlate well with previous findings for similar Pt<sup>II</sup> complexes of 1,3-bis(2-pyridyl)benzene that the emission of the yellow phase is the combination of monomer and excimer-like emissions, while the absorption and emission of the red phase is dominated by MMLCT excited state.<sup>12</sup>

The above data are consistent with the following behaviour. Cooling an isotropic melt of **5** leads to a red, liquid-crystalline phase, the colour of which is indicative of the presence of short Pt<sup>II</sup>···Pt interactions judging by characteristic MMLCT absorption and emission.

It has been argued that such metallophilic interactions are responsible for the induction of mesomorphism in related Pt<sup>II</sup> materials.<sup>8</sup> Whether these interactions actually drive mesophase formation or whether it is the case that the optimum intermolecular aggregation imposed by mesophase formation leads to a geometry that favours the Pt<sup>II</sup>···Pt interactions remains a moot point. However, whatever the mechanism, it is clear that metallophilic Pt<sup>II</sup>···Pt interactions are key to the thermochromic behaviour of these complexes. Because thermodynamically the red mesophase is metastable, it eventually reverts to the yellow crystalline phase in which Pt<sup>II</sup>···Pt interactions are disrupted. In this way, it is possible to switch the Pt<sup>II</sup>···Pt interactions on and off manipulating the sample only thermally without the need for intervention by solvent or mechanical treatment. This binary switch in morphology is in turn translated to a binary switch in both absorption (red/yellow) and emission (NIR/orange) characteristics of the film.

## Notes and references

<sup>a</sup> Department of Applied Sciences, Northumbria University, Newcastle upon Tyne, NE1 8ST, U.K.; E-mail: valery.kozhevnikov@northumbria.ac.uk

<sup>b</sup> Institut de Physique et de Chimie des Matériaux de Strasbourg (IPCMS), UMR 7504 (CNRS-Université de Strasbourg), 23 Rue du Loess BP 43, 67034 Strasbourg Cedex 2, France.

<sup>c</sup> Complex Assemblies of Soft Matter Laboratory (COMPASS), UMI 3254 (CNRS-Solvay-University of Pennsylvania), CRTB, 350 George Patterson Boulevard, Bristol, PA 19007, USA.

<sup>d</sup> Department of Chemistry, University of York, Heslington, York YO10 5DD, UK; E-mail: duncan.bruce@york.ac.uk.

† Electronic Supplementary Information (ESI) available: synthetic procedures, DSC and SAXS data. See DOI: 10.1039/c000000x/

- 1 K. M.-C. Wong, M. M.-Y. Chan, V. W.-W. Yam, *Adv. Mater.* 2014, **26**, 5558; J. R. Kumpfer, S. D. Taylor, W. B. Connick, S. J. Rowan, *J. Mater. Chem.* 2012, **22**, 14196; P. Du, J. Schneider, W. W. Brennessel, R. Eisenberg, *Inorg. Chem.* 2007, **47**, 69; X.-P. Zhang, T. Wu, J. Liu, J.-X. Zhang, C.-H. Li, X.-Z. You, *J. Mater. Chem. C* 2014, **2**, 184; J. Ni, X. Zhang, Y. H. Wu, L. Y. Zhang, Z. N. Chen, *Chem. – Eur. J.*, 2011, **17**, 1171; M. L. Muro, C. A. Daws, F. N. Castellano, *Chem. Commun.* 2008, **46**, 6134.
- 2 B. Donnio and D. W. Bruce, Liquid crystalline *ortho*-palladated complexes, in *Palladacycles - Synthesis, Characterization and Applications*, ed. M. Pfeffer, J. Dupont, Wiley-VCH, Weinheim, 2008.
- 3 A.-Y. Tam, K.-C. Wong, N. Zhu, G. Wang, V. W.-W. Yam, *Langmuir* 2009, **25**, 8685; F. Camerel, R. Ziessel, B. Donnio, C. Bourgogne, D. Guillon, M. Schmutz, C. Iacovita, *Angew. Chem. Int. Ed.*, 2007, **46**, 2659
- 4 W. Lu, V. A. L. Roy, C.-M. Che, *Chem. Commun.* 2006, **38**, 3972; M. Mauro, A. Aliprandi, C. Cebrian, D. Wang, C. Kubel, L. De Cola, *Chem. Commun.* 2014, **50**, 7269.
- 5 C.-C. Kwok, S.-C. Yu, I. H. T. Sham, C.-M. Che, *Chem. Commun.* 2004, **21**, 2512.
- 6 H. J. Coles and M. N. Pivnenko, *Nature*, 2005, **436**, 997.
- 7 S. W. Thomas III, S. Yagi, T. M. Swager, *J. Mater. Chem.* 2005, **15**, 2829; K. Venkatesan, P. H. J. Kouwer, S. Yagi, P. Mueller, T. M. Swager, *J. Mater. Chem.* 2008, **18**, 400; A. Santoro, A. C. Whitwood, J. A. G. Williams, V. N. Kozhevnikov, D. W. Bruce, *Chem. Mater.*, 2009, **21**, 3871; Y. Wang, Y. Liu, J. Luo, H. Qi, X. Li, M. Nin, M. Liu, D. Shi, W. Zhu, Y. Cao, *Dalton Trans.*, 2011, **40**, 5046; M. Spencer, A. Santoro, G. R. Freeman, A. Diez, P. R. Murray, J. Torroba, A. C. Whitwood, L. J. Yellowlees, J. A. G. Williams, D. W. Bruce, *Dalton Trans.*, 2012, **41**, 14244.
- 8 T. Hegmann, J. Kain, S. Diele, B. Schubert, H. Bögel, C. Tschierske, *J. Mater. Chem.*, 2003, **13**, 991. C. Damm, G. Israel, T. Hegmann, C. Tschierske, *J. Mater. Chem.* 2006, **16**, 1808; Y. Wang, Y. Liu, J.

- 
- Luo, H. Qi, X. Li, M. Nin, M. Liu, D. Hu, Y. Cao, *Dalton Trans.*, 2011, **40**, 5046.
- 9 M. Krikorian, S. Liu, T. M. Swager, *J. Am. Chem. Soc.*, 2014, **136**, 2952.
- 10 C. T. Liao, H. H. Chen, H. F. Hsu, A. Poloek, H. H. Yeh, Y. Chi, K. W. Wang, C. H. Lai, G. H. Lee, C. W. Shih, P. T. Chou, *Chem. Eur. J.*, 2011, **17**, 546; Y. Li, A. Y.-Y. Tam, K. Wong, M.-C. W. Li, L. Wu, V. W.-W. Yam, *Chem. Eur. J.*, 2011, **17**, 8048. H.-C. Chang, T. Shiozaki, A. Kamata, K. Kishida, T. Ohmori, D. Kiriya, T. Yamauchi, H. Furukawa, S. Kitagawa, *J. Mater. Chem.*, 2007, **17**, 4136.
- 11 V. N. Kozhevnikov, B. Donnio, D. W. Bruce, *Angew. Chem. Int. Ed* 2008, **47**, 6286.
- 12 E. Rossi, A. Colombo, C. Dragonetti, D. Roberto, F. Demartin, M. Cocchi, P. Brulatti, V. Fattori, J. A. G. Williams, *Chem. Commun.* 2012, 48, 3182.