

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

First Ever Observation of the Intermediate of Phosphonium Salt & Ylide Hydrolysis: *P*-Hydroxytetraorganophosphorane

Peter A. Byrne,^{*,a,b} Yannick Ortin,^a Declan G. Gilheany^a

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

P-hydroxytetraorganophosphorane, the long-postulated intermediate in phosphonium salt & ylide hydrolysis, has been observed and characterised by low temperature NMR, finally definitively establishing its involvement in these reactions.

The results require modification of the previously accepted mechanism for ylide hydrolysis: *P*-hydroxytetraorganophosphorane is generated directly by 4-centre reaction of ylide with water.

The alkaline hydrolysis of phosphonium salts has been afforded extensive attention in the chemical literature, and has been put to good use in the synthesis of tertiary phosphine oxides.^{1,2} Phosphonium ylide hydrolysis too has been employed in diverse synthetic applications.³ Recently, several examples of tandem or domino reactions have been reported^{4,5,6} in which the reactivity of an ylide is utilised (e.g. for the synthesis of dihydrobenzofurans,⁴ or peptidyl ketones⁵) followed by hydrolytic extrusion of phosphine oxide from the resulting adduct (which may be a phosphonium ylide, betaine, or salt). Furthermore, as a result of recent developments in phosphine oxide reduction methodology (and indeed the reduction of the P=O bond in related species),⁷ phosphonium salt hydrolysis followed by phosphine oxide reduction has now become an attractive route to phosphines (and derivatives). This is of particular importance for chiral phosphonium salts, which are hydrolysed stereospecifically⁸ and thus give the possibility of obtaining enantiopure or enantio-enriched chiral phosphines.

The hydrolysis of a phosphonium ylide (by addition of water to an anhydrous solution of the ylide)^{9,10} and that of the parent phosphonium salt under alkaline conditions afford the same products – phosphine oxide and the alkane or arene that results from scission of the P-C bond to the group that gives the most stable carbanion (i.e. the order of ease of P-C bond cleavage is *p*-nitrobenzyl > benzyl > phenyl > alkyl).^{8,11,12,13,14,15,16,17,18,19,20,21} Hydrolysis of phosphonium salts has been shown to be 3rd order (1st order in phosphonium salt, 2nd order in hydroxide) for benzyltriphenylphosphonium,^{11,12,14} benzyltrialkylphosphonium,¹⁴ mixed benzyl/alkyl/phenyl phosphonium, mixed alkyl/phenyl phosphonium,¹⁴ tetraphenylphosphonium,^{12,14} and tetrabenzylphosphonium salts.¹³ 3rd order hydrolysis has also been observed for cyclic phosphonium salts.²² The hydrolysis reactions of *para*-nitrobenzyltriphenylphosphonium bromide (**1**, Chart 1),^{12,23} of

methyltris(pentafluorophenyl)phosphonium triflate,²³ and of phosphonium iodides,²⁰ respectively, are exceptional in that they exhibit 1st order dependence on hydroxide concentration.





Chart 1. Phosphonium salts, ylides, & *P*-hydroxytetraorganohydroxyphosphorane **8**.

The mechanism that has been proposed for alkaline hydrolysis of phosphonium salts is shown in Scheme 1.^{11,12,13,14,16,17,24} Nucleophilic attack of hydroxide at phosphorus gives a hydroxyphosphorane (with apical oxygen^{8,25,26}). This is deprotonated by hydroxide to give an oxyanionic phosphorane, which expels a carbanion (probably protonated in the process of its expulsion) to give phosphine oxide and alkane or arene.²⁷ Largely on the basis that ylide hydrolysis gives the same products as phosphonium salt hydrolysis,¹⁵ it has generally been concluded that ylide hydrolysis proceeds by the same mechanism with one extra (initial) step - protonation of the ylide by water to give phosphonium hydroxide.^{10,28,29}



Scheme 1. The mechanism of alkaline hydrolysis of a phosphonium salt.

The existence of an intermediate in these reactions is a necessity in view of the kinetic order of the reaction (& in particular that it is 2nd order in hydroxide). Based on analogy with many other instances in which phosphoranes are involved,^{25,30,31,32,33,34,35} including the relatively stable *P*-alkoxytetraorganophosphoranes that have been observed spectroscopically,³⁵ it is reasonable to conclude that the intermediate in question is a trigonal bipyramidal (TBP) *P*-hydroxytetraorganophosphorane, as shown in Scheme 1. Consistent with this interpretation are the results of hydrolysis reactions of phosphonium salts in which the phosphorus atom is in a cycle of five or fewer members (see the ESI for additional discussion on this topic).† Certain chiral small ring phosphonium salts (and also compound **2**; see Chart 1)^{24,36} have been shown to undergo hydrolysis with retention of

configuration at phosphorus,³⁷ while other cyclic phosphonium salts undergo ring opening in the course of hydrolysis,^{20,22,38} including dibenzophospholium salts such as **4** (Chart 1).^{18,39} In TBP phosphoranes, the leaving group is obliged to depart from an apical site.^{25,40} However, in a cyclic hydroxyphosphorane, the apical sites are expected to be occupied by hydroxide and one ring P-C bond.^{25,26,30,35,41} Thus, for decomposition of the intermediate to occur, the reaction must proceed either with ring cleavage or with stereomutation by pseudorotation to place the leaving group in an apical site in the transition state.⁴² If starting from a chiral phosphonium salt, the ligand permutation that occurs during pseudorotation can reveal itself in the product phosphine oxide. Racemisation or retention of configuration (the latter is observed in the examples cited above) at phosphorus in the product indicates the intervention of stereomutation in the intermediate. That both ring cleavage and phosphorus ligand permutation (resulting in retention of configuration at phosphorus) are known to occur during phosphonium salt hydrolysis gives strong support to the existence of a TBP intermediate. The hydrolysis of **3**, (in which the phosphorus is not constrained in a ring) proceeds stereospecifically with *inversion* of configuration at P.^{8,43}




Fig. 1. (a) Partial ^{31}P NMR and (b) partial ^1H NMR showing signals for *P*-hydroxyphosphorane **8**.

Despite the strong evidence implicating *P*-hydroxytetraorganophosphoranes formation in phosphonium salt & ylide hydrolysis, this putative intermediate has never


been observed or characterised spectroscopically.^{44,45,46,47,48} Since low temperature NMR analysis of oxaphosphetanes (OPAs)^{32,33,34} and of alkoxyphosphoranes³⁵ was known to be possible, and had proven to be so useful in the analysis of the reactions in which these species are involved, we undertook to determine if spectroscopic observation of *P*-hydroxytetraorganophosphorane(s) could be realised. We also considered that these experiments might allow us to clarify what seemed to us to be an inconsistency in the accepted mechanism of ylide hydrolysis.^{10,15,28} The supposition that the first step of ylide hydrolysis involves formation of phosphonium hydroxide is inconsistent with the relative pK_{a} s of H_2O (31 in DMSO)⁴⁹ and phosphonium ylides (e.g. $\text{pK}_{\text{a}} = 22$ in DMSO for deprotonation of MePh_3P^+),⁵⁰ in organic media. Since stepwise protonation and subsequent hydroxide addition seems therefore to be an impossible process in organic media, we hoped to establish if there exists any evidence for the formation (or otherwise) of phosphonium hydroxide in these reactions.

Ylide hydrolysis reactions are extremely rapid at ambient temperature (20 °C), and all such reactions investigated by us resulted only in the observation of phosphine oxide (by ^{31}P NMR). Indeed, even at -70 °C, rapid conversion of ylide to phosphine oxide was observed by ^{31}P NMR for ylides **5** and **6** (Chart 1), with no appearance of any high field signals that could be attributed to an intermediate.

However, the addition of water to a solution of constrained (cyclic) ylide **7** (Chart 1) at -80 °C resulted in the formation of a compound with a chemical shift of -80.6 ppm in the ^{31}P NMR (see Fig. 1(a); obtained at -70 °C).[†] The high field chemical shift is indicative of a phosphorane structure (electron-rich at phosphorus).⁵¹ This compound was also characterised by ^1H , ^{13}C , gCOSY, gHSQC, ^1H - ^{13}C gHMBC and ^1H - ^{31}P HMBC. The ^1H NMR (Fig. 1(b)) contains a doublet for the methyl group at δ_{H} 0.90, two multiplets arising from the diastereotopic CH_2 protons (δ_{H} 2.32 & 2.68, each of which couples to methyl group by COSY), and a distinctive broadened singlet at δ_{H} 5.11 for the *P*-hydroxyl proton. The hydroxyl proton is shown to be coupled to phosphorus by ^1H - ^{31}P HMBC, and to the methylene carbon by ^1H - ^{13}C HMBC, indicating beyond doubt that this species is indeed *P*-hydroxyphosphorane (**8**, shown in Chart 1). In the ^{13}C NMR of **8**, a value of $^1J_{\text{PC}}$ of 119 Hz is observed for the methylene carbon.[†] The magnitude of the value of $^1J_{\text{PC}}$ observed for the methylene group of **8** is consistent with its occupation of an equatorial position in a pentavalent phosphorane.^{30,31,33,34,35}

This is the first time that a *P*-hydroxytetraorganophosphorane – the proposed intermediate in phosphonium salt and ylide hydrolysis – has been observed spectroscopically. Significantly, a small amount of residual ylide could also be observed in the ^{31}P NMR (δ_{P} -11.6). Since water (present in excess), ylide and *P*-hydroxyphosphorane can be seen from the ^{31}P and ^1H NMR to co-exist in solution, it is apparent that an equilibrium involving these three species is in operation (*cf.* reactions of ylides and alcohols to give alkoxyphosphoranes³⁵). Furthermore, since the pK_{a} of the ylide is too low in organic media to effect deprotonation of water, we conclude that that phosphonium hydroxide (which is

not detected) plays no role in the process of interconversion between ylide and *P*-hydroxytetraorganophosphorane. We propose instead that the phosphorane is formed directly from ylide + H₂O by (reversible) concerted addition of an O-H bond across the P=C bond. Indeed, it is even possible that the hydrolysis of *p*-nitrobenzyl phosphonium salt **1** (Chart 1), which is 1st order in each of phosphonium salt & hydroxide, may proceed by initial formation of ylide, and hence occurs by a different mechanism to that shown in Scheme 1. The sample of *P*-hydroxyphosphorane **8** was stored at -19 °C, and was observed by ¹H & ³¹P NMR (run at 25 °C) to have survived for 5 days at -19 °C. Warming of the sample to 25 °C resulted in the formation of ring-opened phosphine oxide **10** (Scheme 2) almost to the exclusion of the other two possible phosphine oxide products (*cf.* other cyclic phosphonium salts, especially **4**, *vide supra*). The identity of **10** was confirmed unambiguously by comparison with the NMR data of an authentic sample prepared by independent means – α -methylation of phosphine oxide **9** via a phosphinoxy carbanion (see Scheme 2).



Scheme 2. Synthesis of phosphine oxide **10**.

In conclusion, we have succeeded in carrying out the first ever spectroscopic observation of a *P*-hydroxytetraorganophosphorane, the intermediate of phosphonium salt and ylide hydrolysis. The ³¹P NMR chemical shift and associated spectral data reported above establish that the intermediate is a TBP phosphorane, and validates the operation of the proposed mechanism shown in Scheme 1 for phosphonium salt hydrolysis. The establishment of the structure of this intermediate provides a basis for a unifying explanation of the outcomes (ring opening, ring expansion, retention of stereochemistry at phosphorus) of the hydrolysis reactions of the various cyclic phosphonium salts.²⁸ It seems likely that spectroscopic observation of the *P*-hydroxyphosphoranes arising from hydrolysis of other cyclic phosphonium salts (e.g. **4**) should be possible, at least in some cases. Given the relative stability of **8** (it survived under inert atmosphere for several days at -19 °C), it may be possible to pursue novel synthetic avenues by testing the reactivity of **8** and analogues. Finally, we have provided evidence that the first step of ylide hydrolysis is not protonation of the ylide, but rather is concerted addition of an O-H bond across the P=C bond.

Acknowledgements

This work was supported by the Synthesis and Solid State Pharmaceutical Centre (SSPC) and Science Foundation Ireland (SFI) under grant number 12/RC/2275. We also thank SFI for the acquisition at UCD of an NMR spectrometer equipped for low temperature experiments (SFI Infrastructural Award [12/RI/2341(2)]).

Notes and references

- ^a Centre for Synthesis & Chemical Biology, University College Dublin, Belfield, Dublin 4, Ireland. Tel: +353-1-7162308.
- ^b Current address: Department Chemie, Ludwig-Maximilians-Universität München, Butenandtstr. 5-13 (Haus F), 81377 München, Germany. E-mail: peter.byrne@cup.lmu.de.
- [†] Electronic Supplementary Information (ESI) available: Additional discussion of the mechanism of phosphonium salt hydrolysis, experimental procedures & characterisation details for new compounds are available. See DOI: 10.1039/b000000x/
- P. Wyatt, S. Warren, M. McPartlin, T. Woodroffe, *J. Chem. Soc., Perkin Trans. I* 2001, 279; P. Calcagno, B. M. Kariuki, S. J. Kitchin, J. M. A. Robinson, D. Philp, K. D. M. Harris, *Chem. Eur. J.* 2000, **6**, 2338; D. W. Allen, J. P. Mifflin, P. J. Skabara, *J. Organomet. Chem.*, 2000, **601**, 293; F. Zaragoza, *J. Org. Chem.* 2002, **67**, 4963.
- D. Valentine Jr., J. F. Blount, K. Toth, *J. Org. Chem.* 1980, **45**, 3691.
- E. C. Taylor, S. F. Martin *J. Am. Chem. Soc.* 1974, **96**, 8095; M. Schwarz, J. E. Oliver, P. E.; Sonnet, *J. Org. Chem.* 1975, **40**, 2410; K. Y. Lee, J. E. Na, M. J. Lee, J. N. Kim *Tetrahedron Lett.* 2004, **45**, 5977.
- J.-B. Zhu, P. Wang, P.; S. Liao, Y. Tang *Org. Lett.* 2013, **15**, 3054.
- A. El-Dahshan, Ahsanullah, J. Rademann *Biopolymers* 2010, **94**, 220.
- J.-B. Zhu, P. Wang, P.; S. Liao, Y. Tang *Chem. Commun.* 2013, **49**, 4570; D. Duvvuru, P. Retailleau, J.-F. Betzer, A. Marinetti *Eur. J. Org. Chem.* 2012, 897–900.
- Y. Li, S. Das, S. Zhou, K. Junge, M. Beller, *J. Am. Chem. Soc.* 2012, **134**, 9727; Y. Li, L.-Q. Lu, S. Das, S. Pisiewicz, K. Junge, M. Beller, *J. Am. Chem. Soc.* 2012, **134**, 18325; T. Yano, M. Hoshino, M. Kuroboshi, H. Tanaka, *Synlett* 2010, **5**, 801; K. V. Rajendran, D. G. Gilheany *Chem. Commun.* 2012, **48**, 817; P. A. Byrne, K. V. Rajendran, J. Muldoon, D. G. Gilheany, *Org. & Biomol. Chem.* 2012, **10**, 3531; N. P. Kenny, K. V. Rajendran, E. V. Jennings, D. G. Gilheany *Chem. Eur. J.* 2013, **19**, 14210.
- W. E. McEwen, K. F. Kumli, A. Blade-Font, M. Zanger, C. A. VanderWerf, *J. Am. Chem. Soc.* 1964, **86**, 2378.
- F. Ramirez, S. Dershowitz, *J. Org. Chem.* 1957, **22**, 41; A. W. Johnson, *J. Org. Chem.* 1959, **24**, 282; F. Ramirez, O. P. Madan, C. P. Smith, *Tetrahedron Lett.*, 1965, 201; G. Wittig, G. Laib, *Liebigs Ann. Chem.* 1953, **580**, 57; A. W. Johnson, R. B. Lacount, *Tetrahedron*, 1960, **9**, 130; M. P. Cooke, *J. Org. Chem.* 1973, **38**, 4082.
- A. Schnell, J. G. Dawber, J. C. Tebby, *J. Chem. Soc., Perkin Trans. II*, 1976, 633; A. Schnell, J. C. Tebby, *J. Chem. Soc., Chem. Commun.*, 1975, 134; A. Schnell, J. C. Tebby, *J. Chem. Soc., Perkin Trans. I*, 1977, 1883.
- C. A. VanderWerf, W. E. McEwen, M. Zanger, *J. Am. Chem. Soc.* 1959, **81**, 3806.
- G. Asknes, J. Songstad, *Acta. Chem. Scand.* 1962, **16**, 1426.
- W. E. McEwen, G. Axelrad, M. Zanger, C. A. VanderWerf, *J. Am. Chem. Soc.* 1965, **87**, 3948; W. E. McEwen in *Topics in Phosphorus Chemistry*; M. Grayson & E. J. Griffith, Eds.; Interscience: New York, 1965, Vol. 2, pp. 5-9.
- J. G. Dawber, R. G. Skerratt, J. C. Tebby, A. A. C. Waite, *Phosphorus, Sulfur, Silicon Relat. Elem.* 2012, **187**, 1261.
- A. W. Johnson, *Ylides and Imines of Phosphorus*, 1993, Wiley; New York.
- R. R. Holmes, *Pentacoordinated Phosphorus, Volume II*, 1980, American Chemical Society; Washington D. C.; pp. 115-129.
- H. J. Cristau, F. Plénat in *The Chemistry of Organophosphorus Compounds Volume 3*; F. R. Hartley, Ed.; Wiley: Chichester, 1994, Chapter 2; pp. 111-138.
- D. W. Allen, I. T. Miller *J. Chem. Soc. (C)* 1967, 1869; D. W. Allen, I. T. Miller *J. Chem. Soc. (C)* 1969, 252.
- G. W. Fenton, C. K. Ingold, *J. Chem. Soc.* 1929, 2342; M. Schlosser, *Angew. Chem.* 1962, **74**, 291; D. W. Allen, B. G. Huntley, M. T. J. Mellor, *J. Chem. Soc., Perkin Trans. II*, 1974, 1690; L. Hey, C. K. Ingold *J. Chem. Soc.* 1933, 531; D. Q. Allen, B. G. Huntley, M. T. J. Mellor *J. Chem. Soc., Perkin Trans. II* 1972, 63.
- K. Bergeson *Acta Chem. Scand.* 1966, **20**, 899.

- 21 J. G. Dawber, J. C. Tebby, A. A. C. Waite, *J. Chem. Soc., Perkin Trans. I*, 1983, 1923.
- 22 G. Aksnes, K. Bergesen *Acta Chem. Scand.* 1965, **19**, 931.
- 23 B. Siegel *J. Am. Chem. Soc.* 1979, **101**, 2265.
- 24 J. R. Corfield, N. J. De'Ath, S. Trippett, *J. Chem. Soc. (C)* 1971, 1930.
- 25 F. H. Westheimer, *Acc. Chem. Res.* 1968, **1**, 70.
- 26 K. Mislow *Acc. Chem. Res.* 1970, **3**, 321; R. K. Oran, S. Trippett *J. Chem. Soc., Perkin Trans. I* 1973, 1300.
- 27 J. R. Corfield, S. Trippett, *J. Chem. Soc. Chem. Commun.* 1970, 1267.
- 28 U. Schöllkopf *Angew. Chem.* 1959, **71**, 260; H. J. Bestmann *Angew. Chem.* 1965, **77**, 260; D. D. Coffmann, C. S. Marvel *J. Am. Chem. Soc.* 1929, **51**, 3496.
- 29 See ESI for additional discussion on the differences that have been observed between phosphonium salt & ylide hydrolysis.
- 30 P. B. Kay, S. Trippett, *J. Chem. Res. (S)*, 1986, 62
- 31 E.g. *P*-fluorophosphoranes: R. G. Cavell, J. A. Gibson, K. I. The *Inorg. Chem.* 1978, **17**, 2880; R. G. Cavell, J. A. Gibson, K. I. The *J. Am. Chem. Soc.* 1977, **99**, 7841; K. I. The, R. G. Cavell *Inorg. Chem.* 1977, **16**, 2887; K. I. The, R. G. Cavell *J. Chem. Soc., Chem. Commun.* 1975, 279; K. I. The, R. G. Cavell *J. Chem. Soc., Chem. Commun.* 1975, 716; R. G. Cavell, D. D. Poulin, K. I. The, A. J. Tomlinson *J. Chem. Soc., Chem. Commun.* 1974, 19.
- 32 E.g. oxaphosphetanes: T. Fleck, *J. Am. Chem. Soc.* 1989, **111**, 5861; E. Vedejs, C. F. Marth, R. Ruggeri, *J. Am. Chem. Soc.* 1988, **110**, 3940; E. Vedejs, C. F. Marth, *J. Am. Chem. Soc.* 1988, **110**, 3948; E. Vedejs, K. A. J. Snoble, *J. Am. Chem. Soc.* 1973, **95**, 5778; E. Vedejs, G. P. Meier, K. A. J. Snoble, *J. Am. Chem. Soc.* 1981, **103**, 2823; E. Vedejs, C. F. Marth, *J. Am. Chem. Soc.* 1990, **112**, 3905; B. E. Maryanoff, A. B. Reitz, M. S. Mutter, R. R. Inners, H. R. Almond Jr., *J. Am. Chem. Soc.* 1985, **107**, 1068; B. E. Maryanoff, A. B. Reitz, M. S. Mutter, R. R. Inners, H. R. Almond Jr., R. R. Whittle, R. A. Olofson, *J. Am. Chem. Soc.* 1986, **108**, 7664; A. B. Reitz, M. S. Mutter, B. E. Maryanoff, *J. Am. Chem. Soc.* 1984, **106**, 1873; F. Bangerter, M. Karpf, L. A. Meier, P. Rys, P. Skrabal, *J. Am. Chem. Soc.* 1998, **120**, 10653; P. A. Byrne, D. G. Gilheany *Chem. Soc. Rev.* 2013, **42**, 6670.
- 33 E. Vedejs, C. F. Marth, *J. Am. Chem. Soc.* 1989, **111**, 1519.
- 34 P. A. Byrne, D. G. Gilheany *J. Am. Chem. Soc.* 2012, **134**, 9225; P. A. Byrne, J. Muldoon, Y. Ortin, D. G. Gilheany, *Eur. J. Org. Chem.* 2014, 86.
- 35 H. Schmidbaur, H. Stühler, W. Buchner *Chem. Ber.* 1973, **106**, 1238; H. Schmidbaur, W. Buchner, D. Scheutzwow *Chem. Ber.* 1973, **106**, 1251; H. Schmidbaur, W. Buchner, F. H. Köhler *J. Am. Chem. Soc.* 1974, **96**, 6208; H. Schmidbaur, H. Stühler *Chem. Ber.* 1974, **107**, 1420; H. Schmidbaur, P. Holl *Chem. Ber.* 1976, **109**, 3151 (The product in this case is a cyclic 1,2-oxaphospholane - from ylide + epoxide); D. Q. Allen, B. G. Huntley, M. T. J. Mellor *J. Chem. Soc., Perkin Trans. II* 1972, 63; P. A. Byrne, D. G. Gilheany, unpublished results.
- 36 N. J. De'Ath, S. Trippett *J. Chem. Soc. Chem. Commun.* 1969, 172.
- 37 K. L. Marsi *J. Chem. Soc. Chem. Commun.*, 1968, 846; W. Hawes, S. Trippett *J. Chem. Soc. Chem. Commun.*, 1968, 295.
- 38 S. E. Fishwick, J. A. Flint *J. Chem. Soc. Chem. Commun.* 1968, 182; S. E. Fishwick, J. A. Flint, W. Hawes, S. Trippett *J. Chem. Soc. Chem. Commun.* 1967, 1113.
- 39 E. M. Richards, J. C. Tebby *J. Chem. Soc. Chem. Commun.* 1967, 957.
- 40 No exceptions to the apical departure rule for trigonal bipyramidal phosphoranes have yet been reported.^{25,26,31,32,33,34,35}
- 41 Note that where there is a choice between two elements, the most electronegative element occupies the apical site.
- 42 E. P. A. Couzijn, J. C. Slootweg, A. W. Ehlers, K. Lamertsma *J. Am. Chem. Soc.* 2010, **132**, 18127; E. L. Muetterties, *J. Am. Chem. Soc.* 1969, **91**, 4115; E. L. Muetterties *J. Am. Chem. Soc.* 1969, **91**, 1636; J. García López, A. Morán Ramallal, J. González, L. Rocés, S. García-Granda, M. J. Iglesias, P. Oña-Burgos, F. López Ortiz *J. Am. Chem. Soc.* 2012, **134**, 19504.
- 43 The benzyl group is apparently cleaved far faster than ligand permutation can occur at phosphorus in the *P*-hydroxytetraorganophosphorane.
- 44 *P*-hydroxyphosphoranes with multiple P-O bonds have been reported,^{45,46} while a hydroxyphosphorane structure has been proposed for the product of the reaction of arylsulfonylglycyl chlorides with Ph₃P.⁴⁷ However, as pointed out by Chesnut & Quin,⁴⁸ on the basis of the chemical shift of δ_p 18.7 for this entity, it seems likely that it is in fact a phosphonium salt.¹⁵
- 45 Y. Segall, I. Granoth, *J. Am. Chem. Soc.* 1978, **100**, 5130; Y. Segall, I. Granoth, *J. Am. Chem. Soc.* 1979, **101**, 3687; I. Granoth, J. C. Martin *J. Am. Chem. Soc.* 1978, **100**, 5229; F. Ramirez, M. Nowakowski, J. F. Maracek *J. Am. Chem. Soc.* 1977, **99**, 4517.
- 46 I. Granoth, J. C. Martin *J. Am. Chem. Soc.* 1981, **103**, 2711. In this paper, alkoxyhalophosphorane structures are assigned to the products of the reactions of phosphinites with alkyl halides; these entities appear in fact to be alkoxyphosphonium halides based on their low field ³¹P NMR chemical shifts.
- 47 I. I. Yavari, A. Alizadeh *Tetrahedron* 2001, **57**, 9873.
- 48 D. B. Chesnut, L. D. Quin *Tetrahedron* 2005, **62**, 12343.
- 49 W. N. Olmstead, Z. Margolin, F. G. Bordwell, *J. Org. Chem.* 1980, **45**, 3295; F. G. Bordwell, *Acc. Chem. Res.* 1988, **21**, 456.
- 50 X.-M. Zhang, F. G. Bordwell, *J. Am. Chem. Soc.* 1994, **116**, 968.
- 51 The existence of this *P*-hydroxyphosphorane at low temperature may account for some signals that we were previously unable to assign in the high field region of ³¹P NMR spectra of some dibenzophosphole-derived OPAs in reference 34. It is likely that small amounts of *P*-hydroxyphosphorane were produced by reaction of ylide with adventitious water.